

Table 1.103C Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 or Older: Standard Errors of Numbers in Thousands, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	375	387	178	178	180	182
Amphetamine or Methylphenidate Products	323	340	171	173	173	176
Amphetamine Products	305	317	167	167	169	170
Methylphenidate Products	131	157	57	64	57	64
Anorectic (Weight-Loss) Stimulants	140	141	28	42	28	42
Provigil®	56	69	29	28	*	*
Any Other Prescription Stimulant ¹	148	158	26	10	26	10

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.103D Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 or Older: Standard Errors of Percentages, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	0.14	0.14	0.06	0.06	0.87	0.85
Amphetamine or Methylphenidate Products	0.12	0.12	0.06	0.06	1.03	1.02
Amphetamine Products	0.11	0.12	0.06	0.06	1.10	1.09
Methylphenidate Products	0.05	0.06	0.02	0.02	1.67	1.77
Anorectic (Weight-Loss) Stimulants	0.05	0.05	0.01	0.02	1.10	1.79
Provigil®	0.02	0.03	0.01	0.01	*	*
Any Other Prescription Stimulant ¹	0.05	0.06	0.01	0.00	0.96	0.41

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.103P Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 or Older: P Values from Tests of Differences of Percentages, 2019 Versus 2018

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	0.7284		0.4006		0.2427	
Amphetamine or Methylphenidate Products	0.3122		0.5292		0.1325	
Amphetamine Products	0.3955		0.5030		0.1367	
Methylphenidate Products	0.2888		0.5778		0.2204	
Anorectic (Weight-Loss) Stimulants	0.1438		0.2476		0.1111	
Provigil®	0.1810		0.3369		*	
Any Other Prescription Stimulant ¹	0.8355		0.0554		0.0553	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.104C Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 to 17 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 to 17: Standard Errors of Numbers in Thousands, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	62	67	27	37	28	37
Amphetamine or Methylphenidate Products	58	65	27	36	27	37
Amphetamine Products	51	58	25	35	26	35
Methylphenidate Products	35	39	13	15	13	15
Anorectic (Weight-Loss) Stimulants	9	15	3	7	*	*
Provigil®	8	5	3	2	*	*
Any Other Prescription Stimulant ¹	26	31	5	8	5	8

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.104D Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 to 17 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 to 17: Standard Errors of Percentages, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	0.25	0.27	0.11	0.15	1.42	1.65
Amphetamine or Methylphenidate Products	0.23	0.26	0.11	0.14	1.55	1.78
Amphetamine Products	0.21	0.23	0.10	0.14	1.86	2.13
Methylphenidate Products	0.14	0.16	0.05	0.06	2.25	2.27
Anorectic (Weight-Loss) Stimulants	0.04	0.06	0.01	0.03	*	*
Provigil®	0.03	0.02	0.01	0.01	*	*
Any Other Prescription Stimulant ¹	0.10	0.12	0.02	0.03	1.62	2.08

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.104P Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 to 17 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 to 17: P Values from Tests of Differences of Percentages, 2019 Versus 2018

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	0.3495		0.1914		0.2942	
Amphetamine or Methylphenidate Products	0.4057		0.2057		0.2900	
Amphetamine Products	0.3155		0.2164		0.3735	
Methylphenidate Products	0.5662		0.7512		0.8974	
Anorectic (Weight-Loss) Stimulants	0.2199		0.3423		*	
Provigil®	0.2926		0.6540		*	
Any Other Prescription Stimulant ¹	0.0700		0.5687		0.8204	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.105C Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 or Older: Standard Errors of Numbers in Thousands, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse	Misuse	Misuse	Misuse
			in Past Year among Total Population (2018)	in Past Year among Total Population (2019)	in Past Year among Any Users of Stimulant Subtypes (2018)	in Past Year among Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	368	379	175	173	177	177
Amphetamine or Methylphenidate Products	316	330	169	167	170	170
Amphetamine Products	299	305	164	162	166	164
Methylphenidate Products	127	151	57	61	58	61
Anorectic (Weight-Loss) Stimulants	139	140	27	41	27	41
Provigil®	56	69	29	28	*	*
Any Other Prescription Stimulant ¹	146	155	25	7	25	7

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.105D Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 or Older: Standard Errors of Percentages, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	0.15	0.15	0.07	0.07	0.95	0.92
Amphetamine or Methylphenidate Products	0.13	0.13	0.07	0.07	1.14	1.13
Amphetamine Products	0.12	0.12	0.07	0.06	1.21	1.18
Methylphenidate Products	0.05	0.06	0.02	0.02	2.00	2.08
Anorectic (Weight-Loss) Stimulants	0.06	0.06	0.01	0.02	1.11	1.83
Provigil®	0.02	0.03	0.01	0.01	*	*
Any Other Prescription Stimulant ¹	0.06	0.06	0.01	0.00	1.05	0.32

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.105P Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 or Older: P Values from Tests of Differences of Percentages, 2019 Versus 2018

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	0.8457		0.2616		0.1692	
Amphetamine or Methylphenidate Products	0.3748		0.3668		0.0823	
Amphetamine Products	0.4876		0.3514		0.0957	
Methylphenidate Products	0.3379		0.5173		0.1889	
Anorectic (Weight-Loss) Stimulants	0.1134		0.3048		0.1387	
Provigil®	0.1472		0.3557		*	
Any Other Prescription Stimulant ¹	0.5872		0.0251		0.0252	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.106C Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 to 25 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 to 25: Standard Errors of Numbers in Thousands, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	127	134	93	93	104	102
Amphetamine or Methylphenidate Products	124	132	92	92	103	102
Amphetamine Products	118	129	88	92	99	101
Methylphenidate Products	64	61	42	34	43	34
Anorectic (Weight-Loss) Stimulants	24	25	7	9	*	*
Provigil®	12	13	9	4	*	*
Any Other Prescription Stimulant ¹	38	34	5	7	5	7

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.106D Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 to 25 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 to 25: Standard Errors of Percentages, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	0.37	0.40	0.27	0.27	1.39	1.47
Amphetamine or Methylphenidate Products	0.36	0.39	0.27	0.27	1.47	1.52
Amphetamine Products	0.35	0.38	0.26	0.27	1.54	1.60
Methylphenidate Products	0.19	0.18	0.12	0.10	2.84	2.83
Anorectic (Weight-Loss) Stimulants	0.07	0.08	0.02	0.03	*	*
Provigil®	0.04	0.04	0.03	0.01	*	*
Any Other Prescription Stimulant ¹	0.11	0.10	0.02	0.02	1.20	1.93

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.106P Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 to 25 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 to 25: P Values from Tests of Differences of Percentages, 2019 Versus 2018

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	0.5179		0.0505		0.0280	
Amphetamine or Methylphenidate Products	0.6220		0.0658		0.0275	
Amphetamine Products	0.5754		0.0585		0.0254	
Methylphenidate Products	0.6718		0.0858		0.0537	
Anorectic (Weight-Loss) Stimulants	0.8984		0.2956		*	
Provigil®	0.9701		0.0973		*	
Any Other Prescription Stimulant ¹	0.1749		0.8766		0.6734	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.107C Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 26 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 26 or Older: Standard Errors of Numbers in Thousands, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse	Misuse	Misuse	Misuse
			in Past Year among Total Population (2018)	in Past Year among Total Population (2019)	in Past Year among Any Users of Stimulant Subtypes (2018)	in Past Year among Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	329	344	134	138	134	139
Amphetamine or Methylphenidate Products	273	293	126	130	126	132
Amphetamine Products	258	271	125	128	125	129
Methylphenidate Products	109	134	39	49	39	49
Anorectic (Weight-Loss) Stimulants	137	136	27	40	27	40
Provigil®	54	68	27	27	*	*
Any Other Prescription Stimulant ¹	140	151	24	*	24	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.107D Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 26 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 26 or Older: Standard Errors of Percentages, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	0.15	0.16	0.06	0.06	1.05	1.05
Amphetamine or Methylphenidate Products	0.13	0.14	0.06	0.06	1.37	1.38
Amphetamine Products	0.12	0.13	0.06	0.06	1.47	1.47
Methylphenidate Products	0.05	0.06	0.02	0.02	2.48	2.65
Anorectic (Weight-Loss) Stimulants	0.06	0.06	0.01	0.02	1.17	1.97
Provigil®	0.03	0.03	0.01	0.01	*	*
Any Other Prescription Stimulant ¹	0.07	0.07	0.01	*	1.26	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.107P Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 26 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 26 or Older: *P* Values from Tests of Differences of Percentages, 2019 Versus 2018

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	0.5891		0.9846		0.7883	
Amphetamine or Methylphenidate Products	0.1875		0.7924		0.5589	
Amphetamine Products	0.2572		0.7937		0.6269	
Methylphenidate Products	0.1617		0.4861		0.9933	
Anorectic (Weight-Loss) Stimulants	0.1090		0.4180		0.2125	
Provigil®	0.1382		0.5937		*	
Any Other Prescription Stimulant ¹	0.8266		*		*	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.