

**Table 10.6C Co-Occurring Substance Use Disorder and Mental Illness in Past Year among Persons Aged 18 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2015-2019**

<b>Substance Use Disorder (SUD)/ Mental Illness/Age Group</b>	<b>2015</b>	<b>2016</b>	<b>2017</b>	<b>2018</b>	<b>2019</b>
<b>SUD and Any Mental Illness</b>	247	253	246	276	283
18-25	75	83	86	88	85
26 or Older	234	236	226	261	265
26-49	166	179	161	192	192
50 or Older	159	157	158	170	160
<b>SUD and Serious Mental Illness</b>	124	141	152	149	158
18-25	42	49	57	53	54
26 or Older	116	130	138	140	145
26-49	87	98	100	105	111
50 or Older	75	81	95	95	91
<b>SUD and Any Mental Illness Excluding Serious Mental Illness</b>	208	206	194	220	220
18-25	65	68	66	70	69
26 or Older	200	194	180	211	208
26-49	141	140	128	155	150
50 or Older	137	135	131	145	134

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to persons with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.

**Table 10.6D Co-Occurring Substance Use Disorder and Mental Illness in Past Year among Persons Aged 18 or Older, by Age Group: Standard Errors of Percentages, 2015-2019**

<b>Substance Use Disorder (SUD)/ Mental Illness/Age Group</b>	<b>2015</b>	<b>2016</b>	<b>2017</b>	<b>2018</b>	<b>2019</b>
<b>SUD and Any Mental Illness</b>	0.10	0.10	0.10	0.11	0.11
18-25	0.21	0.24	0.25	0.26	0.25
26 or Older	0.11	0.11	0.11	0.12	0.12
26-49	0.17	0.18	0.16	0.19	0.19
50 or Older	0.15	0.14	0.14	0.15	0.14
<b>SUD and Serious Mental Illness</b>	0.05	0.06	0.06	0.06	0.06
18-25	0.12	0.14	0.17	0.16	0.16
26 or Older	0.06	0.06	0.06	0.07	0.07
26-49	0.09	0.10	0.10	0.10	0.11
50 or Older	0.07	0.07	0.08	0.08	0.08
<b>SUD and Any Mental Illness Excluding Serious Mental Illness</b>	0.09	0.08	0.08	0.09	0.09
18-25	0.19	0.20	0.19	0.21	0.20
26 or Older	0.10	0.09	0.08	0.10	0.10
26-49	0.14	0.14	0.13	0.15	0.15
50 or Older	0.13	0.12	0.12	0.13	0.12

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to persons with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.

**Table 10.6P Co-Occurring Substance Use Disorder and Mental Illness in Past Year among Persons Aged 18 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2019 Versus 2015-2018**

<b>Substance Use Disorder (SUD)/ Mental Illness/Age Group</b>	<b>2015</b>	<b>2016</b>	<b>2017</b>	<b>2018</b>	<b>2019</b>
<b>SUD and Any Mental Illness</b>	0.0019	0.0042	0.0181	0.4480	
18-25	0.0000	0.0000	0.0418	0.2106	
26 or Older	0.0570	0.0719	0.0474	0.6556	
26-49	0.0017	0.0119	0.0015	0.5007	
50 or Older	0.7428	0.9544	0.7173	0.9921	
<b>SUD and Serious Mental Illness</b>	0.0000	0.0001	0.0554	0.0836	
18-25	0.0000	0.0003	0.7285	0.2418	
26 or Older	0.0000	0.0020	0.0411	0.1360	
26-49	0.0000	0.0003	0.0078	0.0753	
50 or Older	0.1825	0.4665	0.8094	0.7600	
<b>SUD and Any Mental Illness Excluding Serious Mental Illness</b>	0.8799	0.4776	0.1065	0.7839	
18-25	0.0346	0.0146	0.0226	0.4934	
26 or Older	0.7312	0.9623	0.3071	0.6496	
26-49	0.4459	0.5864	0.0375	0.6862	
50 or Older	0.2291	0.6860	0.5471	0.8507	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) that are used in the comparisons are a subset of estimates of any mental illness (AMI) because SMI is limited to persons with AMI that resulted in serious functional impairment. The mental illness estimates that are used in the comparisons are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015-2019.