

Table 11.13C Type of Treatment Received in Past Year for Depression among Persons Aged 12 to 17 with Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year: Standard Errors of Numbers in Thousands, 2004-2019

| MDE/Type of Treatment | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 |
|--|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| MDE | | | | | | | | | | | | | | | | |
| Saw or Talked to a Health Professional OR Used Prescription Medication | 39 | 38 | 37 | 40 | 35 | 35 | 35 | 35 | 37 | 43 | 50 | 51 | 51 | 54 | 59 | 63 |
| Saw or Talked to a Health Professional BUT Did Not Use Prescription Medication | 30 | 28 | 28 | 27 | 27 | 26 | 25 | 26 | 27 | 31 | 36 | 38 | 37 | 37 | 46 | 42 |
| Used Prescription Medication BUT Did Not See or Talk to a Health Professional | 11 | 10 | 9 | 11 | 10 | 9 | 12 | 10 | 9 | 17 | 14 | 14 | 11 | 13 | 12 | 19 |
| Saw or Talked to a Health Professional AND Used Prescription Medication | 25 | 24 | 21 | 24 | 21 | 20 | 20 | 21 | 23 | 23 | 32 | 30 | 34 | 37 | 33 | 41 |
| MDE with Severe Impairment | | | | | | | | | | | | | | | | |
| Saw or Talked to a Health Professional OR Used Prescription Medication | -- | -- | 34 | 34 | 31 | 31 | 30 | 31 | 34 | 40 | 42 | 45 | 45 | 50 | 52 | 57 |
| Saw or Talked to a Health Professional BUT Did Not Use Prescription Medication | -- | -- | 25 | 23 | 23 | 23 | 22 | 22 | 23 | 28 | 30 | 33 | 30 | 33 | 40 | 38 |
| Used Prescription Medication BUT Did Not See or Talk to a Health Professional | -- | -- | 8 | 9 | 9 | 9 | 9 | 7 | 9 | 17 | 13 | 13 | 10 | 12 | 10 | 17 |
| Saw or Talked to a Health Professional AND Used Prescription Medication | -- | -- | 20 | 22 | 20 | 19 | 18 | 20 | 22 | 23 | 27 | 28 | 32 | 34 | 31 | 38 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown past year treatment for depression data were excluded.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004-2019.

Table 11.13D Type of Treatment Received in Past Year for Depression among Persons Aged 12 to 17 with Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year: Standard Errors of Percentages, 2004-2019

| MDE/Type of Treatment | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 |
|--|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| MDE | | | | | | | | | | | | | | | | |
| Saw or Talked to a Health Professional OR Used Prescription Medication | 1.35 | 1.40 | 1.54 | 1.51 | 1.45 | 1.51 | 1.48 | 1.45 | 1.35 | 1.35 | 1.42 | 1.37 | 1.29 | 1.26 | 1.36 | 1.34 |
| Saw or Talked to a Health Professional BUT Did Not Use Prescription Medication | 1.21 | 1.19 | 1.30 | 1.17 | 1.22 | 1.19 | 1.22 | 1.20 | 1.10 | 1.10 | 1.19 | 1.12 | 1.09 | 1.04 | 1.20 | 1.03 |
| Used Prescription Medication BUT Did Not See or Talk to a Health Professional | 0.47 | 0.45 | 0.47 | 0.52 | 0.48 | 0.47 | 0.63 | 0.50 | 0.40 | 0.65 | 0.50 | 0.47 | 0.36 | 0.41 | 0.36 | 0.50 |
| Saw or Talked to a Health Professional AND Used Prescription Medication | 1.03 | 1.02 | 0.98 | 1.10 | 1.00 | 0.99 | 0.95 | 0.97 | 0.97 | 0.86 | 1.07 | 0.94 | 1.02 | 1.01 | 0.89 | 0.98 |
| MDE with Severe Impairment | | | | | | | | | | | | | | | | |
| Saw or Talked to a Health Professional OR Used Prescription Medication | -- | -- | 1.97 | 1.90 | 1.69 | 1.82 | 1.78 | 1.76 | 1.67 | 1.61 | 1.66 | 1.61 | 1.57 | 1.55 | 1.66 | 1.64 |
| Saw or Talked to a Health Professional BUT Did Not Use Prescription Medication | -- | -- | 1.65 | 1.43 | 1.44 | 1.48 | 1.47 | 1.45 | 1.31 | 1.34 | 1.36 | 1.40 | 1.27 | 1.28 | 1.45 | 1.28 |
| Used Prescription Medication BUT Did Not See or Talk to a Health Professional | -- | -- | 0.61 | 0.66 | 0.61 | 0.63 | 0.63 | 0.51 | 0.54 | 0.87 | 0.63 | 0.59 | 0.45 | 0.51 | 0.43 | 0.61 |
| Saw or Talked to a Health Professional AND Used Prescription Medication | -- | -- | 1.34 | 1.44 | 1.30 | 1.24 | 1.24 | 1.29 | 1.29 | 1.14 | 1.26 | 1.15 | 1.35 | 1.31 | 1.18 | 1.29 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2019 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown past year treatment for depression data were excluded.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004-2019.

Table 11.13P Type of Treatment Received in Past Year for Depression among Persons Aged 12 to 17 with Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year: P Values from Tests of Differences of Percentages, 2019 Versus 2004-2018

| MDE/Type of Treatment | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 |
|--|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|------|
| MDE | | | | | | | | | | | | | | | | |
| Saw or Talked to a Health Professional OR Used Prescription Medication | 0.0653 | 0.0011 | 0.0021 | 0.0097 | 0.0027 | 0.0000 | 0.0014 | 0.0027 | 0.0012 | 0.0055 | 0.1792 | 0.0198 | 0.2718 | 0.2995 | 0.2264 | |
| Saw or Talked to a Health Professional BUT Did Not Use Prescription Medication | 0.2162 | 0.0865 | 0.7946 | 0.0833 | 0.4808 | 0.0745 | 0.2452 | 0.3677 | 0.2636 | 0.5605 | 0.4008 | 0.3258 | 0.0870 | 0.2382 | 0.7130 | |
| Used Prescription Medication BUT Did Not See or Talk to a Health Professional | 0.9555 | 0.2995 | 0.1517 | 0.3223 | 0.5456 | 0.1393 | 0.6542 | 0.2769 | 0.1121 | 0.7330 | 0.3826 | 0.4838 | 0.0194 | 0.1275 | 0.0556 | |
| Saw or Talked to a Health Professional AND Used Prescription Medication | 0.2447 | 0.0357 | 0.0006 | 0.2183 | 0.0023 | 0.0002 | 0.0044 | 0.0101 | 0.0138 | 0.0003 | 0.6138 | 0.0977 | 0.2310 | 0.6004 | 0.2129 | |
| MDE with Severe Impairment | | | | | | | | | | | | | | | | |
| Saw or Talked to a Health Professional OR Used Prescription Medication | -- | -- | 0.0247 | 0.0089 | 0.0014 | 0.0000 | 0.0001 | 0.0022 | 0.0003 | 0.0361 | 0.0198 | 0.0296 | 0.2691 | 0.2915 | 0.1789 | |
| Saw or Talked to a Health Professional BUT Did Not Use Prescription Medication | -- | -- | 0.9606 | 0.1355 | 0.3073 | 0.1074 | 0.1782 | 0.4239 | 0.0643 | 0.7450 | 0.2898 | 0.4858 | 0.0201 | 0.1814 | 0.7979 | |
| Used Prescription Medication BUT Did Not See or Talk to a Health Professional | -- | -- | 0.2526 | 0.3457 | 0.6683 | 0.2410 | 0.1702 | 0.0975 | 0.2829 | 0.3607 | 0.5616 | 0.4992 | 0.0587 | 0.1011 | 0.0526 | |
| Saw or Talked to a Health Professional AND Used Prescription Medication | -- | -- | 0.0129 | 0.1580 | 0.0055 | 0.0000 | 0.0032 | 0.0193 | 0.0238 | 0.0016 | 0.1147 | 0.1204 | 0.1186 | 0.4621 | 0.2430 | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2006 to 2010 estimates that are used in the comparisons may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2019 National Survey on Drug Use and Health: Methodological Summary and Definitions).

NOTE: Respondents with unknown past year treatment for depression data were excluded.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004-2019.