

Table 6.1C Drugs Used with Alcohol or within 2 Hours of Alcohol Use on Most Recent Use of Alcohol in Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group and Underage and Legal Drinking Age Groups: Standard Errors of Numbers in Thousands, 2018 and 2019

| Drug | AGE GROUP | | | | | | | | | | UNDERAGE AND LEGAL DRINKING AGE GROUP | | | |
|---------------------------|-----------------|-----------------|-------------------|-------------------|-----------------|-----------------|-------------------|-------------------|-----------------|-----------------|---------------------------------------|-------------------|-----------------|-----------------|
| | Aged 12+ (2018) | Aged 12+ (2019) | Aged 12-17 (2018) | Aged 12-17 (2019) | Aged 18+ (2018) | Aged 18+ (2019) | Aged 18-25 (2018) | Aged 18-25 (2019) | Aged 26+ (2018) | Aged 26+ (2019) | Aged 12-20 (2018) | Aged 12-20 (2019) | Aged 21+ (2018) | Aged 21+ (2019) |
| Marijuana | 315 | 308 | 30 | 35 | 312 | 307 | 125 | 111 | 278 | 275 | 76 | 75 | 294 | 302 |
| Cocaine (Including Crack) | 66 | 72 | 2 | 7 | 66 | 72 | 22 | 26 | 63 | 67 | 11 | 13 | 64 | 70 |
| Heroin | 29 | 27 | 3 | * | 29 | 27 | 3 | 2 | 29 | 27 | 3 | * | 29 | 27 |
| Hallucinogens | 27 | 37 | 5 | 2 | 27 | 37 | 17 | 10 | 21 | 35 | 8 | 5 | 26 | 36 |
| Inhalants | 13 | 18 | 2 | 5 | 13 | 17 | 6 | 8 | 12 | 16 | 3 | 5 | 13 | 17 |
| Methamphetamine | 53 | 58 | 1 | 1 | 53 | 58 | 6 | 7 | 53 | 57 | 1 | 6 | 53 | 57 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates are based on nonimputed responses to multiple questions; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 6.1D Drugs Used with Alcohol or within 2 Hours of Alcohol Use on Most Recent Use of Alcohol in Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group and Underage and Legal Drinking Age Groups: Standard Errors of Percentages, 2018 and 2019

| Drug | AGE GROUP | | | | | | | | | | UNDERAGE AND LEGAL DRINKING AGE GROUP | | | |
|---------------------------|-----------------|-----------------|-------------------|-------------------|-----------------|-----------------|-------------------|-------------------|-----------------|-----------------|---------------------------------------|-------------------|-----------------|-----------------|
| | Aged 12+ (2018) | Aged 12+ (2019) | Aged 12-17 (2018) | Aged 12-17 (2019) | Aged 18+ (2018) | Aged 18+ (2019) | Aged 18-25 (2018) | Aged 18-25 (2019) | Aged 26+ (2018) | Aged 26+ (2019) | Aged 12-20 (2018) | Aged 12-20 (2019) | Aged 21+ (2018) | Aged 21+ (2019) |
| Marijuana | 0.22 | 0.21 | 1.26 | 1.35 | 0.22 | 0.21 | 0.55 | 0.53 | 0.23 | 0.22 | 0.91 | 0.90 | 0.22 | 0.22 |
| Cocaine (Including Crack) | 0.05 | 0.05 | 0.11 | 0.31 | 0.05 | 0.05 | 0.12 | 0.14 | 0.05 | 0.06 | 0.15 | 0.18 | 0.05 | 0.05 |
| Heroin | 0.02 | 0.02 | 0.12 | * | 0.02 | 0.02 | 0.02 | 0.01 | 0.02 | 0.02 | 0.04 | * | 0.02 | 0.02 |
| Hallucinogens | 0.02 | 0.03 | 0.22 | 0.07 | 0.02 | 0.03 | 0.09 | 0.05 | 0.02 | 0.03 | 0.12 | 0.07 | 0.02 | 0.03 |
| Inhalants | 0.01 | 0.01 | 0.10 | 0.24 | 0.01 | 0.01 | 0.03 | 0.04 | 0.01 | 0.01 | 0.04 | 0.08 | 0.01 | 0.01 |
| Methamphetamine | 0.04 | 0.04 | 0.03 | 0.03 | 0.04 | 0.04 | 0.03 | 0.04 | 0.04 | 0.05 | 0.01 | 0.09 | 0.04 | 0.04 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates are based on nonimputed responses to multiple questions; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 6.1P Drugs Used with Alcohol or within 2 Hours of Alcohol Use on Most Recent Use of Alcohol in Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group and Underage and Legal Drinking Age Groups: P Values from Tests of Differences of Percentages, 2019 Versus 2018

| Drug | AGE GROUP | | | | | | | | | | UNDERAGE AND LEGAL DRINKING AGE GROUP | | | |
|---------------------------|-----------------|-----------------|-------------------|-------------------|-----------------|-----------------|-------------------|-------------------|-----------------|-----------------|---------------------------------------|-------------------|-----------------|-----------------|
| | Aged 12+ (2018) | Aged 12+ (2019) | Aged 12-17 (2018) | Aged 12-17 (2019) | Aged 18+ (2018) | Aged 18+ (2019) | Aged 18-25 (2018) | Aged 18-25 (2019) | Aged 26+ (2018) | Aged 26+ (2019) | Aged 12-20 (2018) | Aged 12-20 (2019) | Aged 21+ (2018) | Aged 21+ (2019) |
| Marijuana | 0.0008 | | 0.0741 | | 0.0014 | | 0.2061 | | 0.0016 | | 0.4986 | | 0.0007 | |
| Cocaine (Including Crack) | 0.7517 | | 0.5522 | | 0.7850 | | 0.5899 | | 0.9200 | | 0.8718 | | 0.7668 | |
| Heroin | 0.6816 | | * | | 0.6330 | | 0.3943 | | 0.5710 | | * | | 0.6074 | |
| Hallucinogens | 0.8141 | | 0.6107 | | 0.7706 | | 0.0644 | | 0.2213 | | 0.3326 | | 0.6513 | |
| Inhalants | 0.6529 | | 0.5192 | | 0.7583 | | 0.6868 | | 0.8605 | | 0.7166 | | 0.7056 | |
| Methamphetamine | 0.1698 | | 0.9874 | | 0.1688 | | 0.5332 | | 0.1473 | | 0.3196 | | 0.1953 | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates that are used in the comparisons are based on nonimputed responses to multiple questions; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.