

**Table 6.5C Source Where Pain Relievers Were Obtained for Most Recent Misuse among Past Year Misusers Aged 12 or Older, by Age Group:  
Standard Errors of Numbers in Thousands, 2018 and 2019**

<b>Source for Most Recent Misuse among Past Year Misusers of Pain Relievers</b>	<b>Aged 12+ (2018)</b>	<b>Aged 12+ (2019)</b>	<b>Aged 12-17 (2018)</b>	<b>Aged 12-17 (2019)</b>	<b>Aged 18+ (2018)</b>	<b>Aged 18+ (2019)</b>	<b>Aged 18-25 (2018)</b>	<b>Aged 18-25 (2019)</b>	<b>Aged 26+ (2018)</b>	<b>Aged 26+ (2019)</b>
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	189	184	24	22	188	182	37	42	184	176
Prescription from One Doctor	180	184	23	20	179	182	35	40	175	177
Prescriptions from More Than One Doctor	40	20	7	7	40	19	7	9	39	16
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	39	22	3	4	39	21	5	9	38	19
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	190	198	26	26	188	197	58	53	177	190
From Friend or Relative for Free	167	169	22	20	166	168	50	44	159	162
Bought from Friend or Relative	79	79	9	12	79	78	23	25	75	74
Took from Friend or Relative without Asking	36	55	10	11	34	53	19	22	28	49
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	62	65	11	8	61	64	22	19	56	61
<b>SOME OTHER WAY<sup>1</sup></b>	53	66	10	7	52	66	17	20	49	64

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.5D Source Where Pain Relievers Were Obtained for Most Recent Misuse among Past Year Misusers Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2018 and 2019**

Source for Most Recent Misuse among Past Year Misusers of Pain Relievers	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	1.53	1.56	3.21	3.59	1.61	1.64	1.81	2.05	1.92	1.95
Prescription from One Doctor	1.49	1.58	3.12	3.45	1.57	1.65	1.73	1.97	1.87	1.97
Prescriptions from More Than One Doctor	0.43	0.22	1.23	1.56	0.45	0.22	0.40	0.58	0.56	0.23
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.42	0.24	0.54	0.85	0.44	0.25	0.29	0.56	0.55	0.27
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	1.51	1.54	3.25	3.51	1.59	1.61	2.06	2.11	1.89	1.94
From Friend or Relative for Free	1.45	1.46	3.14	3.29	1.53	1.52	2.12	2.09	1.84	1.81
Bought from Friend or Relative	0.82	0.82	1.54	2.37	0.87	0.86	1.23	1.45	1.04	1.00
Took from Friend or Relative without Asking	0.38	0.59	1.57	2.27	0.39	0.61	1.09	1.30	0.40	0.69
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.65	0.70	1.72	1.59	0.69	0.73	1.18	1.12	0.79	0.84
<b>SOME OTHER WAY<sup>1</sup></b>	0.57	0.70	1.73	1.58	0.59	0.73	0.93	1.19	0.70	0.88

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.5P Source Where Pain Relievers Were Obtained for Most Recent Misuse among Past Year Misusers Aged 12 or Older, by Age Group:  
P Values from Tests of Differences of Percentages, 2019 Versus 2018**

Source for Most Recent Misuse among Past Year Misusers of Pain Relievers	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	0.9753		0.7710		0.9378		0.0388		0.5078	
Prescription from One Doctor	0.6501		0.9650		0.6590		0.1151		0.9956	
Prescriptions from More Than One Doctor	0.0546		0.4686		0.0352		0.2561		0.0126	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.8352		0.8519		0.8254		0.2705		0.6189	
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	0.8098		0.7173		0.7699		0.0223		0.7260	
From Friend or Relative for Free	0.4398		0.5047		0.4680		0.1238		0.7494	
Bought from Friend or Relative	0.7819		0.3505		0.6695		0.4223		0.8710	
Took from Friend or Relative without Asking	0.0522		0.4549		0.0558		0.6972		0.0136	
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.7132		0.4264		0.7970		0.6334		0.9418	
<b>SOME OTHER WAY<sup>1</sup></b>	0.2852		0.5759		0.2280		0.2419		0.3823	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.7C Source Where Stimulants Were Obtained for Most Recent Misuse among Past Year Misusers Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2018 and 2019**

<b>Source for Most Recent Misuse among Past Year Misusers of Stimulants</b>	<b>Aged 12+ (2018)</b>	<b>Aged 12+ (2019)</b>	<b>Aged 12-17 (2018)</b>	<b>Aged 12-17 (2019)</b>	<b>Aged 18+ (2018)</b>	<b>Aged 18+ (2019)</b>	<b>Aged 18-25 (2018)</b>	<b>Aged 18-25 (2019)</b>	<b>Aged 26+ (2018)</b>	<b>Aged 26+ (2019)</b>
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	58	65	11	16	57	63	26	28	50	56
Prescription from One Doctor	57	64	9	15	55	63	26	28	49	56
Prescriptions from More Than One Doctor	14	9	6	5	13	7	7	3	11	6
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	2	2	*	*	2	*	1	*	2
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	157	150	24	31	155	146	89	86	119	117
From Friend or Relative for Free	128	132	16	27	127	130	66	63	102	108
Bought from Friend or Relative	74	71	15	12	73	69	48	45	54	53
Took from Friend or Relative without Asking	24	29	10	9	21	27	10	15	19	23
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	35	38	7	8	34	37	21	19	26	31
<b>SOME OTHER WAY<sup>1</sup></b>	21	30	3	2	21	30	15	16	14	24

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.7D Source Where Stimulants Were Obtained for Most Recent Misuse among Past Year Misusers Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2018 and 2019**

<b>Source for Most Recent Misuse among Past Year Misusers of Stimulants</b>	<b>Aged 12+ (2018)</b>	<b>Aged 12+ (2019)</b>	<b>Aged 12-17 (2018)</b>	<b>Aged 12-17 (2019)</b>	<b>Aged 18+ (2018)</b>	<b>Aged 18+ (2019)</b>	<b>Aged 18-25 (2018)</b>	<b>Aged 18-25 (2019)</b>	<b>Aged 26+ (2018)</b>	<b>Aged 26+ (2019)</b>
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	1.10	1.21	2.97	3.61	1.16	1.29	1.14	1.32	1.94	2.04
Prescription from One Doctor	1.08	1.20	2.52	3.42	1.14	1.29	1.11	1.32	1.90	2.03
Prescriptions from More Than One Doctor	0.29	0.18	1.71	1.34	0.28	0.16	0.31	0.15	0.45	0.25
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.04	0.05	0.58	*	*	0.06	*	0.04	*	0.09
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	1.32	1.39	3.50	3.81	1.39	1.49	1.54	1.67	2.17	2.38
From Friend or Relative for Free	1.63	1.78	3.90	4.17	1.72	1.92	1.89	2.16	2.66	2.85
Bought from Friend or Relative	1.29	1.39	3.60	2.72	1.36	1.50	1.77	1.93	1.99	2.04
Took from Friend or Relative without Asking	0.47	0.59	2.73	2.08	0.46	0.61	0.46	0.79	0.76	0.91
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.68	0.75	2.02	1.95	0.71	0.80	0.96	0.96	1.05	1.20
<b>SOME OTHER WAY<sup>1</sup></b>	0.42	0.61	0.85	0.56	0.44	0.67	0.68	0.82	0.58	0.95

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.7P Source Where Stimulants Were Obtained for Most Recent Misuse among Past Year Misusers Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2019 Versus 2018**

<b>Source for Most Recent Misuse among Past Year Misusers of Stimulants</b>	<b>Aged 12+ (2018)</b>	<b>Aged 12+ (2019)</b>	<b>Aged 12-17 (2018)</b>	<b>Aged 12-17 (2019)</b>	<b>Aged 18+ (2018)</b>	<b>Aged 18+ (2019)</b>	<b>Aged 18-25 (2018)</b>	<b>Aged 18-25 (2019)</b>	<b>Aged 26+ (2018)</b>	<b>Aged 26+ (2019)</b>
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	0.2608		0.5245		0.3501		0.3980		0.6005	
Prescription from One Doctor	0.2107		0.3224		0.3029		0.3357		0.5644	
Prescriptions from More Than One Doctor	0.5593		0.8039		0.4827		0.5201		0.6491	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.8348		*		*		*		*	
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	0.0845		0.7037		0.1037		0.4745		0.1491	
From Friend or Relative for Free	0.7847		0.0356		0.5500		0.5536		0.6034	
Bought from Friend or Relative	0.1448		0.0161		0.3247		0.6708		0.5170	
Took from Friend or Relative without Asking	0.9424		0.4302		0.8167		0.1787		0.5500	
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.8707		0.9079		0.8698		0.5799		0.5127	
<b>SOME OTHER WAY<sup>1</sup></b>	0.0762		0.4823		0.0596		0.4233		0.0539	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.9C Source Where Tranquilizers Were Obtained for Most Recent Misuse among Past Year Misusers Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2018 and 2019**

<b>Source for Most Recent Misuse among Past Year Misusers of Tranquilizers</b>	<b>Aged 12+ (2018)</b>	<b>Aged 12+ (2019)</b>	<b>Aged 12-17 (2018)</b>	<b>Aged 12-17 (2019)</b>	<b>Aged 18+ (2018)</b>	<b>Aged 18+ (2019)</b>	<b>Aged 18-25 (2018)</b>	<b>Aged 18-25 (2019)</b>	<b>Aged 26+ (2018)</b>	<b>Aged 26+ (2019)</b>
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	115	123	9	11	115	122	16	22	114	118
Prescription from One Doctor	115	119	9	8	115	119	16	21	113	115
Prescriptions from More Than One Doctor	11	27	3	*	10	27	4	7	9	26
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	7	7	2	*	7	4	1	4	7	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	179	146	23	24	177	145	62	55	164	134
From Friend or Relative for Free	139	127	18	19	138	125	50	46	128	117
Bought from Friend or Relative	100	60	12	13	99	59	33	26	92	52
Took from Friend or Relative without Asking	43	39	6	5	43	39	13	16	41	35
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	46	54	16	18	44	50	27	24	35	43
<b>SOME OTHER WAY<sup>1</sup></b>	35	35	8	6	35	34	14	16	32	30

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.9D Source Where Tranquilizers Were Obtained for Most Recent Misuse among Past Year Misusers Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2018 and 2019**

Source for Most Recent Misuse among Past Year Misusers of Tranquilizers	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	1.78	1.95	2.29	2.79	1.89	2.07	1.06	1.61	2.54	2.66
Prescription from One Doctor	1.77	1.93	2.16	2.22	1.89	2.05	1.05	1.52	2.54	2.62
Prescriptions from More Than One Doctor	0.19	0.54	0.71	*	0.20	0.58	0.28	0.56	0.25	0.77
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.13	0.14	0.49	*	0.14	0.08	0.08	0.29	0.19	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	1.83	1.94	3.94	4.42	1.94	2.07	1.92	2.36	2.63	2.64
From Friend or Relative for Free	1.90	1.94	3.80	4.26	2.05	2.05	2.26	2.57	2.74	2.69
Bought from Friend or Relative	1.61	1.14	2.86	3.16	1.71	1.20	1.90	1.89	2.27	1.45
Took from Friend or Relative without Asking	0.77	0.77	1.51	1.43	0.82	0.83	0.87	1.25	1.10	1.03
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.82	1.02	3.53	4.31	0.83	1.04	1.64	1.79	0.94	1.22
<b>SOME OTHER WAY<sup>1</sup></b>	0.63	0.68	2.09	1.58	0.66	0.73	0.87	1.26	0.85	0.89

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.9P Source Where Tranquilizers Were Obtained for Most Recent Misuse among Past Year Misusers Aged 12 or Older, by Age Group:  
P Values from Tests of Differences of Percentages, 2019 Versus 2018**

Source for Most Recent Misuse among Past Year Misusers of Tranquilizers	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	0.1588		0.8445		0.1441		0.0658		0.2953	
Prescription from One Doctor	0.2428		0.6868		0.2173		0.2115		0.3448	
Prescriptions from More Than One Doctor	0.2495		*		0.2231		0.1030		0.4345	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.8723		*		0.4630		0.4836		*	
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	0.0714		0.4829		0.0529		0.2187		0.1159	
From Friend or Relative for Free	0.3117		0.2351		0.2369		0.5148		0.3161	
Bought from Friend or Relative	0.2167		0.6623		0.2345		0.3225		0.4180	
Took from Friend or Relative without Asking	0.8042		0.7263		0.7695		0.4784		0.9602	
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.8863		0.7002		0.8142		0.4497		0.3616	
<b>SOME OTHER WAY<sup>1</sup></b>	0.3502		0.7009		0.3043		0.1548		0.6349	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.11C Source Where Sedatives Were Obtained for Most Recent Misuse among Past Year Misusers Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2018 and 2019**

Source for Most Recent Misuse among Past Year Misusers of Sedatives	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	61	68	*	*	60	68	*	*	60	67
Prescription from One Doctor	60	68	*	*	59	68	*	*	59	67
Prescriptions from More Than One Doctor	11	2	*	*	11	2	*	*	11	2
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	3	2	*	*	3	*	*	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	69	74	*	*	67	73	*	*	63	70
From Friend or Relative for Free	67	67	*	*	65	67	*	*	62	64
Bought from Friend or Relative	11	24	*	*	10	24	*	*	9	22
Took from Friend or Relative without Asking	9	22	*	*	7	19	*	*	7	18
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	9	10	*	*	7	10	*	*	5	*
<b>SOME OTHER WAY<sup>1</sup></b>	14	7	*	*	14	7	*	*	10	0

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.11D Source Where Sedatives Were Obtained for Most Recent Misuse among Past Year Misusers Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2018 and 2019**

Source for Most Recent Misuse among Past Year Misusers of Sedatives	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	4.54	4.75	*	*	4.75	5.07	*	*	5.58	5.91
Prescription from One Doctor	4.54	4.76	*	*	4.75	5.08	*	*	5.58	5.91
Prescriptions from More Than One Doctor	1.05	0.21	*	*	1.12	0.23	*	*	1.33	0.20
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.25	0.20	*	*	0.26	*	*	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	4.46	4.74	*	*	4.67	5.07	*	*	5.50	5.90
From Friend or Relative for Free	4.43	4.79	*	*	4.62	5.02	*	*	5.45	5.83
Bought from Friend or Relative	1.02	2.21	*	*	1.07	2.32	*	*	1.14	2.61
Took from Friend or Relative without Asking	0.87	2.03	*	*	0.74	1.86	*	*	0.82	2.14
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.85	0.98	*	*	0.76	1.03	*	*	0.59	*
<b>SOME OTHER WAY<sup>1</sup></b>	1.31	0.65	*	*	1.39	0.66	*	*	1.21	0.06

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.11P Source Where Sedatives Were Obtained for Most Recent Misuse among Past Year Misusers Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2019 Versus 2018**

Source for Most Recent Misuse among Past Year Misusers of Sedatives	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>										
	0.3966		*		0.3590		*		0.4168	
Prescription from One Doctor	0.2666		*		0.2259		*		0.2830	
Prescriptions from More Than One Doctor	0.1147		*		0.1153		*		0.1154	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.8909		*		*		*		*	
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>										
	0.6621		*		0.5873		*		0.5945	
From Friend or Relative for Free	0.1310		*		0.1613		*		0.2026	
Bought from Friend or Relative	0.1377		*		0.1767		*		0.2539	
Took from Friend or Relative without Asking	0.1229		*		0.2317		*		0.2608	
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>										
	0.7869		*		0.9952		*		*	
<b>SOME OTHER WAY<sup>1</sup></b>										
	0.1064		*		0.0869		*		0.0490	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.