

**Table 6.6C Source Where Friend or Relative Obtained Pain Relievers among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Pain Relievers from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2018 and 2019**

Source Where Friend or Relative Obtained Pain Relievers	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	153	153	17	*	152	152	43	39	146	148
Prescription from One Doctor	150	151	16	*	149	150	42	39	143	147
Prescriptions from More Than One Doctor	20	20	5	*	19	20	10	7	16	19
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	10	8	2	*	10	8	3	3	*	7
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	50	48	6	*	49	48	17	13	46	47
From Friend or Relative for Free	46	38	5	*	46	38	13	12	44	37
Bought from Friend or Relative	14	24	*	*	13	24	7	5	11	23
Took from Friend or Relative without Asking	13	19	1	*	13	18	8	4	10	17
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	19	24	4	*	19	23	8	6	18	22
<b>SOME OTHER WAY<sup>1</sup></b>	35	16	1	*	35	15	7	5	34	14

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.6D Source Where Friend or Relative Obtained Pain Relievers among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Pain Relievers from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Percentages, 2018 and 2019**

Source Where Friend or Relative Obtained Pain Relievers	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	1.72	1.66	4.62	*	1.78	1.68	2.77	2.56	2.15	2.02
Prescription from One Doctor	1.79	1.76	5.53	*	1.85	1.78	2.90	2.79	2.22	2.14
Prescriptions from More Than One Doctor	0.58	0.63	3.33	*	0.58	0.65	1.46	1.15	0.63	0.75
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.30	0.25	1.33	*	0.31	0.26	0.48	0.51	*	0.29
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	1.39	1.44	4.06	*	1.43	1.47	2.38	2.16	1.72	1.78
From Friend or Relative for Free	1.31	1.18	3.07	*	1.35	1.22	1.82	2.00	1.64	1.44
Bought from Friend or Relative	0.40	0.74	*	*	0.40	0.76	0.97	0.90	0.42	0.91
Took from Friend or Relative without Asking	0.37	0.59	0.80	*	0.39	0.57	1.18	0.79	0.38	0.68
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.55	0.75	2.33	*	0.57	0.73	1.19	1.03	0.69	0.86
<b>SOME OTHER WAY<sup>1</sup></b>	1.00	0.51	0.91	*	1.04	0.49	1.04	0.96	1.29	0.56

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.6P Source Where Friend or Relative Obtained Pain Relievers among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Pain Relievers from a Friend or Relative for Free in Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2019 Versus 2018**

Source Where Friend or Relative Obtained Pain Relievers	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	0.9145		*		0.6949		0.6580		0.8150	
Prescription from One Doctor	0.7845		*		0.6722		0.3383		0.9231	
Prescriptions from More Than One Doctor	0.6389		*		0.8705		0.2649		0.7777	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.9279		*		0.9256		0.9613		*	
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	0.7809		*		0.7810		0.8323		0.7091	
From Friend or Relative for Free	0.8480		*		0.9430		0.4958		0.7872	
Bought from Friend or Relative	0.4656		*		0.3600		0.3963		0.2062	
Took from Friend or Relative without Asking	0.6989		*		0.8960		0.3271		0.7549	
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.8048		*		0.8429		0.7128		0.9593	
<b>SOME OTHER WAY<sup>1</sup></b>	0.3556		*		0.2416		0.7803		0.2600	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.8C Source Where Friend or Relative Obtained Stimulants among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Stimulants from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2018 and 2019**

Source Where Friend or Relative Obtained Stimulants	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	113	111	*	*	112	108	57	53	93	89
Prescription from One Doctor	112	111	*	*	111	108	56	53	93	89
Prescriptions from More Than One Doctor	13	12	*	*	13	9	10	8	9	4
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	34	47	*	*	33	46	20	20	26	42
From Friend or Relative for Free	27	38	*	*	26	37	17	16	19	34
Bought from Friend or Relative	20	27	*	*	20	27	8	12	18	24
Took from Friend or Relative without Asking	7	0	*	*	6	*	6	*	*	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	15	15	*	*	15	13	7	2	13	13
<b>SOME OTHER WAY<sup>1</sup></b>	9	14	*	*	9	14	4	4	*	13

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.8D Source Where Friend or Relative Obtained Stimulants among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Stimulants from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Percentages, 2018 and 2019**

Source Where Friend or Relative Obtained Stimulants	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	1.45	1.94	*	*	1.49	2.04	1.97	2.14	2.14	3.04
Prescription from One Doctor	1.54	1.98	*	*	1.58	2.07	2.11	2.31	2.22	3.04
Prescriptions from More Than One Doctor	0.54	0.53	*	*	0.56	0.41	0.94	0.91	0.67	0.32
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	1.32	1.82	*	*	1.35	1.92	1.85	2.10	1.88	2.83
From Friend or Relative for Free	1.04	1.55	*	*	1.07	1.63	1.63	1.78	1.40	2.43
Bought from Friend or Relative	0.79	1.10	*	*	0.83	1.18	0.80	1.28	1.30	1.75
Took from Friend or Relative without Asking	0.30	0.01	*	*	0.24	*	0.57	*	*	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.62	0.62	*	*	0.65	0.62	0.71	0.27	0.97	0.99
<b>SOME OTHER WAY<sup>1</sup></b>	0.37	0.57	*	*	0.38	0.62	0.37	0.50	*	0.96

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.8P Source Where Friend or Relative Obtained Stimulants among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Stimulants from a Friend or Relative for Free in Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2019 Versus 2018**

Source Where Friend or Relative Obtained Stimulants	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	0.2868		*		0.3180		0.9030		0.2156	
Prescription from One Doctor	0.3212		*		0.4975		0.8805		0.3517	
Prescriptions from More Than One Doctor	0.9084		*		0.2878		0.9251		0.1643	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*		*	
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	0.2879		*		0.2545		0.8099		0.2272	
From Friend or Relative for Free	0.2157		*		0.2230		0.9534		0.1404	
Bought from Friend or Relative	0.7177		*		0.7558		0.2967		0.8896	
Took from Friend or Relative without Asking	0.0726		*		*		*		*	
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.8666		*		0.5405		0.1153		0.9305	
<b>SOME OTHER WAY<sup>1</sup></b>	0.5713		*		0.5154		0.7909		*	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.10C Source Where Friend or Relative Obtained Tranquilizers among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Tranquilizers from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2018 and 2019**

Source Where Friend or Relative Obtained Tranquilizers	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	129	112	*	*	128	111	38	37	122	107
Prescription from One Doctor	128	110	*	*	128	109	37	37	121	105
Prescriptions from More Than One Doctor	10	14	*	*	9	14	6	2	7	14
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	10	*	*	4	10	*	3	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	37	40	*	*	36	37	23	20	27	32
From Friend or Relative for Free	27	28	*	*	26	27	14	16	22	22
Bought from Friend or Relative	23	28	*	*	22	26	18	13	13	23
Took from Friend or Relative without Asking	11	6	*	*	11	5	2	*	11	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	22	22	*	*	21	20	16	14	14	15
<b>SOME OTHER WAY<sup>1</sup></b>	10	15	*	*	10	14	5	1	9	14

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.10D Source Where Friend or Relative Obtained Tranquilizers among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Tranquilizers from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Percentages, 2018 and 2019**

Source Where Friend or Relative Obtained Tranquilizers	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	1.72	2.02	*	*	1.74	2.06	3.46	3.68	1.80	2.42
Prescription from One Doctor	1.75	2.19	*	*	1.78	2.25	3.47	3.73	1.85	2.72
Prescriptions from More Than One Doctor	0.39	0.67	*	*	0.39	0.70	0.86	0.45	0.43	0.95
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.17	0.46	*	*	0.18	0.48	*	0.59	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	1.45	1.76	*	*	1.46	1.77	3.03	3.26	1.56	2.07
From Friend or Relative for Free	1.07	1.26	*	*	1.07	1.31	1.99	2.72	1.26	1.47
Bought from Friend or Relative	0.90	1.26	*	*	0.92	1.28	2.53	2.35	0.74	1.52
Took from Friend or Relative without Asking	0.44	0.27	*	*	0.45	0.25	0.29	*	0.62	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.87	1.01	*	*	0.89	1.00	2.29	2.48	0.80	1.02
<b>SOME OTHER WAY<sup>1</sup></b>	0.40	0.67	*	*	0.42	0.70	0.75	0.18	0.50	0.96

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.10P Source Where Friend or Relative Obtained Tranquilizers among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Tranquilizers from a Friend or Relative for Free in Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2019 Versus 2018**

Source Where Friend or Relative Obtained Tranquilizers	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	0.7889		*		0.9250		0.4981		0.4988	
Prescription from One Doctor	0.4607		*		0.5544		0.4219		0.1880	
Prescriptions from More Than One Doctor	0.2378		*		0.2307		0.2752		0.1055	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2848		*		0.2823		*		*	
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	0.9670		*		0.9385		0.8133		0.7335	
From Friend or Relative for Free	0.6647		*		0.8582		0.6672		0.6376	
Bought from Friend or Relative	0.4486		*		0.5757		0.3387		0.1322	
Took from Friend or Relative without Asking	0.5246		*		0.4668		*		*	
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.9974		*		0.7302		0.7043		0.9960	
<b>SOME OTHER WAY<sup>1</sup></b>	0.4210		*		0.5163		0.1241		0.2817	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.12C Source Where Friend or Relative Obtained Sedatives among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Sedatives from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2018 and 2019**

Source Where Friend or Relative Obtained Sedatives	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	61	64	*	*	60	*	*	*	*	*
Prescription from One Doctor	60	64	*	*	60	64	*	*	*	*
Prescriptions from More Than One Doctor	3	5	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	12	17	*	*	11	*	*	*	*	*
From Friend or Relative for Free	10	*	*	*	10	*	*	*	*	*
Bought from Friend or Relative	3	7	*	*	3	*	*	*	*	*
Took from Friend or Relative without Asking	5	3	*	*	*	*	*	*	*	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	*	0	*	*	*	0	*	*	*	*
<b>SOME OTHER WAY<sup>1</sup></b>	*	*	*	*	*	*	*	*	*	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.12D Source Where Friend or Relative Obtained Sedatives among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Sedatives from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Percentages, 2018 and 2019**

Source Where Friend or Relative Obtained Sedatives	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	3.06	3.46	*	*	3.12	*	*	*	*	*
Prescription from One Doctor	3.07	3.62	*	*	3.14	3.69	*	*	*	*
Prescriptions from More Than One Doctor	0.57	1.05	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	2.18	3.46	*	*	2.12	*	*	*	*	*
From Friend or Relative for Free	1.89	*	*	*	1.91	*	*	*	*	*
Bought from Friend or Relative	0.58	1.49	*	*	0.61	*	*	*	*	*
Took from Friend or Relative without Asking	1.00	0.70	*	*	*	*	*	*	*	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	*	0.03	*	*	*	0.04	*	*	*	*
<b>SOME OTHER WAY<sup>1</sup></b>	*	*	*	*	*	*	*	*	*	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 6.12P Source Where Friend or Relative Obtained Sedatives among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Sedatives from a Friend or Relative for Free in Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2019 Versus 2018**

Source Where Friend or Relative Obtained Sedatives	Aged 12+ (2018)	Aged 12+ (2019)	Aged 12-17 (2018)	Aged 12-17 (2019)	Aged 18+ (2018)	Aged 18+ (2019)	Aged 18-25 (2018)	Aged 18-25 (2019)	Aged 26+ (2018)	Aged 26+ (2019)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	0.8915		*		*		*		*	
Prescription from One Doctor	0.9831		*		0.9875		*		*	
Prescriptions from More Than One Doctor	0.5433		*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*		*	
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	0.3797		*		*		*		*	
From Friend or Relative for Free	*		*		*		*		*	
Bought from Friend or Relative	0.3405		*		*		*		*	
Took from Friend or Relative without Asking	0.7969		*		*		*		*	
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	*		*		*		*		*	
<b>SOME OTHER WAY<sup>1</sup></b>	*		*		*		*		*	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.