Table 7.68D Marijuana Use in Past Month among Persons Aged 12 or Older, by Age Group: Standard Errors of Percentages, 1971-2019

Survey Year	Aged 12-17	Aged 18-25	Aged 26-34
1971	0.98	1.74	1.13
1972	1.00	2.02	1.57
1974	1.23	1.86	1.21
1976	1.22	1.82	1.41
1977	1.22	1.44	1.66
1979	2.51	5.03	3.34
1982	1.82	3.95	3.06
1985	1.82	3.15	2.96
1988	0.99	2.38	2.22
1990	0.93	2.01	1.74
1991	0.60	1.82	1.16
1992	0.60	1.62	1.45
1993	0.67	1.70	1.22
1994	0.50	0.81	0.47
1995	0.55	0.71	0.47
1996	0.54	0.80	0.42
1997	0.56	0.88	0.54
1998	0.57	0.68	0.46
1999	0.20	0.38	0.32
2000	0.21	0.34	0.33
2001	0.24	0.39	0.45
2002	0.24	0.36	0.45
2003	0.24	0.37	0.44
2004	0.23	0.37	0.47
2005	0.22	0.37	0.51
2006	0.21	0.35	0.47
2007	0.22	0.37	0.44
2008	0.22	0.37	0.48
2009	0.24	0.38	0.47
2010	0.25	0.38	0.54
2011	0.24	0.39	0.54
2012	0.22	0.39	0.52
2013	0.23	0.39	0.62
2014	0.27	0.45	0.45
2015	0.24	0.40	0.43
2016	0.24	0.42	0.46
2017	0.24	0.46	0.47
2018	0.25	0.47	0.50
2019	0.26	0.44	0.50

^{* =} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates for 1979 through 1993 have been adjusted for comparability with estimates for 1994 through 1998.

NOTE: Estimates for 1999 through 2019 have not been adjusted for comparability. See Appendix C of the *Results from the 2004 National Survey on Drug Use and Health: National Findings* for information on the comparability of the 1999 and later estimates with pre-1999 estimates.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 1971-2019.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2019 National Survey on Drug Use and Health: Methodological Summary and Definitions).

NOTE: Standard errors for 1971 through 1977 were calculated by multiplying the standard error of the percentage (assuming simple random sampling) by the square root of the estimated "design effect" (i.e., 1.25). See Appendix A of the *National Household Survey on Drug Abuse: Main Findings 1979* for further information on these procedures.

NOTE: Estimates for 1971 through 1977 have been adjusted to improve their comparability with estimates based on later versions of the NSDUH instrument.

Table 7.69D Cocaine Use in Past Year among Persons Aged 12 or Older, by Age Group: Standard Errors of Percentages, 1971-2019

Survey Year	Aged 12-17	Aged 18-25	Aged 26-34
1971			
1972	0.48		
1974	0.61	1.09	
1976	0.56	1.01	
1977	0.51	0.91	
1979	0.60	2.11	1.23
1982	0.80	2.10	1.41
1985	0.59	1.72	1.48
1988	0.43	1.54	1.02
1990	0.56	0.98	0.83
1991	0.22	0.82	0.55
1992	0.19	0.75	0.55
1993	0.15	0.64	0.52
1994	0.20	0.37	0.32
1995	0.26	0.44	0.27
1996	0.19	0.43	0.34
1997	0.29	0.42	0.38
1998	0.24	0.41	0.35
1999	0.10	0.21	0.22
2000	0.12	0.18	0.18
2001	0.10	0.23	0.28
2002	0.13	0.24	0.31
2003	0.11	0.23	0.36
2004	0.11	0.25	0.30
2005	0.11	0.23	0.27
2006	0.11	0.24	0.35
2007	0.11	0.24	0.32
2008	0.10	0.21	0.33
2009	0.09	0.20	0.30
2010	0.09	0.20	0.33
2011	0.08	0.21	0.25
2012	0.08	0.21	0.31
2013	0.06	0.21	0.32
2014	0.09	0.23	0.23
2015	0.08	0.26	0.24
2016	0.07	0.26	0.24
2017	0.06	0.26	0.28
2018	0.07	0.26	0.28
2019	0.07	0.24	0.27

^{* =} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates for 1979 through 1993 have been adjusted for comparability with estimates for 1994 through 1998.

NOTE: Estimates for 1999 through 2019 have not been adjusted for comparability. See Appendix C of the *Results from the 2004 National Survey on Drug Use and Health: National Findings* for information on the comparability of the 1999 and later estimates with pre-1999 estimates.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 1971-2019.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the 2019 National Survey on Drug Use and Health: Methodological Summary and Definitions).

NOTE: Standard errors for 1971 through 1977 were calculated by multiplying the standard error of the percentage (assuming simple random sampling) by the square root of the estimated "design effect" (i.e., 1.25). See Appendix A of the *National Household Survey on Drug Abuse: Main Findings 1979* for further information on these procedures.

NOTE: Estimates for 1971 through 1977 have been adjusted to improve their comparability with estimates based on later versions of the NSDUH instrument.