

Table 8.1C Any Mental Illness in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Numbers in Thousands, 2018 and 2019

Age Category	Total (2018)	Total (2019)	Male (2018)	Male (2019)	Female (2018)	Female (2019)
TOTAL	644	631	395	404	500	467
18-25	151	163	99	102	115	117
18-20	106	144	63	76	84	105
21-25	155	165	94	97	116	122
26 or Older	615	599	380	385	477	441
26-49	351	366	228	232	253	266
26-29	187	194	122	120	132	145
30-34	198	213	125	135	139	155
35-39	183	190	122	114	137	148
40-44	154	175	100	102	116	135
45-49	152	160	89	100	123	125
50 or Older	480	460	302	297	371	339
50-54	244	230	169	158	171	171
55-59	244	254	157	164	192	183
60-64	222	207	127	119	176	172
65 or Older	310	295	174	172	254	232

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Any Mental Illness (AMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 8.1D Any Mental Illness in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Percentages, 2018 and 2019

Age Category	Total (2018)	Total (2019)	Male (2018)	Male (2019)	Female (2018)	Female (2019)
TOTAL	0.26	0.25	0.33	0.33	0.39	0.36
18-25	0.44	0.48	0.58	0.60	0.68	0.70
18-20	0.66	0.78	0.81	0.91	1.06	1.15
21-25	0.57	0.60	0.78	0.79	0.84	0.90
26 or Older	0.29	0.28	0.37	0.37	0.43	0.39
26-49	0.35	0.36	0.46	0.46	0.50	0.52
26-29	0.83	0.87	1.15	1.16	1.25	1.27
30-34	0.74	0.78	0.99	1.04	1.03	1.13
35-39	0.72	0.78	1.02	1.02	1.06	1.12
40-44	0.70	0.80	0.95	0.99	1.04	1.17
45-49	0.69	0.74	0.85	0.92	1.08	1.11
50 or Older	0.42	0.40	0.57	0.55	0.61	0.55
50-54	1.07	1.04	1.58	1.45	1.45	1.47
55-59	1.03	1.08	1.44	1.49	1.50	1.47
60-64	0.98	0.94	1.18	1.18	1.54	1.40
65 or Older	0.60	0.56	0.76	0.73	0.90	0.80

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Any Mental Illness (AMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 8.1P Any Mental Illness in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: P Values from Tests of Differences of Percentages, 2019 Versus 2018

Age Category	Total (2018)	Total (2019)	Male (2018)	Male (2019)	Female (2018)	Female (2019)
TOTAL	0.0001		0.0173		0.0007	
18-25	0.0000		0.0083		0.0000	
18-20	0.0007		0.0424		0.0085	
21-25	0.0003		0.0838		0.0008	
26 or Older	0.0024		0.0696		0.0126	
26-49	0.0000		0.0137		0.0000	
26-29	0.0080		0.2738		0.0078	
30-34	0.0087		0.1028		0.0290	
35-39	0.0724		0.6699		0.0071	
40-44	0.0040		0.1751		0.0084	
45-49	0.2505		0.0262		0.8936	
50 or Older	0.8566		0.6412		0.8693	
50-54	0.8395		0.8931		0.6495	
55-59	0.5184		0.7746		0.4683	
60-64	0.7271		0.7481		0.6500	
65 or Older	1.0000		0.3761		0.5287	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Any Mental Illness (AMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. The mental illness estimates that are used in the comparisons are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 8.4C Serious Mental Illness in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Numbers in Thousands, 2018 and 2019

Age Category	Total (2018)	Total (2019)	Male (2018)	Male (2019)	Female (2018)	Female (2019)
TOTAL	300	328	187	203	233	257
18-25	86	90	51	57	70	70
18-20	52	63	29	35	45	48
21-25	74	83	45	48	58	64
26 or Older	282	315	179	193	215	250
26-49	186	195	124	117	141	163
26-29	100	98	72	63	72	74
30-34	95	101	59	65	68	76
35-39	83	96	54	57	65	79
40-44	71	88	45	51	57	71
45-49	78	83	46	48	62	66
50 or Older	211	226	125	148	164	177
50-54	124	125	71	90	93	84
55-59	112	127	75	89	86	93
60-64	97	97	53	41	79	85
65 or Older	108	112	53	73	91	86

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Serious Mental Illness (SMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of SMI are a subset of estimates of any mental illness (AMI) because SMI is limited to persons with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 8.4D Serious Mental Illness in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Percentages, 2018 and 2019

Age Category	Total (2018)	Total (2019)	Male (2018)	Male (2019)	Female (2018)	Female (2019)
TOTAL	0.12	0.13	0.16	0.17	0.18	0.20
18-25	0.25	0.27	0.30	0.34	0.41	0.42
18-20	0.39	0.44	0.41	0.50	0.70	0.69
21-25	0.34	0.37	0.42	0.45	0.51	0.56
26 or Older	0.13	0.15	0.17	0.19	0.19	0.22
26-49	0.18	0.19	0.25	0.23	0.28	0.32
26-29	0.53	0.52	0.76	0.67	0.77	0.80
30-34	0.41	0.43	0.51	0.58	0.59	0.65
35-39	0.37	0.44	0.49	0.55	0.56	0.68
40-44	0.36	0.44	0.46	0.53	0.56	0.69
45-49	0.38	0.40	0.47	0.48	0.60	0.62
50 or Older	0.19	0.20	0.24	0.27	0.27	0.29
50-54	0.59	0.59	0.70	0.87	0.86	0.79
55-59	0.51	0.59	0.73	0.85	0.72	0.84
60-64	0.47	0.46	0.51	0.42	0.76	0.77
65 or Older	0.21	0.21	0.23	0.31	0.32	0.30

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Serious Mental Illness (SMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of SMI are a subset of estimates of any mental illness (AMI) because SMI is limited to persons with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 8.4P Serious Mental Illness in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: P Values from Tests of Differences of Percentages, 2019 Versus 2018

Age Category	Total (2018)	Total (2019)	Male (2018)	Male (2019)	Female (2018)	Female (2019)
TOTAL	0.0001		0.0258		0.0014	
18-25	0.0138		0.3579		0.0181	
18-20	0.0396		0.0591		0.2928	
21-25	0.1636		0.8502		0.0361	
26 or Older	0.0007		0.0350		0.0080	
26-49	0.0006		0.2013		0.0010	
26-29	0.3867		0.8612		0.3085	
30-34	0.1770		0.4228		0.2679	
35-39	0.0340		0.8343		0.0040	
40-44	0.0269		0.2585		0.0648	
45-49	0.3207		0.2794		0.5936	
50 or Older	0.1060		0.0746		0.5389	
50-54	0.5892		0.2127		0.6900	
55-59	0.3185		0.6958		0.2963	
60-64	0.3749		0.6178		0.5425	
65 or Older	0.4116		0.1864		0.9105	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Serious Mental Illness (SMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of SMI that are used in the comparisons are a subset of estimates of any mental illness (AMI) because SMI is limited to persons with AMI that resulted in serious functional impairment. The mental illness estimates that are used in the comparisons are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.