

Table 8.22C Received Substance Use Treatment at a Specialty Facility and/or Mental Health Services in Past Year among Persons Aged 18 or Older, by Past Year Co-Occurring Substance Use Disorder and Level of Mental Illness and Age Group: Standard Errors of Numbers in Thousands, 2018 and 2019

| Co-Occurring Substance Use Disorder (SUD) and Level of Mental Illness/Age Group | Received Substance Use Treatment (SU Tx) at a Specialty Facility OR Mental Health (MH) Services (2018) | Received Substance Use Treatment (SU Tx) at a Specialty Facility OR Mental Health (MH) Services (2019) | Received SU Tx at a Specialty Facility BUT NOT MH Services (2018) | Received SU Tx at a Specialty Facility BUT NOT MH Services (2019) | Received MH Services BUT NOT SU Tx at a Specialty Facility (2018) | Received MH Services BUT NOT SU Tx at a Specialty Facility (2019) | Received SU Tx at a Specialty Facility AND MH Services (2018) | Received SU Tx at a Specialty Facility AND MH Services (2019) |
|--|---|---|--|--|--|--|--|--|
| SUD and Any Mental Illness | 214 | 190 | 75 | 35 | 185 | 168 | 65 | 78 |
| 18-25 | 57 | 62 | 11 | 12 | 52 | 59 | 15 | 21 |
| 26 or Older | 204 | 180 | 74 | 32 | 175 | 158 | 63 | 73 |
| 26-49 | 135 | 127 | 27 | 25 | 120 | 114 | 47 | 56 |
| 50 or Older | 146 | 122 | * | 20 | 121 | 107 | 41 | 48 |
| SUD and Serious Mental Illness | 134 | 132 | 35 | 20 | 122 | 119 | 45 | 59 |
| 18-25 | 40 | 42 | 6 | 8 | 37 | 40 | 12 | 15 |
| 26 or Older | 127 | 121 | 35 | 18 | 116 | 110 | 43 | 57 |
| 26-49 | 86 | 94 | 13 | 18 | 76 | 83 | 36 | 47 |
| 50 or Older | * | * | * | * | * | * | * | * |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Specialty facilities for substance use treatment include hospitals (inpatient only), rehabilitation facilities (inpatient or outpatient), or mental health centers.

NOTE: Mental Health Services for adults includes inpatient treatment/counseling, outpatient treatment/counseling, or use of prescription medication for problems with emotions, nerves, or mental health. Respondents with unknown mental health service information were excluded.

NOTE: Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to persons with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 8.22D Received Substance Use Treatment at a Specialty Facility and/or Mental Health Services in Past Year among Persons Aged 18 or Older, by Past Year Co-Occurring Substance Use Disorder and Level of Mental Illness and Age Group: Standard Errors of Percentages, 2018 and 2019

| Co-Occurring Substance Use Disorder (SUD) and Level of Mental Illness/Age Group | Received Substance Use Treatment (SU Tx) at a Specialty Facility OR Mental Health (MH) Services (2018) | Received Substance Use Treatment (SU Tx) at a Specialty Facility OR Mental Health (MH) Services (2019) | Received SU Tx at a Specialty Facility BUT NOT MH Services (2018) | Received SU Tx at a Specialty Facility BUT NOT MH Services (2019) | Received MH Services BUT NOT SU Tx at a Specialty Facility (2018) | Received MH Services BUT NOT SU Tx at a Specialty Facility (2019) | Received SU Tx at a Specialty Facility AND MH Services (2018) | Received SU Tx at a Specialty Facility AND MH Services (2019) |
|--|---|---|--|--|--|--|--|--|
| SUD and Any Mental Illness | 1.49 | 1.47 | 0.79 | 0.36 | 1.45 | 1.42 | 0.68 | 0.79 |
| 18-25 | 1.71 | 1.77 | 0.45 | 0.48 | 1.68 | 1.79 | 0.62 | 0.80 |
| 26 or Older | 1.93 | 1.85 | 1.07 | 0.45 | 1.88 | 1.79 | 0.91 | 1.01 |
| 26-49 | 1.89 | 1.84 | 0.53 | 0.48 | 1.80 | 1.79 | 0.91 | 1.04 |
| 50 or Older | 4.68 | 4.73 | * | 1.14 | 5.09 | 4.67 | 2.34 | 2.58 |
| SUD and Serious Mental Illness | 2.12 | 2.11 | 1.09 | 0.54 | 2.41 | 2.31 | 1.36 | 1.57 |
| 18-25 | 2.75 | 2.90 | 0.68 | 0.82 | 2.82 | 3.02 | 1.30 | 1.56 |
| 26 or Older | 2.70 | 2.58 | 1.48 | 0.68 | 3.15 | 2.93 | 1.85 | 2.07 |
| 26-49 | 3.06 | 2.65 | 0.75 | 0.89 | 3.09 | 2.99 | 1.99 | 2.20 |
| 50 or Older | * | * | * | * | * | * | * | * |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Specialty facilities for substance use treatment include hospitals (inpatient only), rehabilitation facilities (inpatient or outpatient), or mental health centers.

NOTE: Mental Health Services for adults includes inpatient treatment/counseling, outpatient treatment/counseling, or use of prescription medication for problems with emotions, nerves, or mental health. Respondents with unknown mental health service information were excluded.

NOTE: Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to persons with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 8.22P Received Substance Use Treatment at a Specialty Facility and/or Mental Health Services in Past Year among Persons Aged 18 or Older, by Past Year Co-Occurring Substance Use Disorder and Level of Mental Illness and Age Group: P Values from Tests of Differences of Percentages, 2019 Versus 2018

| Co-Occurring Substance Use Disorder (SUD) and Level of Mental Illness/Age Group | Received Substance Use Treatment (SU Tx) at a Specialty Facility OR Mental Health (MH) Services (2018) | Received Substance Use Treatment (SU Tx) at a Specialty Facility OR Mental Health (MH) Services (2019) | Received SU Tx at a Specialty Facility BUT NOT MH Services (2018) | Received SU Tx at a Specialty Facility BUT NOT MH Services (2019) | Received MH Services BUT NOT SU Tx at a Specialty Facility (2018) | Received MH Services BUT NOT SU Tx at a Specialty Facility (2019) | Received SU Tx at a Specialty Facility AND MH Services (2018) | Received SU Tx at a Specialty Facility AND MH Services (2019) |
|--|---|---|--|--|--|--|--|--|
| SUD and Any Mental Illness | 0.2052 | | 0.1120 | | 0.2768 | | 0.4796 | |
| 18-25 | 0.8534 | | 0.3292 | | 0.6388 | | 0.1850 | |
| 26 or Older | 0.2010 | | 0.1552 | | 0.3250 | | 0.6785 | |
| 26-49 | 0.6001 | | 0.6096 | | 0.5597 | | 0.8100 | |
| 50 or Older | 0.1495 | | * | | 0.4170 | | 0.7102 | |
| SUD and Serious Mental Illness | 0.3399 | | 0.5298 | | 0.2554 | | 0.3992 | |
| 18-25 | 0.9292 | | 0.6497 | | 0.7359 | | 0.5494 | |
| 26 or Older | 0.2640 | | 0.4465 | | 0.2709 | | 0.4964 | |
| 26-49 | 0.8273 | | 0.6551 | | 0.5038 | | 0.5967 | |
| 50 or Older | * | | * | | * | | * | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Specialty facilities for substance use treatment include hospitals (inpatient only), rehabilitation facilities (inpatient or outpatient), or mental health centers.

NOTE: Mental Health Services for adults includes inpatient treatment/counseling, outpatient treatment/counseling, or use of prescription medication for problems with emotions, nerves, or mental health. Respondents with unknown mental health service information were excluded.

NOTE: Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) that are used in the comparisons are a subset of estimates of any mental illness (AMI) because SMI is limited to persons with AMI that resulted in serious functional impairment. The mental illness estimates that are used in the comparisons are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.