

**Table 8.36C Major Depressive Episode in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Numbers in Thousands, 2018 and 2019**

<b>Age Category</b>	<b>Total (2018)</b>	<b>Total (2019)</b>	<b>Male (2018)</b>	<b>Male (2019)</b>	<b>Female (2018)</b>	<b>Female (2019)</b>
<b>TOTAL</b>	379	385	227	240	303	310
<b>18-25</b>	115	124	70	73	94	96
18-20	76	102	43	53	64	76
21-25	102	108	61	62	81	85
<b>26 or Older</b>	356	360	213	230	282	289
26-49	225	221	146	138	166	176
26-29	120	115	81	77	85	84
30-34	114	123	74	73	80	94
35-39	94	110	63	58	72	94
40-44	96	100	59	61	75	77
45-49	87	96	50	60	74	73
50 or Older	274	275	164	183	223	214
50-54	156	146	99	100	111	105
55-59	140	137	89	85	109	108
60-64	122	128	72	69	99	104
65 or Older	153	159	80	111	132	120

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 8.36D Major Depressive Episode in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Percentages, 2018 and 2019**

<b>Age Category</b>	<b>Total (2018)</b>	<b>Total (2019)</b>	<b>Male (2018)</b>	<b>Male (2019)</b>	<b>Female (2018)</b>	<b>Female (2019)</b>
<b>TOTAL</b>	0.15	0.16	0.19	0.20	0.24	0.24
<b>18-25</b>	0.34	0.37	0.42	0.44	0.56	0.58
18-20	0.54	0.64	0.60	0.69	0.93	1.01
21-25	0.44	0.47	0.57	0.58	0.68	0.73
<b>26 or Older</b>	0.17	0.17	0.21	0.22	0.25	0.26
26-49	0.23	0.22	0.30	0.28	0.33	0.35
26-29	0.61	0.60	0.86	0.81	0.90	0.89
30-34	0.49	0.53	0.65	0.64	0.69	0.80
35-39	0.42	0.49	0.58	0.56	0.63	0.79
40-44	0.48	0.50	0.59	0.63	0.73	0.76
45-49	0.43	0.47	0.52	0.59	0.71	0.70
50 or Older	0.24	0.24	0.31	0.34	0.37	0.35
50-54	0.74	0.70	0.98	0.97	1.01	0.98
55-59	0.64	0.64	0.88	0.84	0.91	0.97
60-64	0.58	0.60	0.69	0.70	0.96	0.92
65 or Older	0.30	0.30	0.35	0.47	0.47	0.42

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 8.36P Major Depressive Episode in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2019 Versus 2018**

Age Category	Total (2018)	Total (2019)	Male (2018)	Male (2019)	Female (2018)	Female (2019)
<b>TOTAL</b>	0.0029		0.0125		0.0808	
<b>18-25</b>	0.0039		0.0602		0.0258	
18-20	0.0487		0.0608		0.3470	
21-25	0.0400		0.3668		0.0466	
<b>26 or Older</b>	0.0244		0.0387		0.2466	
26-49	0.0044		0.1417		0.0132	
26-29	0.7013		0.7347		0.8136	
30-34	0.0378		0.1603		0.1061	
35-39	0.0592		0.4396		0.0032	
40-44	0.2250		0.4102		0.3927	
45-49	0.6460		0.1533		0.6508	
50 or Older	0.5426		0.1451		0.6952	
50-54	0.9680		0.8422		0.8291	
55-59	0.8939		0.9407		0.8491	
60-64	0.6610		0.5366		0.9277	
65 or Older	0.4441		0.0666		0.6652	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 8.37C Major Depressive Episode with Severe Impairment in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category:  
Standard Errors of Numbers in Thousands, 2018 and 2019**

Age Category	Total (2018)	Total (2019)	Male (2018)	Male (2019)	Female (2018)	Female (2019)
<b>TOTAL</b>	311	331	184	200	244	259
<b>18-25</b>	93	101	57	60	74	79
18-20	62	73	34	39	51	57
21-25	77	86	48	50	60	69
<b>26 or Older</b>	293	316	175	189	226	246
26-49	189	184	122	116	138	148
26-29	100	95	70	61	69	70
30-34	96	91	64	54	64	75
35-39	76	91	47	50	59	76
40-44	81	85	50	53	62	64
45-49	67	81	41	49	54	63
50 or Older	222	236	125	146	180	184
50-54	133	114	72	79	103	83
55-59	110	119	73	74	84	90
60-64	97	108	50	52	81	92
65 or Older	113	134	56	87	99	103

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 8.37D Major Depressive Episode with Severe Impairment in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category:  
Standard Errors of Percentages, 2018 and 2019**

Age Category	Total (2018)	Total (2019)	Male (2018)	Male (2019)	Female (2018)	Female (2019)
<b>TOTAL</b>	0.13	0.13	0.16	0.17	0.19	0.20
<b>18-25</b>	0.28	0.30	0.34	0.36	0.44	0.48
18-20	0.46	0.50	0.49	0.56	0.78	0.80
21-25	0.34	0.39	0.46	0.48	0.52	0.62
<b>26 or Older</b>	0.14	0.15	0.17	0.18	0.20	0.22
26-49	0.19	0.18	0.25	0.23	0.27	0.29
26-29	0.53	0.51	0.76	0.65	0.73	0.77
30-34	0.41	0.40	0.56	0.49	0.55	0.65
35-39	0.34	0.42	0.44	0.48	0.52	0.66
40-44	0.41	0.43	0.51	0.56	0.61	0.64
45-49	0.34	0.40	0.42	0.49	0.54	0.61
50 or Older	0.20	0.21	0.24	0.27	0.30	0.30
50-54	0.63	0.56	0.71	0.78	0.95	0.79
55-59	0.51	0.56	0.72	0.73	0.71	0.82
60-64	0.47	0.51	0.49	0.54	0.79	0.82
65 or Older	0.22	0.26	0.25	0.37	0.36	0.36

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

**Table 8.37P Major Depressive Episode with Severe Impairment in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category:  
P Values from Tests of Differences of Percentages, 2019 Versus 2018**

Age Category	Total (2018)	Total (2019)	Male (2018)	Male (2019)	Female (2018)	Female (2019)
<b>TOTAL</b>	0.0004		0.0237		0.0071	
<b>18-25</b>	0.0004		0.0458		0.0032	
18-20	0.0885		0.1334		0.3731	
21-25	0.0021		0.1991		0.0033	
<b>26 or Older</b>	0.0076		0.0698		0.0516	
26-49	0.0025		0.2937		0.0023	
26-29	0.5435		0.8637		0.4860	
30-34	0.1066		0.7168		0.0666	
35-39	0.0392		0.7010		0.0044	
40-44	0.1529		0.3312		0.3003	
45-49	0.2922		0.1616		0.7609	
50 or Older	0.2905		0.1312		0.8460	
50-54	0.6304		0.6400		0.3223	
55-59	0.7131		1.0000		0.5859	
60-64	0.3269		0.4446		0.5845	
65 or Older	0.1673		0.0757		0.6575	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.