

Mental Health Annual Report 2013–2018

Use of Mental Health Services: National Client-Level Data

DEPARTMENT OF HEALTH AND HUMAN SERVICES
Substance Abuse and Mental Health Services Administration

ACKNOWLEDGMENTS

This report was prepared for the Substance Abuse and Mental Health Services Administration (SAMHSA), U.S. Department of Health and Human Services (HHS), by Eagle Technologies, Inc., under Contract No. HHSS283201600001C. The Contracting Officer's Representative (COR) at SAMHSA/CBHSQ was Nichele Waller.

SAMHSA complies with applicable federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. SAMHSA *cumple con las leyes federales de derechos civiles aplicables y no discrimina por motivos de raza, color, nacionalidad, edad, discapacidad o sexo.*

PUBLIC DOMAIN NOTICE

All material appearing in this report is in the public domain and may be reproduced or copied without permission from SAMHSA. Citation of the source is appreciated. However, this publication may not be reproduced or distributed for a fee without the specific, written authorization of the Office of Communications, SAMHSA, U.S. Department of Health and Human Services.

RECOMMENDED CITATION

Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. *Mental Health Annual Report: 2013–2018. Use of Mental Health Services: National Client-Level Data*. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2020.

ELECTRONIC ACCESS AND COPIES OF PUBLICATION

This publication may be downloaded at
<https://www.samhsa.gov/data/data-we-collect/mental-health-client-level-data>

ORIGINATING OFFICE

Center for Behavioral Health Statistics and Quality
Substance Abuse and Mental Health Services Administration
5600 Fishers Lane, Room 15SEH03
Rockville, Maryland 20857

June 2020

TABLE OF CONTENTS

List of Tables	vii
List of Figures	xxxix
Highlights	1
Background	5
Introduction	9
Chapter 1. Summary of Annual Data in State Client-Level Data Reporting: 2013–2018	13
Individuals Served	13
Regional Distributions	16
Any-Mention Mental Health Diagnoses.....	17
Chapter 2. Characteristics of Children and Adolescents Aged 17 Years and Younger Receiving Mental Health Services	19
Number Served	19
Gender.....	19
Race	19
Ethnicity.....	20
Living Arrangements	20
SED Status	20
Client Treatment Status at the Start of the Reporting Period.....	20
SED Status, by Service Setting.....	20
Chapter 3. Characteristics of Young Adults Aged 18 to 24 Years Receiving Mental Health Services	21
Number Served	21
Gender.....	21
Race and Ethnicity	21
Education	22
Marital Status.....	22
Living Arrangements	22
SMI Status.....	22
Veteran Status	22
Employment Status	22
Client Treatment Status at the Start of the Reporting Period.....	22

TABLE OF CONTENTS (CONTINUED)

Service Setting	23
SMI Status, by Service Setting	23
Chapter 4. Characteristics of Adults Aged 25 to 54 Years Receiving Mental Health Services	25
Number Served	25
Gender	25
Race	25
Ethnicity	26
Education	26
Marital Status	26
Living Arrangements	26
SMI Status	26
Veteran Status	27
Employment Status	27
Client Treatment Status at the Start of the Reporting Period	27
Service Setting	27
SMI Status, by Service Setting	27
Chapter 5. Characteristics of Older Adults Aged 55 Years and Older Receiving Mental Health Services	29
Number Served	29
Gender	29
Race	29
Ethnicity	30
Education	30
Marital Status	30
Living Arrangements	30
SMI Status	30
Veteran Status	30
Employment Status	31
Client Treatment Status at the Start of the Reporting Period	31
Service Setting	31
SMI Status, by Service Setting	31

TABLE OF CONTENTS (CONTINUED)

Chapter 6. Characteristics of Individuals With Co-occurring Mental Health and Substance Use Disorders Receiving Mental Health Services	33
Number Served	33
Gender.....	33
Age.....	33
Race	34
Ethnicity.....	34
Substance Use Diagnosis	34
Living Arrangements	34
SMI/SED Status.....	34
Veteran Status	34
Employment Status	35
Service Setting.....	35
Client Treatment Status at the Start of the Reporting Period.....	35
Chapter 7. Client Outcomes Among Individuals Receiving Mental Health Services: 2013–2018	37
Living Arrangements	37
Employment Status	37
Transitions in Living Arrangements	38
Transitions in Employment Status	41
Chapter 8. Geographic and Clinical Trends: 2013–2018	45
Chapter 9. Client Characteristics by State or Jurisdiction: 2013–2018.....	47
Chapter 10. Client Characteristics by State or Jurisdiction: 2018.....	49
Tables.....	51
Appendix A. MH-TEDS/MH-CLD Crosswalk Tables.....	773
Appendix B. MH-TEDS and MH-CLD 2011–2018 Reporting Methods and 2018 Reporting Period, by State or Jurisdiction.....	783
Appendix C. MH-TEDS and MH-CLD Data Dictionary of Included Variables	785

TABLE OF CONTENTS (CONTINUED)

Appendix D. SMI/SED Federal Definitions	797
Serious Mental Illness (SMI)	797
Serious Emotional Disturbance (SED).....	797
Appendix E. Mental Health and Substance Use Diagnosis Codes.....	799
Appendix F. Terms and Acronyms	811
Appendix G. List of Contributors	813

LIST OF TABLES

Summary of Annual Data in State Client-Level Data Reporting: 2013–2018

1.1a	Individuals, by mental health diagnosis: Number, 2013–2018	52
1.1b	Individuals, by mental health diagnosis: Column percent distribution, 2013–2018	53
1.2a	Individuals, by gender and age: Number, 2013–2018.....	54
1.2b	Individuals, by gender and age: Column percent distribution, 2013–2018.....	55
1.3a	Individuals, by race and ethnicity: Number, 2013–2018.....	56
1.3b	Individuals, by race and ethnicity: Column percent distribution, 2013–2018.....	57
1.4a	Individuals aged 17 years and younger, by mental health diagnosis: Number, 2013–2018	58
1.4b	Individuals aged 17 years and younger, by mental health diagnosis: Column percent distribution, 2013–2018	59
1.5a	Individuals aged 18 to 24 years, by mental health diagnosis: Number, 2013–2018	60
1.5b	Individuals aged 18 to 24 years, by mental health diagnosis: Column percent distribution, 2013–2018	61
1.6a	Individuals aged 25 to 54 years, by mental health diagnosis: Number, 2013–2018	62
1.6b	Individuals aged 25 to 54 years, by mental health diagnosis: Column percent distribution, 2013–2018	63
1.7a	Individuals aged 55 years and older, by mental health diagnosis: Number, 2013–2018	64
1.7b	Individuals aged 55 years and older, by mental health diagnosis: Column percent distribution, 2013–2018	65
1.8a	Individuals, by marital status and living arrangement: Number, 2013–2018.....	66
1.8b	Individuals, by marital status and living arrangement: Column percent distribution, 2013–2018.....	67
1.9a	Individuals, by SMI/SED status and co-occurring mental health and substance use disorder status: Number, 2013–2018	68

LIST OF TABLES (CONTINUED)

1.9b	Individuals, by SMI/SED status and co-occurring mental health and substance use disorder status: Column percent distribution, 2013–2018.....	69
1.10a	Individuals aged 18 years and older, by employment status, ‘detailed not in labor force,’ education, and veteran status: Number, 2013–2018	70
1.10b	Individuals aged 18 years and older, by employment status, ‘detailed not in labor force,’ education, and veteran status: Column percent distribution, 2013–2018	71
1.11a	Individuals, by timing of admission and mental health service setting: Number, 2013–2018	72
1.11b	Individuals, by timing of admission and mental health service setting: Column percent distribution, 2013–2018	73
1.12a	Individuals, by number of mental health diagnoses and any mention of mental health diagnosis: Number, 2013–2018	74
1.12b	Individuals, by number of mental health diagnoses and any mention of mental health diagnosis: Column percent, 2013–2018.....	75
1.13a	Individuals, by Census region and division: Number, 2013–2018.....	76
1.13b	Individuals, by Census region and division: Column percent distribution, 2013–2018	77
1.14a	Individuals with primary diagnosis of trauma- and stressor-related disorders, by gender, age, race, and ethnicity: Number, 2013–2018.....	78
1.14b	Individuals with primary diagnosis of trauma- and stressor-related disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018.....	80
1.15a	Individuals with primary diagnosis of anxiety disorders, by gender, age, race, and ethnicity: Number, 2013–2018.....	82
1.15b	Individuals with primary diagnosis of anxiety disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018.....	84
1.16a	Individuals with primary diagnosis of attention deficit/hyperactivity disorder, by gender, age, race, and ethnicity: Number, 2013–2018	86

LIST OF TABLES (CONTINUED)

1.16b	Individuals with primary diagnosis of attention deficit/hyperactivity disorder, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018	88
1.17a	Individuals with primary diagnosis of bipolar disorders, by gender, age, race, and ethnicity: Number, 2013–2018.....	90
1.17b	Individuals with primary diagnosis of bipolar disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018.....	92
1.18a	Individuals with primary diagnosis of depressive disorders, by gender, age, race, and ethnicity: Number, 2013–2018	94
1.18b	Individuals with primary diagnosis of depressive disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018	96
1.19a	Individuals with primary diagnosis of schizophrenia and other psychotic disorders, by gender, age, race, and ethnicity: Number, 2013–2018.....	98
1.19b	Individuals with primary diagnosis of schizophrenia and other psychotic disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018.....	100
1.20a	Individuals with co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Number, 2013–2018	102
1.20b	Individuals with co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018	104
1.21a	Individuals without co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Number, 2013–2018.....	106
1.21b	Individuals without co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018.....	108
1.22a	Individuals who received mental health services and U.S. population, by gender, marital status, and age group: Number, 2018	110
1.22b	Individuals who received mental health services and U.S. population, by gender, marital status, and age group: Column percent distribution, 2018	111
1.22c	Individuals who received mental health services and U.S. population, by gender, marital status, and age group: Row percent distribution, 2018	112

LIST OF TABLES (CONTINUED)

1.23a	Individuals who received mental health services and U.S. population, by race, ethnicity and age group: Number, 2018.....	113
1.23b	Individuals who received mental health services and U.S. population, by race, ethnicity and age group: Column percent distribution, 2018.....	114
1.23c	Individuals who received mental health services and U.S. population, by race, ethnicity and age group: Row percent distribution, 2018.....	115

Characteristics of Children and Adolescents Aged 17 Years and Younger Receiving Mental Health Services

2.1a	Individuals aged 17 years and younger, by gender, age, race, ethnicity, and SED status: Number, 2013–2018	116
2.1b	Individuals aged 17 years and younger, by gender, age, race, ethnicity, and SED status: Column percent distribution, 2013–2018	118
2.2a	Individuals aged 17 years and younger with SED, by gender, age, race, and ethnicity: Number, 2013–2018	120
2.2b	Individuals aged 17 years and younger with SED, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018	122
2.3a	Individuals aged 17 years and younger, by mental health diagnosis, gender, race, and ethnicity: Number, 2018	124
2.3b	Individuals aged 17 years and younger, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018.....	125
2.3c	Individuals aged 17 years and younger, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018	126
2.4a	Individuals aged 17 years and younger with SED, by mental health diagnosis, gender, race, and ethnicity: Number, 2018.....	127
2.4b	Individuals aged 17 years and younger with SED, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018.....	128
2.4c	Individuals aged 17 years and younger with SED, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018	129
2.5a	Individuals aged 17 years and younger, by mental health diagnosis, living arrangement, and SED status: Number, 2018.....	130

LIST OF TABLES (CONTINUED)

2.5b	Individuals aged 17 years and younger, by mental health diagnosis, living arrangement, and SED status: Column percent distribution, 2018.....	131
2.5c	Individuals aged 17 years and younger, by mental health diagnosis, living arrangement, and SED status: Row percent distribution, 2018.....	132
2.6a	Individuals aged 17 years and younger with SED, by mental health diagnosis and living arrangement: Number, 2018.....	133
2.6b	Individuals aged 17 years and younger with SED, by mental health diagnosis and living arrangement: Column percent distribution, 2018.....	134
2.6c	Individuals aged 17 years and younger with SED, by mental health diagnosis and living arrangement: Row percent distribution, 2018.....	135
2.7a	Individuals aged 17 years and younger, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018.....	136
2.7b	Individuals aged 17 years and younger, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018.....	137
2.7c	Individuals aged 17 years and younger, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018.....	138
2.8a	Individuals aged 17 years and younger with SED, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018.....	139
2.8b	Individuals aged 17 years and younger with SED, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018.....	140
2.8c	Individuals aged 17 years and younger with SED, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018.....	141
2.9	Individuals aged 17 years and younger, by mental health service setting and SED status: Number and percent distribution, 2018.....	142
Characteristics of Young Adults Aged 18 to 24 Years Receiving Mental Health Services		
3.1a	Individuals aged 18 to 24 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2013–2018.....	143

LIST OF TABLES (CONTINUED)

3.1b	Individuals aged 18 to 24 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2013–2018	145
3.2a	Individuals aged 18 to 24 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Number, 2013–2018	147
3.2b	Individuals aged 18 to 24 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Column percent distribution, 2013–2018.....	149
3.3a	Individuals aged 18 to 24 years, by mental health diagnosis, gender, race, and ethnicity: Number, 2018	151
3.3b	Individuals aged 18 to 24 years, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018	152
3.3c	Individuals aged 18 to 24 years, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018.....	153
3.4a	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Number, 2018	154
3.4b	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018.....	155
3.4c	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018	156
3.5a	Individuals aged 18 to 24 years, by mental health diagnosis, education, and marital status: Number, 2018.....	157
3.5b	Individuals aged 18 to 24 years, by mental health diagnosis, education, and marital status: Column percent distribution, 2018.....	158
3.5c	Individuals aged 18 to 24 years, by mental health diagnosis, education, and marital status: Row percent distribution, 2018	159
3.6a	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, education, and marital status: Number, 2018	160
3.6b	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, education, and marital status: Column percent distribution, 2018	161
3.6c	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, education, marital status: Row percent distribution, 2018	162

LIST OF TABLES (CONTINUED)

3.7a	Individuals aged 18 to 24 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Number, 2018	163
3.7b	Individuals aged 18 to 24 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Column percent distribution, 2018	164
3.7c	Individuals aged 18 to 24 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Row percent distribution, 2018	165
3.8a	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, living arrangement and veteran status: Number, 2018	166
3.8b	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, living arrangement and veteran status: Column percent distribution, 2018	167
3.8c	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, living arrangement and veteran status: Row percent distribution, 2018	168
3.9a	Individuals aged 18 to 24 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2018	169
3.9b	Individuals aged 18 to 24 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2018	170
3.9c	Individuals aged 18 to 24 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2018	171
3.10a	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2018	172
3.10b	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2018.....	173
3.10c	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2018.....	174
3.11a	Individuals aged 18 to 24 years, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018	175
3.11b	Individuals aged 18 to 24 years, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018.....	176

LIST OF TABLES (CONTINUED)

3.11c	Individuals aged 18 to 24 years, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018.....	177
3.12a	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018.....	178
3.12b	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018.....	179
3.12c	Individuals aged 18 to 24 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018.....	180
3.13	Individuals aged 18 to 24 years, by mental health service setting and SMI status: Number and percent distribution, 2018.....	181

Characteristics of Adults Aged 25 to 54 Years Receiving Mental Health Services

4.1a	Individuals aged 25 to 54 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2013–2018	182
4.1b	Individuals aged 25 to 54 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2013–2018	184
4.2a	Individuals aged 25 to 54 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Number, 2013–2018	186
4.2b	Individuals aged 25 to 54 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Column percent distribution, 2013–2018.....	188
4.3a	Individuals aged 25 to 54 years, by mental health diagnosis, gender, race, and ethnicity: Number, 2018	190
4.3b	Individuals aged 25 to 54 years, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018	191
4.3c	Individuals aged 25 to 54 years, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018.....	192
4.4a	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Number, 2018	193
4.4b	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018.....	194

LIST OF TABLES (CONTINUED)

4.4c	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018	195
4.5a	Individuals aged 25 to 54 years, by mental health diagnosis, education, and marital status: Number, 2018	196
4.5b	Individuals aged 25 to 54 years, by mental health diagnosis, education, and marital status: Column percent distribution, 2018	197
4.5c	Individuals aged 25 to 54 years, by mental health diagnosis, education, and marital status: Row percent distribution, 2018	198
4.6a	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, education, and marital status: Number, 2018	199
4.6b	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, education, and marital status: Column percent distribution, 2018	200
4.6c	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, education, and marital status: Row percent distribution, 2018	201
4.7a	Individuals aged 25 to 54 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Number, 2018	202
4.7b	Individuals aged 25 to 54 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Column percent distribution, 2018	203
4.7c	Individuals aged 25 to 54 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Row percent distribution, 2018	204
4.8a	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, living arrangement, and veteran status: Number, 2018	205
4.8b	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, living arrangement, and veteran status: Column percent distribution, 2018	206
4.8c	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, living arrangement, and veteran status: Row percent distribution, 2018	207
4.9a	Individuals aged 25 to 54 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2018	208
4.9b	Individuals aged 25 to 54 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2018	209

LIST OF TABLES (CONTINUED)

4.9c	Individuals aged 25 to 54 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2018	210
4.10a	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2018	211
4.10b	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2018.....	212
4.10c	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2018.....	213
4.11a	Individuals aged 25 to 54 years, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018	214
4.11b	Individuals aged 25 to 54 years, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018.....	215
4.11c	Individuals aged 25 to 54 years, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018.....	216
4.12a	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018.....	217
4.12b	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018.....	218
4.12c	Individuals aged 25 to 54 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018.....	219
4.13	Individuals aged 25 to 54 years, by mental health service setting and SMI status: Number and percent distribution, 2018.....	220
Characteristics of Older Adults Aged 55 Years and Older Receiving Mental Health		
5.1a	Individuals aged 55 years and older, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2013–2018.....	221
5.1b	Individuals aged 55 years and older, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2013–2018	223

LIST OF TABLES (CONTINUED)

5.2a	Individuals aged 55 years and older with SMI, by gender, age, race, ethnicity, education, and veteran status: Number, 2013–2018	225
5.2b	Individuals aged 55 years and older with SMI, by gender, age, race, ethnicity, education, and veteran status: Column percent distribution, 2013–2018	227
5.3a	Individuals aged 55 years and older, by mental health diagnosis, gender, race, and ethnicity: Number, 2018.....	229
5.3b	Individuals aged 55 years and older, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018.....	230
5.3c	Individuals aged 55 years and older, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018.....	231
5.4a	Individuals aged 55 years and older with SMI, by mental health diagnosis, gender, race, and ethnicity: Number, 2018.....	232
5.4b	Individuals aged 55 years and older with SMI, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018.....	233
5.4c	Individuals aged 55 years and older with SMI, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018.....	234
5.5a	Individuals aged 55 years and older, by mental health diagnosis, education, and marital status: Number, 2018	235
5.5b	Individuals aged 55 years and older, by mental health diagnosis, education, and marital status: Column percent distribution, 2018	236
5.5c	Individuals aged 55 years and older, by mental health diagnosis, education, and marital status: Row percent distribution, 2018.....	237
5.6a	Individuals aged 55 years and older with SMI, by mental health diagnosis, education, and marital status: Number, 2018	238
5.6b	Individuals aged 55 years and older with SMI, by mental health diagnosis, education, and marital status: Column percent distribution, 2018	239
5.6c	Individuals aged 55 years and older with SMI, by mental health diagnosis, education, and marital status: Row percent distribution, 2018.....	240
5.7a	Individuals aged 55 years and older, by mental health diagnosis, living arrangement, SMI status, and veteran status: Number, 2018	241

LIST OF TABLES (CONTINUED)

5.7b	Individuals aged 55 years and older, by mental health diagnosis, living arrangement, SMI status, and veteran status: Column percent distribution, 2018	242
5.7c	Individuals aged 55 years and older, by mental health diagnosis, living arrangement, SMI status, and veteran status: Row percent distribution, 2018	243
5.8a	Individuals aged 55 years and older with SMI, by mental health diagnosis, living arrangement, and veteran status: Number, 2018	244
5.8b	Individuals aged 55 years and older with SMI, by mental health diagnosis, living arrangement, and veteran status: Column percent distribution, 2018	245
5.8c	Individuals aged 55 years and older with SMI, by mental health diagnosis, living arrangement, and veteran status: Row percent distribution, 2018	246
5.9a	Individuals aged 55 years and older, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2018	247
5.9b	Individuals aged 55 years and older, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2018.....	248
5.9c	Individuals aged 55 years and older, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2018.....	249
5.10a	Individuals aged 55 years and older with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2018	250
5.10b	Individuals aged 55 years and older with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2018.....	251
5.10c	Individuals aged 55 years and older with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2018.....	252
5.11a	Individuals aged 55 years and older, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018	253
5.11b	Individuals aged 55 years and older, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018.....	254

LIST OF TABLES (CONTINUED)

5.11c	Individuals aged 55 years and older, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018.....	255
5.12a	Individuals aged 55 years and older with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018.....	256
5.12b	Individuals aged 55 years and older with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018.....	257
5.12c	Individuals aged 55 years and older with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018.....	258
5.13	Individuals aged 55 years and older, by mental health service setting and SMI status: Number and percent distribution, 2018.....	259

Characteristics of Individuals With Co-occurring Mental Health and Substance Use Disorders Receiving Mental Health Services

6.1a	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by gender, age, race, ethnicity, education, veteran status, and SMI/SED status: Number, 2013–2018	260
6.1b	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by gender, age, race, ethnicity, education, veteran status, and SMI/SED status: Column percent distribution, 2013–2018	262
6.2a	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, gender, and age: Number, 2018	264
6.2b	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, gender, and age: Column percent distribution, 2018	265
6.2c	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, gender, and age: Row percent distribution, 2018	266
6.3a	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, race, and ethnicity: Number, 2018	267

LIST OF TABLES (CONTINUED)

6.3b	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, race, and ethnicity: Column percent distribution, 2018	268
6.3c	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, race, and ethnicity: Row percent distribution, 2018	269
6.4a	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis and substance use diagnosis: Number, 2018	270
6.4b	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis and substance use diagnosis: Column percent distribution, 2018.....	271
6.4c	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis and substance use diagnosis: Row percent distribution, 2018	272
6.5a	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, education, and marital status: Number, 2018.....	273
6.5b	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, education, and marital status: Column percent distribution, 2018.....	274
6.5c	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, education, and marital status: Row percent distribution, 2018.....	275
6.6a	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, living arrangement, SMI/SED status, and veteran status: Number, 2018.....	276
6.6b	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, living arrangement, SMI/SED status, and veteran status: Column percent distribution, 2018	277
6.6c	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, living arrangement, SMI/SED status, and veteran status: Row percent distribution, 2018.....	278

LIST OF TABLES (CONTINUED)

6.7a	Individuals aged 18 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2018	279
6.7b	Individuals aged 18 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2018	280
6.7c	Individuals aged 18 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2018.....	281
6.8a	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018.....	282
6.8b	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018.....	283
6.8c	Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018.....	284

Client Outcomes Among Individuals Receiving Mental Health Services: 2013–2018

7.1a	Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2015: Number, 2015	285
7.1b	Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2015: Column percent distribution, 2015	288
7.1c	Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2015: Row percent distribution, 2015	291
7.2a	Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2018: Number, 2018	294
7.2b	Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2018: Column percent distribution, 2018	297
7.2c	Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2018: Row percent distribution, 2018	300

LIST OF TABLES (CONTINUED)

7.3a	Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2015: Number, 2015	303
7.3b	Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2015: Column percent distribution, 2018	306
7.3c	Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2015: Row percent distribution, 2018	309
7.4a	Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2018: Number, 2018	312
7.4b	Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2018: Column percent distribution, 2018	315
7.4c	Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2018: Row percent distribution, 2018	318
7.5a	Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and living arrangement in 2015 and 2018: Number	321
7.5b	Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and living arrangement in 2015 and 2018: Row percent distribution.....	323
7.6a	Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and employment status in 2015 and 2018: Number	325
7.6b	Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and employment status in 2015 and 2018: Row percent distribution	326
7.7	Individuals aged 18 years and older served in consecutive reporting periods, by mental health diagnosis and living arrangement: Number, 2013–2018	327
7.8a	Individuals aged 18 years and older served in both 2013 and 2015, by mental health diagnosis in 2013 and living arrangement in 2013 and 2015: Number	329

LIST OF TABLES (CONTINUED)

7.8b	Individuals aged 18 years and older served in both 2013 and 2015, by mental health diagnosis in 2013 and living arrangement in 2013 and 2015: Row percent distribution	331
7.9a	Individuals aged 18 years and older served in both 2013 and 2016, by mental health diagnosis in 2013 and living arrangement in 2013 and 2016: Number	333
7.9b	Individuals aged 18 years and older served in both 2013 and 2016, by mental health diagnosis in 2013 and living arrangement in 2013 and 2016: Row percent distribution	335
7.10a	Individuals aged 18 years and older served in both 2013 and 2017, by mental health diagnosis in 2013 and living arrangement in 2013 and 2017: Number	337
7.10b	Individuals aged 18 years and older served in both 2013 and 2017, by mental health diagnosis in 2013 and living arrangement in 2013 and 2017: Row percent distribution	339
7.11a	Individuals aged 18 years and older served in both 2013 and 2018, by mental health diagnosis in 2013 and living arrangement in 2013 and 2018: Number	341
7.11b	Individuals aged 18 years and older served in both 2013 and 2018, by mental health diagnosis in 2013 and living arrangement in 2013 and 2018: Row percent distribution	343
7.12	Individuals aged 18 years and older served in consecutive reporting periods, by mental health diagnosis and employment status: Number, 2013–2018	345
7.13a	Individuals aged 18 years and older served in both 2013 and 2015, by mental health diagnosis in 2013 and employment status in 2013 and 2015: Number	346
7.13b	Individuals aged 18 years and older served in both 2013 and 2015, by mental health diagnosis in 2013 and employment status in 2013 and 2015: Row percent distribution	347
7.14a	Individuals aged 18 years and older served in both 2013 and 2016, by mental health diagnosis in 2013 and employment status in 2013 and 2016: Number	348

LIST OF TABLES (CONTINUED)

7.14b	Individuals aged 18 years and older served in both 2013 and 2016, by mental health diagnosis in 2013 and employment status in 2013 and 2016: Row percent distribution	349
7.15a	Individuals aged 18 years and older served in both 2013 and 2017, by mental health diagnosis in 2013 and employment status in 2013 and 2017: Number	350
7.15b	Individuals aged 18 years and older served in both 2013 and 2017, by mental health diagnosis in 2013 and employment status in 2013 and 2017: Row percent distribution	351
7.16a	Individuals aged 18 years and older served in both 2013 and 2018, by mental health diagnosis in 2013 and employment status in 2013 and 2018: Number	352
7.16b	Individuals aged 18 years and older served in both 2013 and 2018, by mental health diagnosis in 2013 and employment status in 2013 and 2018: Row percent distribution	353

Geographic and Clinical Trends: 2013–2018

8.1	Individuals, by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2018	354
8.2	Individuals per 100,000 population, by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2018.....	357
8.3	Individuals per 100,000 population, by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2018, adjusted for age, gender, and race.....	360
8.4	Individuals with primary diagnosis of trauma- and stressor-related disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018	363
8.5	Individuals with primary diagnosis of anxiety disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018	366
8.6	Individuals with primary diagnosis of attention deficit/hyperactivity disorder, by Census region, Census division, and state or jurisdiction: Number, 2013–2018	369

LIST OF TABLES (CONTINUED)

8.7	Individuals with primary diagnosis of bipolar disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018	372
8.8	Individuals with primary diagnosis of depressive disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018.....	375
8.9	Individuals with primary diagnosis of schizophrenia and other psychotic disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018	378
8.10	Individuals with co-occurring mental health and substance use disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018	381

Client Characteristics by State or Jurisdiction: 2013–2018

9.1a	Alabama individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	384
9.1b	Alabama individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	386
9.2a	Arizona individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	388
9.2b	Arizona individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	390
9.3a	Arkansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	392
9.3b	Arkansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	394
9.4a	California individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	396
9.4b	California individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	398
9.5a	Colorado individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	400
9.5b	Colorado individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	402

LIST OF TABLES (CONTINUED)

9.6a	Connecticut individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	404
9.6b	Connecticut individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	406
9.7a	Delaware individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	408
9.7b	Delaware individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	410
9.8a	District of Columbia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	412
9.8b	District of Columbia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	414
9.9a	Florida individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	416
9.9b	Florida individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	418
9.10a	Hawaii individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	420
9.10b	Hawaii individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	422
9.11a	Idaho individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	424
9.11b	Idaho individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	426
9.12a	Illinois individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	428
9.12b	Illinois individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	430
9.13a	Indiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	432

LIST OF TABLES (CONTINUED)

9.13b	Indiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	434
9.14a	Iowa individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018	436
9.14b	Iowa individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018	438
9.15a	Kansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	440
9.15b	Kansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	442
9.16a	Kentucky individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	444
9.16b	Kentucky individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	446
9.17a	Louisiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	448
9.17b	Louisiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	450
9.18a	Maine individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018	452
9.18b	Maine individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018	454
9.19a	Maryland individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	456
9.19b	Maryland individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	458
9.20a	Massachusetts individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	460
9.20b	Massachusetts individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	462

LIST OF TABLES (CONTINUED)

9.21a	Michigan individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	464
9.21b	Michigan individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	466
9.22a	Minnesota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	468
9.22b	Minnesota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	470
9.23a	Mississippi individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	472
9.23b	Mississippi individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	474
9.24a	Missouri individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	476
9.24b	Missouri individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	478
9.25a	Montana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	480
9.25b	Montana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	482
9.26a	Nebraska individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	484
9.26b	Nebraska individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	486
9.27a	Nevada individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	488
9.27b	Nevada individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	490
9.28a	New Jersey individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	492

LIST OF TABLES (CONTINUED)

9.28b	New Jersey individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	494
9.29a	New Mexico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	496
9.29b	New Mexico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	498
9.30a	New York individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	500
9.30b	New York individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	502
9.31a	North Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	504
9.31b	North Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	506
9.32a	North Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	508
9.32b	North Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	510
9.33a	Ohio individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018	512
9.33b	Ohio individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018	514
9.34a	Oklahoma individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	516
9.34b	Oklahoma individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	518
9.35a	Oregon individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	520
9.35b	Oregon individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	522

LIST OF TABLES (CONTINUED)

9.36a	Pennsylvania individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	524
9.36b	Pennsylvania individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	526
9.37a	Puerto Rico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	528
9.37b	Puerto Rico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	530
9.38a	Rhode Island individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	532
9.38b	Rhode Island individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	534
9.39a	South Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	536
9.39b	South Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	538
9.40a	South Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	540
9.40b	South Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	542
9.41a	Tennessee individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	544
9.41b	Tennessee individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	546
9.42a	Texas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018	548
9.42b	Texas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018	550
9.43a	Utah individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018	552

LIST OF TABLES (CONTINUED)

9.43b	Utah individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018	554
9.44a	Vermont individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	556
9.44b	Vermont individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	558
9.45a	Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	560
9.45b	Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	562
9.46a	Washington individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	564
9.46b	Washington individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	566
9.47a	West Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	568
9.47b	West Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	570
9.48a	Wisconsin individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	572
9.48b	Wisconsin individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	574
9.49a	Wyoming individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018.....	576
9.49b	Wyoming individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	578
9.50a	Other jurisdictions individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018	580
9.50b	Other jurisdictions individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018.....	582

LIST OF TABLES (CONTINUED)

Client Characteristics by State or Other Jurisdiction: 2018

10.1a	Alabama individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	584
10.1b	Alabama individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	586
10.2a	Arizona individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	588
10.2b	Arizona individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	590
10.3a	Arkansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	592
10.3b	Arkansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	594
10.4a	California individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	596
10.4b	California individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	598
10.5a	Colorado individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	600
10.5b	Colorado individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	602
10.6a	Connecticut individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	604
10.6b	Connecticut individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	606
10.7a	Delaware individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	608
10.7b	Delaware individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	610
10.8a	District of Columbia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	612

LIST OF TABLES (CONTINUED)

10.8b	District of Columbia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	614
10.9a	Florida individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	616
10.9b	Florida individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	618
10.10a	Hawaii individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	620
10.10b	Hawaii individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	622
10.11a	Idaho individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018	624
10.11b	Idaho individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018	626
10.12a	Illinois individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	628
10.12b	Illinois individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	630
10.13a	Indiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	632
10.13b	Indiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	634
10.14a	Iowa individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018	636
10.14b	Iowa individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018	638
10.15a	Kentucky individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	640
10.15b	Kentucky individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	642

LIST OF TABLES (CONTINUED)

10.16a	Louisiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	644
10.16b	Louisiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	646
10.17a	Maryland individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	648
10.17b	Maryland individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	650
10.18a	Massachusetts individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	652
10.18b	Massachusetts individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	654
10.19a	Michigan individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	656
10.19b	Michigan individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	658
10.20a	Minnesota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	660
10.20b	Minnesota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	662
10.21a	Mississippi individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	664
10.21b	Mississippi individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	666
10.22a	Missouri individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	668
10.22b	Missouri individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	670
10.23a	Montana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	672

LIST OF TABLES (CONTINUED)

10.23b	Montana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	674
10.24a	Nebraska individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	676
10.24b	Nebraska individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	678
10.25a	Nevada individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	680
10.25b	Nevada individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	682
10.26a	New Mexico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	684
10.26b	New Mexico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	686
10.27a	New York individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	688
10.27b	New York individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	690
10.28a	North Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	692
10.28b	North Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	694
10.29a	North Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	696
10.29b	North Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	698
10.30a	Ohio individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018	700
10.30b	Ohio individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018	702

LIST OF TABLES (CONTINUED)

10.31a	Oklahoma individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	704
10.31b	Oklahoma individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	706
10.32a	Oregon individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	708
10.32b	Oregon individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	710
10.33a	Pennsylvania individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	712
10.33b	Pennsylvania individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	714
10.34a	Puerto Rico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	716
10.34b	Puerto Rico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	718
10.35a	Rhode Island individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	720
10.35b	Rhode Island individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	722
10.36a	South Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	724
10.36b	South Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	726
10.37a	South Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	728
10.37b	South Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	730
10.38a	Tennessee individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	732

LIST OF TABLES (CONTINUED)

10.38b	Tennessee individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	734
10.39a	Texas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018	736
10.39b	Texas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018	738
10.40a	Utah individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018	740
10.40b	Utah individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018	742
10.41a	Vermont individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	744
10.41b	Vermont individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	746
10.42a	Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	748
10.42b	Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	750
10.43a	Washington individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	752
10.43b	Washington individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	754
10.44a	West Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	756
10.44b	West Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	758
10.45a	Wisconsin individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	760
10.45b	Wisconsin individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	762

LIST OF TABLES (CONTINUED)

10.46a	Wyoming individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	764
10.46b	Wyoming individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	766
10.47a	Other jurisdictions individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018.....	768
10.47b	Other jurisdictions individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018.....	770
Appendix A		
A1	MH-TEDS/MH-CLD Crosswalk.....	773
Appendix B		
B1	MH-TEDS and MH-CLD 2011–2018 reporting methods and 2018 reporting period, by state or jurisdiction	783
Appendix C		
C1	Number of states reporting data, MH-CLD and MH-TEDS 2013–2018	793
C2	Item percentage response rate, by state or jurisdiction: MH-CLD and MH-TEDS 2018	794

LIST OF FIGURES

Client Outcomes Among Individuals Receiving Mental Health Services: 2013–2018

1	Census regions and Census divisions of the United States.....	16
2	Living arrangements in 2018 of individuals aged 18 years and older who reported living in a private residence in 2015, by 2015 mental health diagnosis.....	38
3	Living arrangements in 2018 of individuals aged 18 years and older who reported being homeless in 2015, by 2015 mental health diagnosis.....	39
4	Living arrangements in 2018 of individuals aged 18 years and older who reported being in jail or a correctional facility in 2015, by 2015 mental health diagnosis.....	40
5	Employment status in 2018 of individuals aged 18 years and older who reported being employed in 2015, by 2015 mental health diagnosis.....	41
6	Employment status in 2018 of individuals aged 18 years and older who reported being unemployed in 2015, by 2015 mental health diagnosis.....	42
7	Employment status in 2018 of individuals aged 18 years and older who were not in the labor force in 2015, by 2015 mental health diagnosis	43

HIGHLIGHTS

This report presents results from the Mental Health Client-Level Data (MH-CLD) and Mental Health Treatment Episode Data Set (MH-TEDS) for individuals receiving mental health services from state mental health systems in 2018, as well as selected trends in data collected from such individuals between 2013 and 2018. The report provides information on mental health diagnoses, mental health treatment settings, and demographic and substance use characteristics of individuals in mental health treatment in facilities that reported to individual state administrative data systems.

In the 2018 reporting period, 48 states and other jurisdictions (hereafter, “states”) submitted 6,213,791 records for individuals receiving mental health services. Alaska, American Samoa, the Federated States of Micronesia, Georgia, Guam, Kansas, Maine, Marshall Islands, New Jersey, and the U.S. Virgin Islands did not report data for the 2018 reporting period, as of July 8, 2019 [Table 1.1 and Appendix Table B1].

Of the 6,213,791 eligible client-level records for the 2018 reporting period, 88 percent had valid mental health or substance use diagnoses. Such records form the basis for most tables in this report. However, individual records may not have data for all variables; therefore, the analysis of specific variables may be based on a smaller number of individuals [Tables 1.1a–b].

State Client-Level Data Reporting: 2013–2018

For the 2013 reporting period, 45 states reported that a total of 5,576,659 individuals received mental health services; for the 2014 reporting period, 47 states reported that a total of 5,818,881 individuals received mental health services; for the 2015 reporting period, 45 states reported that a total of 5,311,897 individuals received mental health services; for the 2016 reporting period, 47 states reported that a total of 5,813,339 individuals received mental health services; for the 2017 reporting period, 48 states reported that a total of 5,934,383 individuals received mental health services; and for the 2018 reporting period, 48 states reported that a total of 6,213,791 individuals received mental health services [Table 1.1].

Children and Adolescents Aged 17 Years and Younger Receiving Mental Health Services: 2018

A total of 1,800,613 children and adolescents aged 17 years and younger were served in the 2018 reporting period [Table 2.1a]; 1,630,278 had a valid mental health diagnosis code [Table 2.3a].

Among males aged 17 years and younger served in the 2018 reporting period, the most frequently reported diagnoses were ADHD (31 percent), trauma- and stressor-related disorders (20 percent), anxiety disorders (9 percent), and depressive disorders (9 percent) [Table 2.3c].

Among females aged 17 years and younger served in the 2018 reporting period, the most frequently reported diagnoses were trauma- and stressor-related disorders (28 percent), depressive disorders (20 percent), anxiety disorders (14 percent), and ADHD (14 percent) [Table 2.3c].

Young Adults Aged 18 to 24 Years Receiving Mental Health Services: 2018

A total of 645,452 young adults aged 18 to 24 years were served in 2018 [Table 3.1a]; 557,478 had a valid mental health diagnosis code [Table 3.3a].

Depressive disorders were the most frequently reported diagnoses among individuals aged 18 to 24 years served in the 2018 reporting period, across all selected categories for gender, race, ethnicity, educational attainment, and marital status [Tables 3.3c and 3.5c].

Adults Aged 25 to 54 Years Receiving Mental Health Services: 2018

A total of 2,831,082 adults aged 25 to 54 years were served in 2018 [Table 4.1a]; 2,476,907 had a valid mental health diagnosis code [Table 4.3a].

Among males aged 25 to 54 years served in the 2018 reporting period, the most frequently reported diagnoses were depressive disorders (25 percent), schizophrenia and other psychotic disorders (24 percent), and bipolar disorders (14 percent); among females, the most frequently reported diagnoses were depressive disorders (33 percent), bipolar disorders (18 percent), and anxiety disorders (14 percent) [Table 4.3c].

Older Adults Aged 55 Years and Older Receiving Mental Health Services: 2018

A total of 932,573 adults aged 55 years and older were served in 2018 [Table 5.1a]; 827,060 had a valid mental health diagnosis code [Table 5.3a].

Among males aged 55 years and older served in the 2018 reporting period, the most frequently reported diagnoses were schizophrenia and other psychotic disorders (30 percent), depressive disorders (30 percent), and bipolar disorders (12 percent); among females served during this period, the most frequently reported diagnoses were depressive disorders (39 percent), schizophrenia and other psychotic disorders (19 percent), and bipolar disorders (15 percent) [Table 5.3c].

Individuals With Co-Occurring Mental Health and Substance Use Disorders Receiving Mental Health Services: 2018

A total of 1,026,726 individuals aged 12 years and older with co-occurring mental and substance use disorders were served in 2018 [Table 6.1a].

For males with co-occurring mental and substance use disorders aged 12 years or older served during the 2018 reporting period, the most frequently reported diagnoses were depressive disorders (25 percent), schizophrenia and other psychotic disorders (22 percent), and bipolar disorders (14 percent); among females, the most frequently reported diagnoses were depressive disorders (31 percent), bipolar disorders (19 percent) and anxiety disorders (12 percent) [Table 6.2c].

Client Outcomes Among Individuals Receiving Mental Health Services in 2015 and 2018

Among individuals served in both the 2015 and 2018 reporting periods, 92 percent of those who lived in a private residence in 2015 also reported living in a private residence in 2018; 3 percent transitioned to homelessness; 2 percent transitioned to residential care; and 3 percent, combined, transitioned to other living arrangements, institutional settings, or jail [Table 7.5b].

Among individuals aged 18 years and older who were served in both the 2015 and 2018 reporting periods, 66 percent of those who were employed in 2015 also reported being employed in 2018; 13 percent transitioned to being unemployed, and 21 percent had left the labor force [Table 7.6b].

Individuals Who Received Mental Health Services per 100,000 Population, by State or Jurisdiction in 2018, Adjusted for Age, Gender, and Race

Tables 8.2 and 8.3 present, respectively, the number and adjusted number of individuals per 100,000 population served in the 2018 reporting period, by mental health diagnosis, Census region, Census division, and state or jurisdiction.

Among individuals served in the 2018 reporting period, the primary mental health diagnoses with the largest per 100,000 population ratios (adjusted for age, gender, and ethnicity) were depressive disorders (443 per 100,000 population), trauma- and stressor-related disorders (246 per 100,000 population), and schizophrenia and other psychotic disorders (201 per 100,000 population) [Table 8.3].

Across all Census divisions in the 2018 reporting period, depressive disorders were the primary mental health diagnosis with the highest adjusted ratio per 100,000 population, ranging from 238 per 100,000 population for New England to 695 per 100,000 population for the West North Central division [Table 8.3].

BACKGROUND

Mental health services data are currently collected for and reported to the Substance Abuse and Mental Health Services Administration's (SAMHSA) Center for Behavioral Health Statistics and Quality (CBHSQ). State mental health agencies (SMHAs) are the state entities with primary responsibility for reporting the data in accordance with the reporting terms and conditions of the Behavioral Health Services Information System (BHSIS) Agreements funded by SAMHSA. The BHSIS Agreement stipulates that states and jurisdictions submit Uniform Reporting System (URS) data and client-level data by December 1 each year, in compliance with mental health block grant reporting requirements (42 U.S.C. §300x-6).

URS data reporting started in 2002 in an effort spearheaded by SAMHSA's Center for Mental Health Services (CMHS) to collect both system performance measures and client outcomes at an aggregate level. The collection of client-level data came later.

In 2008, SAMHSA, through CMHS, tested the feasibility of transitioning from aggregate reporting of SAMHSA's mental health national outcome measures (NOMs)¹ to individual-level data reporting. This led to the development of the Mental Health Client-Level Data (MH-CLD) reporting method. The development was supported by data infrastructure grants (DIGs), which were a vehicle through which states and jurisdictions received financial and technical assistance from SAMHSA to assist in their capacity-building efforts toward meeting the block grant reporting requirements.

The general framework for the MH-CLD involves a compilation of the demographic, clinical, and outcome data of individuals served by the SMHA within a state-defined 12-month reporting period. States may choose the calendar year or the state fiscal year as their reporting period. Individuals served is defined as all enrolled individuals who received mental health and support services, including screening, assessment, crisis services, and telemedicine, from programs operated or funded by the SMHA during the reporting period.

Two data sets are submitted each reporting period: Basic Client Information (BCI) is due December 1, and the State Hospital Readmission (SHR) data set is due March 1 of the following year. These two data sets inform five mental health NOMs: access to services/capacity, stability in housing/residential status, 30-day and 180-day readmission to state hospital, adult employment or children's school attendance and education level, and criminal justice involvement. There are five additional NOMs—social connectedness, decreased mental illness symptomatology, perception of care, cost-effectiveness, and use of evidence-based practices—that have not been converted into client-level data reporting and continue to be captured in the URS. Data are reported in accordance with the Health Insurance Portability and Accountability Act (HIPAA) rules for non-protected health information (non-PHI).

In 2010, SAMHSA, through CBHSQ, began a parallel effort of testing the feasibility of building a national behavioral health database that integrates both mental health and substance use data using a uniform platform for data reporting. This effort led to the integration of mental health data into

¹ National outcome measures can be found at https://www.dasis.samhsa.gov/dasis2/mhclld/mh_cld_final_instruction_manual.pdf

the well-established system of reporting for substance use client-level data, known as the Treatment Episode Data Set (TEDS).² To enable reporting of mental health client data to TEDS, the system was modified to incorporate data elements that capture the characteristics unique to the mental health services system and the individuals receiving mental health services. These characteristics include, among others, the extended period of time that an individual may be engaged in treatment, the recovery process, and the simultaneous receipt of different types of services from different service providers. This method of reporting mental health data at the individual level is referred to as the Mental Health Treatment Episode Data Set (MH-TEDS).

Unlike MH-CLD, MH-TEDS is structured around treatment events, that is, admissions and discharges from specific service settings. Admission and discharge records can be linked to track treatment episodes and the treatment services received by individuals. Thus, with MH-TEDS, both the individual client and the treatment episode can serve as a unit of analysis, whereas with MH-CLD the client is the sole unit of analysis. The common TEDS platform in which both the substance use and mental health data are reported enhances the ability to report data for people with co-occurring mental and substance use disorders. It also offers optional data fields for individuals with mental illness that are not captured in MH-CLD, such as referral source, details on criminal justice referral, income sources, and health insurance.

MH-TEDS is a compilation of demographic, substance use, mental health, clinical, legal, and socioeconomic characteristics of individuals who are primarily receiving publicly funded mental health and/or substance use services. MH-TEDS data support program performance measurement and management goals. The same set of mental health NOMs enumerated under MH-CLD is also supported by MH-TEDS. Thus, while both MH-TEDS and MH-CLD collect similar client-level data, the collection methodology differs.

National mental health client-level data reporting was launched in 2011, when MH-CLD was the only reporting method available. The number of states reporting MH-CLD or MH-TEDS has increased from 19 in 2011 to 48 in 2018 (see Appendix B for details). It is important to note that the level and extent of reporting varies. Note also that New York and Tennessee submit data only for a subset of the total number of individuals that they served during a reporting period.

Beginning in the 2014 reporting period, states could choose which method—MH-CLD or MH-TEDS—to use for data reporting. In 2018, 33 states and two other jurisdictions (Commonwealth of the Northern Mariana Islands and the Republic of Palau) used MH-CLD; 11 states (Connecticut, Delaware, Louisiana, Michigan, Mississippi, Nebraska, North Carolina, Oklahoma, Pennsylvania, Vermont, and West Virginia), the District of Columbia, and Puerto Rico used MH-TEDS [Appendix Table B1]. Connecticut reports data through both MH-CLD (for children and adolescents) and MH-TEDS (for adults).

In summary, this report presents the client-level data reported for the 2013 through 2018 reporting periods. Data from all six reporting periods are presented in Chapters 1 through 6. Chapter 7 presents outcomes for individuals served across two or more reporting periods. Chapter 8 presents, for each state or jurisdiction, the number of individuals served in the 2013–2018 reporting periods, according

² Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. *Treatment Episode Data Set (TEDS): 2016. National Admissions to and Discharges from Publicly-Funded Substance Use Treatment*. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2018.

to select mental health diagnoses for age, gender, and race based on the U.S. state-level population. Chapter 9 presents state-level data for the 2013 through 2018 reporting periods, and Chapter 10 presents state-level data only for the 2018 reporting period.

See Appendix A for a crosswalk of data fields collected in MH-TEDS and MH-CLD.

INTRODUCTION

This report presents mental health client-level data submitted for the 2013 through 2018 reporting periods using the Mental Health Client-Level Data (MH-CLD) and the Mental Health Treatment Episode Data Set (MH-TEDS). This report provides information on the demographics, characteristics, and outcomes of individuals served by state mental health agencies (SMHAs) for these six reporting periods, with particular focus on the most recent reporting period. The Substance Abuse and Mental Health Services Administration (SAMHSA) has provided support to states to build capacity to collect and report client-level data.

Methodology

Reconciling data from MH-TEDS and MH-CLD

Both the MH-TEDS and MH-CLD include individuals served through SMHAs during the state-defined 12-month reporting period. Most states elect to use the state fiscal year, which, for the 2018 reporting period, usually ran from July 1, 2017, through June 30, 2018; however, some states define their fiscal year differently. See Appendix B for details.

MH-CLD presented in this report contains one record for each individual served. Data items are populated by the states based on the status of the individual, either at the start and end of the state's reporting period, or the most recently available data. MH-TEDS contains one record for each admission to, and discharge from, a particular service type/setting (referred to as a treatment episode) within the client's treatment continuum during the reporting period. Data collected from MH-TEDS admission records and update/discharge records are stored as two separate data sets that are linked using a HIPAA-compliant, non-protected health information, unique client identifier located in both files. Linking the records allows for the creation of a single client record comparable to the MH-CLD format.

The process of creating an individual record using the MH-TEDS data allows for the reporting of these individuals in both the MH-TEDS and MH-CLD data sets. Tables in all chapters of this report use individuals as the unit of analysis for the combined data from MH-TEDS and MH-CLD.

Selection of reporting variables

MH-TEDS and MH-CLD collect a variety of data items, including demographics (e.g., age, gender, race, and ethnicity), national outcome measures (NOMs, e.g., living arrangements, employment status, criminal justice involvement, and school attendance), and other clinical measures (e.g., co-occurring mental health and substance use disorders). Throughout this report, data for living arrangements and employment are presented according to individuals' status at the end of the given reporting period. Valid data on criminal justice involvement were reported for less than 40 percent of the individuals served for each reporting period; thus, arrest data were excluded from this report. Global Assessment of Functioning (GAF) and Children's Global Assessment (CGAS) scores were dropped from the fifth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) and thus are excluded from this report due to insufficient data. See Appendix C for a description

of the variables; see Appendix D for the federal definitions of serious mental illness (SMI) and serious emotional disturbance (SED).

Mental health diagnoses are reported in the MH-TEDS and MH-CLD as codes from either the International Classification of Diseases, Clinical Modification (ICD-9-CM/ICD-10-CM), or from the DSM-IV/5. Up to three mental health diagnostic codes may be reported for each client. For this report, diagnostic codes were categorized based on a modified version of the Clinical Classification Software (CCS). The mental health section of the CCS has 15 diagnostic categories and 25 subcategories; in all, 12 mental health diagnoses were selected for tabulation, with all other mental health diagnoses grouped as other mental health diagnoses for a given table. See Appendix E for details.

The 12 mental health diagnostic groups used in this report are: trauma- and stressor- related disorders; anxiety disorders; attention deficit/hyperactivity disorder (ADHD); conduct disorder; delirium, dementia, and amnesic and other cognitive disorders (delirium/dementia); bipolar disorders; depressive disorders; oppositional defiant disorder; personality disorders; pervasive developmental disorders; schizophrenia and other psychotic disorders; and alcohol or substance-related disorders.

Records with partially complete data have been retained in this report. For example, an entire client record is not removed from the report because a single data item is missing. When records include missing or invalid data for a specific variable, those records are excluded from tabulations of that variable. The total number of records on which a percentage distribution is based is reported in each table.

Stratification of reporting population and subgroups

In general, this report presents national- and state-level data from the MH-CLD in the 2018 reporting period, as well as trend data from the 2013–2018 reporting periods. There are three major levels of data reporting: (1) the entire population within each age category, (2) the subgroup population of individuals with, or at risk for, serious emotional disturbance (SED) or serious mental illness (SMI), and (3) the entire population within each mental health diagnosis category.

Because different stages of life are associated with distinct forms of mental and behavioral disorders and with distinctive treatment service utilization patterns, this report examines the characteristics of individuals receiving mental health services by the following age groups: children and adolescents (aged 17 years and younger), young adults (aged 18 to 24 years), adults (aged 25 to 54 years), and older adults (aged 55 years and older). Accordingly, Chapters 2 through 5, which focus on specific age groups, include age-specific sets of mental health diagnoses.³

Additionally, this report includes the subpopulation of individuals with co-occurring mental and substance use disorders. For this subpopulation, the report includes individuals aged 12 years and older to match the population used in the substance use TEDS reports. Determination of individuals with co-occurring mental and substance use disorders was based on the variable in the data set called *co-occurring mental and substance use disorders*.

³ For example, individuals with a primary mental health diagnosis of schizophrenia and other psychotic disorders are grouped with *other* for persons aged 17 years and younger in Chapter 2. Although the subject of recent interest, this diagnostic category is not reported separately for this age group. Less than 1 percent of all individuals served in this age group reported schizophrenia and other psychotic disorders as a diagnosis.

Identification of individuals with a serious mental illness (SMI) or serious emotional disturbance (SED) is based on each state's definitions for SMI and SED. This is reported as a single variable: SMI/SED status. Individuals aged 17 years and younger who are indicated in the data set as having SED, or being at risk of SED, are included in the SED subpopulation, while individuals aged 18 years and older who are indicated as having an SMI are included in the SMI subpopulation. Exceptions may exist for a few states where young adults (aged 18 to 21 years) continue to receive mental health services from the state's child mental health system. They are included in the SED subpopulation as reported by the state.

Individual episode-level analysis

Chapter 7 presents data on cohorts of individuals who received mental health services through SMHAs during two or more reporting periods. While the narrative of Chapter 7 focuses specifically on a cohort of individuals served in both the 2015 and 2018 reporting periods, additional tables for Chapter 7 encompass cohorts of individuals served across various other combinations of reporting periods between 2013 and 2018. Individuals were selected for analysis if records with the same unique identification number, from the same state, indicated that they were served in both of the given reporting periods.

Limitations

For both MH-TEDS and MH-CLD, states report only individuals served through SMHAs during the state-defined 12-month reporting period. These data do not represent the total national demand for mental health treatment or describe the mental health status of the national population. In addition, New York and Tennessee report only a subset of the total individuals served through their SMHAs.

Except for Tables 1.12a–b, presented in Chapter 1, this report uses the first-mention (primary) mental health diagnosis as the variable by which individuals are grouped. About 88 percent of all individuals in the 2018 data have at least one mental health diagnosis, while 22 percent have two mental health diagnoses, and 9 percent have three mental health diagnoses [Tables 1.12a–b].

CHAPTER 1

SUMMARY OF ANNUAL DATA IN STATE CLIENT-LEVEL DATA REPORTING: 2013–2018

This chapter provides a brief background on state reporting of mental health client-level data and a summary of the data submitted for the 2013 through 2018 reporting periods.⁴ Data on NOMs and clinical measures are presented by demographic characteristics and mental health diagnoses⁵ for all individuals.

Caution must be used when comparing data over time. The number of states submitting client-level data has varied for each reporting period: 45 states reported 2013 data, 47 states reported 2014 data, 45 states reported 2015 data, 47 states reported 2016 data, 48 states reported 2017 data, and 48 states reported 2018 data.

All data tables represent sums of individuals who received publicly funded mental health services—individuals served—through the auspices of state mental health agencies (SMHAs).

Individuals Served

Tables 1.1a–b. For the 2013 reporting period, 45 states reported that a total of 5,576,659 individuals received mental health services; for the 2014 reporting period, 47 states reported that a total of 5,818,881 individuals received mental health services; for the 2015 reporting period, 45 states reported that a total of 5,311,897 individuals received mental health services; for the 2016 reporting period, 47 states reported that a total of 5,813,339 individuals received mental health services; for the 2017 reporting period, 48 states reported that a total of 5,934,383 individuals received mental health services; and for the 2018 reporting period, 48 states reported that a total of 6,213,791 individuals received mental health services.⁶

Gender

Tables 1.2a–b. Females accounted for between 51 and 52 percent of all individuals served in every reporting period between 2013 and 2018.

⁴ Hereafter, “states” may refer to any of the 50 states, District of Columbia, the Commonwealth of Puerto Rico, and the combined jurisdictions of the Republic of Palau and the Commonwealth of the Northern Mariana Islands.

⁵ Except where otherwise noted, throughout this report, the term “diagnosis” refers to the first reported mental health diagnosis on the client-level data record. This is only the first of up to three diagnoses (per client) that qualify as a mental health diagnosis, as recorded by the reporting state. While first-mentioned mental health diagnoses are mutually exclusive, a client with a first-mention diagnosis of, for example, a depressive disorder, may also have a second-mention diagnosis of anxiety disorder, and a third-mention diagnosis of schizophrenia or other psychotic disorder.

⁶ Data were not submitted or were incomplete for one or more years in some states or jurisdictions.

Age group

Tables 1.2a–b. In all six reporting periods from 2013 through 2018, between 28 and 30 percent of individuals served were aged 17 years and younger; between 10 and 11 percent were aged 18 to 24 years; 46 percent were aged 25 to 54 years; and between 14 and 15 percent were aged 55 years and older. The most frequently reported diagnoses among specific age groups were relatively consistent across the six reporting periods [Tables 1.4a–1.7b].

*Race and ethnicity*⁷

Tables 1.3a–b. For each of the reporting periods from 2013 through 2018, Whites represented between 67 and 68 percent of the individuals served; Blacks or African Americans represented between 20 and 21 percent; and American Indians or Alaska Natives, Asians, Native Hawaiians or Other Pacific Islanders, and other groups, combined, represented between 12 and 13 percent.⁸

Hispanics accounted for between 15 and 17 percent of all individuals served in each of the six reporting periods from 2013 through 2018.

Table 1.23b. Blacks or African Americans accounted for 20 percent of individuals served in the 2018 reporting period but just 13 percent of the 2018 U.S. population.⁹

Marital status

Tables 1.8a–b. Among individuals aged 18 years and older in each reporting period from 2013 through 2018, more than half reported never being married (ranging between 55 and 60 percent); those who were married at the time of reporting ranged between 15 and 18 percent.

Tables 1.22a–c. Among individuals aged 25 to 54 years served in the 2018 reporting period, 16 percent were currently married. In contrast, 57 percent of the 2018 U.S. population in that age group were currently married.¹⁰

Living arrangements

Tables 1.8a–b. Individuals living in a private residence accounted for between 83 and 85 percent of those served in the six reporting periods from 2013 through 2018; between 3 and 4 percent reported being homeless.

SMI/SED status

Tables 1.9a–b. Individuals with serious mental illness (SMI) accounted for between 72 and 78 percent of individuals aged 18 years and older who were served in any of the six reporting periods from 2013 through 2018. Children and adolescents aged 17 years and younger who had, or were at risk for,

⁷ Data on specific Hispanic or Latino origin (hereafter referred to as “Hispanic origin”) are presented for a subset of states that reported information on individuals’ specific Hispanic or Latino origin.

⁸ *Other* includes individuals who reported two or more races or some other race.

⁹ Source: U.S. Census Bureau, Population Estimates Program, SC-EST2018-alldata6: “Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2018.”

¹⁰ Source: U.S. Census Bureau, Current Population Survey, 2018 Annual Social and Economic Supplement (ASEC), March 2018, Table A1. <https://www.census.gov/data/tables/2018/demo/families/cps-2018.html>

serious emotional disturbance (SED) accounted for between 70 and 76 percent of individuals aged 17 years and younger who were served in any of the six reporting periods from 2013 through 2018.

Employment

Tables 1.10a–b. Among individuals aged 18 years and older served in any of the six reporting periods from 2013 through 2018, the proportion employed ranged between 17 and 23 percent, the proportion unemployed ranged between 27 and 31 percent, and the proportion not in the labor force ranged between 48 and 52 percent.

Education

Tables 1.10a–b. Individuals who completed Grade 12 or a general equivalency diploma (GED) accounted for between 46 and 49 percent of individuals aged 18 years and older who were served in any of the six reporting periods from 2013–2018; between 19 and 24 percent completed at least a year of education beyond Grade 12.

Veteran status

Tables 1.10a–b. In both the 2017 and 2018 reporting periods, veterans accounted for 3 percent of individuals aged 18 years and older served.

Client treatment status at the start of the reporting period

Tables 1.11a–b. Among individuals served in any of the six reporting periods from 2013 through 2018, those newly admitted to treatment¹¹ ranged between 46 to 54 percent, and those continuing treatment¹² ranged between 46 and 54 percent.

Service setting

Tables 1.11a–b. For the 2013 to 2018 reporting periods, the proportion of individuals who received mental health services from community-based programs ranged between 97 and 98 percent. Further analysis of the settings in which individuals received mental health services during the 2018 reporting period is presented in Chapters 2 through 6.

Demographic characteristics of individuals with selected primary mental health diagnoses

Tables 1.14a–1.19b present the gender, age, race, and ethnicity of individuals with the following primary diagnoses: trauma- and stressor-related disorders [Tables 1.14a–b], anxiety disorders [Tables 1.15a–b], attention deficit/hyperactivity disorder [Tables 1.16a–b], bipolar disorders [Tables 1.17a–b], depressive disorders [Tables 1.18a–b], and schizophrenia and other psychotic disorders [Tables 1.19a–b].

¹¹ Individuals who were newly admitted to treatment either (a) had not previously received a service and started receiving services during the reporting year, or (b) had previously received a service, were discharged or inactive for an extended period of time, and resumed receiving services during the reporting period.

¹² Individuals continuing treatment consists of those who were admitted prior to the start of the reporting period and who received a service during the reporting period.


Demographic characteristics, by status for co-occurring mental health and substance use disorders

Tables 1.20a–1.21b present the gender, age, race, and ethnicity of individuals grouped by the presence or absence of co-occurring mental health and substance use disorders.

Regional Distributions

Tables 1.13a–b and Figure 1. About a third (between 31 and 36 percent) of all individuals who received mental health services in the 2013 through 2018 reporting periods were in the South. The Census regions with the next largest proportions of individuals who received mental health services in the 2013 through 2018 reporting periods were the West (between 27 and 31 percent), the Midwest (between 18 and 27 percent), and the Northeast (between 11 and 23 percent).

Figure 1. Census regions and Census divisions of the United States



SOURCE: U.S. Census Bureau.

Any-Mention Mental Health Diagnoses

Tables 1.12a–b present the number and percentage distribution of individuals who received any mention—whether first, second, or third—of the listed mental health diagnoses for the 2013, 2014, 2015, 2016, 2017, and 2018 reporting periods. Differences among reporting periods should be interpreted with great caution; variation may be more a function of the mix and number of reporting states than of actual differences.

CHAPTER 2

CHARACTERISTICS OF CHILDREN AND ADOLESCENTS AGED 17 YEARS AND YOUNGER RECEIVING MENTAL HEALTH SERVICES

This chapter presents data on children and adolescents aged 17 years and younger who received mental health services through SMHAs in the 2013–2018 reporting periods, with particular focus on the 2018 reporting period. Data on NOMs and clinical measures are presented by demographic characteristics and mental health diagnoses for all children and adolescents. Characteristics for the subpopulation of children and adolescents with, or at risk for, SED are presented where the percentages notably differ from the characteristics of all children and adolescents.

Number Served

Tables 2.1a–b present the total number of individuals aged 17 years and younger served in the 2013 through 2018 reporting periods, by demographic characteristics.

The number of individuals served who were aged 17 years and younger increased from 1,650,728 in the 2013 reporting period to 1,800,613 in the 2018 reporting period.

Tables 2.3a–2.4c. In the 2018 reporting period, 1,630,278 individuals aged 17 years and younger had a valid mental health diagnosis code, of which 1,174,041 were reported to have, or be at risk of, SED.

Only individuals with valid mental health diagnoses are included in these and all subsequent tabulations of mental health diagnoses in this chapter.

Gender

Tables 2.3a–2.4c. Tables 2.3a–c present the most frequently reported diagnoses, by gender, among individuals aged 17 years and younger, while Tables 2.4a–c present the most frequently reported diagnoses, by gender, for individuals aged 17 years and younger with SED. Males accounted for 57 percent and females accounted for 43 percent of individuals aged 17 years and younger who were served in the 2018 reporting period. The most frequently reported diagnoses differed by gender.

Race

Tables 2.3a–2.4c. The most frequently reported diagnoses among individuals aged 17 years and younger served in the 2018 reporting period varied by race. Trauma- and stressor-related disorders were reported most frequently among American Indians or Alaska Natives (30 percent), Native Hawaiian or Other Pacific Islanders (27 percent), Whites (24 percent), and Asians (23 percent); ADHD was the most frequently reported diagnosis among Blacks or African Americans (32 percent).

Ethnicity

Tables 2.3a–2.4c. Among Hispanic individuals aged 17 years and younger served in the 2018 reporting period, the most frequently reported diagnoses were trauma- and stressor-related disorders (23 percent); next were ADHD (20 percent) and depressive disorders (17 percent). Among non-Hispanics, the most frequently reported diagnoses were ADHD (25 percent), trauma- and stressor-related disorders (23 percent), and depressive disorders (13 percent).

Living Arrangements

Tables 2.5a–2.6c. Among individuals aged 17 years and younger served in the 2018 reporting period, trauma- and stressor-related disorders were the most frequently reported diagnoses among those who were in foster care (47 percent), were homeless (39 percent), were in other living arrangements (25 percent), or were in residential care (21 percent). During the same period, individuals aged 17 years and younger living in a private residence were most frequently reported with an ADHD diagnosis (24 percent).

SED Status

Tables 2.5a–2.6c. Among individuals aged 17 years and younger with SED served in the 2018 reporting period, the most frequently reported diagnosis was ADHD (23 percent); among those without SED, the most frequently reported diagnoses were trauma- and stressor-related disorders (27 percent).

Client Treatment Status at the Start of the Reporting Period

Tables 2.7a–2.8c. Among individuals aged 17 years and younger who were continuing treatment during the 2018 reporting period, the most frequently reported diagnosis was ADHD (29 percent); among those who were newly admitted to treatment in the 2018 reporting period, the most frequently reported diagnoses were trauma- and stressor-related disorders (27 percent).

SED Status, by Service Setting

Table 2.9. SED status varied by service setting among individuals aged 17 years and younger served in the 2018 reporting period. It should be noted that individuals may have received services in more than one setting and therefore may be counted in multiple rows of Table 2.9.

CHAPTER 3

CHARACTERISTICS OF YOUNG ADULTS AGED 18 TO 24 YEARS RECEIVING MENTAL HEALTH SERVICES

This chapter presents data on young adults aged 18 to 24 years who received mental health services through SMHAs in the 2013–2018 reporting periods, with particular focus on the 2018 reporting period. Data on NOMs and clinical measures for all young adults are presented by demographic characteristics and mental health diagnoses. Characteristics for the subpopulation of young adults with an SMI are presented where the percentages notably differ from all young adults who received mental health services.

Number Served

Tables 3.1a–b present the total number of individuals aged 18 to 24 years served in the 2013 through 2018 reporting periods, by demographic characteristics.

The number of individuals served who were aged 18 to 24 years increased from 594,733 in the 2013 reporting period to 645,452 in the 2018 reporting period.

Tables 3.3a–3.4c. In the 2018 reporting period, 557,478 individuals aged 18 to 24 years had a valid mental health diagnosis code, of which 380,549 were reported to have SMI.

Only individuals with valid mental health diagnoses are included in these and all subsequent tabulations of mental health diagnoses in this chapter.

Gender

Tables 3.3a–c. Females accounted for 54 percent of individuals aged 18 to 24 years who were served during the 2018 reporting period. The most frequently reported diagnoses among males were depressive disorders (24 percent), schizophrenia and other psychotic disorders (14 percent), anxiety disorders (11 percent), and bipolar disorders (11 percent). Among females aged 18 to 24 years, the most frequently reported diagnoses were depressive disorders (35 percent), anxiety disorders (17 percent), and trauma- and stressor-related disorders (15 percent); schizophrenia and other psychotic disorders were reported in 5 percent of females aged 18 to 24 years.

Race and Ethnicity

Tables 3.3a–c. Depressive disorders were the most frequently reported diagnoses among individuals of every racial and ethnic group aged 18 to 24 years served during the 2018 reporting period, ranging from 27 percent among Blacks or African Americans to 31 percent among Native Hawaiians or Other Pacific Islanders and Whites, respectively.

Education

Tables 3.5a–c. Among individuals of every education level aged 18 to 24 years served during the 2018 reporting period, the most frequently reported diagnoses were depressive disorders, ranging from 24 percent of individuals who completed Grade 8 or less to 38 percent among individuals who completed at least a year of education beyond Grade 12.

Marital Status

Tables 3.5a–3.6c. Among individuals across all marital status categories aged 18 to 24 years who were served during the 2018 reporting period, the most frequently reported diagnoses were depressive disorders, ranging from 29 percent among widowed individuals to 38 percent among separated individuals.

Living Arrangements

Tables 3.7a–3.8c. Among individuals aged 18 to 24 years who were served during the 2018 reporting period, depressive disorders were the most frequently reported diagnoses among those who lived in a private residence (32 percent), those who had other living arrangements (29 percent), those who were homeless (27 percent), those who were in foster care (21 percent), and those in residential care (20 percent). Trauma- and stressor-related disorders were the most frequently reported diagnoses for those who were in jail or correctional facilities (22 percent). Among those in institutional settings, schizophrenia and other psychotic disorders were the most frequently reported diagnoses (32 percent).

SMI Status

Tables 3.7a–3.8c. Among individuals aged 18 to 24 years served during the 2018 reporting period, depressive disorders were the most frequently reported diagnoses for those who had SMI (34 percent). Trauma- and stressor-related disorders were the most frequently reported diagnosis for those who did not have SMI (20 percent).

Veteran Status

Tables 3.7a–3.8c. Among individuals aged 18 to 24 years served during the 2018 reporting period, depressive disorders were the most frequently reported diagnoses for those who were veterans (37 percent) and those who were not veterans (30 percent).

Employment Status

Tables 3.9a–3.10c. Among individuals aged 18 to 24 years served in the 2018 reporting period, 27 percent were employed, 28 percent were unemployed, and 45 percent were not in the labor force; of those not in the labor force, 45 percent were students.

Client Treatment Status at the Start of the Reporting Period

Tables 3.11a–3.12c. Among individuals aged 18 to 24 years served during the 2018 reporting period, the most frequently reported diagnoses were depressive disorders, regardless of individuals' treatment status at the start of the reporting period (32 percent among individuals who were newly admitted to treatment and 26 percent among individuals who were continuing treatment).

Service Setting

Tables 3.11a–3.12c. The most frequently reported diagnoses varied by mental health service setting. Individuals may have received services in more than one service setting during the reporting period and therefore may be counted in multiple rows of Tables 3.11a–c and Tables 3.12a–c.

Among individuals aged 18 to 24 years served in the 2018 reporting period, the most frequently reported diagnoses were schizophrenia and other psychotic disorders for those in state psychiatric hospitals (41 percent) and residential treatment centers (23 percent); depressive disorders were the most frequently reported diagnoses among those served in other psychiatric inpatient facilities (31 percent) and community-based programs (30 percent). Trauma- and stressor-related disorders were the most frequently reported diagnoses among those in institutions under the justice system (24 percent).

SMI Status, by Service Setting

Table 3.13. Among individuals aged 18 to 24 years served during the 2018 reporting period, SMI status varied by service setting. It should be noted that individuals may have received services in more than one setting and therefore may be counted in multiple rows of Table 3.13.

CHAPTER 4

CHARACTERISTICS OF ADULTS AGED 25 TO 54 YEARS RECEIVING MENTAL HEALTH SERVICES

This chapter presents data on adults aged 25 to 54 years who received mental health services through SMHAs in the 2013–2018 reporting periods, with particular focus on the 2018 reporting period. Data on NOMs and clinical measures for all adults aged 25 to 54 years old are presented by demographic characteristics and mental health diagnoses. Characteristics for the subpopulation of adults with SMI are presented where the percentages notably differ from the characteristics of all adults aged 25 to 54 years old.

Number Served

Tables 4.1a–b present the total of individuals aged 25 to 54 years reported in the 2013 through 2018 reporting periods, by demographic characteristics.

The number of individuals served who were aged 25 to 54 years increased from 2,558,630 in the 2013 reporting period to 2,831,082 in the 2018 reporting period.

Tables 4.3a–4.4c. In the 2018 reporting period, 2,476,907 individuals aged 25 to 54 years had a valid mental health diagnosis code, of which 1,848,515 were reported to have SMI.

Only individuals with valid mental health diagnoses are included in these and all subsequent tabulations of mental health diagnoses in this chapter.

Gender

Tables 4.3a–4.4c. Females accounted for 55 percent of individuals aged 25 to 54 years who were served during the 2018 reporting period. The most frequently reported diagnoses among females were depressive disorders (33 percent), bipolar disorders (18 percent), and trauma- and stressor-related disorders (13 percent). Among males, the most frequently reported diagnoses were depressive disorders (25 percent), schizophrenia and other psychotic disorders (24 percent), and bipolar disorders (14 percent).

Race

Tables 4.3a–4.4c. Among individuals of all races, except Asians, who were aged 25 to 54 years and were served in the 2018 reporting period, depressive disorders were the most frequently reported diagnoses; depressive disorders were reported among 30 percent of Whites, 29 percent of Blacks or African Americans, 27 percent of American Indian or Alaska Natives, and 27 percent of Native Hawaiians or Other Pacific Islanders. Among Asians aged 25 to 54 years served in the 2018 reporting period, schizophrenia and other psychotic disorders were the most frequently reported diagnoses (30 percent).

Ethnicity

Tables 4.3a–4.4c. Among individuals aged 25 to 54 years served in the 2018 reporting period, depressive disorders were the most frequently reported diagnoses for both those of Hispanic origin (31 percent) and non-Hispanic origin (30 percent).

Education

Tables 4.5a–4.6c. Among individuals aged 25 to 54 years served in the 2018 reporting period, depressive disorders were the most frequently reported diagnoses, ranging from 28 percent among those who completed at most Grade 8 to 32 percent among those who completed at least a year of education beyond Grade 12.

Marital Status

Tables 4.5a–4.6c. Depressive disorders were the most frequently reported diagnoses among individuals aged 25 to 54 years served during the 2018 reporting period across all marital status categories (31 percent). Among individuals in this age group with schizophrenia and other psychotic disorders, 76 percent were never married.

Living Arrangements

Tables 4.7a–4.8c. Among individuals aged 25 to 54 years served during the 2018 reporting period who were homeless, lived in a private residence, or had other living arrangements,¹³ the most frequently reported diagnoses were depressive disorders, ranging from 25 percent of individuals with other living arrangements to 31 percent of individuals living in a private residence.

Among individuals aged 25 to 54 years served in the 2018 reporting period, schizophrenia and other psychotic disorders were the most frequently reported diagnoses among those living in an institutional setting (47 percent), in residential care (40 percent), in foster care (28 percent), or in jails or correctional facilities (20 percent).

SMI Status

Tables 4.7a–4.8c. Among individuals with SMI who were aged 25 to 54 years and were served in the 2018 reporting period, the most frequently reported diagnoses were depressive disorders (32 percent); the next most frequent diagnoses for those with SMI were schizophrenia and other psychotic disorders (20 percent) and bipolar disorders (18 percent). Among those without SMI, the most frequent diagnoses were depressive disorders (21 percent) and trauma- and stressor-related disorders (21 percent).

Among individuals with schizophrenia and other psychotic disorders who were aged 25 to 54 years and were served in the 2018 reporting period, 93 percent were reported to have SMI.

¹³ “Other living arrangements” includes crisis residences and other undefined residences.

Veteran Status

Tables 4.7a–4.8c. Among individuals aged 25 to 54 years served during the 2018 reporting period, depressive disorders were the most frequently reported diagnoses both for those who were veterans (29 percent) and those who were not veterans (29 percent).

Among those who were veterans, 16 percent had diagnoses of trauma- and stressor-related disorders; 12 percent of non-veterans aged 25 to 54 years who were served in the 2018 reporting period had diagnoses of trauma- and stressor-related disorders.

Employment Status

Tables 4.9a–4.10c. Among individuals aged 25 to 54 years served in the 2018 reporting period, 26 percent were employed, 31 percent were unemployed, and 43 percent were not in the labor force. The most frequent diagnoses across all employment statuses were depressive disorders.

Client Treatment Status at the Start of the Reporting Period

Tables 4.11a–4.12c. Among individuals aged 25 to 54 years served in the 2018 reporting period, depressive disorders were the most frequently reported diagnoses, regardless of individuals' treatment status at the start of the reporting period; depressive disorders were reported for 31 percent of individuals who were newly admitted to treatment and for 28 percent of individuals continuing treatment. The second most frequently reported diagnoses for individuals continuing treatment were schizophrenia and other psychotic disorders (22 percent); for those newly admitted to treatment, the second most frequently reported diagnoses were anxiety disorders (14 percent).

Service Setting

Tables 4.11a–4.12c. It should be noted that individuals may have received services in more than one service setting during the reporting period and therefore may be counted in multiple rows of Tables 4.11a–c and 4.12a–c.

Community-based programs accounted for 97 percent of individuals aged 25 to 54 years served during the 2018 reporting period. Among individuals aged 25 to 54 years served in community-based programs during the 2018 reporting period, the most frequent diagnoses were depressive disorders (30 percent), bipolar disorders (16 percent), and schizophrenia and other psychotic disorders (16 percent). Depressive disorders were also the most frequently reported diagnoses among those aged 25 to 54 years who were served during the 2018 reporting period in other psychiatric inpatient settings (30 percent). Schizophrenia and other psychotic disorders were the most frequently reported diagnoses among individuals aged 25 to 54 years who were served during the 2018 reporting period in state psychiatric hospitals (53 percent), residential treatment centers (37 percent), and institutions under the justice system (21 percent).

SMI Status, by Service Setting

Table 4.13 presents the number and percentage of individuals aged 25 to 54 years served during the 2018 reporting period, by mental health service setting and SMI status. It should be noted that individuals may have received services in more than one setting and therefore may be counted in multiple rows of Table 4.13.

CHAPTER 5

CHARACTERISTICS OF OLDER ADULTS AGED 55 YEARS AND OLDER RECEIVING MENTAL HEALTH SERVICES

This chapter presents data on individuals aged 55 years and older who received mental health services through SMHAs in the 2013–2018 reporting periods, with particular focus on the 2018 reporting period. Data on NOMs and clinical measures for older adults are presented by diagnostic categories and mental health diagnoses. Characteristics for the subpopulation of older adults with SMI are presented where the percentages notably differ from the characteristics of all individuals aged 55 and older.

Number Served

Tables 5.1a–b present the total number of individuals aged 55 years and older served in the 2013 through 2018 reporting periods, by demographic characteristics.

The number of individuals served who were aged 55 years and older increased from 769,081 in the 2013 reporting period to 932,573 in the 2018 reporting period.

Tables 5.3a–5.4c. In the 2018 reporting period, 827,060 individuals aged 55 years and older had a valid mental health diagnosis code, of which 655,241 were reported to have SMI.

Only individuals with valid mental health diagnoses are included in these and all subsequent tabulations of mental health diagnoses in this chapter.

Gender

Tables 5.3a–5.4c. Females accounted for 59 percent of individuals aged 55 years and older who were served in the 2018 reporting period. Among females, the most frequently reported diagnoses were depressive disorders (39 percent), schizophrenia and other psychotic disorders (19 percent), and bipolar disorders (15 percent). Among males, the most frequently reported diagnoses were schizophrenia and other psychotic disorders (30 percent), depressive disorders (30 percent), and bipolar disorders (12 percent).

Race

Tables 5.3a–5.4c. Among individuals of all races, except for Blacks or African Americans, who were aged 55 years and older and were served in the 2018 reporting period, the most frequently reported diagnoses were depressive disorders, ranging from 34 percent of American Indians or Alaska Natives to 42 percent of Asians; among Blacks or African Americans, the two most frequently reported diagnoses were schizophrenia and other psychotic disorders (38 percent) and depressive disorders (32 percent).

Ethnicity

Tables 5.3a–5.4c. Among individuals aged 55 years and older served during the 2018 reporting period, the most frequently reported diagnoses were depressive disorders both for those of Hispanic origin (43 percent) and non-Hispanic origin (35 percent). Schizophrenia and other psychotic disorders were the second most frequently reported diagnoses both for Hispanics and non-Hispanics (21 percent and 24 percent, respectively).

Education

Tables 5.5a–5.6c. Among individuals across all levels of educational attainment who were aged 55 years or older and were served in the 2018 reporting period, the most frequently reported diagnoses were depressive disorders.

Marital Status

Tables 5.5a–5.6c. Among individuals aged 55 years and older served in the 2018 reporting period, the most frequently reported diagnoses across all marital statuses, except those who were never married, were depressive disorders, ranging from 38 percent of those who were divorced to 44 percent of those who were widowed. Among never-married individuals aged 55 years and older who were served in the 2018 reporting period, the most frequently reported diagnoses were schizophrenia and other psychotic disorders (37 percent).

Living Arrangements

Tables 5.7a–5.8c. Among individuals aged 55 years and older served in the 2018 reporting period who were living in a private residence, were homeless, or had other living arrangements, the most frequently reported diagnoses were depressive disorders, ranging from 29 percent among those with other living arrangements to 38 percent among those who were living in a private residence.

Among individuals aged 55 years and older served in the 2018 reporting period, schizophrenia and other psychotic disorders were the most frequently reported diagnoses among those living in residential care (54 percent), foster care (44 percent), an institutional setting (39 percent), or a jail or correctional facility (29 percent).

SMI Status

Tables 5.7a–5.8c. Among individuals aged 55 years and older served in the 2018 reporting period, the most frequently reported diagnoses were depressive disorders, both among those who had SMI (37 percent) and those who did not have SMI (28 percent). For those with SMI, the second most frequently reported diagnoses were schizophrenia and other psychotic disorders (27 percent); among those without SMI, the second most frequently reported diagnoses were anxiety disorders (16 percent).

Veteran Status

Tables 5.7a–5.8c. Among individuals aged 55 years and older served during the 2018 reporting period, depressive disorders were the most frequently reported diagnoses both for those who were veterans (31 percent) and those who were not veterans (35 percent).

Schizophrenia and other psychotic disorders were reported among 28 percent of veterans aged 55 years and older who were served in the 2018 reporting period; 24 percent of non-veterans aged 55 years and older had diagnoses of schizophrenia and other psychotic disorders.

Employment Status

Tables 5.9a–5.10c. Among individuals aged 55 years and older served in the 2018 reporting period, 12 percent were employed, 18 percent were unemployed, and 70 percent were not in the labor force (of which 61 percent were disabled and 13 percent were retired). Among individuals across all employment statuses who were aged 55 years and older and were served in the 2018 reporting period, the most frequently reported diagnoses were depressive disorders, ranging from 34 percent among those not in the labor force to 40 percent among those unemployed.

Client Treatment Status at the Start of the Reporting Period

Tables 5.11a–5.12c. Among individuals aged 55 years and older served in the 2018 reporting period, depressive disorders were the most frequently reported diagnoses, regardless of individuals' treatment status at the start of the reporting period; depressive disorders were reported for 38 percent of those newly admitted to treatment in the 2018 reporting period and for 33 percent of individuals who were continuing treatment. Schizophrenia and other psychotic disorders were the second most frequently reported diagnoses in the same period for individuals aged 55 years and older who were continuing treatment (29 percent) and for those newly admitted to treatment (15 percent).

Service Setting

Tables 5.11a–5.12c. Individuals may have received services in more than one service setting and therefore may be counted in multiple rows of Tables 5.11a–c and 5.12a–c.

Community-based programs accounted for 98 percent of individuals aged 55 years and older served during the 2018 reporting period. Depressive disorders were the most frequent diagnoses among individuals aged 55 years and older who were served during the 2018 reporting period in community-based programs (36 percent) and other psychiatric inpatient settings (34 percent). Schizophrenia and other psychotic disorders were the most frequently reported diagnoses among individuals aged 55 years and older served during the same period in state psychiatric hospitals (59 percent), residential treatment centers (51 percent), and institutions under the justice system (30 percent).

SMI Status, by Service Setting

Table 5.13 presents the number and percentage distribution of individuals aged 55 years and older who were served in the 2018 reporting period by mental health service setting and SMI status. It should be noted that individuals may have received services in more than one setting during the reporting period and therefore may be counted in multiple columns of Table 5.13.

During the 2018 reporting period, community-based programs reported the lowest proportion among all service settings of individuals aged 55 years and older who had SMI (79 percent); institutions under the justice system and state psychiatric hospitals had the highest proportions of individuals aged 55 years and older with SMI (92 and 91 percent each).

CHAPTER 6

CHARACTERISTICS OF INDIVIDUALS WITH CO-OCCURRING MENTAL HEALTH AND SUBSTANCE USE DISORDERS RECEIVING MENTAL HEALTH SERVICES

This chapter presents data on individuals aged 12 years and older reported to have co-occurring mental health and substance use disorders who received mental health services through SMHAs during the 2013–2018 reporting periods, with particular focus on the 2018 reporting period. Data on NOMs and clinical measures for all individuals are presented by demographic characteristics and mental health diagnoses. It should be noted that the definition for co-occurring mental health and substance use disorders includes those who have a reported, identified substance use disorder but may not have a formal substance use diagnosis in addition to a reported mental health diagnosis. It should also be noted that data reported in this chapter exclude individuals whose only mental health diagnoses were alcohol or substance-related disorders.

Number Served

Tables 1.2a and 6.1a. States reported a total of 1,026,726 individuals with co-occurring mental health and substance use disorders aged 12 years and older who were served in the 2018 reporting period, accounting for 17 percent of all individuals served during this period.

Mental health diagnoses for individuals with co-occurring mental health and substance use disorders aged 12 years and older who were served in the 2018 reporting period differed somewhat by gender, age group, race, ethnicity, living arrangements, employment, not in labor force details, service setting, and timing of admission. Mental health diagnoses for this cohort were largely similar across SMI/SED status.

Gender

Tables 6.2a–c. For males with co-occurring mental health and substance use disorders aged 12 years or older served during the 2018 reporting period, the most frequently reported diagnoses were depressive disorders (25 percent), schizophrenia and other psychotic disorders (22 percent), and bipolar disorders (14 percent). Among their female counterparts, the most frequently reported diagnoses were depressive disorders (31 percent), bipolar disorders (19 percent) and anxiety disorders (12 percent).

Age

Tables 6.2a–c. Among individuals with co-occurring mental health and substance use disorders served in the 2018 reporting period, for all age categories, the most frequently reported diagnoses were depressive disorders, ranging between 21 percent in the 12 to 14 years age group and 33 percent in the 55 to 59 years age group, as well as 33 percent in the 60 to 64 years age group. Schizophrenia

and other psychotic disorders were reported almost as frequently as depressive disorders in the 65 years and older age group (29 percent).

Race

Tables 6.3a–c. Among individuals with co-occurring mental health and substance use disorders served during the 2018 reporting period, the most frequently reported diagnoses were depressive disorders for Whites (29 percent) and American Indian or Alaska Natives (29 percent). Schizophrenia and other psychotic disorders were the most frequently reported diagnoses for Asians (27 percent), Blacks or African Americans (29 percent). The most frequently reported diagnoses among Hawaiians or Other Pacific Islanders were schizophrenia and other psychotic disorders (24 percent).

Ethnicity

Tables 6.3a–c. Among individuals with co-occurring mental health and substance use disorders served in the 2018 reporting period, the most frequently reported diagnoses were depressive disorders for both those of Hispanic origin (26 percent) and non-Hispanic origin (29 percent).

Substance Use Diagnosis

Tables 6.4a–c. Among individuals with co-occurring mental health and substance use disorders served during the 2018 reporting period, mental health diagnoses showed no major differences by substance use diagnosis. Depressive disorders were the most frequently reported diagnoses for individuals who had co-occurring alcohol dependence (33 percent), opioid dependence (33 percent), cocaine dependence (30 percent), and marijuana dependence (25 percent). Depressive disorders were also the most frequently reported mental health diagnoses for those who had co-occurring non-dependent use of opioids (34 percent), cocaine (28 percent), alcohol (29 percent), and marijuana (25 percent). Depressive disorders were also the most frequently reported diagnoses for individuals with alcohol intoxication (34 percent), alcoholic psychoses (32 percent), and drug psychoses (24 percent).

Living Arrangements

Tables 6.6a–c. Among individuals with co-occurring mental health and substance use diagnoses served during the 2018 reporting period, depressive disorders were the most frequently reported diagnoses for those living in private residences (30 percent), those who were homeless (29 percent), and among those in other living arrangements (27 percent). Schizophrenia and other psychotic disorders were the most frequently reported diagnoses among those in residential care (37 percent), institutional settings (34 percent), and foster care (28 percent).

SMI/SED Status

Tables 6.6a–c. Among individuals who had co-occurring mental health and substance use disorders and were served during the 2018 reporting period, depressive disorders were the most frequently reported diagnoses among those with SMI (29 percent), among those with SED (25 percent), and among those without SMI/SED (25 percent).

Veteran Status

Tables 6.6a–c. Among individuals who had co-occurring mental health and substance use disorders and who were served during the 2018 reporting period, depressive disorders were the most frequently reported diagnoses both for those who were veterans (32 percent) and those who were not (30 percent).

Among individuals who had co-occurring mental health and substance use disorders who were served during the 2018 reporting period, schizophrenia and other psychotic disorders were reported among 22 percent of veterans and 17 percent of non-veterans.

Employment Status

Tables 6.7a–c. Among individuals who had co-occurring mental health and substance use disorders, who were aged 18 years and older, and who were served during the 2018 reporting period, 21 percent were employed, 30 percent were unemployed, and 48 percent were not in the labor force; the most frequently reported diagnoses were depressive disorders across all employment statuses (34 percent among the unemployed, 32 percent among the employed, and 27 percent among those not in the labor force).

Service Setting

Tables 6.8a–c. Individuals may have received services in more than one service setting during the reporting period and therefore may be counted in multiple rows of Tables 6.8a–c.

Among individuals with co-occurring mental health and substance use disorders served during the 2018 reporting period, the most frequently reported diagnosis varied by service setting: Schizophrenia and other psychotic disorders were the most frequently reported diagnoses for those in state psychiatric hospitals (49 percent), residential treatment centers (31 percent), and institutions under the justice system (19 percent). Depressive disorders were the most frequently reported diagnoses for those in other psychiatric inpatient settings (32 percent) and community-based programs (29 percent).

Client Treatment Status at the Start of the Reporting Period

Tables 6.8a–c. Among individuals who had co-occurring mental health and substance use disorders and who were served during the 2018 reporting period, depressive disorders were the most frequently reported diagnoses, regardless of individuals' treatment status at the start of the reporting period; depressive disorders were reported for 32 percent of those newly admitted to treatment in the 2018 reporting period and for 25 percent of individuals who were continuing treatment.

CHAPTER 7

CLIENT OUTCOMES AMONG INDIVIDUALS RECEIVING MENTAL HEALTH SERVICES: 2013–2018

This chapter presents data on cohorts of individuals who received mental health services through SMHAs during two or more reporting periods. While the following narrative focuses specifically on a cohort of individuals served in both the 2015 and 2018 reporting periods [Tables 7.1a–7.6b], additional tables for Chapter 7 [Tables 7.7–7.16b] encompass cohorts of individuals served across various other combinations of reporting periods between 2013 and 2018. The length of treatment was not taken into consideration. NOMs are presented by demographic characteristics, mental health diagnoses, and clinical measures for all individuals, based on the last reported status in each period.

Individuals were selected for analysis by matching any individual records in 2015 with any in 2018 from the same state and with the same unique identification number; other individuals served in the 2015 and 2018 reporting periods were excluded from analysis in this chapter. The resulting data set contained 1,800,068 unique clients [Table 7.1a].

Living Arrangements

Tables 7.1a–7.2c. States reported a total of 1,019,988 individuals served in both the 2015 and 2018 reporting periods who had known living arrangements in both reporting years.

Among individuals served in both the 2015 and 2018 reporting periods, those living in a private residence accounted for 83 percent in both 2015 and 2018; those living in residential care accounted for 5 percent in 2015 and 6 percent in 2018; those who were homeless accounted for 3 percent in 2015 and 4 percent in 2018; those who were in institutional settings accounted for 2 percent in both 2015 and 2018; those in jails/correctional facilities accounted for 1 percent in 2015 and 2 percent in 2018; and those in foster care accounted for 2 percent in 2015 and 1 percent in 2018.

Employment Status


Tables 7.3a–7.4c. States reported a total of 636,139 individuals aged 18 or older in 2015 who were served in both the 2015 and 2018 reporting periods who had known employment status in both years. Among individuals aged 18 years and older in 2018 who were served in both the 2015 and 2018 reporting periods, those who were not in the labor force accounted for 62 percent in both 2015 and 2018; those who were unemployed accounted for 20 percent in 2015 and 19 percent in 2018; and those who were employed accounted for 18 percent in 2015 and 19 percent in 2018.

Transitions in Living Arrangements

Individuals living in a private residence in 2015

Tables 7.5a–b and Figure 2. Among individuals served in both the 2015 and 2018 reporting periods, 92 percent of those who lived in a private residence in 2015 also reported living in a private residence in 2018; 3 percent transitioned to homelessness; 2 percent transitioned to residential care; and 3 percent, combined, transitioned to other living arrangements, institutional settings, or jail.

Figure 2. Living arrangements in 2018 of individuals aged 18 years and older who reported living in a private residence in 2015, by 2015 mental health diagnosis




NOTES: Percentages may not sum to 100 percent due to rounding. Less than 1 percent of individuals served in 2015 and 2018 moved into foster homes or foster care in 2018.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods.

Individuals who were homeless in 2015

Tables 7.5a–b and Figure 3. Among individuals aged 18 years and older served in both the 2015 and 2018 reporting periods, 43 percent who were homeless in 2015 also reported being homeless in 2018; 44 percent transitioned to a private residence, 6 percent to residential care, 4 percent to other living arrangements, 2 percent to institutional settings, and 2 percent to jail or correctional facilities.

Figure 3. Living arrangements in 2018 of individuals aged 18 years and older who reported being homeless in 2015, by 2015 mental health diagnosis




NOTES: Percentages may not sum to 100 percent due to rounding. Less than 1 percent of individuals served in 2015 and 2018 moved into foster homes or foster care in 2018.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods.

Individuals in jail or correctional facilities in 2015

Tables 7.5a–b and Figure 4. Among individuals aged 18 years and older served in both the 2015 and 2018 reporting periods, 60 percent of those who were in jail or correctional facilities in 2015 also reported being in jail or correctional facilities in 2018; 24 percent transitioned to private residences, 6 percent to homelessness, 5 percent to residential care, and 3 percent to institutional settings.

Figure 4. Living arrangements in 2018 of individuals aged 18 years and older who reported being in jail or a correctional facility in 2015, by 2015 mental health diagnosis



NOTES: Percentages may not sum to 100 percent due to rounding. Less than 1 percent of individuals served in 2015 and 2018 moved into foster homes or foster care in 2018.


SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods.

Transitions in Employment Status

Individuals who were employed in 2015

Tables 7.6a–b and Figure 5. Among individuals aged 18 years and older who were served in both the 2015 and 2018 reporting periods, 66 percent of those who were employed in 2015 also reported being employed in 2018; 13 percent transitioned to being unemployed, and 21 percent had left the labor force.

Figure 5. Employment status in 2018 of individuals aged 18 years and older who reported being employed in 2015, by 2015 mental health diagnosis




NOTES: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods.

Individuals who were unemployed in 2015

Tables 7.6a–b and Figure 6. Among individuals served in both the 2015 and 2018 reporting periods, 56 percent of those who were unemployed in 2015 also reported being unemployed in 2018; 30 percent of those unemployed in 2015 reported in 2018 that they were not in the labor force, and 13 percent reported that they were employed.

Figure 6. Employment status in 2018 of individuals aged 18 years and older who reported being unemployed in 2015, by 2015 mental health diagnosis




NOTES: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods.

Individuals who were not in the labor force in 2015

Tables 7.6a–b and Figure 7. Among individuals served in both the 2015 and 2018 reporting periods overall, 84 percent of those who were not in the labor force in 2015 also reported not being in the labor force in 2018; 9 percent transitioned to unemployment, and 7 percent transitioned to employment in 2018. The pattern of transition from not in the labor force to other employment statuses was similar among individuals served with bipolar and depressive disorders compared with individuals across all selected mental health diagnoses combined.

Figure 7. Employment status in 2018 of individuals aged 18 years and older who were not in the labor force in 2015, by 2015 mental health diagnosis



NOTES: Percentages may not sum to 100 percent due to rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods.

CHAPTER 8

GEOGRAPHIC AND CLINICAL TRENDS: 2013–2018

It is important to note that differences among reporting periods should be interpreted with great caution; variation may be more a function of the mix and number of reporting states than of actual differences. Throughout this chapter, states/jurisdictions may refer to any of the 50 states, the District of Columbia, the Commonwealth of Puerto Rico, and the combined jurisdictions of the Republic of Palau and the Commonwealth of the Northern Mariana Islands.

Table 8.1 presents the number of individuals served in the 2018 reporting period, by mental health diagnosis, Census region, Census division, and state or jurisdiction.

Tables 8.2 and 8.3 present, respectively, the number and adjusted number of individuals per 100,000 population served in the 2018 reporting period, by mental health diagnosis, Census region, Census division, and state or jurisdiction.¹⁴

Among individuals served in the 2018 reporting period, the primary mental health diagnoses with the largest per 100,000 population ratios (adjusted for age, gender, and ethnicity) were depressive disorders (443 per 100,000 population), trauma- and stressor-related disorders (246 per 100,000 population), and schizophrenia and other psychotic disorders (201 per 100,000 population) [Table 8.3].

Across all Census divisions in the 2018 reporting period, depressive disorders were the primary mental health diagnosis with the highest adjusted ratio per 100,000 population, ranging from 238 per 100,000 population for New England to 695 per 100,000 population for the West North Central division [Table 8.3].

In 35 of 45 states and jurisdictions for which age, gender, and race adjustment could be performed, depressive disorders were the primary diagnosis group with the highest adjusted ratio of individuals per 100,000 population. For Hawaii, Idaho, Massachusetts, Nevada, and New York, schizophrenia and other psychotic disorders had the highest adjusted ratio of individuals per 100,000 population. In New Mexico, Oregon, South Dakota, and Vermont, trauma- and stressor-related disorders had the highest adjusted ratio of individuals per 100,000 population. In Montana, the diagnosis with the highest adjusted ratio of individuals per 100,000 population was anxiety disorders [Table 8.3].

Tables 8.4–8.9 present—by Census region, Census division, and state or jurisdiction—the number of individuals served in the 2013 through 2018 reporting periods, grouped by the following primary diagnoses: trauma- and stressor-related disorders [Table 8.4], anxiety disorders [Table 8.5], attention

¹⁴ The adjusted number of individuals per 100,000 population served in the 2018 reporting period is calculated based on age, sex, and race of the mid-year 2010 population (according to vintage 2018 estimates). States that submitted insufficient data for inclusion in this report, as well as their reference populations, are excluded from these calculations. U.S. Census Bureau, Population Estimates Program, SC-EST2018-alldata6: Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2018. <https://www2.census.gov/programs-surveys/popest/tables/2010-2018/state/asrh/sc-est2018-alldata6.csv>

deficit/hyperactivity disorder [Table 8.6], bipolar disorders [Table 8.7], depressive disorders [Table 8.8], and schizophrenia and other psychotic disorders [Table 8.9].

Table 8.10 presents the number of individuals with co-occurring mental health and substance use disorders who were served in the 2013 through 2018 reporting periods, by Census region, Census division, and state or jurisdiction. It should be noted that the definition for co-occurring mental health and substance use disorders includes those who have a reported, identified substance use disorder but may not have a formal substance use diagnosis in addition to a reported mental health diagnosis. It should also be noted that data reported in this chapter exclude individuals whose only mental health diagnoses were substance use disorders.

CHAPTER 9
CLIENT CHARACTERISTICS BY STATE OR JURISDICTION:
2013–2018

Tables 9.1a–9.50b present, for each state or jurisdiction, the number and percent distribution of individuals served in the 2013 through 2018 reporting periods, according to mental health diagnosis, gender, age, race, ethnicity, and veteran status.

CHAPTER 10

CLIENT CHARACTERISTICS BY STATE OR JURISDICTION: 2018

Tables 10.1a–10.47b present, for each state or jurisdiction, the number and percent distribution of individuals served in the 2018 reporting period, according to mental health diagnosis, gender, age, race, ethnicity, and veteran status.

States that submitted no or insufficient data for the 2018 reporting period—Kansas, Maine, and New Jersey—do not appear in tables for this chapter.

TABLES

Table 1.1a. Individuals, by mental health diagnosis: Number, 2013–2018

<i>Mental health diagnosis¹</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total	5,576,659	5,818,881	5,311,897	5,813,339	5,934,383	6,213,791
Trauma- and stressor-related disorders	645,568	660,241	637,555	664,355	734,330	793,054
Anxiety disorders	376,408	409,035	427,333	493,641	579,070	654,028
Attention deficit/hyperactivity disorder+	442,351	420,745	351,786	406,362	429,499	461,444
Conduct disorder+	134,084	128,790	108,805	101,972	92,296	86,573
Delirium/dementia	25,278	26,010	20,292	15,298	11,421	11,463
Mood disorders	2,121,517	2,207,515	2,013,579	1,921,176	1,922,966	2,011,300
Bipolar disorders	854,076	880,161	758,549	663,415	607,293	601,023
Depressive disorders	1,267,441	1,327,354	1,255,030	1,257,761	1,315,673	1,410,277
Oppositional defiant disorder+	120,659	114,832	97,732	97,031	100,927	103,714
Pervasive developmental disorders+	58,322	67,896	44,635	69,443	64,777	65,578
Personality disorders	29,089	31,118	31,330	37,771	41,677	45,746
Schizophrenia and other psychotic disorders	630,500	656,286	593,824	576,500	626,384	660,521
Alcohol or substance-related disorders	161,155	189,290	115,824	112,563	124,679	144,759
Other mental disorders ²	204,065	213,013	206,063	284,349	380,782	455,832
No/deferred diagnosis ³	627,663	694,110	663,139	1,032,878	825,575	719,779

+ Indicates diagnoses typically identified in childhood (age < 18 years).

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.1b. Individuals, by mental health diagnosis: Column percent distribution, 2013–2018

<i>Mental health diagnosis¹</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total	100.0	100.0	100.0	100.0	100.0	100.0
Trauma- and stressor-related disorders	11.6	11.3	12.0	11.4	12.4	12.8
Anxiety disorders	6.7	7.0	8.0	8.5	9.8	10.5
Attention deficit/hyperactivity disorder+	7.9	7.2	6.6	7.0	7.2	7.4
Conduct disorder+	2.4	2.2	2.0	1.8	1.6	1.4
Delirium/dementia	†	†	†	†	†	†
Mood disorders	38.0	37.9	37.9	33.0	32.4	32.4
Bipolar disorders	15.3	15.1	14.3	11.4	10.2	9.7
Depressive disorders	22.7	22.8	23.6	21.6	22.2	22.7
Oppositional defiant disorder+	2.2	2.0	1.8	1.7	1.7	1.7
Pervasive developmental disorders+	1.0	1.2	†	1.2	1.1	1.1
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	11.3	11.3	11.2	9.9	10.6	10.6
Alcohol or substance-related disorders	2.9	3.3	2.2	1.9	2.1	2.3
Other mental disorders ²	3.7	3.7	3.9	4.9	6.4	7.3
No/deferred diagnosis ³	11.3	11.9	12.5	17.8	13.9	11.6

+ Indicates diagnoses typically identified in childhood (age < 18 years); † Less than 1 percent of individuals.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.2a. Individuals, by gender and age: Number, 2013–2018

<i>Gender and age</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	47	45	47	48	48
Total	5,576,659	5,818,881	5,311,897	5,813,339	5,934,383	6,213,791
Gender						
Male	2,713,018	2,828,069	2,570,572	2,799,851	2,873,064	3,021,655
Female	2,857,645	2,981,225	2,732,474	2,994,143	3,037,899	3,175,592
Total	5,570,663	5,809,294	5,303,046	5,793,994	5,910,963	6,197,247
Age¹						
0 to 11 years	812,119	849,907	731,351	811,404	812,159	892,484
12 to 14 years	403,736	417,244	376,750	401,746	401,160	438,967
15 to 17 years	434,873	439,431	411,136	444,858	443,765	469,162
18 to 20 years	268,477	259,939	234,919	260,296	268,132	288,195
21 to 24 years	326,256	336,104	304,520	333,579	345,253	357,257
25 to 29 years	441,976	463,364	432,045	487,419	527,755	557,653
30 to 34 years	446,072	471,411	438,256	483,106	511,309	534,107
35 to 39 years	383,178	406,379	388,512	435,995	467,430	494,186
40 to 44 years	407,261	415,026	369,047	387,645	389,997	400,992
45 to 49 years	435,990	442,877	390,827	415,729	413,334	414,109
50 to 54 years	444,153	469,914	428,470	452,852	438,901	430,035
55 to 59 years	337,373	368,375	350,060	385,959	389,871	402,710
60 to 64 years	203,503	225,348	220,758	249,173	255,186	273,170
65 years and older	228,205	250,048	231,583	259,691	263,860	256,693
Total	5,573,172	5,815,367	5,308,234	5,809,452	5,928,112	6,209,720

¹ Age is based on the individual's age at the midpoint of the reporting period.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.2b. Individuals, by gender and age: Column percent distribution, 2013–2018

<i>Gender and age</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	47	45	47	48	48
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	48.7	48.7	48.5	48.3	48.6	48.8
Female	51.3	51.3	51.5	51.7	51.4	51.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	14.6	14.6	13.8	14.0	13.7	14.4
12 to 14 years	7.2	7.2	7.1	6.9	6.8	7.1
15 to 17 years	7.8	7.6	7.7	7.7	7.5	7.6
18 to 20 years	4.8	4.5	4.4	4.5	4.5	4.6
21 to 24 years	5.9	5.8	5.7	5.7	5.8	5.8
25 to 29 years	7.9	8.0	8.1	8.4	8.9	9.0
30 to 34 years	8.0	8.1	8.3	8.3	8.6	8.6
35 to 39 years	6.9	7.0	7.3	7.5	7.9	8.0
40 to 44 years	7.3	7.1	7.0	6.7	6.6	6.5
45 to 49 years	7.8	7.6	7.4	7.2	7.0	6.7
50 to 54 years	8.0	8.1	8.1	7.8	7.4	6.9
55 to 59 years	6.1	6.3	6.6	6.6	6.6	6.5
60 to 64 years	3.7	3.9	4.2	4.3	4.3	4.4
65 years and older	4.1	4.3	4.4	4.5	4.5	4.1
Total	100.0	100.0	100.0	100.0	100.0	100.0

¹ Age is based on the individual's age at the midpoint of the reporting period.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.3a. Individuals, by race and ethnicity: Number, 2013–2018

<i>Race and ethnicity</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	47	45	47	48	48
Total	5,576,659	5,818,881	5,311,897	5,813,339	5,934,383	6,213,791
Race						
American Indian or Alaska Native	62,079	73,160	73,483	76,232	79,740	89,566
Asian ¹	60,040	69,140	67,106	78,385	82,820	91,274
Black or African American	1,051,934	1,106,235	993,314	1,070,717	1,090,821	1,138,522
Native Hawaiian or Other Pacific Islander	13,130	14,158	15,587	17,693	12,649	15,296
White	3,493,575	3,621,667	3,256,197	3,596,852	3,667,144	3,844,577
Other	467,957	488,936	440,646	491,417	537,599	579,001
Total	5,148,715	5,373,296	4,846,333	5,331,296	5,470,773	5,758,236
Ethnicity						
Not of Hispanic or Latino origin	3,891,620	3,995,971	3,593,763	3,912,972	4,464,374	4,603,454
Hispanic or Latino origin	703,437	750,208	725,197	800,551	856,460	893,469
Total	4,595,057	4,746,179	4,318,960	4,713,523	5,320,834	5,496,923
Specific Hispanic or Latino origin						
Mexican	32,588	45,719	54,002	42,618	52,074	49,932
Puerto Rican	29,990	49,184	51,790	44,068	51,139	27,716
Cuban	9,951	9,323	9,503	8,850	8,816	9,489
Other specific	45,395	56,850	51,609	51,118	59,058	59,964
Total	117,924	161,076	166,904	146,654	171,087	147,101

¹ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.3b. Individuals, by race and ethnicity: Column percent distribution, 2013–2018

<i>Race and ethnicity</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	47	45	47	48	48
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	1.2	1.4	1.5	1.4	1.5	1.6
Asian ¹	1.2	1.3	1.4	1.5	1.5	1.6
Black or African American	20.4	20.6	20.5	20.1	19.9	19.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	67.9	67.4	67.2	67.5	67.0	66.8
Other	9.1	9.1	9.1	9.2	9.8	10.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	84.7	84.2	83.2	83.0	83.9	83.7
Hispanic or Latino origin	15.3	15.8	16.8	17.0	16.1	16.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	27.6	28.4	32.4	29.1	30.4	33.9
Puerto Rican	25.4	30.5	31.0	30.0	29.9	18.8
Cuban	8.4	5.8	5.7	6.0	5.2	6.5
Other specific	38.5	35.3	30.9	34.9	34.5	40.8
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.4a. Individuals aged 17 years and younger, by mental health diagnosis: Number, 2013–2018

<i>Mental health diagnosis</i> ¹	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals aged 0 to 17 years	1,650,728	1,706,582	1,519,237	1,658,008	1,657,084	1,800,613
Trauma- and stressor-related disorders	324,173	329,697	300,043	325,701	343,048	386,335
Anxiety disorders	110,703	124,465	127,468	146,398	163,507	186,058
Attention deficit/hyperactivity disorder+	383,840	360,448	293,813	340,594	357,649	384,329
Conduct disorder	123,355	119,363	99,232	92,855	80,193	75,786
Mood disorders	312,644	331,301	298,143	262,988	233,013	250,271
Bipolar disorders	129,415	132,404	105,092	67,516	33,221	26,214
Depressive disorders	183,229	198,897	193,051	195,472	199,792	224,057
Oppositional defiant disorder	109,717	105,328	88,511	91,303	94,179	98,418
Pervasive developmental disorders	45,801	54,063	32,315	53,911	47,823	46,303
Alcohol or substance-related disorders	9,101	15,194	8,736	9,331	10,174	9,005
Other mental disorders ²	87,924	99,499	90,696	131,433	152,557	193,773
No/deferred diagnosis ³	143,470	167,224	180,280	203,494	174,941	170,335

+ Indicates diagnoses typically identified in childhood (age < 18 years).

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.4b. Individuals aged 17 years and younger, by mental health diagnosis: Column percent distribution, 2013–2018

<i>Mental health diagnosis</i> ¹	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals aged 0 to 17 years	100.0	100.0	100.0	100.0	100.0	100.0
Trauma- and stressor-related disorders	19.6	19.3	19.7	19.6	20.7	21.5
Anxiety disorders	6.7	7.3	8.4	8.8	9.9	10.3
Attention deficit/hyperactivity disorder+	23.3	21.1	19.3	20.5	21.6	21.3
Conduct disorder	7.5	7.0	6.5	5.6	4.8	4.2
Mood disorders	18.9	19.4	19.6	15.9	14.1	13.9
Bipolar disorders	7.8	7.8	6.9	4.1	2.0	1.5
Depressive disorders	11.1	11.7	12.7	11.8	12.1	12.4
Oppositional defiant disorder	6.6	6.2	5.8	5.5	5.7	5.5
Pervasive developmental disorders	2.8	3.2	2.1	3.3	2.9	2.6
Alcohol or substance-related disorders	†	†	†	†	†	†
Other mental disorders ²	5.3	5.8	6.0	7.9	9.2	10.8
No/deferred diagnosis ³	8.7	9.8	11.9	12.3	10.6	9.5

+ Indicates diagnoses typically identified in childhood (age < 18 years); † Less than 1 percent of individuals.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.5a. Individuals aged 18 to 24 years, by mental health diagnosis: Number, 2013–2018

<i>Mental health diagnosis</i> ¹	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals aged 18 to 24 years	594,733	596,043	539,439	593,875	613,385	645,452
Trauma- and stressor-related disorders	62,731	59,530	58,196	59,427	66,457	70,164
Anxiety disorders	44,296	46,445	49,330	58,684	69,054	80,312
Attention deficit/hyperactivity disorder+	30,898	28,485	26,193	28,535	30,203	31,550
Mood disorders	245,158	245,601	221,704	211,884	210,810	226,862
Bipolar disorders	110,862	109,516	91,184	74,882	62,950	61,603
Depressive disorders	134,296	136,085	130,520	137,002	147,860	165,259
Personality disorders	3,666	3,797	3,761	4,713	5,681	6,454
Schizophrenia and other psychotic disorders	50,060	52,510	47,444	46,390	49,489	50,990
Alcohol or substance-related disorders	24,858	27,942	14,908	16,317	17,126	18,475
Other mental disorders ²	49,718	45,092	42,486	52,010	66,807	72,671
No/deferred diagnosis ³	83,348	86,641	75,417	115,915	97,758	87,974

+ Indicates diagnoses typically identified in childhood (age < 18 years).

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.5b. Individuals aged 18 to 24 years, by mental health diagnosis: Column percent distribution, 2013–2018

<i>Mental health diagnosis</i> ¹	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals aged 18 to 24 years	100.0	100.0	100.0	100.0	100.0	100.0
Trauma- and stressor-related disorders	10.5	10.0	10.8	10.0	10.8	10.9
Anxiety disorders	7.4	7.8	9.1	9.9	11.3	12.4
Attention deficit/hyperactivity disorder+	5.2	4.8	4.9	4.8	4.9	4.9
Mood disorders	41.2	41.2	41.1	35.7	34.4	35.1
Bipolar disorders	18.6	18.4	16.9	12.6	10.3	9.5
Depressive disorders	22.6	22.8	24.2	23.1	24.1	25.6
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	8.4	8.8	8.8	7.8	8.1	7.9
Alcohol or substance-related disorders	4.2	4.7	2.8	2.7	2.8	2.9
Other mental disorders ²	8.4	7.6	7.9	8.8	10.9	11.3
No/deferred diagnosis ³	14.0	14.5	14.0	19.5	15.9	13.6

+ Indicates diagnoses typically identified in childhood (age < 18 years); † Less than 1 percent of individuals.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.6a. Individuals aged 25 to 54 years, by mental health diagnosis: Number, 2013–2018

<i>Mental health diagnosis</i> ¹	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals aged 25 to 54 years	2,558,630	2,668,971	2,447,157	2,662,746	2,748,726	2,831,082
Trauma- and stressor-related disorders	213,648	222,216	228,296	232,527	263,676	276,239
Anxiety disorders	179,033	191,692	201,280	233,557	277,618	311,401
Attention deficit/hyperactivity disorder+	25,230	29,104	29,493	34,105	37,511	41,662
Mood disorders	1,210,553	1,251,869	1,127,292	1,090,202	1,098,760	1,127,106
Bipolar disorders	507,354	523,618	453,661	416,657	401,908	398,050
Depressive disorders	703,199	728,251	673,631	673,545	696,852	729,056
Personality disorders	19,753	21,088	21,286	25,254	27,942	30,448
Schizophrenia and other psychotic disorders	401,111	412,619	369,050	359,050	386,146	405,426
Alcohol or substance-related disorders	110,461	126,431	75,816	75,497	83,268	100,093
Other mental disorders ²	95,670	95,623	94,559	116,745	162,025	184,532
No/deferred diagnosis ³	303,171	318,329	300,085	495,809	411,780	354,175

+ Indicates diagnoses typically identified in childhood (age < 18 years).

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.6b. Individuals aged 25 to 54 years, by mental health diagnosis: Column percent distribution, 2013–2018

<i>Mental health diagnosis</i> ¹	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals aged 25 to 54 years	100.0	100.0	100.0	100.0	100.0	100.0
Trauma- and stressor-related disorders	8.4	8.3	9.3	8.7	9.6	9.8
Anxiety disorders	7.0	7.2	8.2	8.8	10.1	11.0
Attention deficit/hyperactivity disorder+	†	1.1	1.2	1.3	1.4	1.5
Mood disorders	47.3	46.9	46.1	40.9	40.0	39.8
Bipolar disorders	19.8	19.6	18.5	15.6	14.6	14.1
Depressive disorders	27.5	27.3	27.5	25.3	25.4	25.8
Personality disorders	†	†	†	†	1.0	1.1
Schizophrenia and other psychotic disorders	15.7	15.5	15.1	13.5	14.0	14.3
Alcohol or substance-related disorders	4.3	4.7	3.1	2.8	3.0	3.5
Other mental disorders ²	3.7	3.6	3.9	4.4	5.9	6.5
No/deferred diagnosis ³	11.8	11.9	12.3	18.6	15.0	12.5

+ Indicates diagnoses typically identified in childhood (age < 18 years); † Less than 1 percent of individuals.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.7a. Individuals aged 55 years and older, by mental health diagnosis: Number, 2013–2018

<i>Mental health diagnosis</i> ¹	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals aged 55 years and older	769,081	843,771	802,401	894,823	908,917	932,573
Trauma- and stressor-related disorders	44,797	48,596	50,789	46,360	60,431	59,957
Anxiety disorders	42,248	46,186	48,972	54,486	68,329	76,046
Delirium/dementia	18,812	18,841	14,385	10,283	7,156	6,966
Mood disorders	352,574	378,230	365,877	355,259	379,184	406,375
Bipolar disorders	106,239	114,407	108,361	104,009	108,904	115,042
Depressive disorders	246,335	263,823	257,516	251,250	270,280	291,333
Personality disorders	4,536	5,031	5,081	5,873	6,632	7,301
Schizophrenia and other psychotic disorders	168,732	180,237	168,101	162,135	181,172	194,943
Alcohol or substance-related disorders	16,430	19,195	15,829	11,033	13,554	17,176
Other mental disorders ²	24,264	26,404	27,037	32,457	52,904	58,296
No/deferred diagnosis ³	96,688	121,051	106,330	216,937	139,555	105,513

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.7b. Individuals aged 55 years and older, by mental health diagnosis: Column percent distribution, 2013–2018

<i>Mental health diagnosis¹</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	47	45	47	48	48
Total individuals aged 55 years and older	100.0	100.0	100.0	100.0	100.0	100.0
Trauma- and stressor-related disorders	5.8	5.8	6.3	5.2	6.6	6.4
Anxiety disorders	5.5	5.5	6.1	6.1	7.5	8.2
Delirium/dementia	2.4	2.2	1.8	1.1	†	†
Mood disorders	45.8	44.8	45.6	39.7	41.7	43.6
Bipolar disorders	13.8	13.6	13.5	11.6	12.0	12.3
Depressive disorders	32.0	31.3	32.1	28.1	29.7	31.2
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	21.9	21.4	20.9	18.1	19.9	20.9
Alcohol or substance-related disorders	2.1	2.3	2.0	1.2	1.5	1.8
Other mental disorders ²	3.2	3.1	3.4	3.6	5.8	6.3
No/deferred diagnosis ³	12.6	14.3	13.3	24.2	15.4	11.3

† Less than 1 percent of individuals.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ No/deferred diagnosis includes unknown, not collected, missing, invalid, or other non-mental health diagnoses.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.8a. Individuals, by marital status and living arrangement: Number, 2013–2018

<i>Marital status and living arrangement</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	47	45	47	48	48
Total	5,576,659	5,818,881	5,311,897	5,813,339	5,934,383	6,213,791
Marital status						
Total individuals aged 18 years and older	3,922,444	4,108,785	3,788,997	4,151,444	4,271,028	4,409,107
Never married	1,151,441	1,309,306	1,085,021	1,308,840	1,401,547	1,460,309
Currently married	360,996	382,420	343,635	365,641	355,269	360,589
Separated	139,869	164,834	134,090	153,837	147,973	146,589
Divorced	343,970	377,560	332,624	356,402	379,029	368,856
Widowed	65,792	71,529	62,120	69,912	68,006	77,430
Total	2,062,068	2,305,649	1,957,490	2,254,632	2,351,824	2,413,773
Living arrangement						
Homeless	116,735	116,751	120,379	121,442	139,704	158,941
Foster home/foster care	57,624	60,545	49,442	42,927	50,753	51,597
Residential care	115,787	124,623	121,488	114,333	132,081	131,152
Institutional setting	49,214	57,994	54,821	50,157	52,450	58,126
Jail/correctional facility	69,640	66,226	62,898	59,301	64,401	67,992
Private residence	2,845,956	2,768,395	2,783,266	2,710,042	2,846,909	3,024,152
Other	111,065	153,824	126,319	145,677	147,854	89,414
Total	3,366,021	3,348,358	3,318,613	3,243,879	3,434,152	3,581,374

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.8b. Individuals, by marital status and living arrangement: Column percent distribution, 2013–2018

<i>Marital status and living arrangement</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	47	45	47	48	48
Total	100.0	100.0	100.0	100.0	100.0	100.0
Marital status						
Total individuals aged 18 years and older	100.0	100.0	100.0	100.0	100.0	100.0
Never married	55.8	56.8	55.4	58.1	59.6	60.5
Currently married	17.5	16.6	17.6	16.2	15.1	14.9
Separated	6.8	7.1	6.9	6.8	6.3	6.1
Divorced	16.7	16.4	17.0	15.8	16.1	15.3
Widowed	3.2	3.1	3.2	3.1	2.9	3.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement						
Homeless	3.5	3.5	3.6	3.7	4.1	4.4
Foster home/foster care	1.7	1.8	1.5	1.3	1.5	1.4
Residential care	3.4	3.7	3.7	3.5	3.8	3.7
Institutional setting	1.5	1.7	1.7	1.5	1.5	1.6
Jail/correctional facility	2.1	2.0	1.9	1.8	1.9	1.9
Private residence	84.5	82.7	83.9	83.5	82.9	84.4
Other	3.3	4.6	3.8	4.5	4.3	2.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.9a. Individuals, by SMI/SED status and co-occurring mental health and substance use disorder status: Number, 2013–2018

<i>SMI/SED status and co-occurring mental health and substance use disorder status</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	47	45	47	48	48
Total	5,576,659	5,818,881	5,311,897	5,813,339	5,934,383	6,213,791
SMI status (individuals aged 18 years and older)¹						
SMI	2,345,633	2,482,833	2,297,480	2,392,154	2,617,478	2,884,305
Not SMI	899,626	942,735	862,460	792,535	826,034	812,698
Total	3,245,259	3,425,568	3,159,940	3,184,689	3,443,512	3,697,003
SED status (individuals aged 17 years and younger)¹						
SED	993,409	1,121,791	951,957	990,041	1,044,419	1,174,041
Not SED	421,613	358,730	346,904	406,609	389,453	407,630
Total	1,415,022	1,480,521	1,298,861	1,396,650	1,433,872	1,581,671
Co-occurring mental health and substance use disorders (individuals aged 12 years and older)²						
Yes	918,368	929,294	847,413	865,802	1,117,677	1,260,328
No	3,514,423	3,411,905	3,166,793	3,395,252	3,282,243	3,441,194
Total³	4,432,791	4,341,199	4,014,206	4,261,054	4,399,920	4,701,522

SED = serious emotional disturbance; SMI = serious mental illness.

¹ The federal definitions for SED and SMI status can be found in Appendix D.

² Co-occurring mental health and substance use disorders are only presented for individuals aged 12 years and older.

³ Total number of individuals aged 12 years and older for whom co-occurring mental health and substance use disorders were reported.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 1.9b. Individuals, by SMI/SED status and co-occurring mental health and substance use disorder status:
Column percent distribution, 2013–2018**

<i>SMI/SED status and co-occurring mental health and substance use disorder status</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	47	45	47	48	48
Total	100.0	100.0	100.0	100.0	100.0	100.0
SMI status (individuals aged 18 years and older)¹						
SMI	72.3	72.5	72.7	75.1	76.0	78.0
Not SMI	27.7	27.5	27.3	24.9	24.0	22.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
SED status (individuals aged 17 years and younger)¹						
SED	70.2	75.8	73.3	70.9	72.8	74.2
Not SED	29.8	24.2	26.7	29.1	27.2	25.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Co-occurring mental health and substance use disorders (individuals aged 12 years and older)²						
Yes	20.7	21.4	21.1	20.3	25.4	26.8
No	79.3	78.6	78.9	79.7	74.6	73.2
Total³	100.0	100.0	100.0	100.0	100.0	100.0

SED = serious emotional disturbance; SMI = serious mental illness.

¹ The federal definitions for SED and SMI status can be found in Appendix D.

² Co-occurring mental health and substance use disorders are only presented for individuals aged 12 years and older.

³ Total percentage of individuals aged 12 years and older for whom co-occurring mental health and substance use disorders were reported.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.10a. Individuals aged 18 years and older, by employment status, ‘detailed not in labor force,’ education, and veteran status: Number, 2013–2018

<i>Employment status, ‘detailed not in labor force,’ education, and veteran status</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals aged 18 years and older	3,922,444	4,108,785	3,788,997	4,151,444	4,271,028	4,409,107
Employment status						
Employed	334,469	411,923	422,812	422,244	482,087	530,803
Full time	146,315	205,018	211,222	216,618	242,892	268,250
Part time	126,285	136,512	145,495	140,807	168,149	180,041
Unemployed	601,533	593,550	582,538	515,856	649,126	643,175
Not in labor force (see detail below)	1,025,423	1,100,959	1,041,928	984,140	1,069,552	1,102,235
Total	1,961,425	2,106,432	2,047,278	1,922,240	2,200,765	2,276,213
Detailed not in labor force						
Total individuals aged 18 years and older not in labor force	1,025,423	1,100,959	1,041,928	984,140	1,069,552	1,102,235
Disabled	382,004	438,537	442,592	445,055	505,381	537,604
Retired	29,624	34,320	38,514	40,119	47,770	50,054
Student	105,604	95,331	80,558	76,447	86,261	87,836
Homemaker	44,771	47,810	48,334	48,574	51,315	44,872
Sheltered/non-competitive employment	8,564	13,826	11,760	9,833	13,690	13,629
Other	454,856	471,135	420,170	364,112	365,135	368,240
Highest school grade completed						
0 to 8	201,729	215,153	222,981	227,302	197,102	187,072
9 to 11	317,008	325,958	329,782	332,483	337,189	349,251
12 (or GED)	680,587	774,709	803,771	839,793	851,624	961,697
More than 12	278,891	326,127	342,885	367,417	432,449	479,701
Total	1,478,215	1,641,947	1,699,419	1,766,995	1,818,364	1,977,721
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	50,439	46,860
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	1,486,721	1,519,278
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	1,537,160	1,566,138

n/c Not collected.

GED = general equivalency diploma.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.10b. Individuals aged 18 years and older, by employment status, ‘detailed not in labor force,’ education, and veteran status: Column percent distribution, 2013–2018

<i>Employment status, ‘detailed not in labor force,’ education, and veteran status</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals aged 18 years and older	100.0	100.0	100.0	100.0	100.0	100.0
Employment status						
Employed	17.1	19.6	20.7	22.0	21.9	23.3
Full time	7.5	9.7	10.3	11.3	11.0	11.8
Part time	6.4	6.5	7.1	7.3	7.6	7.9
Unemployed	30.7	28.2	28.5	26.8	29.5	28.3
Not in labor force (see detail below)	52.3	52.3	50.9	51.2	48.6	48.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force						
Total individuals aged 18 years and older not in labor force	100.0	100.0	100.0	100.0	100.0	100.0
Disabled	37.3	39.8	42.5	45.2	47.3	48.8
Retired	2.9	3.1	3.7	4.1	4.5	4.5
Student	10.3	8.7	7.7	7.8	8.1	8.0
Homemaker	4.4	4.3	4.6	4.9	4.8	4.1
Sheltered/non-competitive employment	†	1.3	1.1	†	1.3	1.2
Other	44.4	42.8	40.3	37.0	34.1	33.4
Highest school grade completed						
0 to 8	13.6	13.1	13.1	12.9	10.8	9.5
9 to 11	21.4	19.9	19.4	18.8	18.5	17.7
12 (or GED)	46.0	47.2	47.3	47.5	46.8	48.6
More than 12	18.9	19.9	20.2	20.8	23.8	24.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	3.3	3.0
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	96.7	97.0
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

GED = general equivalency diploma.

NOTES: Percentages may not sum to 100 percent because of rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.11a. Individuals, by timing of admission and mental health service setting: Number, 2013–2018

<i>Timing of admission and mental health service setting</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Timing of admission						
New individuals	2,686,660	2,677,733	2,592,375	3,143,256	3,177,990	3,329,865
Continuing individuals	2,889,999	3,141,148	2,719,522	2,670,083	2,756,393	2,883,926
Total	5,576,659	5,818,881	5,311,897	5,813,339	5,934,383	6,213,791
Mental health service setting^{1,2}						
State psychiatric hospital	118,761	129,391	119,336	126,139	120,797	119,126
Community-based program	5,474,015	5,695,871	5,183,290	5,665,264	5,788,778	6,081,713
Residential treatment center	49,109	47,615	43,202	56,415	60,154	60,657
Other psychiatric inpatient	148,961	157,679	177,132	234,056	233,774	248,065
Institutions under the justice system	48,724	67,231	70,512	69,039	70,735	69,113
Total	5,576,659	5,818,881	5,311,897	5,813,338	5,934,383	6,213,791

¹ Numbers may not sum to total, because individuals could receive services in more than one service setting during the reporting period.

² Mental health service setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.11b. Individuals, by timing of admission and mental health service setting: Column percent, 2013–2018

<i>Timing of admission and mental health service setting</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Timing of admission						
New individuals	48.2	46.0	48.8	54.1	53.6	53.6
Continuing individuals	51.8	54.0	51.2	45.9	46.4	46.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Mental health service setting^{1,2}						
State psychiatric hospital	2.1	2.2	2.2	2.2	2.0	1.9
Community-based program	98.2	97.9	97.6	97.5	97.5	97.9
Residential treatment center	†	†	†	†	1.0	†
Other psychiatric inpatient	2.7	2.7	3.3	4.0	3.9	4.0
Institutions under the justice system	†	1.2	1.3	1.2	1.2	1.1
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

² Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages for timing of admission may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.12a. Individuals, by number of mental health diagnoses and any mention of mental health diagnosis: Number, 2013–2018

<i>Number of diagnoses and any mention of mental health diagnosis</i>		<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting		45	47	45	47	48	48
	Total	5,576,659	5,818,881	5,311,897	5,813,339	5,934,383	6,213,791
Number of mental health diagnoses							
None		627,663	694,110	663,139	1,032,878	825,575	719,779
1		2,823,691	2,933,818	2,523,146	3,017,455	3,441,870	3,535,468
2		1,567,462	1,578,192	1,501,638	1,227,814	1,207,297	1,373,094
3		557,843	612,761	623,974	535,192	459,641	585,450
Any mention of Mental health diagnosis^{1,2}							
Trauma- and stressor-related disorders		853,416	904,066	888,806	915,822	1,024,429	1,145,889
Anxiety disorders		680,938	761,164	797,941	888,831	1,085,457	1,266,831
Attention deficit/hyperactivity disorder		609,826	611,789	534,056	584,682	625,134	693,471
Conduct disorder		177,055	175,986	152,736	142,337	134,277	131,077
Mood disorders		2,374,017	2,485,295	2,301,620	2,206,154	2,243,965	2,394,353
Bipolar disorders		971,605	998,673	874,681	759,967	704,533	713,319
Depressive disorders		1,452,613	1,546,242	1,487,935	1,494,381	1,584,468	1,734,691
Oppositional defiant disorder		200,431	193,031	168,355	170,161	184,070	196,012
Pervasive developmental disorders		83,515	94,564	71,302	96,830	96,651	105,479
Schizophrenia and other psychotic disorders		676,837	707,491	644,141	625,605	692,868	735,029
Personality disorders		178,540	186,681	170,276	147,202	159,207	168,380
Alcohol or substance-related disorders		326,369	350,750	270,757	238,747	214,613	239,033
Delirium/dementia		38,514	42,024	35,059	25,569	18,412	20,442
Other mental disorders ³		1,288,282	1,251,922	1,201,118	866,039	609,623	749,130
No/deferred diagnosis		627,663	694,110	663,139	1,032,878	825,575	719,779

¹ Numbers may not sum to the total, because up to three diagnoses may be reported for an individual.

² Up to three diagnoses can be reported for each individual; “any mention” includes all diagnoses reported. For this reason, the sum of any mention of bipolar and depressive disorders may be larger than any mention of mood disorders. Mental health diagnostic category definitions can be found in Appendix E.

³ “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.12b. Individuals, by number of mental health diagnoses and any mention of mental health diagnosis: Column percent, 2013–2018

<i>Number of diagnoses and any mention of mental health diagnosis</i>		2013	2014	2015	2016	2017	2018
Number of states reporting		45	47	45	47	48	48
	Total	100.0	100.0	100.0	100.0	100.0	100.0
Number of mental health diagnoses							
None		11.3	11.9	12.5	17.8	13.9	11.6
1		50.6	50.4	47.5	51.9	58.0	56.9
2		28.1	27.1	28.3	21.1	20.3	22.1
3		10.0	10.5	11.7	9.2	7.7	9.4
Any mention of Mental health diagnosis^{1,2}							
Trauma- and stressor-related disorders		15.3	15.5	16.7	15.8	17.3	18.4
Anxiety disorders		12.2	13.1	15.0	15.3	18.3	20.4
Attention deficit/hyperactivity disorder		10.9	10.5	10.1	10.1	10.5	11.2
Conduct disorder		3.2	3.0	2.9	2.4	2.3	2.1
Mood disorders		42.6	42.7	43.3	37.9	37.8	38.5
Bipolar disorders		17.4	17.2	16.5	13.1	11.9	11.5
Depressive disorders		26.0	26.6	28.0	25.7	26.7	27.9
Oppositional defiant disorder		3.6	3.3	3.2	2.9	3.1	3.2
Pervasive developmental disorders		1.5	1.6	1.3	1.7	1.6	1.7
Schizophrenia and other psychotic disorders		12.1	12.2	12.1	10.8	11.7	11.8
Personality disorders		3.2	3.2	3.2	2.5	2.7	2.7
Alcohol or substance-related disorders		5.9	6.0	5.1	4.1	3.6	3.8
Delirium/dementia		†	†	†	†	†	†
Other mental disorders ³		23.1	21.5	22.6	14.9	10.3	12.1
No/deferred diagnosis		11.3	11.9	12.5	17.8	13.9	11.6

† Less than 1 percent of individuals.

¹ Percentages may not sum to 100 percent, because up to three diagnoses may be reported for an individual.

² Up to three diagnoses can be reported for each individual; “any mention” includes all diagnoses reported. For this reason, the sum of any mention of bipolar and depressive disorders may be larger than any mention of mood disorders. Mental health diagnostic category definitions can be found in Appendix E.

³ “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.13a. Individuals, by Census region and division: Number, 2013–2018

<i>Census region and Census division</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	47	45	47	48	48
Total	5,576,453	5,797,312	5,288,528	5,795,396	5,908,465	6,210,481
Region¹						
Northeast	1,270,672	1,173,111	560,450	1,115,347	871,878	801,388
Midwest	1,005,850	1,267,305	1,220,167	1,223,143	1,484,457	1,654,571
South	1,786,971	1,814,022	1,882,068	1,831,834	1,854,663	1,928,812
West	1,512,960	1,542,874	1,625,843	1,625,072	1,697,467	1,825,710
Division¹						
New England	246,846	179,096	180,824	191,369	240,399	178,047
Middle Atlantic	1,023,826	994,015	379,626	923,978	631,479	623,341
East North Central	687,866	742,524	752,154	721,586	982,915	1,045,726
West North Central	317,984	524,781	468,013	501,557	501,542	608,845
South Atlantic	835,400	847,031	878,731	800,461	812,765	849,366
East South Central	441,818	449,787	467,125	461,375	441,287	463,160
West South Central	509,753	517,204	536,212	569,998	600,611	616,286
Mountain	520,428	520,164	565,460	592,591	650,505	696,587
Pacific	992,532	1,022,710	1,060,383	1,032,481	1,046,962	1,129,123

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.13b. Individuals, by Census region and division: Column percent distribution, 2013–2018

<i>Census region and Census division</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total	100.0	100.0	100.0	100.0	100.0	100.0
Region¹						
Northeast	22.8	20.2	10.6	19.2	14.8	12.9
Midwest	18.0	21.9	23.1	21.1	25.1	26.6
South	32.0	31.3	35.6	31.6	31.4	31.1
West	27.1	26.6	30.7	28.0	28.7	29.4
Total						
Division¹	4.4	3.1	3.4	3.3	4.1	2.9
Middle Atlantic	18.4	17.1	7.2	15.9	10.7	10.0
East North Central	12.3	12.8	14.2	12.5	16.6	16.8
West North Central	5.7	9.1	8.8	8.7	8.5	9.8
South Atlantic	15.0	14.6	16.6	13.8	13.8	13.7
East South Central	7.9	7.8	8.8	8.0	7.5	7.5
West South Central	9.1	8.9	10.1	9.8	10.2	9.9
Mountain	9.3	9.0	10.7	10.2	11.0	11.2
Pacific	17.8	17.6	20.1	17.8	17.7	18.2

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.14a. Individuals with primary diagnosis of trauma- and stressor-related disorders, by gender, age, race, and ethnicity: Number, 2013–2018

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals with primary diagnosis of trauma- and stressor-related disorders	645,568	660,241	637,555	664,355	734,330	793,054
Gender						
Male	268,543	275,355	266,187	278,916	312,326	340,879
Female	376,237	383,945	370,248	384,020	419,667	450,131
Total	644,780	659,300	636,435	662,936	731,993	791,010
Age¹						
0 to 11 years	186,479	193,741	173,421	190,198	200,730	227,728
12 to 14 years	73,090	73,966	67,643	71,585	75,072	85,042
15 to 17 years	64,604	61,990	58,979	63,918	67,246	73,565
18 to 20 years	30,443	27,762	26,897	28,011	30,967	33,991
21 to 24 years	32,288	31,768	31,299	31,416	35,490	36,173
25 to 29 years	42,549	43,890	45,407	47,363	55,270	57,846
30 to 34 years	42,515	44,281	45,353	47,131	53,727	56,434
35 to 39 years	34,711	37,027	39,535	41,303	47,726	51,277
40 to 44 years	33,698	34,252	34,153	33,811	37,564	39,710
45 to 49 years	31,812	32,556	32,623	32,449	35,863	36,910
50 to 54 years	28,363	30,210	31,225	30,470	33,526	34,062
55 to 59 years	19,797	21,340	22,793	22,818	26,667	27,990
60 to 64 years	11,523	12,475	13,435	12,946	15,697	17,042
65 years and older	13,477	14,781	14,561	10,596	18,067	14,925
Total	645,349	660,039	637,324	664,015	733,612	792,695

Continued. See notes at end of table.

Table 1.14a. Individuals with primary diagnosis of trauma- and stressor-related disorders, by gender, age, race, and ethnicity: Number, 2013–2018 (continued)

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	10,474	12,662	13,033	14,399	15,095	15,951
Asian ²	7,080	8,008	7,794	9,398	9,811	10,242
Black or African American	101,401	107,328	100,392	99,418	118,467	126,593
Native Hawaiian or Other Pacific Islander	1,559	1,629	1,828	2,075	2,013	2,668
White	401,023	409,678	392,734	418,045	453,204	482,619
Other	64,007	63,235	58,711	62,225	73,156	82,317
Total	585,544	602,540	574,492	605,560	671,746	720,390
Ethnicity						
Not of Hispanic or Latino origin	419,387	422,278	418,316	451,153	552,535	573,300
Hispanic or Latino origin	98,823	102,236	99,768	111,576	119,793	124,482
Total	518,210	524,514	518,084	562,729	672,328	697,782
Specific Hispanic or Latino origin						
Mexican	5,790	9,036	9,534	7,292	12,405	11,814
Puerto Rican	3,822	3,656	3,580	2,989	3,382	2,998
Cuban	623	613	692	692	903	789
Other specific	5,840	8,901	7,466	7,081	11,125	10,658
Total	16,075	22,206	21,272	18,054	27,815	26,259

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.14b. Individuals with primary diagnosis of trauma- and stressor-related disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018

<i>Gender, age, race, and ethnicity</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	47	45	47	48	48
Total individuals with primary diagnosis of trauma- and stressor-related disorders	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	41.6	41.8	41.8	42.1	42.7	43.1
Female	58.4	58.2	58.2	57.9	57.3	56.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	28.9	29.4	27.2	28.6	27.4	28.7
12 to 14 years	11.3	11.2	10.6	10.8	10.2	10.7
15 to 17 years	10.0	9.4	9.3	9.6	9.2	9.3
18 to 20 years	4.7	4.2	4.2	4.2	4.2	4.3
21 to 24 years	5.0	4.8	4.9	4.7	4.8	4.6
25 to 29 years	6.6	6.6	7.1	7.1	7.5	7.3
30 to 34 years	6.6	6.7	7.1	7.1	7.3	7.1
35 to 39 years	5.4	5.6	6.2	6.2	6.5	6.5
40 to 44 years	5.2	5.2	5.4	5.1	5.1	5.0
45 to 49 years	4.9	4.9	5.1	4.9	4.9	4.7
50 to 54 years	4.4	4.6	4.9	4.6	4.6	4.3
55 to 59 years	3.1	3.2	3.6	3.4	3.6	3.5
60 to 64 years	1.8	1.9	2.1	1.9	2.1	2.1
65 years and older	2.1	2.2	2.3	1.6	2.5	1.9
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 1.14b. Individuals with primary diagnosis of trauma- and stressor-related disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	1.8	2.1	2.3	2.4	2.2	2.2
Asian ²	1.2	1.3	1.4	1.6	1.5	1.4
Black or African American	17.3	17.8	17.5	16.4	17.6	17.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	68.5	68.0	68.4	69.0	67.5	67.0
Other	10.9	10.5	10.2	10.3	10.9	11.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	80.9	80.5	80.7	80.2	82.2	82.2
Hispanic or Latino origin	19.1	19.5	19.3	19.8	17.8	17.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	36.0	40.7	44.8	40.4	44.6	45.0
Puerto Rican	23.8	16.5	16.8	16.6	12.2	11.4
Cuban	3.9	2.8	3.3	3.8	3.2	3.0
Other specific	36.3	40.1	35.1	39.2	40.0	40.6
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Percentages may not sum to 100 percent because of rounding. This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.15a. Individuals with primary diagnosis of anxiety disorders, by gender, age, race, and ethnicity: Number, 2013–2018

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	46	48	48
Total individuals with primary diagnosis of anxiety disorders	376,408	409,035	427,333	493,641	579,070	654,028
Gender						
Male	152,340	167,615	176,308	196,875	229,128	254,741
Female	223,852	241,113	250,591	288,233	348,550	398,151
Total	376,192	408,728	426,899	485,108	577,678	652,892
Age¹						
0 to 11 years	57,145	63,171	62,006	68,419	75,555	84,127
12 to 14 years	26,061	29,832	31,162	35,999	40,597	47,217
15 to 17 years	27,497	31,462	34,300	41,980	47,355	54,714
18 to 20 years	19,493	20,374	21,491	26,066	30,565	36,802
21 to 24 years	24,803	26,071	27,839	32,618	38,489	43,510
25 to 29 years	37,044	39,599	42,377	50,530	61,334	69,649
30 to 34 years	36,488	39,411	41,937	48,799	58,661	66,030
35 to 39 years	28,830	31,401	33,796	40,663	49,594	56,685
40 to 44 years	26,961	28,309	28,879	32,459	37,479	42,253
45 to 49 years	25,699	26,741	27,159	30,737	35,662	39,444
50 to 54 years	24,011	26,231	27,132	30,369	34,888	37,340
55 to 59 years	17,957	19,765	21,154	24,560	29,207	33,172
60 to 64 years	10,978	12,327	13,393	15,740	18,943	22,334
65 years and older	13,313	14,094	14,425	14,186	20,179	20,540
Total	376,280	408,788	427,050	493,125	578,508	653,817

Continued. See notes at end of table.

Table 1.15a. Individuals with primary diagnosis of anxiety disorders, by gender, age, race, and ethnicity: Number, 2013–2018 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	4,524	5,780	6,651	7,541	9,665	11,316
Asian ²	3,013	4,035	4,637	6,119	7,664	8,336
Black or African American	36,279	41,039	42,759	46,901	57,810	64,167
Native Hawaiian or Other Pacific Islander	839	986	1,243	1,677	1,392	1,959
White	272,565	290,326	298,679	344,367	408,660	455,853
Other	31,897	37,640	38,046	43,890	48,537	58,146
Total	349,117	379,806	392,015	450,495	533,728	599,777
Ethnicity						
Not of Hispanic or Latino origin	265,784	281,602	298,441	348,928	452,986	474,463
Hispanic or Latino origin	48,995	56,346	61,375	72,561	80,841	90,629
Total	314,779	337,948	359,816	421,489	533,827	565,092
Specific Hispanic or Latino origin						
Mexican	2,619	3,862	5,033	3,872	6,018	5,639
Puerto Rican	1,790	1,648	1,736	1,786	1,897	1,990
Cuban	593	638	720	762	745	712
Other specific	3,107	4,396	4,278	4,399	5,656	5,836
Total	8,109	10,544	11,767	10,819	14,316	14,177

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.15b. Individuals with primary diagnosis of anxiety disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	46	48	48
Total individuals with primary diagnosis of anxiety disorders	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	40.5	41.0	41.3	40.6	39.7	39.0
Female	59.5	59.0	58.7	59.4	60.3	61.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	15.2	15.5	14.5	13.9	13.1	12.9
12 to 14 years	6.9	7.3	7.3	7.3	7.0	7.2
15 to 17 years	7.3	7.7	8.0	8.5	8.2	8.4
18 to 20 years	5.2	5.0	5.0	5.3	5.3	5.6
21 to 24 years	6.6	6.4	6.5	6.6	6.7	6.7
25 to 29 years	9.8	9.7	9.9	10.2	10.6	10.7
30 to 34 years	9.7	9.6	9.8	9.9	10.1	10.1
35 to 39 years	7.7	7.7	7.9	8.2	8.6	8.7
40 to 44 years	7.2	6.9	6.8	6.6	6.5	6.5
45 to 49 years	6.8	6.5	6.4	6.2	6.2	6.0
50 to 54 years	6.4	6.4	6.4	6.2	6.0	5.7
55 to 59 years	4.8	4.8	5.0	5.0	5.0	5.1
60 to 64 years	2.9	3.0	3.1	3.2	3.3	3.4
65 years and older	3.5	3.4	3.4	2.9	3.5	3.1
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 1.15b. Individuals with primary diagnosis of anxiety disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018 (continued)

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	1.3	1.5	1.7	1.7	1.8	1.9
Asian ²	†	1.1	1.2	1.4	1.4	1.4
Black or African American	10.4	10.8	10.9	10.4	10.8	10.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	78.1	76.4	76.2	76.4	76.6	76.0
Other	9.1	9.9	9.7	9.7	9.1	9.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	84.4	83.3	82.9	82.8	84.9	84.0
Hispanic or Latino origin	15.6	16.7	17.1	17.2	15.1	16.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	32.3	36.6	42.8	35.8	42.0	39.8
Puerto Rican	22.1	15.6	14.8	16.5	13.3	14.0
Cuban	7.3	6.1	6.1	7.0	5.2	5.0
Other specific	38.3	41.7	36.4	40.7	39.5	41.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Percentages may not sum to 100 percent because of rounding. This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.16a. Individuals with primary diagnosis of attention deficit/hyperactivity disorder, by gender, age, race, and ethnicity: Number, 2013–2018

<i>Gender, age, race, and ethnicity</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	47	45	46	47	48
Total individuals with primary diagnosis of attention deficit/hyperactivity disorder	442,351	420,745	351,786	406,362	429,499	461,444
Gender						
Male	316,134	298,232	248,457	284,399	301,185	325,382
Female	125,645	121,791	102,403	120,934	126,826	134,576
Total	441,779	420,023	350,860	405,333	428,011	459,958
Age¹						
0 to 11 years	234,350	224,878	175,823	204,785	215,412	232,353
12 to 14 years	91,750	83,526	71,694	82,395	85,927	93,125
15 to 17 years	57,740	52,044	46,296	53,414	56,310	58,851
18 to 20 years	20,592	17,857	16,331	18,403	19,397	20,333
21 to 24 years	10,306	10,628	9,862	10,132	10,806	11,217
25 to 29 years	8,776	9,945	10,272	10,929	12,038	13,428
30 to 34 years	6,076	7,238	7,385	8,323	9,111	10,184
35 to 39 years	3,836	4,404	4,558	5,690	6,513	7,306
40 to 44 years	2,859	3,218	3,048	3,837	4,097	4,570
45 to 49 years	2,081	2,434	2,330	2,955	3,264	3,597
50 to 54 years	1,602	1,865	1,900	2,371	2,488	2,577
55 to 59 years	928	1,173	1,191	1,586	1,807	2,007
60 to 64 years	468	607	696	935	1,057	1,230
65 years and older	906	875	361	565	1,147	648
Total	442,270	420,692	351,747	406,320	429,374	461,426

Continued. See notes at end of table.

Table 1.16a. Individuals with primary diagnosis of attention deficit/hyperactivity disorder, by gender, age, race, and ethnicity: Number, 2013–2018 (continued)

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	4,345	4,829	4,307	4,244	4,134	4,702
Asian ²	1,828	1,871	1,706	2,265	2,335	3,025
Black or African American	107,683	99,897	88,294	96,287	103,260	105,300
Native Hawaiian or Other Pacific Islander	617	662	653	773	633	851
White	256,925	250,570	204,688	240,648	252,104	270,418
Other	42,746	37,424	27,907	38,653	46,040	47,784
Total	414,144	395,253	327,555	382,870	408,506	432,080
Ethnicity						
Not of Hispanic or Latino origin	316,080	298,095	249,365	292,329	335,168	338,159
Hispanic or Latino origin	67,463	60,920	46,557	62,876	71,199	72,374
Total	383,543	359,015	295,922	355,205	406,367	410,533
Specific Hispanic or Latino origin						
Mexican	3,743	4,448	4,724	3,047	3,836	4,000
Puerto Rican	2,679	2,045	1,931	1,585	2,290	2,004
Cuban	572	396	393	369	385	392
Other specific	4,278	4,479	3,887	3,615	4,221	4,175
Total	11,272	11,368	10,935	8,616	10,732	10,571

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.16b. Individuals with primary diagnosis of attention deficit/hyperactivity disorder, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	46	47	48
Total individuals with primary diagnosis of attention deficit/hyperactivity disorder	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	71.6	71.0	70.8	70.2	70.4	70.7
Female	28.4	29.0	29.2	29.8	29.6	29.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	53.0	53.5	50.0	50.4	50.2	50.4
12 to 14 years	20.7	19.9	20.4	20.3	20.0	20.2
15 to 17 years	13.1	12.4	13.2	13.1	13.1	12.8
18 to 20 years	4.7	4.2	4.6	4.5	4.5	4.4
21 to 24 years	2.3	2.5	2.8	2.5	2.5	2.4
25 to 29 years	2.0	2.4	2.9	2.7	2.8	2.9
30 to 34 years	1.4	1.7	2.1	2.0	2.1	2.2
35 to 39 years	†	1.0	1.3	1.4	1.5	1.6
40 to 44 years	†	†	†	†	†	†
45 to 49 years	†	†	†	†	†	†
50 to 54 years	†	†	†	†	†	†
55 to 59 years	†	†	†	†	†	†
60 to 64 years	†	†	†	†	†	†
65 years and older	†	†	†	†	†	†
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 1.16b. Individuals with primary diagnosis of attention deficit/hyperactivity disorder, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	1.0	1.2	1.3	1.1	1.0	1.1
Asian ²	†	†	†	†	†	†
Black or African American	26.0	25.3	27.0	25.1	25.3	24.4
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	62.0	63.4	62.5	62.9	61.7	62.6
Other	10.3	9.5	8.5	10.1	11.3	11.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	82.4	83.0	84.3	82.3	82.5	82.4
Hispanic or Latino origin	17.6	17.0	15.7	17.7	17.5	17.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	33.2	39.1	43.2	35.4	35.7	37.8
Puerto Rican	23.8	18.0	17.7	18.4	21.3	19.0
Cuban	5.1	3.5	3.6	4.3	3.6	3.7
Other specific	38.0	39.4	35.5	42.0	39.3	39.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Percentages may not sum to 100 percent because of rounding. This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.17a. Individuals with primary diagnosis of bipolar disorders, by gender, age, race, and ethnicity: Number, 2013–2018

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	46	45	46	47	48
Total individuals with primary diagnosis of bipolar disorders	854,076	880,161	758,549	663,415	607,293	601,023
Gender						
Male	357,922	372,043	322,508	273,345	240,041	235,506
Female	495,459	506,778	434,726	388,706	366,061	364,430
Total	853,381	878,821	757,234	662,051	606,102	599,936
Age¹						
0 to 11 years	35,526	36,678	28,906	16,730	6,380	4,045
12 to 14 years	38,773	39,965	31,108	19,287	8,975	6,966
15 to 17 years	55,116	55,761	45,078	31,499	17,866	15,203
18 to 20 years	46,470	44,454	36,556	29,559	22,946	22,589
21 to 24 years	64,392	65,062	54,628	45,323	40,004	39,014
25 to 29 years	87,928	91,116	79,428	71,469	68,109	67,763
30 to 34 years	92,271	94,869	81,734	74,269	70,739	70,656
35 to 39 years	80,701	83,970	74,826	70,768	70,321	70,976
40 to 44 years	84,373	85,340	71,839	64,517	61,945	61,056
45 to 49 years	84,005	85,507	72,580	67,812	65,303	64,113
50 to 54 years	78,076	82,816	73,254	67,822	65,491	63,486
55 to 59 years	52,865	57,014	52,852	51,406	52,770	55,072
60 to 64 years	29,221	31,651	30,753	30,195	31,318	33,997
65 years and older	24,153	25,742	24,756	22,408	24,816	25,973
Total	853,870	879,945	758,298	663,064	606,983	600,909

Continued. See notes at end of table.

Table 1.17a. Individuals with primary diagnosis of bipolar disorders, by gender, age, race, and ethnicity: Number, 2013–2018 (continued)

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	8,009	9,181	8,392	6,518	5,853	5,800
Asian ²	5,852	6,522	5,939	5,539	5,201	5,607
Black or African American	134,677	143,798	124,360	106,639	99,024	99,061
Native Hawaiian or Other Pacific Islander	1,989	2,089	2,071	1,737	960	1,065
White	597,036	608,576	518,366	462,005	433,581	427,450
Other	52,484	54,254	46,884	44,251	33,327	37,235
Total	800,047	824,420	706,012	626,689	577,946	576,218
Ethnicity						
Not of Hispanic or Latino origin	651,506	662,812	566,995	506,262	496,797	490,968
Hispanic or Latino origin	87,620	89,909	76,090	72,616	56,517	57,112
Total	739,126	752,721	643,085	578,878	553,314	548,080
Specific Hispanic or Latino origin						
Mexican	3,498	5,391	4,795	2,587	3,242	2,974
Puerto Rican	5,027	4,677	4,369	3,175	3,206	2,989
Cuban	1,369	1,318	1,283	1,073	915	949
Other specific	6,304	8,010	6,177	4,503	4,698	4,858
Total	16,198	19,396	16,624	11,338	12,061	11,770

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.17b. Individuals with primary diagnosis of bipolar disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	46	45	46	47	48
Total individuals with primary diagnosis of bipolar disorders	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	41.9	42.3	42.6	41.3	39.6	39.3
Female	58.1	57.7	57.4	58.7	60.4	60.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	4.2	4.2	3.8	2.5	1.1	†
12 to 14 years	4.5	4.5	4.1	2.9	1.5	1.2
15 to 17 years	6.5	6.3	5.9	4.8	2.9	2.5
18 to 20 years	5.4	5.1	4.8	4.5	3.8	3.8
21 to 24 years	7.5	7.4	7.2	6.8	6.6	6.5
25 to 29 years	10.3	10.4	10.5	10.8	11.2	11.3
30 to 34 years	10.8	10.8	10.8	11.2	11.7	11.8
35 to 39 years	9.5	9.5	9.9	10.7	11.6	11.8
40 to 44 years	9.9	9.7	9.5	9.7	10.2	10.2
45 to 49 years	9.8	9.7	9.6	10.2	10.8	10.7
50 to 54 years	9.1	9.4	9.7	10.2	10.8	10.6
55 to 59 years	6.2	6.5	7.0	7.8	8.7	9.2
60 to 64 years	3.4	3.6	4.1	4.6	5.2	5.7
65 years and older	2.8	2.9	3.3	3.4	4.1	4.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 1.17b. Individuals with primary diagnosis of bipolar disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018 (continued)

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	1.0	1.1	1.2	1.0	1.0	1.0
Asian ²	†	†	†	†	†	†
Black or African American	16.8	17.4	17.6	17.0	17.1	17.2
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	74.6	73.8	73.4	73.7	75.0	74.2
Other	6.6	6.6	6.6	7.1	5.8	6.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	88.1	88.1	88.2	87.5	89.8	89.6
Hispanic or Latino origin	11.9	11.9	11.8	12.5	10.2	10.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	21.6	27.8	28.8	22.8	26.9	25.3
Puerto Rican	31.0	24.1	26.3	28.0	26.6	25.4
Cuban	8.5	6.8	7.7	9.5	7.6	8.1
Other specific	38.9	41.3	37.2	39.7	39.0	41.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Percentages may not sum to 100 percent because of rounding. This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 1.18a. Individuals with primary diagnosis of depressive disorders, by gender, age, race, and ethnicity:
Number, 2013–2018**

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals with primary diagnosis of depressive disorders	1,267,441	1,327,354	1,255,030	1,257,761	1,315,673	1,410,277
Gender						
Male	448,787	473,878	454,264	458,297	483,700	522,635
Female	817,585	851,797	799,005	797,366	829,651	885,445
Total	1,266,372	1,325,675	1,253,269	1,255,663	1,313,351	1,408,080
Age¹						
0 to 11 years	33,134	34,526	31,092	30,111	30,555	33,950
12 to 14 years	58,869	66,493	64,146	63,407	64,445	73,268
15 to 17 years	91,226	97,878	97,813	101,954	104,792	116,839
18 to 20 years	59,889	59,676	57,705	61,641	66,889	76,711
21 to 24 years	74,407	76,409	72,815	75,361	80,971	88,548
25 to 29 years	105,402	109,754	105,550	110,660	120,788	131,091
30 to 34 years	110,001	115,140	108,854	110,687	117,433	124,637
35 to 39 years	101,918	106,305	101,583	104,264	111,333	118,321
40 to 44 years	115,230	116,610	104,037	100,216	100,297	104,196
45 to 49 years	130,451	132,587	117,431	114,934	115,726	118,304
50 to 54 years	140,197	147,855	136,176	132,784	131,275	132,507
55 to 59 years	111,740	120,611	115,869	116,267	119,779	128,185
60 to 64 years	68,419	73,733	73,486	74,116	78,313	87,352
65 years and older	66,176	69,479	68,161	60,867	72,188	75,796
Total	1,267,059	1,327,056	1,254,718	1,257,269	1,314,784	1,409,705

Continued. See notes at end of table.

**Table 1.18a. Individuals with primary diagnosis of depressive disorders, by gender, age, race, and ethnicity:
Number, 2013–2018 (continued)**

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	13,869	16,973	17,523	16,982	18,047	19,312
Asian ²	16,832	20,503	20,106	20,270	20,518	22,533
Black or African American	212,743	227,689	214,273	209,395	222,751	238,281
Native Hawaiian or Other Pacific Islander	3,218	3,593	4,039	4,325	3,015	3,609
White	820,377	852,465	798,234	814,506	852,467	901,982
Other	109,419	113,017	100,505	110,722	117,684	135,823
Total	1,176,458	1,234,240	1,154,680	1,176,200	1,234,482	1,321,540
Ethnicity						
Not of Hispanic or Latino origin	899,265	926,552	881,406	893,898	1,005,219	1,053,960
Hispanic or Latino origin	171,882	179,341	164,383	185,380	191,368	204,991
Total	1,071,147	1,105,893	1,045,789	1,079,278	1,196,587	1,258,951
Specific Hispanic or Latino origin						
Mexican	9,249	12,364	14,421	9,259	13,264	13,009
Puerto Rican	8,400	7,589	7,370	6,149	6,910	7,148
Cuban	3,541	3,297	3,156	2,989	2,776	2,743
Other specific	12,516	14,754	13,542	12,341	14,791	15,208
Total	33,706	38,004	38,489	30,738	37,741	38,108

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 1.18b. Individuals with primary diagnosis of depressive disorders, by gender, age, race, and ethnicity:
Column percent distribution, 2013–2018**

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals with primary diagnosis of depressive disorders	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	35.4	35.7	36.2	36.5	36.8	37.1
Female	64.6	64.3	63.8	63.5	63.2	62.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	2.6	2.6	2.5	2.4	2.3	2.4
12 to 14 years	4.6	5.0	5.1	5.0	4.9	5.2
15 to 17 years	7.2	7.4	7.8	8.1	8.0	8.3
18 to 20 years	4.7	4.5	4.6	4.9	5.1	5.4
21 to 24 years	5.9	5.8	5.8	6.0	6.2	6.3
25 to 29 years	8.3	8.3	8.4	8.8	9.2	9.3
30 to 34 years	8.7	8.7	8.7	8.8	8.9	8.8
35 to 39 years	8.0	8.0	8.1	8.3	8.5	8.4
40 to 44 years	9.1	8.8	8.3	8.0	7.6	7.4
45 to 49 years	10.3	10.0	9.4	9.1	8.8	8.4
50 to 54 years	11.1	11.1	10.9	10.6	10.0	9.4
55 to 59 years	8.8	9.1	9.2	9.2	9.1	9.1
60 to 64 years	5.4	5.6	5.9	5.9	6.0	6.2
65 years and older	5.2	5.2	5.4	4.8	5.5	5.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 1.18b. Individuals with primary diagnosis of depressive disorders, by gender, age, race, and ethnicity:
Column percent distribution, 2013–2018 (continued)**

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	1.2	1.4	1.5	1.4	1.5	1.5
Asian ²	1.4	1.7	1.7	1.7	1.7	1.7
Black or African American	18.1	18.4	18.6	17.8	18.0	18.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	69.7	69.1	69.1	69.2	69.1	68.3
Other	9.3	9.2	8.7	9.4	9.5	10.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	84.0	83.8	84.3	82.8	84.0	83.7
Hispanic or Latino origin	16.0	16.2	15.7	17.2	16.0	16.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	27.4	32.5	37.5	30.1	35.1	34.1
Puerto Rican	24.9	20.0	19.1	20.0	18.3	18.8
Cuban	10.5	8.7	8.2	9.7	7.4	7.2
Other specific	37.1	38.8	35.2	40.1	39.2	39.9
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Percentages may not sum to 100 percent because of rounding. This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.19a. Individuals with primary diagnosis of schizophrenia and other psychotic disorders, by gender, age, race, and ethnicity: Number, 2013–2018

<i>Gender, age, race, and ethnicity</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	47	45	46	47	48
Total individuals with primary diagnosis of schizophrenia and other psychotic disorders	630,500	656,286	593,824	576,500	626,384	660,521
Gender						
Male	383,632	402,328	367,935	359,909	392,731	414,976
Female	246,365	252,549	224,958	215,651	232,794	244,877
Total	629,997	654,877	592,893	575,560	625,525	659,853
Age¹						
0 to 11 years	2,056	2,139	1,746	1,587	1,707	1,445
12 to 14 years	2,520	2,656	2,118	2,026	2,127	2,088
15 to 17 years	5,754	5,810	5,040	4,849	5,058	5,088
18 to 20 years	15,644	15,809	13,575	13,000	13,971	14,165
21 to 24 years	34,416	36,701	33,869	33,390	35,518	36,825
25 to 29 years	53,851	56,582	52,662	53,263	59,867	64,675
30 to 34 years	60,578	64,902	59,800	58,976	64,220	68,071
35 to 39 years	55,652	58,887	55,895	57,034	64,257	70,024
40 to 44 years	63,219	63,610	55,432	53,041	56,727	60,432
45 to 49 years	77,522	76,398	65,090	61,500	64,490	66,022
50 to 54 years	90,289	92,240	80,171	75,236	76,585	76,202
55 to 59 years	74,752	79,370	72,439	70,758	76,715	80,504
60 to 64 years	46,585	50,736	48,151	47,902	53,920	59,181
65 years and older	47,395	50,131	47,511	43,475	50,537	55,258
Total	630,233	655,971	593,499	576,037	625,699	659,980

Continued. See notes at end of table.

Table 1.19a. Individuals with primary diagnosis of schizophrenia and other psychotic disorders, by gender, age, race, and ethnicity: Number, 2013–2018 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	5,491	6,191	5,840	5,868	6,234	6,711
Asian ²	14,399	15,638	14,700	14,685	15,028	16,929
Black or African American	191,698	198,981	182,612	176,797	193,713	205,237
Native Hawaiian or Other Pacific Islander	2,728	2,801	2,770	2,840	1,765	1,958
White	323,116	335,206	295,827	296,143	322,396	334,030
Other	39,487	40,446	39,130	39,777	47,401	58,000
Total	576,919	599,263	540,879	536,110	586,537	622,865
Ethnicity						
Not of Hispanic or Latino origin	455,984	470,108	419,138	430,365	481,585	511,034
Hispanic or Latino origin	64,601	65,807	60,082	65,343	69,746	78,311
Total	520,585	535,915	479,220	495,708	551,331	589,345
Specific Hispanic or Latino origin						
Mexican	2,052	3,106	3,059	2,595	3,700	3,446
Puerto Rican	3,434	3,213	3,102	2,630	3,411	3,739
Cuban	2,123	1,413	1,380	1,343	1,436	1,414
Other specific	5,133	5,347	4,713	4,658	5,363	5,798
Total	12,742	13,079	12,254	11,226	13,910	14,397

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.19b. Individuals with primary diagnosis of schizophrenia and other psychotic disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	46	47	48
Total individuals with primary diagnosis of schizophrenia and other psychotic disorders	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	60.9	61.4	62.1	62.5	62.8	62.9
Female	39.1	38.6	37.9	37.5	37.2	37.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	†	†	†	†	†	†
12 to 14 years	†	†	†	†	†	†
15 to 17 years	†	†	†	†	†	†
18 to 20 years	2.5	2.4	2.3	2.3	2.2	2.1
21 to 24 years	5.5	5.6	5.7	5.8	5.7	5.6
25 to 29 years	8.5	8.6	8.9	9.2	9.6	9.8
30 to 34 years	9.6	9.9	10.1	10.2	10.3	10.3
35 to 39 years	8.8	9.0	9.4	9.9	10.3	10.6
40 to 44 years	10.0	9.7	9.3	9.2	9.1	9.2
45 to 49 years	12.3	11.6	11.0	10.7	10.3	10.0
50 to 54 years	14.3	14.1	13.5	13.1	12.2	11.5
55 to 59 years	11.9	12.1	12.2	12.3	12.3	12.2
60 to 64 years	7.4	7.7	8.1	8.3	8.6	9.0
65 years and older	7.5	7.6	8.0	7.5	8.1	8.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 1.19b. Individuals with primary diagnosis of schizophrenia and other psychotic disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	1.0	1.1	1.1	1.1	1.1
Asian ²	2.5	2.6	2.7	2.7	2.6	2.7
Black or African American	33.2	33.2	33.8	33.0	33.0	33.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	56.0	55.9	54.7	55.2	55.0	53.6
Other	6.8	6.7	7.2	7.4	8.1	9.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	87.6	87.7	87.5	86.8	87.3	86.7
Hispanic or Latino origin	12.4	12.3	12.5	13.2	12.7	13.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	16.1	23.7	25.0	23.1	26.6	23.9
Puerto Rican	27.0	24.6	25.3	23.4	24.5	26.0
Cuban	16.7	10.8	11.3	12.0	10.3	9.8
Other specific	40.3	40.9	38.5	41.5	38.6	40.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTES: Percentages may not sum to 100 percent because of rounding. This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.20a. Individuals with co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Number, 2013–2018

<i>Gender, age, race, and ethnicity</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	46	44	44	45	45
Total individuals with co-occurring mental health and substance use disorders¹	782,501	777,946	758,082	690,705	905,849	1,051,340
Gender						
Male	435,055	432,519	421,047	380,465	482,415	560,368
Female	346,874	344,573	336,592	309,714	420,103	489,086
Total	781,929	777,092	757,639	690,179	902,518	1,049,454
Age²						
0 to 11 years	6,121	3,697	2,024	1,999	22,887	23,623
12 to 14 years	11,944	10,332	8,185	7,686	19,366	18,691
15 to 17 years	47,758	42,145	36,170	33,985	46,671	49,196
18 to 20 years	37,008	33,790	31,459	28,832	36,981	42,764
21 to 24 years	61,377	59,646	57,960	51,362	64,208	73,139
25 to 29 years	93,623	91,810	91,808	85,553	111,611	133,007
30 to 34 years	96,545	96,053	96,626	89,039	113,299	134,479
35 to 39 years	78,764	79,878	81,582	78,199	102,536	123,782
40 to 44 years	83,691	80,715	75,723	66,148	80,857	94,824
45 to 49 years	89,350	86,840	81,933	71,483	85,767	97,095
50 to 54 years	85,375	89,034	86,199	75,850	89,182	99,167
55 to 59 years	52,925	59,017	60,008	56,210	71,686	86,024
60 to 64 years	23,990	27,981	30,161	28,747	38,208	47,587
65 years and older	13,480	16,295	17,413	14,731	21,062	26,971
Total	781,951	777,233	757,251	689,824	904,321	1,050,349

Continued. See notes at end of table.

Table 1.20a. Individuals with co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Number, 2013–2018 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	12,184	13,610	14,290	13,774	15,647	19,130
Asian ³	5,557	6,157	6,301	7,192	7,652	9,541
Black or African American	141,181	142,675	134,840	121,760	169,829	199,925
Native Hawaiian or Other Pacific Islander	2,042	2,279	2,264	1,811	1,798	2,215
White	509,870	501,387	489,318	449,537	586,527	672,893
Other	57,575	59,347	59,549	56,357	75,902	95,273
Total	728,409	725,455	706,562	650,431	857,355	998,977
Ethnicity						
Not of Hispanic or Latino origin	585,348	580,194	570,041	541,710	743,946	849,744
Hispanic or Latino origin	95,393	97,213	93,600	92,863	101,805	112,120
Total	680,741	677,407	663,641	634,573	845,751	961,864
Specific Hispanic or Latino origin						
Mexican	3,719	4,908	5,677	2,410	4,556	5,154
Puerto Rican	3,685	7,462	6,946	5,049	6,611	6,917
Cuban	1,279	843	830	618	877	931
Other specific	5,954	9,534	7,332	5,401	6,877	7,292
Total	14,637	22,747	20,785	13,478	18,921	20,294

¹ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.20b. Individuals with co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	46	44	44	45	45
Total individuals with co-occurring mental health and substance use disorders¹	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	55.6	55.7	55.6	55.1	53.5	53.4
Female	44.4	44.3	44.4	44.9	46.5	46.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age²						
0 to 11 years	†	†	†	†	2.5	2.2
12 to 14 years	1.5	1.3	1.1	1.1	2.1	1.8
15 to 17 years	6.1	5.4	4.8	4.9	5.2	4.7
18 to 20 years	4.7	4.3	4.2	4.2	4.1	4.1
21 to 24 years	7.8	7.7	7.7	7.4	7.1	7.0
25 to 29 years	12.0	11.8	12.1	12.4	12.3	12.7
30 to 34 years	12.3	12.4	12.8	12.9	12.5	12.8
35 to 39 years	10.1	10.3	10.8	11.3	11.3	11.8
40 to 44 years	10.7	10.4	10.0	9.6	8.9	9.0
45 to 49 years	11.4	11.2	10.8	10.4	9.5	9.2
50 to 54 years	10.9	11.5	11.4	11.0	9.9	9.4
55 to 59 years	6.8	7.6	7.9	8.1	7.9	8.2
60 to 64 years	3.1	3.6	4.0	4.2	4.2	4.5
65 years and older	1.7	2.1	2.3	2.1	2.3	2.6
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 1.20b. Individuals with co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	1.7	1.9	2.0	2.1	1.8	1.9
Asian ³	†	†	†	1.1	†	†
Black or African American	19.4	19.7	19.1	18.7	19.8	20.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	70.0	69.1	69.3	69.1	68.4	67.4
Other	7.9	8.2	8.4	8.7	8.9	9.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	86.0	85.6	85.9	85.4	88.0	88.3
Hispanic or Latino origin	14.0	14.4	14.1	14.6	12.0	11.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	25.4	21.6	27.3	17.9	24.1	25.4
Puerto Rican	25.2	32.8	33.4	37.5	34.9	34.1
Cuban	8.7	3.7	4.0	4.6	4.6	4.6
Other specific	40.7	41.9	35.3	40.1	36.3	35.9
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.21a. Individuals without co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Number, 2013–2018

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	46	44	43	45	45
Total individuals without co-occurring mental health and substance use disorders	3,891,997	3,849,896	3,466,494	3,563,762	3,641,596	3,897,679
Gender						
Male	1,810,514	1,788,302	1,598,082	1,652,215	1,677,409	1,793,313
Female	2,077,439	2,055,895	1,861,506	1,903,078	1,955,509	2,095,698
Total	3,887,953	3,844,197	3,459,588	3,555,293	3,632,918	3,889,011
Age¹						
0 to 11 years	724,293	722,033	609,207	664,155	646,508	725,022
12 to 14 years	343,135	338,428	299,439	315,770	312,921	350,089
15 to 17 years	324,154	316,308	292,189	314,115	313,584	339,749
18 to 20 years	180,514	165,974	151,934	161,427	167,616	185,694
21 to 24 years	196,729	192,953	178,782	180,851	187,370	198,958
25 to 29 years	260,796	262,288	247,118	256,895	272,328	290,383
30 to 34 years	266,824	269,239	248,429	254,181	261,923	273,414
35 to 39 years	237,507	238,994	225,245	233,190	244,321	258,125
40 to 44 years	252,935	245,892	216,290	212,050	211,901	219,788
45 to 49 years	272,107	262,863	227,757	225,873	226,825	231,482
50 to 54 years	285,329	282,806	252,744	247,049	245,126	245,105
55 to 59 years	230,040	231,793	215,695	216,172	224,699	237,545
60 to 64 years	146,594	148,951	143,212	143,969	154,551	171,075
65 years and older	169,291	169,759	156,961	135,998	168,788	169,960
Total	3,890,248	3,848,281	3,465,002	3,561,695	3,638,461	3,896,389

Continued. See notes at end of table.

Table 1.21a. Individuals without co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Number, 2013–2018 (continued)

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	40,618	48,741	48,596	46,135	49,619	54,207
Asian ²	47,238	54,275	51,913	55,928	58,782	64,290
Black or African American	739,796	709,690	616,524	599,610	626,260	646,057
Native Hawaiian or Other Pacific Islander	9,980	10,462	11,744	14,254	8,843	11,154
White	2,386,578	2,368,939	2,100,581	2,219,422	2,285,242	2,408,643
Other	348,473	352,131	310,109	347,123	365,866	409,311
Total	3,572,683	3,544,238	3,139,467	3,282,472	3,394,612	3,593,662
Ethnicity						
Not of Hispanic or Latino origin	2,700,658	2,641,448	2,348,812	2,485,821	2,785,447	2,834,871
Hispanic or Latino origin	521,073	518,320	474,815	553,332	558,425	602,231
Total	3,221,731	3,159,768	2,823,627	3,039,153	3,343,872	3,437,102
Specific Hispanic or Latino origin						
Mexican	23,913	32,352	35,452	26,860	39,357	37,325
Puerto Rican	21,346	11,903	11,364	9,236	8,507	8,799
Cuban	5,495	1,502	1,565	1,374	1,766	1,561
Other specific	30,677	31,000	25,768	23,163	31,534	31,525
Total	81,431	76,757	74,149	60,633	81,164	79,210

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.21b. Individuals without co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018

<i>Gender, age, race, and ethnicity</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	46	44	43	45	45
Total individuals without co-occurring mental health and substance use disorders	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	46.6	46.5	46.2	46.5	46.2	46.1
Female	53.4	53.5	53.8	53.5	53.8	53.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 11 years	18.6	18.8	17.6	18.6	17.8	18.6
12 to 14 years	8.8	8.8	8.6	8.9	8.6	9.0
15 to 17 years	8.3	8.2	8.4	8.8	8.6	8.7
18 to 20 years	4.6	4.3	4.4	4.5	4.6	4.8
21 to 24 years	5.1	5.0	5.2	5.1	5.1	5.1
25 to 29 years	6.7	6.8	7.1	7.2	7.5	7.5
30 to 34 years	6.9	7.0	7.2	7.1	7.2	7.0
35 to 39 years	6.1	6.2	6.5	6.5	6.7	6.6
40 to 44 years	6.5	6.4	6.2	6.0	5.8	5.6
45 to 49 years	7.0	6.8	6.6	6.3	6.2	5.9
50 to 54 years	7.3	7.3	7.3	6.9	6.7	6.3
55 to 59 years	5.9	6.0	6.2	6.1	6.2	6.1
60 to 64 years	3.8	3.9	4.1	4.0	4.2	4.4
65 years and older	4.4	4.4	4.5	3.8	4.6	4.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 1.21b. Individuals without co-occurring mental health and substance use disorders, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018 (continued)

<i>Gender, age, race, and ethnicity</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	1.1	1.4	1.5	1.4	1.5	1.5
Asian ²	1.3	1.5	1.7	1.7	1.7	1.8
Black or African American	20.7	20.0	19.6	18.3	18.4	18.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	66.8	66.8	66.9	67.6	67.3	67.0
Other	9.8	9.9	9.9	10.6	10.8	11.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	83.8	83.6	83.2	81.8	83.3	82.5
Hispanic or Latino origin	16.2	16.4	16.8	18.2	16.7	17.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	29.4	42.1	47.8	44.3	48.5	47.1
Puerto Rican	26.2	15.5	15.3	15.2	10.5	11.1
Cuban	6.7	2.0	2.1	2.3	2.2	2.0
Other specific	37.7	40.4	34.8	38.2	38.9	39.8
Total	100.0	100.0	100.0	100.0	100.0	100.0

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 1.22a. Individuals who received mental health services and U.S. population, by gender, marital status, and age group: Number, 2018

Gender and marital status	Age group				Total
	0–17 years	18–24 years	25–54 years	55 years and older	
<i>Individuals who received mental health services¹</i>					
Gender					
Male	1,013,374	305,519	1,306,842	393,913	3,019,648
Female	781,621	337,271	1,518,595	537,368	3,174,855
Total	1,794,995	642,790	2,825,437	931,281	6,194,503
Marital status					
Never married	1,214,487	322,108	931,694	206,507	2,674,796
Currently married	633	15,811	254,377	90,401	361,222
Separated	351	4,232	109,433	32,924	146,940
Divorced	184	2,548	225,515	140,793	369,040
Widowed	45	354	28,741	48,335	77,475
Total	1,215,700	345,053	1,549,760	518,960	3,629,473
<i>U.S. population</i>					
Gender					
Male	37,490,849	15,592,638	64,339,511	43,705,681	161,128,679
Female	35,908,493	14,864,264	64,267,850	50,998,148	166,038,755
Total	73,399,342	30,456,902	128,607,361	94,703,829	327,167,434
Marital status					
Never married	<i>n/a</i>	26,449,000	38,060,000	7,295,000	71,804,000
Currently married	<i>n/a</i>	2,424,000	72,004,000	56,869,000	131,297,000
Separated	<i>n/a</i>	268,000	3,191,000	1,570,000	5,029,000
Divorced	<i>n/a</i>	170,000	12,290,000	13,408,000	25,868,000
Widowed	<i>n/a</i>	36,000	1,460,000	13,581,000	15,077,000
Total	<i>n/a</i>	29,347,000	127,005,000	92,723,000	249,075,000

n/a Population detail required for marital status was not available.

¹ Age is based on the age of the individual who received mental health services at the midpoint of the reporting period.

SOURCES: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019; U.S. population: U.S. Census Bureau, Current Population Survey, 2018 Annual Social and Economic Supplement (ASEC), March 2018, Table A1. <https://www.census.gov/data/tables/2018/demo/families/cps-2018.html>; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2018-alldata6: "Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2018."

Table 1.22b. Individuals who received mental health services and U.S. population, by gender, marital status, and age group: Column percent distribution, 2018

Gender and marital status	Age group				
	0–17 years	18–24 years	25–54 years	55 years and older	Total
<i>Individuals who received mental health services¹</i>					
Gender					
Male	56.5	47.5	46.3	42.3	48.7
Female	43.5	52.5	53.7	57.7	51.3
Total	100.0	100.0	100.0	100.0	100.0
Marital status					
Never married	99.9	93.4	60.1	39.8	73.7
Currently married	†	4.6	16.4	17.4	10.0
Separated	†	1.2	7.1	6.3	4.0
Divorced	†	†	14.6	27.1	10.2
Widowed	†	†	1.9	9.3	2.1
Total	100.0	100.0	100.0	100.0	100.0
<i>U.S. population</i>					
Gender					
Male	51.1	51.2	50.0	46.1	49.2
Female	48.9	48.8	50.0	53.9	50.8
Total	100.0	100.0	100.0	100.0	100.0
Marital status					
Never married	<i>n/a</i>	90.1	30.0	7.9	28.8
Currently married	<i>n/a</i>	8.3	56.7	61.3	52.7
Separated	<i>n/a</i>	†	2.5	1.7	2.0
Divorced	<i>n/a</i>	†	9.7	14.5	10.4
Widowed	<i>n/a</i>	†	1.1	14.6	6.1
Total	<i>n/a</i>	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; *n/a* Population detail required for marital status was not available.

¹ Age is based on the age of the individual who received mental health services at the midpoint of the reporting period.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCES: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019; U.S. population: U.S. Census Bureau, Current Population Survey, 2018 Annual Social and Economic Supplement (ASEC), March 2018, Table A1. <https://www.census.gov/data/tables/2018/demo/families/cps-2018.html>; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2018-alldata6: "Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2018."

Table 1.22c. Individuals who received mental health services and U.S. population, by gender, marital status, and age group: Row percent distribution, 2018

Gender and marital status	Age group				Total
	0–17 years	18–24 years	25–54 years	55 years and older	
<i>Individuals who received mental health services¹</i>					
Gender					
Male	33.6	10.1	43.3	13.0	100.0
Female	24.6	10.6	47.8	16.9	100.0
Total	29.0	10.4	45.6	15.0	100.0
Marital status					
Never married	45.4	12.0	34.8	7.7	100.0
Currently married	†	4.4	70.4	25.0	100.0
Separated	†	2.9	74.5	22.4	100.0
Divorced	†	†	61.1	38.2	100.0
Widowed	†	†	37.1	62.4	100.0
Total	33.5	9.5	42.7	14.3	100.0
<i>U.S. population</i>					
Gender					
Male	23.3	9.7	39.9	27.1	100.0
Female	21.6	9.0	38.7	30.7	100.0
Total	22.4	9.3	39.3	28.9	100.0
Marital status					
Never married	<i>n/a</i>	36.8	53.0	10.2	100.0
Currently married	<i>n/a</i>	1.8	54.8	43.3	100.0
Separated	<i>n/a</i>	5.3	63.5	31.2	100.0
Divorced	<i>n/a</i>	†	47.5	51.8	100.0
Widowed	<i>n/a</i>	†	9.7	90.1	100.0
Total	<i>n/a</i>	11.8	51.0	37.2	100.0

† Less than 1 percent of individuals; *n/a* Population detail required for marital status was not available.

¹ Age is based on the age of the individual who received mental health services at the midpoint of the reporting period.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCES: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019; U.S. population: U.S. Census Bureau, Current Population Survey, 2018 Annual Social and Economic Supplement (ASEC), March 2018, Table A1. <https://www.census.gov/data/tables/2018/demo/families/cps-2018.html>; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2018-alldata6: "Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2018."

Table 1.23a. Individuals who received mental health services and U.S. population, by race, ethnicity, and age group: Number, 2018

Race and ethnicity	Age group				Total
	0–17 years	18–24 years	25–54 years	55 years and older	
<i>Individuals who received mental health services¹</i>					
Race					
American Indian or Alaska Native	28,223	10,042	42,440	8,849	89,554
Asian ²	18,667	9,542	43,607	19,385	91,201
Black or African American	322,049	119,419	521,214	175,479	1,138,161
Native Hawaiian or Other Pacific Islander	4,427	1,908	7,178	1,776	15,289
White	1,017,941	387,978	1,824,351	613,288	3,843,558
Other	249,786	67,959	206,253	54,359	578,357
Total	1,641,093	596,848	2,645,043	873,136	5,756,120
Ethnicity					
Not of Hispanic or Latino origin	1,213,304	465,569	2,184,247	738,800	4,601,920
Hispanic or Latino origin	383,530	103,558	322,753	82,984	892,825
Total	1,596,834	569,127	2,507,000	821,784	5,494,745
<i>U.S. population</i>					
Race					
American Indian or Alaska Native	1,166,440	456,752	1,709,898	814,431	4,147,521
Asian	3,933,986	1,811,082	9,059,773	4,525,759	19,330,600
Black or African American	11,075,827	4,673,598	18,011,391	10,043,503	43,804,319
Native Hawaiian or Other Pacific Islander	218,738	83,988	354,783	141,909	799,418
White	53,006,956	22,257,040	96,696,261	78,178,839	250,139,096
Other	3,997,395	1,174,442	2,775,255	999,388	8,946,480
Total	73,399,342	30,456,902	128,607,361	94,703,829	327,167,434
Ethnicity					
Not of Hispanic or Latino origin	54,698,158	23,636,702	103,565,518	85,395,310	267,295,688
Hispanic or Latino origin	18,701,184	6,820,200	25,041,843	9,308,519	59,871,746
Total	73,399,342	30,456,902	128,607,361	94,703,829	327,167,434

¹ Age is based on the age of the individual who received mental health services at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

SOURCES: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2018-alldata6: "Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2018."

Table 1.23b. Individuals who received mental health services and U.S. population, by race, ethnicity, and age group: Column percent distribution, 2018

Race and ethnicity	Age group				Total
	0–17 years	18–24 years	25–54 years	55 years and older	
<i>Individuals who received mental health services¹</i>					
Race					
American Indian or Alaska Native	1.7	1.7	1.6	1.0	1.6
Asian ²	1.1	1.6	1.6	2.2	1.6
Black or African American	19.6	20.0	19.7	20.1	19.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†
White	62.0	65.0	69.0	70.2	66.8
Other	15.2	11.4	7.8	6.2	10.0
Total	100.0	100.0	100.0	100.0	100.0
Ethnicity					
Not of Hispanic or Latino origin	76.0	81.8	87.1	89.9	83.8
Hispanic or Latino origin	24.0	18.2	12.9	10.1	16.2
Total	100.0	100.0	100.0	100.0	100.0
<i>U.S. population</i>					
Race					
American Indian or Alaska Native	1.6	1.5	1.3	†	1.3
Asian	5.4	5.9	7.0	4.8	5.9
Black or African American	15.1	15.3	14.0	10.6	13.4
Native Hawaiian or Other Pacific Islander	†	†	†	†	†
White	72.2	73.1	75.2	82.6	76.5
Other	5.4	3.9	2.2	1.1	2.7
Total	100.0	100.0	100.0	100.0	100.0
Ethnicity					
Not of Hispanic or Latino origin	74.5	77.6	80.5	90.2	81.7
Hispanic or Latino origin	25.5	22.4	19.5	9.8	18.3
Total	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ Age is based on the age of the individual who received mental health services at the midpoint of the reporting period.

² For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCES: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2018-alldata6: “Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2018.”

Table 1.23c. Individuals who received mental health services and U.S. population, by race, ethnicity, and age group: Row percent distribution, 2018

Race and ethnicity	Age group				Total
	0–17 years	18–24 years	25–54 years	55 years and older	
<i>Individuals who received mental health services¹</i>					
Race					
American Indian or Alaska Native	31.5	11.2	47.4	9.9	100.0
Asian ²	20.5	10.5	47.8	21.3	100.0
Black or African American	28.3	10.5	45.8	15.4	100.0
Native Hawaiian or Other Pacific Islander	29.0	12.5	46.9	11.6	100.0
White	26.5	10.1	47.5	16.0	100.0
Other	43.2	11.8	35.7	9.4	100.0
Total	28.5	10.4	46.0	15.2	100.0
Ethnicity					
Not of Hispanic or Latino origin	26.4	10.1	47.5	16.1	100.0
Hispanic or Latino origin	43.0	11.6	36.1	9.3	100.0
Total	29.1	10.4	45.6	15.0	100.0
<i>U.S. population</i>					
Race					
American Indian or Alaska Native	28.1	11.0	41.2	19.6	100.0
Asian	20.4	9.4	46.9	23.4	100.0
Black or African American	25.3	10.7	41.1	22.9	100.0
Native Hawaiian or Other Pacific Islander	27.4	10.5	44.4	17.8	100.0
White	21.2	8.9	38.7	31.3	100.0
Other	44.7	13.1	31.0	11.2	100.0
Total	22.4	9.3	39.3	28.9	100.0
Ethnicity					
Not of Hispanic or Latino origin	20.5	8.8	38.7	31.9	100.0
Hispanic or Latino origin	31.2	11.4	41.8	15.5	100.0
Total	22.4	9.3	39.3	28.9	100.0

¹ Age is based on the age of the individual who received mental health services at the midpoint of the reporting period.

² For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCES: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2018-alldata6: “Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2018.”

Table 2.1a. Individuals aged 17 years and younger, by gender, age, race, ethnicity, and SED status: Number, 2013–2018

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	47	45	47	48	48
Total individuals aged 0 to 17 years	1,650,728	1,706,582	1,519,237	1,658,008	1,657,084	1,800,613
Gender						
Male	970,220	989,646	862,910	941,055	933,267	1,013,374
Female	678,593	714,820	653,697	710,853	719,071	781,621
Total	1,648,813	1,704,466	1,516,607	1,651,908	1,652,338	1,794,995
Age¹						
0 to 5 years	168,748	186,262	147,180	165,445	164,824	180,312
6 to 11 years	643,371	663,645	584,171	645,959	647,335	712,172
12 to 14 years	403,736	417,244	376,750	401,746	401,160	438,967
15 to 17 years	434,873	439,431	411,136	444,858	443,765	469,162
Total	1,650,728	1,706,582	1,519,237	1,658,008	1,657,084	1,800,613
Race						
American Indian or Alaska Native	20,981	24,785	23,689	23,759	24,849	28,223
Asian ²	11,659	13,434	12,460	15,436	17,131	18,667
Black or African American	329,634	339,316	296,493	307,222	311,754	322,049
Native Hawaiian or Other Pacific Islander	3,540	3,883	4,382	5,077	3,564	4,427
White	924,813	955,187	826,003	925,604	944,085	1,017,941
Other	221,144	230,267	210,638	226,469	238,763	249,786
Total	1,511,771	1,566,872	1,373,665	1,503,567	1,540,146	1,641,093

Continued. See notes at end of table.

Table 2.1a. Individuals aged 17 years and younger, by gender, age, race, ethnicity, and SED status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Ethnicity						
Not of Hispanic or Latino origin	1,071,376	1,080,124	944,777	1,034,716	1,177,242	1,213,304
Hispanic or Latino origin	318,053	337,472	321,093	345,317	362,162	383,530
Total	1,389,429	1,417,596	1,265,870	1,380,033	1,539,404	1,596,834
Specific Hispanic or Latino origin						
Mexican	17,640	24,177	27,100	19,982	25,095	24,239
Puerto Rican	10,541	11,757	12,892	10,418	9,602	6,342
Cuban	1,731	1,360	1,329	1,221	1,327	1,364
Other specific	16,778	21,950	19,108	17,820	20,644	20,911
Total	46,690	59,244	60,429	49,441	56,668	52,856
SED status³						
SED	1,058,643	1,188,108	1,030,923	1,064,195	1,119,810	1,241,052
Not SED	443,847	384,801	387,164	476,599	458,841	481,155
Total	1,502,490	1,572,909	1,418,087	1,540,794	1,578,651	1,722,207

SED = serious emotional disturbance.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SED status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 2.1b. Individuals aged 17 years and younger, by gender, age, race, ethnicity, and SED status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals aged 0 to 17 years	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	58.8	58.1	56.9	57.0	56.5	56.5
Female	41.2	41.9	43.1	43.0	43.5	43.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
0 to 5 years	10.2	10.9	9.7	10.0	9.9	10.0
6 to 11 years	39.0	38.9	38.5	39.0	39.1	39.6
12 to 14 years	24.5	24.4	24.8	24.2	24.2	24.4
15 to 17 years	26.3	25.7	27.1	26.8	26.8	26.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	1.4	1.6	1.7	1.6	1.6	1.7
Asian ²	†	†	†	1.0	1.1	1.1
Black or African American	21.8	21.7	21.6	20.4	20.2	19.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	61.2	61.0	60.1	61.6	61.3	62.0
Other	14.6	14.7	15.3	15.1	15.5	15.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 2.1b. Individuals aged 17 years and younger, by gender, age, race, ethnicity, and SED status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Ethnicity						
Not of Hispanic or Latino origin	77.1	76.2	74.6	75.0	76.5	76.0
Hispanic or Latino origin	22.9	23.8	25.4	25.0	23.5	24.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	37.8	40.8	44.8	40.4	44.3	45.9
Puerto Rican	22.6	19.8	21.3	21.1	16.9	12.0
Cuban	3.7	2.3	2.2	2.5	2.3	2.6
Other specific	35.9	37.1	31.6	36.0	36.4	39.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
SED status³						
SED	70.5	75.5	72.7	69.1	70.9	72.1
Not SED	29.5	24.5	27.3	30.9	29.1	27.9
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

SED = serious emotional disturbance.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SED status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 2.2a. Individuals aged 17 years and younger with SED, by gender, age, race, and ethnicity: Number, 2013–2018

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	44	47	45	47	47	48
Total individuals with SED aged 0 to 17 years¹	1,058,643	1,188,108	1,030,923	1,064,195	1,119,810	1,241,052
Gender						
Male	637,507	698,562	591,747	616,275	635,806	697,259
Female	419,818	488,060	437,304	445,786	481,118	541,077
Total	1,057,325	1,186,622	1,029,051	1,062,061	1,116,924	1,238,336
Age²						
0 to 5 years	97,016	119,083	83,798	83,303	92,458	100,026
6 to 11 years	420,809	468,316	402,002	419,302	442,550	494,112
12 to 14 years	264,485	294,452	260,949	266,144	277,655	311,360
15 to 17 years	276,333	306,257	284,174	295,446	307,147	335,554
Total	1,058,643	1,188,108	1,030,923	1,064,195	1,119,810	1,241,052
Race						
American Indian or Alaska Native	10,293	13,013	12,997	12,865	15,122	17,592
Asian ³	7,674	9,416	8,354	10,096	11,052	12,658
Black or African American	231,703	251,258	220,655	208,697	212,951	228,662
Native Hawaiian or Other Pacific Islander	2,439	2,860	3,134	3,542	2,471	3,267
White	573,901	652,532	544,553	583,111	606,818	656,551
Other	159,195	180,985	162,024	169,125	196,396	204,991
Total	985,205	1,110,064	951,717	987,436	1,044,810	1,123,721

Continued. See notes at end of table.

Table 2.2a. Individuals aged 17 years and younger with SED, by gender, age, race, and ethnicity: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Ethnicity						
Not of Hispanic or Latino origin	699,459	770,640	649,792	679,721	763,785	797,327
Hispanic or Latino origin	214,012	245,809	228,406	246,301	270,641	279,667
Total	913,471	1,016,449	878,198	926,022	1,034,426	1,076,994
Specific Hispanic or Latino origin						
Mexican	9,321	13,470	16,076	12,851	17,496	16,681
Puerto Rican	8,187	7,486	7,962	7,095	5,713	5,013
Cuban	1,426	1,122	1,147	1,059	1,159	1,204
Other specific	12,422	15,238	14,138	13,169	15,227	15,538
Total	31,356	37,316	39,323	34,174	39,595	38,436

SED = serious emotional disturbance.

¹ The federal definition for SED status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 2.2b. Individuals aged 17 years and younger with SED, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	44	47	45	47	47	48
Total individuals with SED aged 0 to 17 years¹	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	60.3	58.9	57.5	58.0	56.9	56.3
Female	39.7	41.1	42.5	42.0	43.1	43.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age²						
0 to 5 years	9.2	10.0	8.1	7.8	8.3	8.1
6 to 11 years	39.7	39.4	39.0	39.4	39.5	39.8
12 to 14 years	25.0	24.8	25.3	25.0	24.8	25.1
15 to 17 years	26.1	25.8	27.6	27.8	27.4	27.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	1.0	1.2	1.4	1.3	1.4	1.6
Asian ³	†	†	†	1.0	1.1	1.1
Black or African American	23.5	22.6	23.2	21.1	20.4	20.3
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	58.3	58.8	57.2	59.1	58.1	58.4
Other	16.2	16.3	17.0	17.1	18.8	18.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 2.2b. Individuals aged 17 years and younger with SED, by gender, age, race, and ethnicity: Column percent distribution, 2013–2018
(continued)**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Ethnicity						
Not of Hispanic or Latino origin	76.6	75.8	74.0	73.4	73.8	74.0
Hispanic or Latino origin	23.4	24.2	26.0	26.6	26.2	26.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	29.7	36.1	40.9	37.6	44.2	43.4
Puerto Rican	26.1	20.1	20.2	20.8	14.4	13.0
Cuban	4.5	3.0	2.9	3.1	2.9	3.1
Other specific	39.6	40.8	36.0	38.5	38.5	40.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

SED = serious emotional disturbance.

¹ The federal definition for SED status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 2.3a. Individuals aged 17 years and younger, by mental health diagnosis, gender, race, and ethnicity: Number, 2018

Gender, race, and ethnicity	Mental health diagnosis ¹									
	Total	Trauma- and stressor-related disorders				Mood disorders		ODD	PDD	Other mental disorders ¹
		Anxiety disorders	ADHD	Conduct disorder	Bipolar disorders	Depressive disorders				
Total individuals aged 0 to 17 years	1,630,278	386,335	186,058	384,329	75,786	26,214	224,057	98,418	46,303	202,778
Gender										
Male	919,754	188,407	84,280	281,784	53,848	12,744	79,703	66,138	37,455	115,395
Female	705,984	196,771	101,365	101,280	21,675	13,420	143,975	31,945	8,757	86,796
Total	1,625,738	385,178	185,645	383,064	75,523	26,164	223,678	98,083	46,212	202,191
Race										
American Indian or Alaska Native	25,435	7,622	3,240	3,754	751	234	3,613	1,181	428	4,612
Asian ²	15,302	3,469	2,090	2,225	528	211	3,043	441	857	2,438
Black or African American	293,417	60,382	17,855	95,254	20,813	5,021	29,852	26,068	5,852	32,320
Native Hawaiian or Other Pacific Islander	3,997	1,098	622	684	178	34	684	148	89	460
White	914,346	220,941	114,591	214,802	35,114	16,633	126,544	50,150	30,408	105,163
Other	233,318	52,924	28,063	43,443	11,475	2,922	40,071	13,603	5,726	35,091
Total	1,485,815	346,436	166,461	360,162	68,859	25,055	203,807	91,591	43,360	180,084
Ethnicity										
Not of Hispanic or Latino origin	1,106,883	258,876	123,263	278,417	52,227	20,239	141,906	72,025	36,124	123,806
Hispanic or Latino origin	338,031	78,263	40,323	66,238	14,915	4,231	58,094	16,550	7,023	52,394
Total	1,444,914	337,139	163,586	344,655	67,142	24,470	200,000	88,575	43,147	176,200
Specific Hispanic or Latino origin										
Mexican	23,689	7,453	2,663	3,674	928	191	4,948	1,215	346	2,271
Puerto Rican	5,759	1,085	446	1,748	379	91	885	318	142	665
Cuban	1,319	291	116	303	51	15	200	48	34	261
Other specific	19,970	5,518	2,022	3,615	1,061	219	3,787	1,184	278	2,286
Total	50,737	14,347	5,247	9,340	2,419	516	9,820	2,765	800	5,483

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes substance use diagnoses. See Appendix E. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.3b. Individuals aged 17 years and younger, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018

Gender, race, and ethnicity	Mental health diagnosis ¹									
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals aged 0 to 17 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	56.6	48.9	45.4	73.6	71.3	48.7	35.6	67.4	81.1	57.1
Female	43.4	51.1	54.6	26.4	28.7	51.3	64.4	32.6	18.9	42.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	1.7	2.2	1.9	1.0	1.1	†	1.8	1.3	†	2.6
Asian ²	1.0	1.0	1.3	†	†	†	1.5	†	2.0	1.4
Black or African American	19.7	17.4	10.7	26.4	30.2	20.0	14.6	28.5	13.5	17.9
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	†
White	61.5	63.8	68.8	59.6	51.0	66.4	62.1	54.8	70.1	58.4
Other	15.7	15.3	16.9	12.1	16.7	11.7	19.7	14.9	13.2	19.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity										
Not of Hispanic or Latino origin	76.6	76.8	75.4	80.8	77.8	82.7	71.0	81.3	83.7	70.3
Hispanic or Latino origin	23.4	23.2	24.6	19.2	22.2	17.3	29.0	18.7	16.3	29.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican	46.7	51.9	50.8	39.3	38.4	37.0	50.4	43.9	43.3	41.4
Puerto Rican	11.4	7.6	8.5	18.7	15.7	17.6	9.0	11.5	17.8	12.1
Cuban	2.6	2.0	2.2	3.2	2.1	2.9	2.0	1.7	4.3	4.8
Other specific	39.4	38.5	38.5	38.7	43.9	42.4	38.6	42.8	34.8	41.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes substance use diagnoses. See Appendix E. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.3c. Individuals aged 17 years and younger, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals aged 0 to 17 years	100.0	23.7	11.4	23.6	4.6	1.6	13.7	6.0	2.8	12.4
Gender										
Male	100.0	20.5	9.2	30.6	5.9	1.4	8.7	7.2	4.1	12.5
Female	100.0	27.9	14.4	14.3	3.1	1.9	20.4	4.5	1.2	12.3
Total	100.0	23.7	11.4	23.6	4.6	1.6	13.8	6.0	2.8	12.4
Race										
American Indian or Alaska Native	100.0	30.0	12.7	14.8	3.0	†	14.2	4.6	1.7	18.1
Asian ²	100.0	22.7	13.7	14.5	3.5	1.4	19.9	2.9	5.6	15.9
Black or African American	100.0	20.6	6.1	32.5	7.1	1.7	10.2	8.9	2.0	11.0
Native Hawaiian or Other Pacific Islander	100.0	27.5	15.6	17.1	4.5	†	17.1	3.7	2.2	11.5
White	100.0	24.2	12.5	23.5	3.8	1.8	13.8	5.5	3.3	11.5
Other	100.0	22.7	12.0	18.6	4.9	1.3	17.2	5.8	2.5	15.0
Total	100.0	23.3	11.2	24.2	4.6	1.7	13.7	6.2	2.9	12.1
Ethnicity										
Not of Hispanic or Latino origin	100.0	23.4	11.1	25.2	4.7	1.8	12.8	6.5	3.3	11.2
Hispanic or Latino origin	100.0	23.2	11.9	19.6	4.4	1.3	17.2	4.9	2.1	15.5
Total	100.0	23.3	11.3	23.9	4.6	1.7	13.8	6.1	3.0	12.2
Specific Hispanic or Latino origin										
Mexican	100.0	31.5	11.2	15.5	3.9	†	20.9	5.1	1.5	9.6
Puerto Rican	100.0	18.8	7.7	30.4	6.6	1.6	15.4	5.5	2.5	11.5
Cuban	100.0	22.1	8.8	23.0	3.9	1.1	15.2	3.6	2.6	19.8
Other specific	100.0	27.6	10.1	18.1	5.3	1.1	19.0	5.9	1.4	11.4
Total	100.0	28.3	10.3	18.4	4.8	1.0	19.4	5.4	1.6	10.8

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes substance use diagnoses. See Appendix E. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.4a. Individuals aged 17 years and younger with SED, by mental health diagnosis, gender, race, and ethnicity: Number, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years²	1,174,041	263,176	141,028	274,171	53,403	21,667	184,567	70,420	29,603	136,006
Gender										
Male	658,175	129,109	64,806	201,620	38,073	10,498	65,856	47,462	24,181	76,570
Female	513,246	133,517	76,001	71,760	15,197	11,125	118,492	22,748	5,370	59,036
Total	1,171,421	262,626	140,807	273,380	53,270	21,623	184,348	70,210	29,551	135,606
Race										
American Indian or Alaska Native	17,050	5,674	2,349	2,582	512	179	2,684	919	288	1,863
Asian ³	11,834	2,429	1,708	1,714	422	189	2,694	353	600	1,725
Black or African American	216,736	39,708	13,858	72,617	14,770	4,286	25,276	18,217	3,654	24,350
Native Hawaiian or Other Pacific Islander	3,044	849	461	487	146	28	563	129	40	341
White	619,027	140,405	82,021	145,211	22,442	13,281	99,282	34,033	19,314	63,038
Other	194,307	42,942	24,975	31,710	9,630	2,709	37,073	11,071	3,603	30,594
Total	1,061,998	232,007	125,372	254,321	47,922	20,672	167,572	64,722	27,499	121,911
Ethnicity										
Not of Hispanic or Latino origin	756,581	159,127	88,892	191,266	33,968	16,647	112,494	48,680	23,136	82,371
Hispanic or Latino origin	259,019	61,199	32,740	46,047	11,745	3,509	50,449	12,689	4,441	36,200
Total	1,015,600	220,326	121,632	237,313	45,713	20,156	162,943	61,369	27,577	118,571
Specific Hispanic or Latino origin										
Mexican	16,498	5,052	1,714	2,797	710	153	3,474	914	163	1,521
Puerto Rican	4,593	786	346	1,564	285	84	731	203	74	520
Cuban	1,162	250	100	280	46	14	180	39	19	234
Other specific	15,010	3,827	1,441	3,003	786	188	2,981	868	158	1,758
Total	37,263	9,915	3,601	7,644	1,827	439	7,366	2,024	414	4,033

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.4b. Individuals aged 17 years and younger with SED, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018

Gender, race, and ethnicity	Mental health diagnosis ¹									
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	56.2	49.2	46.0	73.8	71.5	48.6	35.7	67.6	81.8	56.5
Female	43.8	50.8	54.0	26.2	28.5	51.4	64.3	32.4	18.2	43.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	1.6	2.4	1.9	1.0	1.1	†	1.6	1.4	1.0	1.5
Asian ³	1.1	1.0	1.4	†	†	†	1.6	†	2.2	1.4
Black or African American	20.4	17.1	11.1	28.6	30.8	20.7	15.1	28.1	13.3	20.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	†
White	58.3	60.5	65.4	57.1	46.8	64.2	59.2	52.6	70.2	51.7
Other	18.3	18.5	19.9	12.5	20.1	13.1	22.1	17.1	13.1	25.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity										
Not of Hispanic or Latino origin	74.5	72.2	73.1	80.6	74.3	82.6	69.0	79.3	83.9	69.5
Hispanic or Latino origin	25.5	27.8	26.9	19.4	25.7	17.4	31.0	20.7	16.1	30.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin										
Mexican	44.3	51.0	47.6	36.6	38.9	34.9	47.2	45.2	39.4	37.7
Puerto Rican	12.3	7.9	9.6	20.5	15.6	19.1	9.9	10.0	17.9	12.9
Cuban	3.1	2.5	2.8	3.7	2.5	3.2	2.4	1.9	4.6	5.8
Other specific	40.3	38.6	40.0	39.3	43.0	42.8	40.5	42.9	38.2	43.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.4c. Individuals aged 17 years and younger with SED, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years²	100.0	22.4	12.0	23.4	4.5	1.8	15.7	6.0	2.5	11.6
Gender										
Male	100.0	19.6	9.8	30.6	5.8	1.6	10.0	7.2	3.7	11.6
Female	100.0	26.0	14.8	14.0	3.0	2.2	23.1	4.4	1.0	11.5
Total	100.0	22.4	12.0	23.3	4.5	1.8	15.7	6.0	2.5	11.6
Race										
American Indian or Alaska Native	100.0	33.3	13.8	15.1	3.0	1.0	15.7	5.4	1.7	10.9
Asian ³	100.0	20.5	14.4	14.5	3.6	1.6	22.8	3.0	5.1	14.6
Black or African American	100.0	18.3	6.4	33.5	6.8	2.0	11.7	8.4	1.7	11.2
Native Hawaiian or Other Pacific Islander	100.0	27.9	15.1	16.0	4.8	†	18.5	4.2	1.3	11.2
White	100.0	22.7	13.2	23.5	3.6	2.1	16.0	5.5	3.1	10.2
Other	100.0	22.1	12.9	16.3	5.0	1.4	19.1	5.7	1.9	15.7
Total	100.0	21.8	11.8	23.9	4.5	1.9	15.8	6.1	2.6	11.5
Ethnicity										
Not of Hispanic or Latino origin	100.0	21.0	11.7	25.3	4.5	2.2	14.9	6.4	3.1	10.9
Hispanic or Latino origin	100.0	23.6	12.6	17.8	4.5	1.4	19.5	4.9	1.7	14.0
Total	100.0	21.7	12.0	23.4	4.5	2.0	16.0	6.0	2.7	11.7
Specific Hispanic or Latino origin										
Mexican	100.0	30.6	10.4	17.0	4.3	†	21.1	5.5	†	9.2
Puerto Rican	100.0	17.1	7.5	34.1	6.2	1.8	15.9	4.4	1.6	11.3
Cuban	100.0	21.5	8.6	24.1	4.0	1.2	15.5	3.4	1.6	20.1
Other specific	100.0	25.5	9.6	20.0	5.2	1.3	19.9	5.8	1.1	11.7
Total	100.0	26.6	9.7	20.5	4.9	1.2	19.8	5.4	1.1	10.8

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.5a. Individuals aged 17 years and younger, by mental health diagnosis, living arrangement, and SED status: Number, 2018

Living arrangement and SED status	Mental health diagnosis ¹									
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals aged 0 to 17 years	1,630,278	386,335	186,058	384,329	75,786	26,214	224,057	98,418	46,303	202,778
Living arrangement										
Homeless	4,780	1,877	439	656	182	77	669	250	40	590
Foster home/foster care	35,535	16,696	3,290	4,609	1,137	436	3,015	1,598	374	4,380
Residential care	14,163	3,045	1,038	2,088	991	610	2,394	957	199	2,841
Private residence	850,319	195,868	101,405	200,789	38,148	14,570	129,536	52,955	21,286	95,762
Other	20,483	5,084	2,087	2,480	1,591	537	3,743	1,136	213	3,612
Total	925,280	222,570	108,259	210,622	42,049	16,230	139,357	56,896	22,112	107,185
SED status²										
SED	1,174,041	263,176	141,028	274,171	53,403	21,667	184,567	70,420	29,603	136,006
Not SED	407,630	110,549	38,581	98,189	20,818	3,947	33,970	26,551	13,296	61,729
Total	1,581,671	373,725	179,609	372,360	74,221	25,614	218,537	96,971	42,899	197,735

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.5b. Individuals aged 17 years and younger, by mental health diagnosis, living arrangement, and SED status: Column percent distribution, 2018

Living arrangement and SED status	Mental health diagnosis ¹									
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals aged 0 to 17 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement										
Homeless	†	†	†	†	†	†	†	†	†	†
Foster home/foster care	3.8	7.5	3.0	2.2	2.7	2.7	2.2	2.8	1.7	4.1
Residential care	1.5	1.4	†	†	2.4	3.8	1.7	1.7	†	2.7
Private residence	91.9	88.0	93.7	95.3	90.7	89.8	93.0	93.1	96.3	89.3
Other	2.2	2.3	1.9	1.2	3.8	3.3	2.7	2.0	†	3.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SED status²										
SED	74.2	70.4	78.5	73.6	72.0	84.6	84.5	72.6	69.0	68.8
Not SED	25.8	29.6	21.5	26.4	28.0	15.4	15.5	27.4	31.0	31.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.5c. Individuals aged 17 years and younger, by mental health diagnosis, living arrangement, and SED status: Row percent distribution, 2018

<i>Living arrangement and SED status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>								
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Conduct disorder</i>	<i>Mood disorders</i>		<i>ODD</i>	<i>PDD</i>	<i>Other mental disorders¹</i>
						<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 0 to 17 years	100.0	23.7	11.4	23.6	4.6	1.6	13.7	6.0	2.8	12.4
Living arrangement										
Homeless	100.0	39.3	9.2	13.7	3.8	1.6	14.0	5.2	†	12.3
Foster home/foster care	100.0	47.0	9.3	13.0	3.2	1.2	8.5	4.5	1.1	12.3
Residential care	100.0	21.5	7.3	14.7	7.0	4.3	16.9	6.8	1.4	20.1
Private residence	100.0	23.0	11.9	23.6	4.5	1.7	15.2	6.2	2.5	11.3
Other	100.0	24.8	10.2	12.1	7.8	2.6	18.3	5.5	1.0	17.6
Total	100.0	24.1	11.7	22.8	4.5	1.8	15.1	6.1	2.4	11.6
SED status²										
SED	100.0	22.4	12.0	23.4	4.5	1.8	15.7	6.0	2.5	11.6
Not SED	100.0	27.1	9.5	24.1	5.1	†	8.3	6.5	3.3	15.1
Total	100.0	23.6	11.4	23.5	4.7	1.6	13.8	6.1	2.7	12.5

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.6a. Individuals aged 17 years and younger with SED, by mental health diagnosis and living arrangement: Number, 2018

Living arrangement	Mental health diagnosis ¹									
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years²	1,174,041	263,176	141,028	274,171	53,403	21,667	184,567	70,420	29,603	136,006
Living arrangement										
Homeless	3,966	1,557	355	589	156	60	567	205	23	454
Foster home/foster care	32,249	15,112	3,100	4,254	1,043	387	2,808	1,413	273	3,859
Residential care	13,041	2,755	954	1,969	929	587	2,282	867	148	2,550
Private residence	701,104	154,768	81,803	175,796	32,377	12,913	110,948	44,374	11,658	76,467
Other	18,104	4,411	1,872	2,189	1,360	469	3,401	970	177	3,255
Total	768,464	178,603	88,084	184,797	35,865	14,416	120,006	47,829	12,279	86,585

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.6b. Individuals aged 17 years and younger with SED, by mental health diagnosis and living arrangement: Column percent distribution, 2018

Living arrangement	Mental health diagnosis ¹									
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement										
Homeless	†	†	†	†	†	†	†	†	†	†
Foster home/foster care	4.2	8.5	3.5	2.3	2.9	2.7	2.3	3.0	2.2	4.5
Residential care	1.7	1.5	1.1	1.1	2.6	4.1	1.9	1.8	1.2	2.9
Private residence	91.2	86.7	92.9	95.1	90.3	89.6	92.5	92.8	94.9	88.3
Other	2.4	2.5	2.1	1.2	3.8	3.3	2.8	2.0	1.4	3.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.6c. Individuals aged 17 years and younger with SED, by mental health diagnosis and living arrangement: Row percent distribution, 2018

Living arrangement	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years²	100.0	22.4	12.0	23.4	4.5	1.8	15.7	6.0	2.5	11.6
Living arrangement										
Homeless	100.0	39.3	9.0	14.9	3.9	1.5	14.3	5.2	†	11.4
Foster home/foster care	100.0	46.9	9.6	13.2	3.2	1.2	8.7	4.4	†	12.0
Residential care	100.0	21.1	7.3	15.1	7.1	4.5	17.5	6.6	1.1	19.6
Private residence	100.0	22.1	11.7	25.1	4.6	1.8	15.8	6.3	1.7	10.9
Other	100.0	24.4	10.3	12.1	7.5	2.6	18.8	5.4	†	18.0
Total	100.0	23.2	11.5	24.0	4.7	1.9	15.6	6.2	1.6	11.3

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.7a. Individuals aged 17 years and younger, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018

Timing of admission and mental health service setting	Mental health diagnosis ¹									
	Total	Trauma- and stressor-related disorders				Mood disorders		ODD	PDD	Other mental disorders ¹
		Anxiety disorders	ADHD	Conduct disorder	Bipolar disorder	Depressive disorder				
Total individuals aged 0 to 17 years	1,630,278	386,335	186,058	384,329	75,786	26,214	224,057	98,418	46,303	202,778
Timing of admission										
New individuals	899,736	247,043	102,958	172,281	42,103	10,427	136,844	49,418	16,217	122,445
Continuing individuals	730,542	139,292	83,100	212,048	33,683	15,787	87,213	49,000	30,086	80,333
Mental health service setting^{2,3}										
State psychiatric hospital	7,793	892	459	758	360	601	2,213	333	102	2,075
Community-based program	1,617,278	383,401	185,415	383,248	74,790	25,585	220,888	97,621	46,178	200,152
Residential treatment center	17,760	3,685	1,061	2,368	1,244	1,079	3,516	1,418	223	3,166
Other psychiatric inpatient	33,811	4,750	2,890	4,343	847	1,988	10,592	1,836	498	6,067
Institutions under the justice system	5,062	914	313	427	518	174	829	323	15	1,549

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.7b. Individuals aged 17 years and younger, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018

Timing of admission and mental health service setting	Mental health diagnosis ¹									
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals aged 0 to 17 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission										
New individuals	55.2	63.9	55.3	44.8	55.6	39.8	61.1	50.2	35.0	60.4
Continuing individuals	44.8	36.1	44.7	55.2	44.4	60.2	38.9	49.8	65.0	39.6
Mental health service setting^{2,3}										
State psychiatric hospital	†	†	†	†	†	2.3	†	†	†	1.0
Community-based program	99.2	99.2	99.7	99.7	98.7	97.6	98.6	99.2	99.7	98.7
Residential treatment center	1.1	†	†	†	1.6	4.1	1.6	1.4	†	1.6
Other psychiatric inpatient	2.1	1.2	1.6	1.1	1.1	7.6	4.7	1.9	1.1	3.0
Institutions under the justice system	†	†	†	†	†	†	†	†	†	†

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages for timing of admission may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.7c. Individuals aged 17 years and younger, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals aged 0 to 17 years	100.0	23.7	11.4	23.6	4.6	1.6	13.7	6.0	2.8	12.4
Timing of admission										
New individuals	100.0	27.5	11.4	19.1	4.7	1.2	15.2	5.5	1.8	13.6
Continuing individuals	100.0	19.1	11.4	29.0	4.6	2.2	11.9	6.7	4.1	11.0
Mental health service setting^{2,3}										
State psychiatric hospital	100.0	11.4	5.9	9.7	4.6	7.7	28.4	4.3	1.3	26.6
Community-based program	100.0	23.7	11.5	23.7	4.6	1.6	13.7	6.0	2.9	12.4
Residential treatment center	100.0	20.7	6.0	13.3	7.0	6.1	19.8	8.0	1.3	17.8
Other psychiatric inpatient	100.0	14.0	8.5	12.8	2.5	5.9	31.3	5.4	1.5	17.9
Institutions under the justice system	100.0	18.1	6.2	8.4	10.2	3.4	16.4	6.4	†	30.6

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.8a. Individuals aged 17 years and younger with SED, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018

Timing of admission and mental health service setting	Mental health diagnosis ¹									
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years²	1,174,041	263,176	141,028	274,171	53,403	21,667	184,567	70,420	29,603	136,006
Timing of admission										
New individuals	621,983	164,205	75,013	120,567	28,516	8,528	110,190	34,273	8,748	71,943
Continuing individuals	552,058	98,971	66,015	153,604	24,887	13,139	74,377	36,147	20,855	64,063
Mental health service setting^{3,4}										
State psychiatric hospital	6,789	740	386	670	281	575	1,998	221	91	1,827
Community-based program	1,163,934	261,006	140,506	273,490	52,591	21,176	181,898	69,952	29,519	133,796
Residential treatment center	13,592	2,679	866	1,660	984	792	3,027	954	181	2,449
Other psychiatric inpatient	25,356	3,168	2,238	2,882	534	1,458	9,091	1,143	324	4,518
Institutions under the justice system	4,979	884	308	425	508	171	812	320	13	1,538

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.8b. Individuals aged 17 years and younger with SED, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018

Timing of admission and mental health service setting	Mental health diagnosis ¹									
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission										
New individuals	53.0	62.4	53.2	44.0	53.4	39.4	59.7	48.7	29.6	52.9
Continuing individuals	47.0	37.6	46.8	56.0	46.6	60.6	40.3	51.3	70.4	47.1
Mental health service setting^{3,4}										
State psychiatric hospital	†	†	†	†	†	2.7	1.1	†	†	1.3
Community-based program	99.1	99.2	99.6	99.8	98.5	97.7	98.6	99.3	99.7	98.4
Residential treatment center	1.2	1.0	†	†	1.8	3.7	1.6	1.4	†	1.8
Other psychiatric inpatient	2.2	1.2	1.6	1.1	†	6.7	4.9	1.6	1.1	3.3
Institutions under the justice system	†	†	†	†	†	†	†	†	†	1.1

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

³ Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages for timing of admission may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.8c. Individuals aged 17 years and younger with SED, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Conduct disorder	Mood disorders		ODD	PDD	Other mental disorders ¹
						Bipolar disorders	Depressive disorders			
Total individuals with SED aged 0 to 17 years²	100.0	22.4	12.0	23.4	4.5	1.8	15.7	6.0	2.5	11.6
Timing of admission										
New individuals	100.0	26.4	12.1	19.4	4.6	1.4	17.7	5.5	1.4	11.6
Continuing individuals	100.0	17.9	12.0	27.8	4.5	2.4	13.5	6.5	3.8	11.6
Mental health service setting^{3,4}										
State psychiatric hospital	100.0	10.9	5.7	9.9	4.1	8.5	29.4	3.3	1.3	26.9
Community-based program	100.0	22.4	12.1	23.5	4.5	1.8	15.6	6.0	2.5	11.5
Residential treatment center	100.0	19.7	6.4	12.2	7.2	5.8	22.3	7.0	1.3	18.0
Other psychiatric inpatient	100.0	12.5	8.8	11.4	2.1	5.8	35.9	4.5	1.3	17.8
Institutions under the justice system	100.0	17.8	6.2	8.5	10.2	3.4	16.3	6.4	†	30.9

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; PDD = pervasive developmental disorders; SED = serious emotional disturbance.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SED status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 2.9. Individuals aged 17 years and younger, by mental health service setting and SED status: Number and percent distribution, 2018

<i>Mental health service setting</i>	<i>SED status¹</i>								
	<i>Number of individuals</i>			<i>Column percent of individuals</i>			<i>Row percent distribution of individuals</i>		
	<i>Total</i>	<i>SED</i>	<i>Not SED</i>	<i>Total</i>	<i>SED</i>	<i>Not SED</i>	<i>Total</i>	<i>SED</i>	<i>Not SED</i>
Total individuals aged 0 to 17 years	1,722,207	1,241,052	481,155	100.0	100.0	100.0	100.0	72.1	27.9
Mental health service setting^{2,3}									
State psychiatric hospital	8,036	7,108	928	†	†	†	100.0	88.5	11.5
Community-based program	1,708,661	1,230,049	478,612	99.2	99.1	99.5	100.0	72.0	28.0
Residential treatment center	17,499	13,772	3,727	1.0	1.1	†	100.0	78.7	21.3
Other psychiatric inpatient	35,754	26,130	9,624	2.1	2.1	2.0	100.0	73.1	26.9
Institutions under the justice system	5,986	5,888	98	†	†	†	100.0	98.4	1.6

† Less than 1 percent of individuals.

SED = serious emotional disturbance.

¹ The federal definition for SED status can be found in Appendix D.

² Individuals could receive services in more than one mental health service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

**Table 3.1a. Individuals aged 18 to 24 years, by gender, age, race, ethnicity, education, veteran status, and SMI status:
Number, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals aged 18 to 24 years	594,733	596,043	539,439	593,875	613,385	645,452
Gender						
Male	292,762	294,354	268,232	286,861	295,369	305,519
Female	301,153	300,224	269,807	304,277	315,454	337,271
Total	593,915	594,578	538,039	591,138	610,823	642,790
Age¹						
18 to 20 years	268,477	259,939	234,919	260,296	268,132	288,195
21 to 24 years	326,256	336,104	304,520	333,579	345,253	357,257
Total	594,733	596,043	539,439	593,875	613,385	645,452
Race						
American Indian or Alaska Native	7,250	8,304	8,332	8,705	8,891	10,042
Asian ²	5,719	6,366	6,580	7,814	8,641	9,542
Black or African American	117,471	118,834	106,406	114,641	116,267	119,419
Native Hawaiian or Other Pacific Islander	1,272	1,366	1,530	1,746	1,596	1,908
White	363,434	362,010	319,329	354,346	368,443	387,978
Other	53,456	54,166	49,945	57,120	61,412	67,959
Total	548,602	551,046	492,122	544,372	565,250	596,848
Ethnicity						
Not of Hispanic or Latino origin	415,140	410,879	364,574	395,350	453,185	465,569
Hispanic or Latino origin	77,857	80,979	80,301	90,885	98,132	103,558
Total	492,997	491,858	444,875	486,235	551,317	569,127
Specific Hispanic or Latino origin						
Mexican	3,324	4,731	6,097	5,098	6,489	6,383
Puerto Rican	2,823	4,302	4,563	4,017	4,787	2,764
Cuban	558	589	614	605	645	680
Other specific	4,876	6,504	5,986	6,054	7,208	7,204
Total	11,581	16,126	17,260	15,774	19,129	17,031

Continued. See notes at end of table.

**Table 3.1a. Individuals aged 18 to 24 years, by gender, age, race, ethnicity, education, veteran status, and SMI status:
Number, 2013–2018 (continued)**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Highest school grade completed						
0 to 8	31,400	31,182	32,851	31,666	27,800	25,707
9 to 11	67,877	66,687	66,177	65,693	71,659	76,769
12 (or GED)	85,242	97,470	98,554	101,779	114,033	134,201
More than 12	21,703	25,157	26,336	28,004	34,079	39,919
Total	206,222	220,496	223,918	227,142	247,571	276,596
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2,483	2,114
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	211,336	219,814
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	213,819	221,928
SMI status³						
SMI	336,335	357,509	318,241	351,981	375,498	412,774
Not SMI	180,265	174,672	163,636	177,970	179,004	176,025
Total	516,600	532,181	481,877	529,951	554,502	588,799

n/c Not collected.

SMI = serious mental illness; GED = general equivalency diploma.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 3.1b. Individuals aged 18 to 24 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals aged 18 to 24 years	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	49.3	49.5	49.9	48.5	48.4	47.5
Female	50.7	50.5	50.1	51.5	51.6	52.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
18 to 20 years	45.1	43.6	43.5	43.8	43.7	44.7
21 to 24 years	54.9	56.4	56.5	56.2	56.3	55.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	1.3	1.5	1.7	1.6	1.6	1.7
Asian ²	1.0	1.2	1.3	1.4	1.5	1.6
Black or African American	21.4	21.6	21.6	21.1	20.6	20.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	66.2	65.7	64.9	65.1	65.2	65.0
Other	9.7	9.8	10.1	10.5	10.9	11.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	84.2	83.5	81.9	81.3	82.2	81.8
Hispanic or Latino origin	15.8	16.5	18.1	18.7	17.8	18.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	28.7	29.3	35.3	32.3	33.9	37.5
Puerto Rican	24.4	26.7	26.4	25.5	25.0	16.2
Cuban	4.8	3.7	3.6	3.8	3.4	4.0
Other specific	42.1	40.3	34.7	38.4	37.7	42.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 3.1b. Individuals aged 18 to 24 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Highest school grade completed						
0 to 8	15.2	14.1	14.7	13.9	11.2	9.3
9 to 11	32.9	30.2	29.6	28.9	28.9	27.8
12 (or GED)	41.3	44.2	44.0	44.8	46.1	48.5
More than 12	10.5	11.4	11.8	12.3	13.8	14.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	1.2	†
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	98.8	99.0
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0
SMI status³						
SMI	65.1	67.2	66.0	66.4	67.7	70.1
Not SMI	34.9	32.8	34.0	33.6	32.3	29.9
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

SMI = serious mental illness; GED = general equivalency diploma.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 3.2a. Individuals aged 18 to 24 years with SMI, by gender, age, race, ethnicity, education, and veteran status:
Number, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	44	47	45	46	46	47
Total individuals with SMI aged 18 to 24 years¹	336,335	357,509	318,241	351,981	375,498	412,774
Gender						
Male	164,928	176,593	159,011	174,320	182,370	194,162
Female	170,830	180,105	158,496	176,532	192,079	217,643
Total	335,758	356,698	317,507	350,852	374,449	411,805
Age²						
18 to 20 years	142,538	149,085	130,146	149,760	160,666	182,989
21 to 24 years	193,797	208,424	188,095	202,221	214,832	229,785
Total	336,335	357,509	318,241	351,981	375,498	412,774
Race						
American Indian or Alaska Native	2,919	4,197	4,212	4,434	5,048	5,615
Asian ³	3,579	4,245	4,184	4,754	5,803	6,427
Black or African American	70,715	74,925	66,542	69,768	71,207	77,456
Native Hawaiian or Other Pacific Islander	809	896	992	1,135	1,000	1,179
White	200,362	210,001	182,265	208,494	219,348	239,691
Other	33,374	36,276	33,548	36,941	46,521	52,417
Total	311,758	330,540	291,743	325,526	348,927	382,785
Ethnicity						
Not of Hispanic or Latino origin	228,879	242,690	211,638	233,056	263,334	282,615
Hispanic or Latino origin	48,707	54,702	53,255	60,300	70,424	75,938
Total	277,586	297,392	264,893	293,356	333,758	358,553
Specific Hispanic or Latino origin						
Mexican	1,795	2,783	3,540	2,955	4,065	4,118
Puerto Rican	1,745	2,004	2,100	2,162	2,409	1,941
Cuban	424	406	418	521	556	615
Other specific	3,353	4,055	3,758	4,163	4,949	5,303
Total	7,317	9,248	9,816	9,801	11,979	11,977

Continued. See notes at end of table.

**Table 3.2a. Individuals aged 18 to 24 years with SMI, by gender, age, race, ethnicity, education, and veteran status:
Number, 2013–2018 (continued)**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Highest school grade completed						
0 to 8	12,970	15,426	14,598	14,476	15,012	15,944
9 to 11	35,728	41,499	39,881	42,971	49,124	55,832
12 (or GED)	50,045	63,927	64,504	70,314	80,283	96,856
More than 12	12,564	16,610	17,335	19,534	24,162	28,742
Total	111,307	137,462	136,318	147,295	168,581	197,374
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	1,694	1,498
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	116,037	144,204
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	117,731	145,702

n/c Not collected.

SMI = serious mental illness; GED = general equivalency diploma.

¹ The federal definition for SMI status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 3.2b. Individuals aged 18 to 24 years with SMI, by gender, age, race, ethnicity, education, and veteran status:
Column percent distribution, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	44	47	45	46	46	47
Total individuals with SMI aged 18 to 24 years¹	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	49.1	49.5	50.1	49.7	48.7	47.1
Female	50.9	50.5	49.9	50.3	51.3	52.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age²						
18 to 20 years	42.4	41.7	40.9	42.5	42.8	44.3
21 to 24 years	57.6	58.3	59.1	57.5	57.2	55.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	†	1.3	1.4	1.4	1.4	1.5
Asian ³	1.1	1.3	1.4	1.5	1.7	1.7
Black or African American	22.7	22.7	22.8	21.4	20.4	20.2
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	64.3	63.5	62.5	64.0	62.9	62.6
Other	10.7	11.0	11.5	11.3	13.3	13.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	82.5	81.6	79.9	79.4	78.9	78.8
Hispanic or Latino origin	17.5	18.4	20.1	20.6	21.1	21.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	24.5	30.1	36.1	30.1	33.9	34.4
Puerto Rican	23.8	21.7	21.4	22.1	20.1	16.2
Cuban	5.8	4.4	4.3	5.3	4.6	5.1
Other specific	45.8	43.8	38.3	42.5	41.3	44.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 3.2b. Individuals aged 18 to 24 years with SMI, by gender, age, race, ethnicity, education, and veteran status:
Column percent distribution, 2013–2018 (continued)**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Highest school grade completed						
0 to 8	11.7	11.2	10.7	9.8	8.9	8.1
9 to 11	32.1	30.2	29.3	29.2	29.1	28.3
12 (or GED)	45.0	46.5	47.3	47.7	47.6	49.1
More than 12	11.3	12.1	12.7	13.3	14.3	14.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	1.4	1.0
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	98.6	99.0
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

SMI = serious mental illness; GED = general equivalency diploma.

¹ The federal definition for SMI status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 3.3a. Individuals aged 18 to 24 years, by mental health diagnosis, gender, race, and ethnicity: Number, 2018

Gender, race, and ethnicity	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
		Anxiety disorders	ADHD	Bipolar disorders	Depressive disorders				
Total individuals aged 18 to 24 years	557,478	70,164	80,312	31,550	61,603	165,259	6,454	50,990	91,146
Gender									
Male	257,941	26,546	29,451	21,021	27,095	60,682	2,746	37,137	53,263
Female	297,454	43,279	50,647	10,383	34,185	104,052	3,672	13,718	37,518
Total	555,395	69,825	80,098	31,404	61,280	164,734	6,418	50,855	90,781
Race									
American Indian or Alaska Native	8,905	1,511	1,397	353	615	2,675	88	687	1,579
Asian ²	8,082	1,005	1,189	283	630	2,415	105	1,164	1,291
Black or African American	101,874	12,947	8,082	6,500	12,275	27,961	1,374	16,493	16,242
Native Hawaiian or Other Pacific Islander	1,736	301	288	59	120	539	21	198	210
White	335,230	41,073	55,485	19,942	39,947	102,804	3,957	21,111	50,911
Other	61,670	7,240	7,376	2,471	5,278	17,927	543	7,575	13,260
Total	517,497	64,077	73,817	29,608	58,865	154,321	6,088	47,228	83,493
Ethnicity									
Not of Hispanic or Latino origin	403,478	51,015	57,128	23,800	48,287	119,402	5,045	35,455	63,346
Hispanic or Latino origin	89,110	10,980	11,850	3,345	8,053	27,852	683	10,337	16,010
Total	492,588	61,995	68,978	27,145	56,340	147,254	5,728	45,792	79,356
Specific Hispanic or Latino origin									
Mexican	6,050	1,120	843	164	461	2,088	45	578	751
Puerto Rican	2,438	351	230	156	272	692	16	307	414
Cuban	633	68	70	44	74	156	2	79	140
Other specific	6,605	1,114	711	274	680	2,082	56	708	980
Total	15,726	2,653	1,854	638	1,487	5,018	119	1,672	2,285

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.3b. Individuals aged 18 to 24 years, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018

Gender, race, and ethnicity	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 18 to 24 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	46.4	38.0	36.8	66.9	44.2	36.8	42.8	73.0	58.7
Female	53.6	62.0	63.2	33.1	55.8	63.2	57.2	27.0	41.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	1.7	2.4	1.9	1.2	1.0	1.7	1.4	1.5	1.9
Asian ²	1.6	1.6	1.6	†	1.1	1.6	1.7	2.5	1.5
Black or African American	19.7	20.2	10.9	22.0	20.9	18.1	22.6	34.9	19.5
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	64.8	64.1	75.2	67.4	67.9	66.6	65.0	44.7	61.0
Other	11.9	11.3	10.0	8.3	9.0	11.6	8.9	16.0	15.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity									
Not of Hispanic or Latino origin	81.9	82.3	82.8	87.7	85.7	81.1	88.1	77.4	79.8
Hispanic or Latino origin	18.1	17.7	17.2	12.3	14.3	18.9	11.9	22.6	20.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	38.5	42.2	45.5	25.7	31.0	41.6	37.8	34.6	32.9
Puerto Rican	15.5	13.2	12.4	24.5	18.3	13.8	13.4	18.4	18.1
Cuban	4.0	2.6	3.8	6.9	5.0	3.1	1.7	4.7	6.1
Other specific	42.0	42.0	38.3	42.9	45.7	41.5	47.1	42.3	42.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.3c. Individuals aged 18 to 24 years, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 18 to 24 years	100.0	12.6	14.4	5.7	11.1	29.6	1.2	9.1	16.3
Gender									
Male	100.0	10.3	11.4	8.1	10.5	23.5	1.1	14.4	20.6
Female	100.0	14.5	17.0	3.5	11.5	35.0	1.2	4.6	12.6
Total	100.0	12.6	14.4	5.7	11.0	29.7	1.2	9.2	16.3
Race									
American Indian or Alaska Native	100.0	17.0	15.7	4.0	6.9	30.0	†	7.7	17.7
Asian ²	100.0	12.4	14.7	3.5	7.8	29.9	1.3	14.4	16.0
Black or African American	100.0	12.7	7.9	6.4	12.0	27.4	1.3	16.2	15.9
Native Hawaiian or Other Pacific Islander	100.0	17.3	16.6	3.4	6.9	31.0	1.2	11.4	12.1
White	100.0	12.3	16.6	5.9	11.9	30.7	1.2	6.3	15.2
Other	100.0	11.7	12.0	4.0	8.6	29.1	†	12.3	21.5
Total	100.0	12.4	14.3	5.7	11.4	29.8	1.2	9.1	16.1
Ethnicity									
Not of Hispanic or Latino origin	100.0	12.6	14.2	5.9	12.0	29.6	1.3	8.8	15.7
Hispanic or Latino origin	100.0	12.3	13.3	3.8	9.0	31.3	†	11.6	18.0
Total	100.0	12.6	14.0	5.5	11.4	29.9	1.2	9.3	16.1
Specific Hispanic or Latino origin									
Mexican	100.0	18.5	13.9	2.7	7.6	34.5	†	9.6	12.4
Puerto Rican	100.0	14.4	9.4	6.4	11.2	28.4	†	12.6	17.0
Cuban	100.0	10.7	11.1	7.0	11.7	24.6	†	12.5	22.1
Other specific	100.0	16.9	10.8	4.1	10.3	31.5	†	10.7	14.8
Total	100.0	16.9	11.8	4.1	9.5	31.9	†	10.6	14.5

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.4a. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Number, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years²	380,549	37,179	47,073	14,853	50,397	129,048	4,333	45,676	51,990
Gender									
Male	176,785	13,849	17,359	9,938	22,365	47,804	1,808	33,327	30,335
Female	202,865	23,243	29,643	4,853	27,865	81,009	2,511	12,259	21,482
Total	379,650	37,092	47,002	14,791	50,230	128,813	4,319	45,586	51,817
Race									
American Indian or Alaska Native	5,378	798	759	157	452	1,863	53	553	743
Asian ³	6,055	582	757	132	561	1,989	53	1,077	904
Black or African American	71,731	5,861	4,487	3,059	10,501	23,054	1,024	14,752	8,993
Native Hawaiian or Other Pacific Islander	1,116	170	152	29	93	370	8	165	129
White	219,331	21,231	31,396	9,014	32,015	77,847	2,566	18,591	26,671
Other	49,467	4,943	5,279	1,286	4,676	15,617	395	7,197	10,074
Total	353,078	33,585	42,830	13,677	48,298	120,740	4,099	42,335	47,514
Ethnicity									
Not of Hispanic or Latino origin	262,631	24,528	31,098	9,819	39,012	90,422	3,415	31,355	32,982
Hispanic or Latino origin	67,763	6,971	7,148	1,506	6,960	23,285	459	9,655	11,779
Total	330,394	31,499	38,246	11,325	45,972	113,707	3,874	41,010	44,761
Specific Hispanic or Latino origin									
Mexican	4,062	614	375	80	394	1,779	33	514	273
Puerto Rican	1,836	215	162	99	244	604	11	291	210
Cuban	576	53	55	40	70	154	2	79	123
Other specific	5,045	714	412	166	623	1,812	37	663	618
Total	11,519	1,596	1,004	385	1,331	4,349	83	1,547	1,224

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.4b. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	46.6	37.3	36.9	67.2	44.5	37.1	41.9	73.1	58.5
Female	53.4	62.7	63.1	32.8	55.5	62.9	58.1	26.9	41.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	1.5	2.4	1.8	1.1	†	1.5	1.3	1.3	1.6
Asian ³	1.7	1.7	1.8	†	1.2	1.6	1.3	2.5	1.9
Black or African American	20.3	17.5	10.5	22.4	21.7	19.1	25.0	34.8	18.9
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	62.1	63.2	73.3	65.9	66.3	64.5	62.6	43.9	56.1
Other	14.0	14.7	12.3	9.4	9.7	12.9	9.6	17.0	21.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity									
Not of Hispanic or Latino origin	79.5	77.9	81.3	86.7	84.9	79.5	88.2	76.5	73.7
Hispanic or Latino origin	20.5	22.1	18.7	13.3	15.1	20.5	11.8	23.5	26.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	35.3	38.5	37.4	20.8	29.6	40.9	39.8	33.2	22.3
Puerto Rican	15.9	13.5	16.1	25.7	18.3	13.9	13.3	18.8	17.2
Cuban	5.0	3.3	5.5	10.4	5.3	3.5	2.4	5.1	10.0
Other specific	43.8	44.7	41.0	43.1	46.8	41.7	44.6	42.9	50.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.4c. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years²	100.0	9.8	12.4	3.9	13.2	33.9	1.1	12.0	13.7
Gender									
Male	100.0	7.8	9.8	5.6	12.7	27.0	1.0	18.9	17.2
Female	100.0	11.5	14.6	2.4	13.7	39.9	1.2	6.0	10.6
Total	100.0	9.8	12.4	3.9	13.2	33.9	1.1	12.0	13.6
Race									
American Indian or Alaska Native	100.0	14.8	14.1	2.9	8.4	34.6	†	10.3	13.8
Asian ³	100.0	9.6	12.5	2.2	9.3	32.8	†	17.8	14.9
Black or African American	100.0	8.2	6.3	4.3	14.6	32.1	1.4	20.6	12.5
Native Hawaiian or Other Pacific Islander	100.0	15.2	13.6	2.6	8.3	33.2	†	14.8	11.6
White	100.0	9.7	14.3	4.1	14.6	35.5	1.2	8.5	12.2
Other	100.0	10.0	10.7	2.6	9.5	31.6	†	14.5	20.4
Total	100.0	9.5	12.1	3.9	13.7	34.2	1.2	12.0	13.5
Ethnicity									
Not of Hispanic or Latino origin	100.0	9.3	11.8	3.7	14.9	34.4	1.3	11.9	12.6
Hispanic or Latino origin	100.0	10.3	10.5	2.2	10.3	34.4	†	14.2	17.4
Total	100.0	9.5	11.6	3.4	13.9	34.4	1.2	12.4	13.5
Specific Hispanic or Latino origin									
Mexican	100.0	15.1	9.2	2.0	9.7	43.8	†	12.7	6.7
Puerto Rican	100.0	11.7	8.8	5.4	13.3	32.9	†	15.8	11.4
Cuban	100.0	9.2	9.5	6.9	12.2	26.7	†	13.7	21.4
Other specific	100.0	14.2	8.2	3.3	12.3	35.9	†	13.1	12.2
Total	100.0	13.9	8.7	3.3	11.6	37.8	†	13.4	10.6

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.5a. Individuals aged 18 to 24 years, by mental health diagnosis, education, and marital status: Number, 2018

<i>Education and marital status</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 18 to 24 years	557,478	70,164	80,312	31,550	61,603	165,259	6,454	50,990	91,146
Highest school grade completed									
0 to 8	24,286	3,407	4,389	2,403	2,095	5,710	252	1,628	4,402
9 to 11	71,170	10,027	8,556	4,709	8,209	21,325	845	6,807	10,692
12 (or GED)	123,564	15,307	15,523	5,528	16,502	40,343	1,520	12,608	16,233
More than 12	36,787	4,559	5,608	1,172	4,813	14,107	421	3,006	3,101
Total	255,807	33,300	34,076	13,812	31,619	81,485	3,038	24,049	34,428
Marital status									
Never married	281,209	33,973	33,534	16,575	36,107	88,086	2,726	27,281	42,927
Currently married	13,596	2,287	1,940	343	1,882	4,977	167	563	1,437
Separated	3,501	598	351	50	583	1,323	44	189	363
Divorced	2,234	377	218	38	354	820	40	130	257
Widowed	317	68	35	6	46	92	3	30	37
Total	300,857	37,303	36,078	17,012	38,972	95,298	2,980	28,193	45,021

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.5b. Individuals aged 18 to 24 years, by mental health diagnosis, education, and marital status: Column percent distribution, 2018

<i>Education and marital status</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 18 to 24 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Highest school grade completed									
0 to 8	9.5	10.2	12.9	17.4	6.6	7.0	8.3	6.8	12.8
9 to 11	27.8	30.1	25.1	34.1	26.0	26.2	27.8	28.3	31.1
12 (or GED)	48.3	46.0	45.6	40.0	52.2	49.5	50.0	52.4	47.2
More than 12	14.4	13.7	16.5	8.5	15.2	17.3	13.9	12.5	9.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marital status									
Never married	93.5	91.1	92.9	97.4	92.6	92.4	91.5	96.8	95.3
Currently married	4.5	6.1	5.4	2.0	4.8	5.2	5.6	2.0	3.2
Separated	1.2	1.6	†	†	1.5	1.4	1.5	†	†
Divorced	†	1.0	†	†	†	†	1.3	†	†
Widowed	†	†	†	†	†	†	†	†	†
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.5c. Individuals aged 18 to 24 years, by mental health diagnosis, education, and marital status: Row percent distribution, 2018

<i>Education and marital status</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 18 to 24 years	100.0	12.6	14.4	5.7	11.1	29.6	1.2	9.1	16.3
Highest school grade completed									
0 to 8	100.0	14.0	18.1	9.9	8.6	23.5	1.0	6.7	18.1
9 to 11	100.0	14.1	12.0	6.6	11.5	30.0	1.2	9.6	15.0
12 (or GED)	100.0	12.4	12.6	4.5	13.4	32.6	1.2	10.2	13.1
More than 12	100.0	12.4	15.2	3.2	13.1	38.3	1.1	8.2	8.4
Total	100.0	13.0	13.3	5.4	12.4	31.9	1.2	9.4	13.5
Marital status									
Never married	100.0	12.1	11.9	5.9	12.8	31.3	†	9.7	15.3
Currently married	100.0	16.8	14.3	2.5	13.8	36.6	1.2	4.1	10.6
Separated	100.0	17.1	10.0	1.4	16.7	37.8	1.3	5.4	10.4
Divorced	100.0	16.9	9.8	1.7	15.8	36.7	1.8	5.8	11.5
Widowed	100.0	21.5	11.0	1.9	14.5	29.0	†	9.5	11.7
Total	100.0	12.4	12.0	5.7	13.0	31.7	†	9.4	15.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.6a. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, education, and marital status: Number, 2018

<i>Education and marital status</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
		<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Bipolar disorders</i>	<i>Depressive disorders</i>				
Total individuals with SMI aged 18 to 24 years²	380,549	37,179	47,073	14,853	50,397	129,048	4,333	45,676	51,990
Highest school grade completed									
0 to 8	15,487	2,186	2,603	1,267	1,634	4,150	197	1,437	2,013
9 to 11	53,300	6,650	5,741	2,883	7,219	17,837	717	6,288	5,965
12 (or GED)	92,377	9,519	9,774	2,833	14,740	34,741	1,238	11,595	7,937
More than 12	27,692	2,621	3,311	540	4,359	12,162	327	2,765	1,607
Total	188,856	20,976	21,429	7,523	27,952	68,890	2,479	22,085	17,522
Marital status									
Never married	193,169	16,187	19,863	6,914	30,946	73,069	2,072	24,410	19,708
Currently married	9,593	1,311	1,173	171	1,621	4,043	125	499	650
Separated	2,642	332	226	23	535	1,143	29	175	179
Divorced	1,655	228	131	22	309	692	29	118	126
Widowed	217	52	14	4	37	68	3	25	14
Total	207,276	18,110	21,407	7,134	33,448	79,015	2,258	25,227	20,677

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.6b. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, education, and marital status: Column percent distribution, 2018

Education and marital status	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
		Anxiety disorders	ADHD	Bipolar disorders	Depressive disorders				
Total individuals with SMI aged 18 to 24 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Highest school grade completed									
0 to 8	8.2	10.4	12.1	16.8	5.8	6.0	7.9	6.5	11.5
9 to 11	28.2	31.7	26.8	38.3	25.8	25.9	28.9	28.5	34.0
12 (or GED)	48.9	45.4	45.6	37.7	52.7	50.4	49.9	52.5	45.3
More than 12	14.7	12.5	15.5	7.2	15.6	17.7	13.2	12.5	9.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marital status									
Never married	93.2	89.4	92.8	96.9	92.5	92.5	91.8	96.8	95.3
Currently married	4.6	7.2	5.5	2.4	4.8	5.1	5.5	2.0	3.1
Separated	1.3	1.8	1.1	†	1.6	1.4	1.3	†	†
Divorced	†	1.3	†	†	†	†	1.3	†	†
Widowed	†	†	†	†	†	†	†	†	†
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma; SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.6c. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, education, marital status: Row percent distribution, 2018

Education and marital status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years²	100.0	9.8	12.4	3.9	13.2	33.9	1.1	12.0	13.7
Highest school grade completed									
0 to 8	100.0	14.1	16.8	8.2	10.6	26.8	1.3	9.3	13.0
9 to 11	100.0	12.5	10.8	5.4	13.5	33.5	1.3	11.8	11.2
12 (or GED)	100.0	10.3	10.6	3.1	16.0	37.6	1.3	12.6	8.6
More than 12	100.0	9.5	12.0	2.0	15.7	43.9	1.2	10.0	5.8
Total	100.0	11.1	11.3	4.0	14.8	36.5	1.3	11.7	9.3
Marital status									
Never married	100.0	8.4	10.3	3.6	16.0	37.8	1.1	12.6	10.2
Currently married	100.0	13.7	12.2	1.8	16.9	42.1	1.3	5.2	6.8
Separated	100.0	12.6	8.6	†	20.2	43.3	1.1	6.6	6.8
Divorced	100.0	13.8	7.9	1.3	18.7	41.8	1.8	7.1	7.6
Widowed	100.0	24.0	6.5	1.8	17.1	31.3	1.4	11.5	6.5
Total	100.0	8.7	10.3	3.4	16.1	38.1	1.1	12.2	10.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma; SMI = serious mental illness

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.7a. Individuals aged 18 to 24 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Number, 2018

<i>Living arrangement, SMI status, and veteran status</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 18 to 24 years	557,478	70,164	80,312	31,550	61,603	165,259	6,454	50,990	91,146
Living arrangement									
Homeless	12,875	1,543	937	190	1,781	3,533	181	2,279	2,431
Foster home/foster care	3,180	660	328	264	261	666	27	215	759
Residential care	9,509	955	744	386	1,403	1,882	110	1,732	2,297
Institutional setting	3,292	225	201	73	371	620	124	1,065	613
Jail/correctional facility	8,851	1,946	640	235	698	1,441	570	1,171	2,150
Private residence	285,143	34,139	37,618	14,784	33,735	90,351	3,019	27,092	44,405
Other	10,965	1,897	1,893	425	983	3,146	72	846	1,703
Total	333,815	41,365	42,361	16,357	39,232	101,639	4,103	34,400	54,358
SMI status²									
SMI	380,549	37,179	47,073	14,853	50,397	129,048	4,333	45,676	51,990
Not SMI	141,746	28,342	25,731	13,943	9,024	27,217	1,731	3,981	31,777
Total	522,295	65,521	72,804	28,796	59,421	156,265	6,064	49,657	83,767
Veteran status									
Veteran	1,849	266	226	69	232	681	7	196	172
Not a veteran	200,732	27,198	25,882	10,413	22,681	60,173	1,663	18,386	34,336
Total	202,581	27,464	26,108	10,482	22,913	60,854	1,670	18,582	34,508

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.7b. Individuals aged 18 to 24 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Column percent distribution, 2018

<i>Living arrangement, SMI status, and veteran status</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 18 to 24 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement									
Homeless	3.9	3.7	2.2	1.2	4.5	3.5	4.4	6.6	4.5
Foster home/foster care	†	1.6	†	1.6	†	†	†	†	1.4
Residential care	2.8	2.3	1.8	2.4	3.6	1.9	2.7	5.0	4.2
Institutional setting	†	†	†	†	†	†	3.0	3.1	1.1
Jail/correctional facility	2.7	4.7	1.5	1.4	1.8	1.4	13.9	3.4	4.0
Private residence	85.4	82.5	88.8	90.4	86.0	88.9	73.6	78.8	81.7
Other	3.3	4.6	4.5	2.6	2.5	3.1	1.8	2.5	3.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SMI status²									
SMI	72.9	56.7	64.7	51.6	84.8	82.6	71.5	92.0	62.1
Not SMI	27.1	43.3	35.3	48.4	15.2	17.4	28.5	8.0	37.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status									
Veteran	†	†	†	†	1.0	1.1	†	1.1	†
Not a veteran	99.1	99.0	99.1	99.3	99.0	98.9	99.6	98.9	99.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.7c. Individuals aged 18 to 24 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Row percent distribution, 2018

Living arrangement, SMI status, and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 18 to 24 years	100.0	12.6	14.4	5.7	11.1	29.6	1.2	9.1	16.3
Living arrangement									
Homeless	100.0	12.0	7.3	1.5	13.8	27.4	1.4	17.7	18.9
Foster home/foster care	100.0	20.8	10.3	8.3	8.2	20.9	†	6.8	23.9
Residential care	100.0	10.0	7.8	4.1	14.8	19.8	1.2	18.2	24.2
Institutional setting	100.0	6.8	6.1	2.2	11.3	18.8	3.8	32.4	18.6
Jail/correctional facility	100.0	22.0	7.2	2.7	7.9	16.3	6.4	13.2	24.3
Private residence	100.0	12.0	13.2	5.2	11.8	31.7	1.1	9.5	15.6
Other	100.0	17.3	17.3	3.9	9.0	28.7	†	7.7	15.5
Total	100.0	12.4	12.7	4.9	11.8	30.4	1.2	10.3	16.3
SMI status²									
SMI	100.0	9.8	12.4	3.9	13.2	33.9	1.1	12.0	13.7
Not SMI	100.0	20.0	18.2	9.8	6.4	19.2	1.2	2.8	22.4
Total	100.0	12.5	13.9	5.5	11.4	29.9	1.2	9.5	16.0
Veteran status									
Veteran	100.0	14.4	12.2	3.7	12.5	36.8	†	10.6	9.3
Not a veteran	100.0	13.5	12.9	5.2	11.3	30.0	†	9.2	17.1
Total	100.0	13.6	12.9	5.2	11.3	30.0	†	9.2	17.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.8a. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, living arrangement and veteran status: Number, 2018

<i>Living arrangement and veteran status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 18 to 24 years²	380,549	37,179	47,073	14,853	50,397	129,048	4,333	45,676	51,990
Living arrangement									
Homeless	10,602	1,140	635	119	1,587	3,103	157	2,121	1,740
Foster home/foster care	2,414	470	270	151	226	568	23	201	505
Residential care	7,358	631	500	173	1,297	1,666	97	1,642	1,352
Institutional setting	2,234	127	138	43	276	401	59	889	301
Jail/correctional facility	6,266	971	449	122	561	1,183	522	1,046	1,412
Private residence	212,945	21,828	23,796	8,040	29,742	76,180	2,429	25,115	25,815
Other	8,330	961	1,591	138	899	2,843	64	795	1,039
Total	250,149	26,128	27,379	8,786	34,588	85,944	3,351	31,809	32,164
Veteran status									
Veteran	1,472	182	147	34	215	620	5	177	92
Not a veteran	140,850	16,883	15,103	5,175	19,520	50,672	1,248	16,438	15,811
Total	142,322	17,065	15,250	5,209	19,735	51,292	1,253	16,615	15,903

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.8b. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, living arrangement and veteran status: Column percent distribution, 2018

<i>Living arrangement and veteran status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 18 to 24 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement									
Homeless	4.2	4.4	2.3	1.4	4.6	3.6	4.7	6.7	5.4
Foster home/foster care	†	1.8	†	1.7	†	†	†	†	1.6
Residential care	2.9	2.4	1.8	2.0	3.7	1.9	2.9	5.2	4.2
Institutional setting	†	†	†	†	†	†	1.8	2.8	†
Jail/correctional facility	2.5	3.7	1.6	1.4	1.6	1.4	15.6	3.3	4.4
Private residence	85.1	83.5	86.9	91.5	86.0	88.6	72.5	79.0	80.3
Other	3.3	3.7	5.8	1.6	2.6	3.3	1.9	2.5	3.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status									
Veteran	1.0	1.1	†	†	1.1	1.2	†	1.1	†
Not a veteran	99.0	98.9	99.0	99.3	98.9	98.8	99.6	98.9	99.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.8c. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, living arrangement and veteran status: Row percent distribution, 2018

<i>Living arrangement and veteran status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 18 to 24 years²	100.0	9.8	12.4	3.9	13.2	33.9	1.1	12.0	13.7
Living arrangement									
Homeless	100.0	10.8	6.0	1.1	15.0	29.3	1.5	20.0	16.4
Foster home/foster care	100.0	19.5	11.2	6.3	9.4	23.5	†	8.3	20.9
Residential care	100.0	8.6	6.8	2.4	17.6	22.6	1.3	22.3	18.4
Institutional setting	100.0	5.7	6.2	1.9	12.4	17.9	2.6	39.8	13.5
Jail/correctional facility	100.0	15.5	7.2	1.9	9.0	18.9	8.3	16.7	22.5
Private residence	100.0	10.3	11.2	3.8	14.0	35.8	1.1	11.8	12.1
Other	100.0	11.5	19.1	1.7	10.8	34.1	†	9.5	12.5
Total	100.0	10.4	10.9	3.5	13.8	34.4	1.3	12.7	12.9
Veteran status									
Veteran	100.0	12.4	10.0	2.3	14.6	42.1	†	12.0	6.3
Not a veteran	100.0	12.0	10.7	3.7	13.9	36.0	†	11.7	11.2
Total	100.0	12.0	10.7	3.7	13.9	36.0	†	11.7	11.2

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.9a. Individuals aged 18 to 24 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2018

Employment status and ‘detailed not in labor force’	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 18 to 24 years	557,478	70,164	80,312	31,550	61,603	165,259	6,454	50,990	91,146
Employment status									
Employed	81,616	11,605	12,269	3,371	9,585	29,981	1,115	4,153	9,537
Full time	35,400	5,824	6,188	1,528	3,273	11,957	582	1,425	4,623
Part time	34,571	4,906	5,129	1,555	3,636	12,542	450	1,895	4,458
Unemployed	82,674	9,233	8,689	2,430	11,881	27,789	890	10,108	11,654
Not in labor force (see detail below)	136,320	15,937	17,983	9,080	15,088	37,581	1,229	13,824	25,598
Total	300,610	36,775	38,941	14,881	36,554	95,351	3,234	28,085	46,789
Detailed not in labor force									
Total individuals aged 18 to 24 years not in labor force	136,320	15,937	17,983	9,080	15,088	37,581	1,229	13,824	25,598
Disabled	22,901	1,429	1,862	1,039	3,758	4,279	336	4,693	5,505
Retired	173	19	12	5	19	49	6	28	35
Student	61,512	7,973	7,853	5,306	4,655	17,710	303	3,800	13,912
Homemaker	3,336	578	511	48	379	1,326	43	96	355
Sheltered/non-competitive employment	1,291	167	124	44	147	335	5	157	312
Other	47,107	5,771	7,621	2,638	6,130	13,882	536	5,050	5,479

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTES: *Employed* includes “full-time/part-time not differentiated,” which is not presented separately; total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.9b. Individuals aged 18 to 24 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 18 to 24 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Employment status									
Employed	27.2	31.6	31.5	22.7	26.2	31.4	34.5	14.8	20.4
Full time	11.8	15.8	15.9	10.3	9.0	12.5	18.0	5.1	9.9
Part time	11.5	13.3	13.2	10.4	9.9	13.2	13.9	6.7	9.5
Unemployed	27.5	25.1	22.3	16.3	32.5	29.1	27.5	36.0	24.9
Not in labor force (see detail below)	45.3	43.3	46.2	61.0	41.3	39.4	38.0	49.2	54.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force									
Total individuals aged 18 to 24 years not in labor force	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Disabled	16.8	9.0	10.4	11.4	24.9	11.4	27.3	33.9	21.5
Retired	†	†	†	†	†	†	†	†	†
Student	45.1	50.0	43.7	58.4	30.9	47.1	24.7	27.5	54.3
Homemaker	2.4	3.6	2.8	†	2.5	3.5	3.5	†	1.4
Sheltered/non-competitive employment	†	1.0	†	†	†	†	†	1.1	1.2
Other	34.6	36.2	42.4	29.1	40.6	36.9	43.6	36.5	21.4

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTES: Percentages may not sum to 100 percent because of rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.9c. Individuals aged 18 to 24 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 18 to 24 years	100.0	12.6	14.4	5.7	11.1	29.6	1.2	9.1	16.3
Employment status									
Employed	100.0	14.2	15.0	4.1	11.7	36.7	1.4	5.1	11.7
Full time	100.0	16.5	17.5	4.3	9.2	33.8	1.6	4.0	13.1
Part time	100.0	14.2	14.8	4.5	10.5	36.3	1.3	5.5	12.9
Unemployed	100.0	11.2	10.5	2.9	14.4	33.6	1.1	12.2	14.1
Not in labor force (see detail below)	100.0	11.7	13.2	6.7	11.1	27.6	†	10.1	18.8
Total	100.0	12.2	13.0	5.0	12.2	31.7	1.1	9.3	15.6
Detailed not in labor force									
Total individuals aged 18 to 24 years not in labor force	100.0	11.7	13.2	6.7	11.1	27.6	†	10.1	18.8
Disabled	100.0	6.2	8.1	4.5	16.4	18.7	1.5	20.5	24.0
Retired	100.0	11.0	6.9	2.9	11.0	28.3	3.5	16.2	20.2
Student	100.0	13.0	12.8	8.6	7.6	28.8	†	6.2	22.6
Homemaker	100.0	17.3	15.3	1.4	11.4	39.7	1.3	2.9	10.6
Sheltered/non-competitive employment	100.0	12.9	9.6	3.4	11.4	25.9	†	12.2	24.2
Other	100.0	12.3	16.2	5.6	13.0	29.5	1.1	10.7	11.6

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTES: Percentages may not sum to 100 percent because of rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.10a. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 18 to 24 years²	380,549	37,179	47,073	14,853	50,397	129,048	4,333	45,676	51,990
Employment status									
Employed	60,576	7,274	7,758	1,852	8,391	25,230	891	3,809	5,371
Full time	24,743	3,659	3,890	790	2,680	9,535	464	1,261	2,464
Part time	25,954	3,220	3,462	942	3,091	10,503	352	1,727	2,657
Unemployed	63,480	5,769	5,634	1,231	10,612	24,029	699	9,355	6,151
Not in labor force (see detail below)	100,308	10,339	11,878	5,104	13,180	31,312	986	12,846	14,663
Total	224,364	23,382	25,270	8,187	32,183	80,571	2,576	26,010	26,185
Detailed not in labor force									
Total individuals with SMI aged 18 to 24 years not in labor force	100,308	10,339	11,878	5,104	13,180	31,312	986	12,846	14,663
Disabled	17,367	1,077	1,324	667	3,336	3,703	304	4,437	2,519
Retired	131	13	6	2	18	39	6	24	23
Student	45,372	5,204	5,440	3,164	3,971	14,934	229	3,555	8,875
Homemaker	2,462	373	327	23	329	1,079	33	89	209
Sheltered/non-competitive employment	1,094	154	101	20	139	317	5	150	208
Other	33,882	3,518	4,680	1,228	5,387	11,240	409	4,591	2,829

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTES: *Employed* includes “full-time/part-time not differentiated,” which is not presented separately; total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.10b. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 18 to 24 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Employment status									
Employed	27.0	31.1	30.7	22.6	26.1	31.3	34.6	14.6	20.5
Full time	11.0	15.6	15.4	9.6	8.3	11.8	18.0	4.8	9.4
Part time	11.6	13.8	13.7	11.5	9.6	13.0	13.7	6.6	10.1
Unemployed	28.3	24.7	22.3	15.0	33.0	29.8	27.1	36.0	23.5
Not in labor force (see detail below)	44.7	44.2	47.0	62.3	41.0	38.9	38.3	49.4	56.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force									
Total individuals with SMI aged 18 to 24 years not in labor force	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Disabled	17.3	10.4	11.1	13.1	25.3	11.8	30.8	34.5	17.2
Retired	†	†	†	†	†	†	†	†	†
Student	45.2	50.3	45.8	62.0	30.1	47.7	23.2	27.7	60.5
Homemaker	2.5	3.6	2.8	†	2.5	3.4	3.3	†	1.4
Sheltered/non-competitive employment	1.1	1.5	†	†	1.1	1.0	†	1.2	1.4
Other	33.8	34.0	39.4	24.1	40.9	35.9	41.5	35.7	19.3

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTES: Percentages may not sum to 100 percent because of rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.10c. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 18 to 24 years²	100.0	9.8	12.4	3.9	13.2	33.9	1.1	12.0	13.7
Employment status									
Employed	100.0	12.0	12.8	3.1	13.9	41.7	1.5	6.3	8.9
Full time	100.0	14.8	15.7	3.2	10.8	38.5	1.9	5.1	10.0
Part time	100.0	12.4	13.3	3.6	11.9	40.5	1.4	6.7	10.2
Unemployed	100.0	9.1	8.9	1.9	16.7	37.9	1.1	14.7	9.7
Not in labor force (see detail below)	100.0	10.3	11.8	5.1	13.1	31.2	†	12.8	14.6
Total	100.0	10.4	11.3	3.6	14.3	35.9	1.1	11.6	11.7
Detailed not in labor force									
Total individuals with SMI aged 18 to 24 years not in labor force	100.0	10.3	11.8	5.1	13.1	31.2	†	12.8	14.6
Disabled	100.0	6.2	7.6	3.8	19.2	21.3	1.8	25.5	14.5
Retired	100.0	9.9	4.6	1.5	13.7	29.8	4.6	18.3	17.6
Student	100.0	11.5	12.0	7.0	8.8	32.9	†	7.8	19.6
Homemaker	100.0	15.2	13.3	†	13.4	43.8	1.3	3.6	8.5
Sheltered/non-competitive employment	100.0	14.1	9.2	1.8	12.7	29.0	†	13.7	19.0
Other	100.0	10.4	13.8	3.6	15.9	33.2	1.2	13.5	8.3

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTES: Percentages may not sum to 100 percent because of rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.11a. Individuals aged 18 to 24 years, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018

Timing of admission and mental health service setting	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 18 to 24 years	557,478	70,164	80,312	31,550	61,603	165,259	6,454	50,990	91,146
Timing of admission									
New individuals	329,494	46,474	50,476	13,593	32,968	105,801	4,037	25,635	50,510
Continuing individuals	227,984	23,690	29,836	17,957	28,635	59,458	2,417	25,355	40,636
Mental health service setting^{2,3}									
State psychiatric hospital	12,867	666	272	140	2,080	2,258	436	5,234	1,781
Community-based program	542,513	68,196	79,817	31,326	60,016	162,010	6,199	47,614	87,335
Residential treatment center	5,971	595	285	225	782	1,221	75	1,376	1,412
Other psychiatric inpatient	32,465	2,660	2,364	588	5,376	10,206	415	6,849	4,007
Institutions under the justice system	9,038	2,213	441	118	436	1,090	573	1,223	2,944

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.11b. Individuals aged 18 to 24 years, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018

Timing of admission and mental health service setting	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 18 to 24 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission									
New individuals	59.1	66.2	62.8	43.1	53.5	64.0	62.6	50.3	55.4
Continuing individuals	40.9	33.8	37.2	56.9	46.5	36.0	37.4	49.7	44.6
Mental health service setting^{2,3}									
State psychiatric hospital	2.3	†	†	†	3.4	1.4	6.8	10.3	2.0
Community-based program	97.3	97.2	99.4	99.3	97.4	98.0	96.0	93.4	95.8
Residential treatment center	1.1	†	†	†	1.3	†	1.2	2.7	1.5
Other psychiatric inpatient	5.8	3.8	2.9	1.9	8.7	6.2	6.4	13.4	4.4
Institutions under the justice system	1.6	3.2	†	†	†	†	8.9	2.4	3.2

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages for timing of admission may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.11c. Individuals aged 18 to 24 years, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018

<i>Timing of admission and mental health service setting</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 18 to 24 years	100.0	12.6	14.4	5.7	11.1	29.6	1.2	9.1	16.3
Timing of admission									
New individuals	100.0	14.1	15.3	4.1	10.0	32.1	1.2	7.8	15.3
Continuing individuals	100.0	10.4	13.1	7.9	12.6	26.1	1.1	11.1	17.8
Mental health service setting^{2,3}									
State psychiatric hospital	100.0	5.2	2.1	1.1	16.2	17.5	3.4	40.7	13.8
Community-based program	100.0	12.6	14.7	5.8	11.1	29.9	1.1	8.8	16.1
Residential treatment center	100.0	10.0	4.8	3.8	13.1	20.4	1.3	23.0	23.6
Other psychiatric inpatient	100.0	8.2	7.3	1.8	16.6	31.4	1.3	21.1	12.3
Institutions under the justice system	100.0	24.5	4.9	1.3	4.8	12.1	6.3	13.5	32.6

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.12a. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018

<i>Timing of admission and mental health service setting</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 18 to 24 years²	380,549	37,179	47,073	14,853	50,397	129,048	4,333	45,676	51,990
Timing of admission									
New individuals	220,856	24,552	29,541	6,519	26,763	81,262	2,453	22,341	27,425
Continuing individuals	159,693	12,627	17,532	8,334	23,634	47,786	1,880	23,335	24,565
Mental health service setting^{3,4}									
State psychiatric hospital	10,093	440	189	97	1,729	1,792	313	4,588	945
Community-based program	368,867	35,719	46,694	14,720	49,172	126,413	4,171	42,833	49,145
Residential treatment center	4,737	405	200	113	693	1,070	63	1,339	854
Other psychiatric inpatient	25,633	1,474	1,498	374	4,543	8,506	339	6,242	2,657
Institutions under the justice system	7,865	1,497	394	103	400	1,013	563	1,204	2,691

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.12b. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018

Timing of admission and mental health service setting	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission									
New individuals	58.0	66.0	62.8	43.9	53.1	63.0	56.6	48.9	52.8
Continuing individuals	42.0	34.0	37.2	56.1	46.9	37.0	43.4	51.1	47.2
Mental health service setting^{3,4}									
State psychiatric hospital	2.7	1.2	†	†	3.4	1.4	7.2	10.0	1.8
Community-based program	96.9	96.1	99.2	99.1	97.6	98.0	96.3	93.8	94.5
Residential treatment center	1.2	1.1	†	†	1.4	†	1.5	2.9	1.6
Other psychiatric inpatient	6.7	4.0	3.2	2.5	9.0	6.6	7.8	13.7	5.1
Institutions under the justice system	2.1	4.0	†	†	†	†	13.0	2.6	5.2

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages for timing of admission may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.12c. Individuals aged 18 to 24 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 18 to 24 years²	100.0	9.8	12.4	3.9	13.2	33.9	1.1	12.0	13.7
Timing of admission									
New individuals	100.0	11.1	13.4	3.0	12.1	36.8	1.1	10.1	12.4
Continuing individuals	100.0	7.9	11.0	5.2	14.8	29.9	1.2	14.6	15.4
Mental health service setting^{3,4}									
State psychiatric hospital	100.0	4.4	1.9	†	17.1	17.8	3.1	45.5	9.4
Community-based program	100.0	9.7	12.7	4.0	13.3	34.3	1.1	11.6	13.3
Residential treatment center	100.0	8.5	4.2	2.4	14.6	22.6	1.3	28.3	18.0
Other psychiatric inpatient	100.0	5.8	5.8	1.5	17.7	33.2	1.3	24.4	10.4
Institutions under the justice system	100.0	19.0	5.0	1.3	5.1	12.9	7.2	15.3	34.2

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 3.13. Individuals aged 18 to 24 years, by mental health service setting and SMI status: Number and percent distribution, 2018

<i>Mental health service setting</i>	<i>SMI status¹</i>								
	<i>Number of individuals</i>			<i>Column percent of individuals</i>			<i>Row percent distribution of individuals</i>		
	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>
Total individuals aged 18 to 24 years	588,799	412,774	176,025	100.0	100.0	100.0	100.0	70.1	29.9
Mental health service setting^{2,3}									
State psychiatric hospital	12,615	11,189	1,426	2.1	2.7	†	100.0	88.7	11.3
Community-based program	573,158	399,615	173,543	97.3	96.8	98.6	100.0	69.7	30.3
Residential treatment center	6,158	4,973	1,185	1.0	1.2	†	100.0	80.8	19.2
Other psychiatric inpatient	34,186	26,780	7,406	5.8	6.5	4.2	100.0	78.3	21.7
Institutions under the justice system	9,796	8,596	1,200	1.7	2.1	†	100.0	87.8	12.2

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ The federal definition for SMI status can be found in Appendix D.

² Individuals could receive services in more than one mental health service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.1a. Individuals aged 25 to 54 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2013–2018

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	47	45	47	48	48
Total individuals aged 25 to 54 years	2,558,630	2,668,971	2,447,157	2,662,746	2,748,726	2,831,082
Gender						
Male	1,140,092	1,198,924	1,108,380	1,201,253	1,260,733	1,306,842
Female	1,416,613	1,466,001	1,335,446	1,454,755	1,479,627	1,518,595
Total	2,556,705	2,664,925	2,443,826	2,656,008	2,740,360	2,825,437
Age¹						
25 to 34 years	888,048	934,775	870,301	970,525	1,039,064	1,091,760
35 to 39 years	383,178	406,379	388,512	435,995	467,430	494,186
40 to 49 years	843,251	857,903	759,874	803,374	803,331	815,101
50 to 54 years	444,153	469,914	428,470	452,852	438,901	430,035
Total	2,558,630	2,668,971	2,447,157	2,662,746	2,748,726	2,831,082
Race						
American Indian or Alaska Native	27,952	32,991	33,940	35,422	37,927	42,440
Asian ²	29,501	34,000	32,967	37,817	39,415	43,607
Black or African American	475,872	502,942	451,937	491,525	498,813	521,214
Native Hawaiian or Other Pacific Islander	6,407	6,852	7,444	8,348	5,935	7,178
White	1,679,857	1,736,676	1,577,075	1,721,495	1,763,177	1,824,351
Other	157,096	162,284	141,843	163,405	185,930	206,253
Total	2,376,685	2,475,745	2,245,206	2,458,012	2,531,197	2,645,043
Ethnicity						
Not of Hispanic or Latino origin	1,864,413	1,918,453	1,739,420	1,873,069	2,122,681	2,184,247
Hispanic or Latino origin	250,298	265,246	257,265	289,042	312,087	322,753
Total	2,114,711	2,183,699	1,996,685	2,162,111	2,434,768	2,507,000
Specific Hispanic or Latino origin						
Mexican	10,123	14,543	18,017	15,014	17,528	16,358
Puerto Rican	12,917	22,841	23,314	19,870	23,920	13,600
Cuban	4,420	4,243	4,257	3,869	3,773	3,716
Other specific	18,736	22,654	20,814	20,990	24,285	24,517
Total	46,196	64,281	66,402	59,743	69,506	58,191

Continued. See notes at end of table.

**Table 4.1a. Individuals aged 25 to 54 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2013–2018
(continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Highest school grade completed						
0 to 8	116,667	122,035	125,806	128,928	113,798	107,573
9 to 11	200,789	206,519	206,850	206,338	206,819	210,449
12 (or GED)	453,633	511,403	524,167	541,308	560,871	628,911
More than 12	197,179	229,392	239,127	252,844	297,725	329,891
Total	968,268	1,069,349	1,095,950	1,129,418	1,179,213	1,276,824
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	29,928	26,951
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	971,795	986,258
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	1,001,723	1,013,209
SMI status³						
SMI	1,695,602	1,802,197	1,658,579	1,767,390	1,861,640	1,973,914
Not SMI	659,682	683,562	638,759	697,206	679,194	681,038
Total	2,355,284	2,485,759	2,297,338	2,464,596	2,540,834	2,654,952

n/c Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 4.1b. Individuals aged 25 to 54 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals aged 25 to 54 years	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	44.6	45.0	45.4	45.2	46.0	46.3
Female	55.4	55.0	54.6	54.8	54.0	53.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
25 to 34 years	34.7	35.0	35.6	36.4	37.8	38.6
35 to 39 years	15.0	15.2	15.9	16.4	17.0	17.5
40 to 49 years	33.0	32.1	31.1	30.2	29.2	28.8
50 to 54 years	17.4	17.6	17.5	17.0	16.0	15.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	1.2	1.3	1.5	1.4	1.5	1.6
Asian ²	1.2	1.4	1.5	1.5	1.6	1.6
Black or African American	20.0	20.3	20.1	20.0	19.7	19.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	70.7	70.1	70.2	70.0	69.7	69.0
Other	6.6	6.6	6.3	6.6	7.3	7.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	88.2	87.9	87.1	86.6	87.2	87.1
Hispanic or Latino origin	11.8	12.1	12.9	13.4	12.8	12.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	21.9	22.6	27.1	25.1	25.2	28.1
Puerto Rican	28.0	35.5	35.1	33.3	34.4	23.4
Cuban	9.6	6.6	6.4	6.5	5.4	6.4
Other specific	40.6	35.2	31.3	35.1	34.9	42.1
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 4.1b. Individuals aged 25 to 54 years, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Highest school grade completed						
0 to 8	12.0	11.4	11.5	11.4	9.7	8.4
9 to 11	20.7	19.3	18.9	18.3	17.5	16.5
12 (or GED)	46.8	47.8	47.8	47.9	47.6	49.3
More than 12	20.4	21.5	21.8	22.4	25.2	25.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	3.0	2.7
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	97.0	97.3
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0
SMI status³						
SMI	72.0	72.5	72.2	71.7	73.3	74.3
Not SMI	28.0	27.5	27.8	28.3	26.7	25.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 4.2a. Individuals aged 25 to 54 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	44	47	45	46	46	47
Total individuals with SMI aged 25 to 54 years¹	1,695,602	1,802,197	1,658,579	1,767,390	1,861,640	1,973,914
Gender						
Male	752,268	807,204	751,951	801,862	853,831	902,699
Female	942,396	992,790	904,921	963,473	1,005,824	1,069,201
Total	1,694,664	1,799,994	1,656,872	1,765,335	1,859,655	1,971,900
Age²						
25 to 34 years	555,231	597,091	559,282	609,819	662,797	719,342
35 to 39 years	251,534	271,244	260,029	286,629	311,879	339,671
40 to 49 years	574,596	596,373	530,058	550,404	565,575	590,231
50 to 54 years	314,241	337,489	309,210	320,538	321,389	324,670
Total	1,695,602	1,802,197	1,658,579	1,767,390	1,861,640	1,973,914
Race						
American Indian or Alaska Native	14,884	19,454	19,588	20,237	23,832	25,869
Asian ³	21,725	25,822	24,596	26,567	29,440	32,823
Black or African American	338,659	358,061	330,499	344,882	357,445	384,904
Native Hawaiian or Other Pacific Islander	5,040	5,568	5,832	6,238	4,163	4,992
White	1,095,790	1,147,831	1,044,245	1,132,943	1,182,336	1,240,710
Other	110,467	119,873	106,572	117,635	147,698	166,823
Total	1,586,565	1,676,609	1,531,332	1,648,502	1,744,914	1,856,121
Ethnicity						
Not of Hispanic or Latino origin	1,248,071	1,303,957	1,192,596	1,271,404	1,406,931	1,481,783
Hispanic or Latino origin	179,132	198,462	187,416	211,459	239,407	253,621
Total	1,427,203	1,502,419	1,380,012	1,482,863	1,646,338	1,735,404
Specific Hispanic or Latino origin						
Mexican	6,688	10,186	12,705	10,396	12,488	11,937
Puerto Rican	9,824	15,876	15,893	13,969	15,855	11,013
Cuban	3,898	3,952	3,981	3,621	3,463	3,503
Other specific	14,404	17,661	16,315	16,066	18,751	19,519
Total	34,814	47,675	48,894	44,052	50,557	45,972

Continued. See notes at end of table.

**Table 4.2a. Individuals aged 25 to 54 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Number, 2013–2018
(continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Highest school grade completed						
0 to 8	68,957	73,791	72,283	76,393	79,665	76,695
9 to 11	138,507	150,991	148,858	155,151	165,255	170,168
12 (or GED)	297,602	356,801	363,653	393,499	429,055	490,256
More than 12	132,569	170,124	175,902	194,205	231,575	257,668
Total	637,635	751,707	760,696	819,248	905,550	994,787
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	20,908	20,351
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	641,461	724,366
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	662,369	744,717

n/c Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ The federal definition for SMI status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 4.2b. Individuals aged 25 to 54 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	44	47	45	46	46	47
Total individuals with SMI aged 25 to 54 years¹	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	44.4	44.8	45.4	45.4	45.9	45.8
Female	55.6	55.2	54.6	54.6	54.1	54.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age²						
25 to 34 years	32.7	33.1	33.7	34.5	35.6	36.4
35 to 39 years	14.8	15.1	15.7	16.2	16.8	17.2
40 to 49 years	33.9	33.1	32.0	31.1	30.4	29.9
50 to 54 years	18.5	18.7	18.6	18.1	17.3	16.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	†	1.2	1.3	1.2	1.4	1.4
Asian ³	1.4	1.5	1.6	1.6	1.7	1.8
Black or African American	21.3	21.4	21.6	20.9	20.5	20.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	69.1	68.5	68.2	68.7	67.8	66.8
Other	7.0	7.1	7.0	7.1	8.5	9.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	87.4	86.8	86.4	85.7	85.5	85.4
Hispanic or Latino origin	12.6	13.2	13.6	14.3	14.5	14.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	19.2	21.4	26.0	23.6	24.7	26.0
Puerto Rican	28.2	33.3	32.5	31.7	31.4	24.0
Cuban	11.2	8.3	8.1	8.2	6.8	7.6
Other specific	41.4	37.0	33.4	36.5	37.1	42.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 4.2b. Individuals aged 25 to 54 years with SMI, by gender, age, race, ethnicity, education, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Highest school grade completed						
0 to 8	10.8	9.8	9.5	9.3	8.8	7.7
9 to 11	21.7	20.1	19.6	18.9	18.2	17.1
12 (or GED)	46.7	47.5	47.8	48.0	47.4	49.3
More than 12	20.8	22.6	23.1	23.7	25.6	25.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	3.2	2.7
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	96.8	97.3
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ The federal definition for SMI status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 4.3a. Individuals aged 25 to 54 years, by mental health diagnosis, gender, race, and ethnicity: Number, 2018

Gender, race, and ethnicity	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	2,476,907	276,239	311,401	41,662	398,050	729,056	30,448	405,426	284,625
Gender									
Male	1,113,169	102,542	114,675	20,851	155,689	279,108	14,559	268,615	157,130
Female	1,360,144	173,225	196,315	20,746	241,745	448,982	15,849	136,410	126,872
Total	2,473,313	275,767	310,990	41,597	397,434	728,090	30,408	405,025	284,002
Race									
American Indian or Alaska Native	37,873	5,858	5,723	553	4,064	10,274	451	4,533	6,417
Asian ²	37,788	3,976	3,787	503	3,754	10,170	330	11,380	3,888
Black or African American	451,298	43,770	30,528	3,290	65,671	131,950	6,207	128,184	41,698
Native Hawaiian or Other Pacific Islander	6,618	1,092	909	103	755	1,797	66	1,297	599
White	1,599,860	180,863	229,170	32,491	284,062	478,100	19,884	199,345	175,945
Other	189,023	18,956	18,867	1,757	23,738	56,952	1,803	37,175	29,775
Total	2,322,460	254,515	288,984	38,697	382,044	689,243	28,741	381,914	258,322
Ethnicity									
Not of Hispanic or Latino origin	1,911,723	214,814	235,084	32,752	325,832	564,706	24,952	312,064	201,519
Hispanic or Latino origin	283,126	30,439	32,033	2,602	36,867	87,583	2,505	50,283	40,814
Total	2,194,849	245,253	267,117	35,354	362,699	652,289	27,457	362,347	242,333
Specific Hispanic or Latino origin									
Mexican	15,702	2,912	1,869	146	1,994	4,879	125	2,134	1,643
Puerto Rican	11,747	1,297	998	94	2,027	3,716	108	2,429	1,078
Cuban	3,359	296	294	35	543	994	14	713	470
Other specific	22,415	3,445	2,475	247	3,164	6,764	175	3,784	2,361
Total	53,223	7,950	5,636	522	7,728	16,353	422	9,060	5,552

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.3b. Individuals aged 25 to 54 years, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	45.0	37.2	36.9	50.1	39.2	38.3	47.9	66.3	55.3
Female	55.0	62.8	63.1	49.9	60.8	61.7	52.1	33.7	44.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	1.6	2.3	2.0	1.4	1.1	1.5	1.6	1.2	2.5
Asian ²	1.6	1.6	1.3	1.3	†	1.5	1.1	3.0	1.5
Black or African American	19.4	17.2	10.6	8.5	17.2	19.1	21.6	33.6	16.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	68.9	71.1	79.3	84.0	74.4	69.4	69.2	52.2	68.1
Other	8.1	7.4	6.5	4.5	6.2	8.3	6.3	9.7	11.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity									
Not of Hispanic or Latino origin	87.1	87.6	88.0	92.6	89.8	86.6	90.9	86.1	83.2
Hispanic or Latino origin	12.9	12.4	12.0	7.4	10.2	13.4	9.1	13.9	16.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	29.5	36.6	33.2	28.0	25.8	29.8	29.6	23.6	29.6
Puerto Rican	22.1	16.3	17.7	18.0	26.2	22.7	25.6	26.8	19.4
Cuban	6.3	3.7	5.2	6.7	7.0	6.1	3.3	7.9	8.5
Other specific	42.1	43.3	43.9	47.3	40.9	41.4	41.5	41.8	42.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.3c. Individuals aged 25 to 54 years, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	100.0	11.2	12.6	1.7	16.1	29.4	1.2	16.4	11.5
Gender									
Male	100.0	9.2	10.3	1.9	14.0	25.1	1.3	24.1	14.1
Female	100.0	12.7	14.4	1.5	17.8	33.0	1.2	10.0	9.3
Total	100.0	11.1	12.6	1.7	16.1	29.4	1.2	16.4	11.5
Race									
American Indian or Alaska Native	100.0	15.5	15.1	1.5	10.7	27.1	1.2	12.0	16.9
Asian ²	100.0	10.5	10.0	1.3	9.9	26.9	†	30.1	10.3
Black or African American	100.0	9.7	6.8	†	14.6	29.2	1.4	28.4	9.2
Native Hawaiian or Other Pacific Islander	100.0	16.5	13.7	1.6	11.4	27.2	†	19.6	9.1
White	100.0	11.3	14.3	2.0	17.8	29.9	1.2	12.5	11.0
Other	100.0	10.0	10.0	†	12.6	30.1	†	19.7	15.8
Total	100.0	11.0	12.4	1.7	16.4	29.7	1.2	16.4	11.1
Ethnicity									
Not of Hispanic or Latino origin	100.0	11.2	12.3	1.7	17.0	29.5	1.3	16.3	10.5
Hispanic or Latino origin	100.0	10.8	11.3	†	13.0	30.9	†	17.8	14.4
Total	100.0	11.2	12.2	1.6	16.5	29.7	1.3	16.5	11.0
Specific Hispanic or Latino origin									
Mexican	100.0	18.5	11.9	†	12.7	31.1	†	13.6	10.5
Puerto Rican	100.0	11.0	8.5	†	17.3	31.6	†	20.7	9.2
Cuban	100.0	8.8	8.8	1.0	16.2	29.6	†	21.2	14.0
Other specific	100.0	15.4	11.0	1.1	14.1	30.2	†	16.9	10.5
Total	100.0	14.9	10.6	†	14.5	30.7	†	17.0	10.4

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.4a. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Number, 2018

Gender, race, and ethnicity	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years²	1,848,515	152,970	186,603	17,206	338,814	592,722	21,997	370,975	167,228
Gender									
Male	837,084	54,671	68,005	8,577	132,135	225,624	10,364	246,050	91,658
Female	1,009,582	98,164	118,481	8,611	206,336	366,593	11,623	124,616	75,158
Total	1,846,666	152,835	186,486	17,188	338,471	592,217	21,987	370,666	166,816
Race									
American Indian or Alaska Native	24,779	3,403	3,227	233	3,276	7,313	292	3,890	3,145
Asian ³	31,346	2,526	2,516	220	3,451	8,651	222	10,878	2,882
Black or African American	359,915	21,461	17,929	1,289	58,276	113,494	4,947	118,121	24,398
Native Hawaiian or Other Pacific Islander	4,763	646	465	41	617	1,348	40	1,217	389
White	1,158,357	99,506	134,345	13,150	239,265	380,272	13,970	180,299	97,550
Other	158,811	12,804	13,663	790	21,319	50,707	1,455	35,286	22,787
Total	1,737,971	140,346	172,145	15,723	326,204	561,785	20,926	349,691	151,151
Ethnicity									
Not of Hispanic or Latino origin	1,391,995	111,222	132,225	11,668	275,238	449,512	18,056	283,524	110,550
Hispanic or Latino origin	231,633	20,032	20,386	1,102	33,148	76,477	1,950	47,664	30,874
Total	1,623,628	131,254	152,611	12,770	308,386	525,989	20,006	331,188	141,424
Specific Hispanic or Latino origin									
Mexican	11,773	1,755	953	60	1,837	4,511	79	1,968	610
Puerto Rican	10,140	878	794	60	1,934	3,446	80	2,349	599
Cuban	3,179	256	260	29	533	964	10	700	427
Other specific	18,502	2,290	1,633	149	2,985	6,338	130	3,613	1,364
Total	43,594	5,179	3,640	298	7,289	15,259	299	8,630	3,000

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.4b. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	45.3	35.8	36.5	49.9	39.0	38.1	47.1	66.4	54.9
Female	54.7	64.2	63.5	50.1	61.0	61.9	52.9	33.6	45.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	1.4	2.4	1.9	1.5	1.0	1.3	1.4	1.1	2.1
Asian ³	1.8	1.8	1.5	1.4	1.1	1.5	1.1	3.1	1.9
Black or African American	20.7	15.3	10.4	8.2	17.9	20.2	23.6	33.8	16.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	66.6	70.9	78.0	83.6	73.3	67.7	66.8	51.6	64.5
Other	9.1	9.1	7.9	5.0	6.5	9.0	7.0	10.1	15.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity									
Not of Hispanic or Latino origin	85.7	84.7	86.6	91.4	89.3	85.5	90.3	85.6	78.2
Hispanic or Latino origin	14.3	15.3	13.4	8.6	10.7	14.5	9.7	14.4	21.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	27.0	33.9	26.2	20.1	25.2	29.6	26.4	22.8	20.3
Puerto Rican	23.3	17.0	21.8	20.1	26.5	22.6	26.8	27.2	20.0
Cuban	7.3	4.9	7.1	9.7	7.3	6.3	3.3	8.1	14.2
Other specific	42.4	44.2	44.9	50.0	41.0	41.5	43.5	41.9	45.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.4c. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years²	100.0	8.3	10.1	†	18.3	32.1	1.2	20.1	9.0
Gender									
Male	100.0	6.5	8.1	1.0	15.8	27.0	1.2	29.4	10.9
Female	100.0	9.7	11.7	†	20.4	36.3	1.2	12.3	7.4
Total	100.0	8.3	10.1	†	18.3	32.1	1.2	20.1	9.0
Race									
American Indian or Alaska Native	100.0	13.7	13.0	†	13.2	29.5	1.2	15.7	12.7
Asian ³	100.0	8.1	8.0	†	11.0	27.6	†	34.7	9.2
Black or African American	100.0	6.0	5.0	†	16.2	31.5	1.4	32.8	6.8
Native Hawaiian or Other Pacific Islander	100.0	13.6	9.8	†	13.0	28.3	†	25.6	8.2
White	100.0	8.6	11.6	1.1	20.7	32.8	1.2	15.6	8.4
Other	100.0	8.1	8.6	†	13.4	31.9	†	22.2	14.3
Total	100.0	8.1	9.9	†	18.8	32.3	1.2	20.1	8.7
Ethnicity									
Not of Hispanic or Latino origin	100.0	8.0	9.5	†	19.8	32.3	1.3	20.4	7.9
Hispanic or Latino origin	100.0	8.6	8.8	†	14.3	33.0	†	20.6	13.3
Total	100.0	8.1	9.4	†	19.0	32.4	1.2	20.4	8.7
Specific Hispanic or Latino origin									
Mexican	100.0	14.9	8.1	†	15.6	38.3	†	16.7	5.2
Puerto Rican	100.0	8.7	7.8	†	19.1	34.0	†	23.2	5.9
Cuban	100.0	8.1	8.2	†	16.8	30.3	†	22.0	13.4
Other specific	100.0	12.4	8.8	†	16.1	34.3	†	19.5	7.4
Total	100.0	11.9	8.3	†	16.7	35.0	†	19.8	6.9

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.5a. Individuals aged 25 to 54 years, by mental health diagnosis, education, and marital status: Number, 2018

<i>Education and marital status</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 25 to 54 years	2,476,907	276,239	311,401	41,662	398,050	729,056	30,448	405,426	284,625
Highest school grade completed									
0 to 8	101,437	13,743	14,337	2,130	13,414	28,445	1,221	17,029	11,118
9 to 11	195,027	19,714	17,483	1,827	33,879	57,173	3,195	44,945	16,811
12 (or GED)	587,601	65,529	65,895	8,598	103,462	180,492	7,916	99,998	55,711
More than 12	312,084	39,689	39,679	5,452	60,454	99,679	4,072	39,300	23,759
Total	1,196,149	138,675	137,394	18,007	211,209	365,789	16,404	201,272	107,399
Marital status									
Never married	824,352	75,910	80,419	12,671	140,516	233,858	8,855	182,491	89,632
Currently married	227,144	31,042	29,693	3,256	41,892	80,389	2,263	19,339	19,270
Separated	97,733	12,610	9,603	890	20,427	35,019	899	10,427	7,858
Divorced	204,661	24,524	19,829	2,006	44,828	70,756	2,170	24,285	16,263
Widowed	26,227	2,988	2,858	220	4,987	9,863	219	2,666	2,426
Total	1,380,117	147,074	142,402	19,043	252,650	429,885	14,406	239,208	135,449

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.5b. Individuals aged 25 to 54 years, by mental health diagnosis, education, and marital status: Column percent distribution, 2018

<i>Education and marital status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 25 to 54 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Highest school grade completed									
0 to 8	8.5	9.9	10.4	11.8	6.4	7.8	7.4	8.5	10.4
9 to 11	16.3	14.2	12.7	10.1	16.0	15.6	19.5	22.3	15.7
12 (or GED)	49.1	47.3	48.0	47.7	49.0	49.3	48.3	49.7	51.9
More than 12	26.1	28.6	28.9	30.3	28.6	27.3	24.8	19.5	22.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marital status									
Never married	59.7	51.6	56.5	66.5	55.6	54.4	61.5	76.3	66.2
Currently married	16.5	21.1	20.9	17.1	16.6	18.7	15.7	8.1	14.2
Separated	7.1	8.6	6.7	4.7	8.1	8.1	6.2	4.4	5.8
Divorced	14.8	16.7	13.9	10.5	17.7	16.5	15.1	10.2	12.0
Widowed	1.9	2.0	2.0	1.2	2.0	2.3	1.5	1.1	1.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.5c. Individuals aged 25 to 54 years, by mental health diagnosis, education, and marital status: Row percent distribution, 2018

<i>Education and marital status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 25 to 54 years	100.0	11.2	12.6	1.7	16.1	29.4	1.2	16.4	11.5
Highest school grade completed									
0 to 8	100.0	13.5	14.1	2.1	13.2	28.0	1.2	16.8	11.0
9 to 11	100.0	10.1	9.0	†	17.4	29.3	1.6	23.0	8.6
12 (or GED)	100.0	11.2	11.2	1.5	17.6	30.7	1.3	17.0	9.5
More than 12	100.0	12.7	12.7	1.7	19.4	31.9	1.3	12.6	7.6
Total	100.0	11.6	11.5	1.5	17.7	30.6	1.4	16.8	9.0
Marital status									
Never married	100.0	9.2	9.8	1.5	17.0	28.4	1.1	22.1	10.9
Currently married	100.0	13.7	13.1	1.4	18.4	35.4	†	8.5	8.5
Separated	100.0	12.9	9.8	†	20.9	35.8	†	10.7	8.0
Divorced	100.0	12.0	9.7	†	21.9	34.6	1.1	11.9	7.9
Widowed	100.0	11.4	10.9	†	19.0	37.6	†	10.2	9.3
Total	100.0	10.7	10.3	1.4	18.3	31.1	1.0	17.3	9.8

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.6a. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, education, and marital status: Number, 2018

<i>Education and marital status</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
		<i>disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 25 to 54 years²	1,848,515	152,970	186,603	17,206	338,814	592,722	21,997	370,975	167,228
Highest school grade completed									
0 to 8	73,890	9,200	8,435	835	11,511	22,397	944	15,621	4,947
9 to 11	162,279	13,730	11,767	937	30,970	50,480	2,803	42,463	9,129
12 (or GED)	469,342	43,334	41,885	4,066	94,376	157,792	6,653	93,817	27,419
More than 12	249,244	26,050	25,542	2,507	55,339	87,295	3,353	36,898	12,260
Total	954,755	92,314	87,629	8,345	192,196	317,964	13,753	188,799	53,755
Marital status									
Never married	642,727	38,522	50,526	4,440	126,491	205,788	7,043	170,169	39,748
Currently married	169,815	16,599	18,424	1,280	36,918	68,062	1,813	17,636	9,083
Separated	78,460	7,011	6,745	334	18,757	31,395	705	9,776	3,737
Divorced	165,766	15,119	13,672	838	41,125	62,371	1,733	22,570	8,338
Widowed	19,344	1,842	1,526	73	4,089	8,313	158	2,428	915
Total	1,076,112	79,093	90,893	6,965	227,380	375,929	11,452	222,579	61,821

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.6b. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, education, and marital status: Column percent distribution, 2018

<i>Education and marital status</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 25 to 54 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Highest school grade completed									
0 to 8	7.7	10.0	9.6	10.0	6.0	7.0	6.9	8.3	9.2
9 to 11	17.0	14.9	13.4	11.2	16.1	15.9	20.4	22.5	17.0
12 (or GED)	49.2	46.9	47.8	48.7	49.1	49.6	48.4	49.7	51.0
More than 12	26.1	28.2	29.1	30.0	28.8	27.5	24.4	19.5	22.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marital status									
Never married	59.7	48.7	55.6	63.7	55.6	54.7	61.5	76.5	64.3
Currently married	15.8	21.0	20.3	18.4	16.2	18.1	15.8	7.9	14.7
Separated	7.3	8.9	7.4	4.8	8.2	8.4	6.2	4.4	6.0
Divorced	15.4	19.1	15.0	12.0	18.1	16.6	15.1	10.1	13.5
Widowed	1.8	2.3	1.7	1.0	1.8	2.2	1.4	1.1	1.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma; SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.6c. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, education, and marital status: Row percent distribution, 2018

Education and marital status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years²	100.0	8.3	10.1	†	18.3	32.1	1.2	20.1	9.0
Highest school grade completed									
0 to 8	100.0	12.5	11.4	1.1	15.6	30.3	1.3	21.1	6.7
9 to 11	100.0	8.5	7.3	†	19.1	31.1	1.7	26.2	5.6
12 (or GED)	100.0	9.2	8.9	†	20.1	33.6	1.4	20.0	5.8
More than 12	100.0	10.5	10.2	1.0	22.2	35.0	1.3	14.8	4.9
Total	100.0	9.7	9.2	†	20.1	33.3	1.4	19.8	5.6
Marital status									
Never married	100.0	6.0	7.9	†	19.7	32.0	1.1	26.5	6.2
Currently married	100.0	9.8	10.8	†	21.7	40.1	1.1	10.4	5.3
Separated	100.0	8.9	8.6	†	23.9	40.0	†	12.5	4.8
Divorced	100.0	9.1	8.2	†	24.8	37.6	1.0	13.6	5.0
Widowed	100.0	9.5	7.9	†	21.1	43.0	†	12.6	4.7
Total	100.0	7.3	8.4	†	21.1	34.9	1.1	20.7	5.7

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma; SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.7a. Individuals aged 25 to 54 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Number, 2018

<i>Living arrangement, SMI status, and veteran status</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
		<i>disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 25 to 54 years	2,476,907	276,239	311,401	41,662	398,050	729,056	30,448	405,426	284,625
Living arrangement									
Homeless	103,214	10,677	6,654	571	16,483	28,713	1,236	22,855	16,025
Foster home/foster care	5,362	409	602	125	538	862	106	1,502	1,218
Residential care	60,916	2,885	3,080	468	8,582	10,434	621	24,450	10,396
Institutional setting	21,287	1,083	1,025	134	2,520	3,172	540	10,017	2,796
Jail/correctional facility	38,006	6,922	2,838	250	3,900	6,427	3,207	7,530	6,932
Private residence	1,244,616	135,442	145,140	19,227	217,113	383,601	14,690	200,993	128,410
Other	46,788	8,269	7,652	905	5,456	11,501	341	6,986	5,678
Total	1,520,189	165,687	166,991	21,680	254,592	444,710	20,741	274,333	171,455
SMI status²									
SMI	1,848,515	152,970	186,603	17,206	338,814	592,722	21,997	370,975	167,228
Not SMI	525,651	109,826	102,406	21,339	49,999	110,035	7,164	27,003	97,879
Total	2,374,166	262,796	289,009	38,545	388,813	702,757	29,161	397,978	265,107
Veteran status									
Veteran	24,580	3,938	2,343	335	3,935	7,218	208	4,483	2,120
Not a veteran	913,104	109,620	103,437	12,956	148,416	268,959	8,551	146,548	114,617
Total	937,684	113,558	105,780	13,291	152,351	276,177	8,759	151,031	116,737

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.7b. Individuals aged 25 to 54 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Column percent distribution, 2018

Living arrangement, SMI status, and veteran status	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement									
Homeless	6.8	6.4	4.0	2.6	6.5	6.5	6.0	8.3	9.3
Foster home/foster care	†	†	†	†	†	†	†	†	†
Residential care	4.0	1.7	1.8	2.2	3.4	2.3	3.0	8.9	6.1
Institutional setting	1.4	†	†	†	†	†	2.6	3.7	1.6
Jail/correctional facility	2.5	4.2	1.7	1.2	1.5	1.4	15.5	2.7	4.0
Private residence	81.9	81.7	86.9	88.7	85.3	86.3	70.8	73.3	74.9
Other	3.1	5.0	4.6	4.2	2.1	2.6	1.6	2.5	3.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SMI status²									
SMI	77.9	58.2	64.6	44.6	87.1	84.3	75.4	93.2	63.1
Not SMI	22.1	41.8	35.4	55.4	12.9	15.7	24.6	6.8	36.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status									
Veteran	2.6	3.5	2.2	2.5	2.6	2.6	2.4	3.0	1.8
Not a veteran	97.4	96.5	97.8	97.5	97.4	97.4	97.6	97.0	98.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.7c. Individuals aged 25 to 54 years, by mental health diagnosis, living arrangement, SMI status, and veteran status: Row percent distribution, 2018

Living arrangement, SMI status, and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	100.0	11.2	12.6	1.7	16.1	29.4	1.2	16.4	11.5
Living arrangement									
Homeless	100.0	10.3	6.4	†	16.0	27.8	1.2	22.1	15.5
Foster home/foster care	100.0	7.6	11.2	2.3	10.0	16.1	2.0	28.0	22.7
Residential care	100.0	4.7	5.1	†	14.1	17.1	1.0	40.1	17.1
Institutional setting	100.0	5.1	4.8	†	11.8	14.9	2.5	47.1	13.1
Jail/correctional facility	100.0	18.2	7.5	†	10.3	16.9	8.4	19.8	18.2
Private residence	100.0	10.9	11.7	1.5	17.4	30.8	1.2	16.1	10.3
Other	100.0	17.7	16.4	1.9	11.7	24.6	†	14.9	12.1
Total	100.0	10.9	11.0	1.4	16.7	29.3	1.4	18.0	11.3
SMI status²									
SMI	100.0	8.3	10.1	†	18.3	32.1	1.2	20.1	9.0
Not SMI	100.0	20.9	19.5	4.1	9.5	20.9	1.4	5.1	18.6
Total	100.0	11.1	12.2	1.6	16.4	29.6	1.2	16.8	11.2
Veteran status									
Veteran	100.0	16.0	9.5	1.4	16.0	29.4	†	18.2	8.6
Not a veteran	100.0	12.0	11.3	1.4	16.3	29.5	†	16.0	12.6
Total	100.0	12.1	11.3	1.4	16.2	29.5	†	16.1	12.4

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.8a. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, living arrangement, and veteran status: Number, 2018

<i>Living arrangement and veteran status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 25 to 54 years²	1,848,515	152,970	186,603	17,206	338,814	592,722	21,997	370,975	167,228
Living arrangement									
Homeless	87,984	8,104	4,681	295	15,057	25,724	1,001	21,480	11,642
Foster home/foster care	3,396	268	291	41	367	615	85	1,288	441
Residential care	51,762	2,003	2,093	225	8,035	9,414	542	23,586	5,864
Institutional setting	16,056	698	666	76	1,923	2,207	306	8,782	1,398
Jail/correctional facility	28,494	3,449	2,027	128	3,420	5,352	3,048	6,902	4,168
Private residence	1,000,892	90,672	93,090	9,012	197,605	333,077	11,932	189,605	75,899
Other	37,182	4,492	6,355	138	5,090	10,599	284	6,566	3,658
Total	1,225,766	109,686	109,203	9,915	231,497	386,988	17,198	258,209	103,070
Veteran status									
Veteran	19,950	2,787	1,478	143	3,589	6,493	165	4,239	1,056
Not a veteran	705,573	72,882	62,330	5,384	132,629	234,169	6,750	134,722	56,707
Total	725,523	75,669	63,808	5,527	136,218	240,662	6,915	138,961	57,763

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.8b. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, living arrangement, and veteran status: Column percent distribution, 2018

<i>Living arrangement and veteran status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 25 to 54 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement									
Homeless	7.2	7.4	4.3	3.0	6.5	6.6	5.8	8.3	11.3
Foster home/foster care	†	†	†	†	†	†	†	†	†
Residential care	4.2	1.8	1.9	2.3	3.5	2.4	3.2	9.1	5.7
Institutional setting	1.3	†	†	†	†	†	1.8	3.4	1.4
Jail/correctional facility	2.3	3.1	1.9	1.3	1.5	1.4	17.7	2.7	4.0
Private residence	81.7	82.7	85.2	90.9	85.4	86.1	69.4	73.4	73.6
Other	3.0	4.1	5.8	1.4	2.2	2.7	1.7	2.5	3.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status									
Veteran	2.7	3.7	2.3	2.6	2.6	2.7	2.4	3.1	1.8
Not a veteran	97.3	96.3	97.7	97.4	97.4	97.3	97.6	96.9	98.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.8c. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, living arrangement, and veteran status: Row percent distribution, 2018

<i>Living arrangement and veteran status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 25 to 54 years²	100.0	8.3	10.1	†	18.3	32.1	1.2	20.1	9.0
Living arrangement									
Homeless	100.0	9.2	5.3	†	17.1	29.2	1.1	24.4	13.2
Foster home/foster care	100.0	7.9	8.6	1.2	10.8	18.1	2.5	37.9	13.0
Residential care	100.0	3.9	4.0	†	15.5	18.2	1.0	45.6	11.3
Institutional setting	100.0	4.3	4.1	†	12.0	13.7	1.9	54.7	8.7
Jail/correctional facility	100.0	12.1	7.1	†	12.0	18.8	10.7	24.2	14.6
Private residence	100.0	9.1	9.3	†	19.7	33.3	1.2	18.9	7.6
Other	100.0	12.1	17.1	†	13.7	28.5	†	17.7	9.8
Total	100.0	8.9	8.9	†	18.9	31.6	1.4	21.1	8.4
Veteran status									
Veteran	100.0	14.0	7.4	†	18.0	32.5	†	21.2	5.3
Not a veteran	100.0	10.3	8.8	†	18.8	33.2	†	19.1	8.0
Total	100.0	10.4	8.8	†	18.8	33.2	†	19.2	8.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.9a. Individuals aged 25 to 54 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 25 to 54 years	2,476,907	276,239	311,401	41,662	398,050	729,056	30,448	405,426	284,625
Employment status									
Employed	348,291	50,051	53,252	8,219	54,593	114,711	4,397	27,904	35,164
Full time	183,124	31,503	33,627	5,401	22,137	57,218	2,744	9,979	20,515
Part time	113,256	14,803	15,480	2,176	18,166	35,944	1,417	12,319	12,951
Unemployed	420,435	45,974	41,679	4,632	75,253	140,141	4,160	62,575	46,021
Not in labor force (see detail below)	571,672	53,137	57,882	7,060	104,447	157,168	7,866	130,318	53,794
Total	1,340,398	149,162	152,813	19,911	234,293	412,020	16,423	220,797	134,979
Detailed not in labor force									
Total individuals aged 25 to 54 years not in labor force	571,672	53,137	57,882	7,060	104,447	157,168	7,866	130,318	53,794
Disabled	293,779	21,594	22,979	2,258	56,976	73,030	4,559	85,665	26,718
Retired	3,300	401	270	17	491	958	36	714	413
Student	20,736	2,388	2,562	641	2,660	5,444	160	3,162	3,719
Homemaker	32,410	5,255	5,202	354	4,789	11,695	317	1,911	2,887
Sheltered/non-competitive employment	8,579	647	698	143	1,102	1,858	34	1,631	2,466
Other	212,868	22,852	26,171	3,647	38,429	64,183	2,760	37,235	17,591

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTES: *Employed* includes “full-time/part-time not differentiated,” which is not presented separately; total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.9b. Individuals aged 25 to 54 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
		<i>disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 25 to 54 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Employment status									
Employed	26.0	33.6	34.8	41.3	23.3	27.8	26.8	12.6	26.1
Full time	13.7	21.1	22.0	27.1	9.4	13.9	16.7	4.5	15.2
Part time	8.4	9.9	10.1	10.9	7.8	8.7	8.6	5.6	9.6
Unemployed	31.4	30.8	27.3	23.3	32.1	34.0	25.3	28.3	34.1
Not in labor force (see detail below)	42.6	35.6	37.9	35.5	44.6	38.1	47.9	59.0	39.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force									
Total individuals aged 25 to 54 years not in labor force	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Disabled	51.4	40.6	39.7	32.0	54.6	46.5	58.0	65.7	49.7
Retired	†	†	†	†	†	†	†	†	†
Student	3.6	4.5	4.4	9.1	2.5	3.5	2.0	2.4	6.9
Homemaker	5.7	9.9	9.0	5.0	4.6	7.4	4.0	1.5	5.4
Sheltered/non-competitive employment	1.5	1.2	1.2	2.0	1.1	1.2	†	1.3	4.6
Other	37.2	43.0	45.2	51.7	36.8	40.8	35.1	28.6	32.7

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTES: Percentages may not sum to 100 percent because of rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.9c. Individuals aged 25 to 54 years, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 25 to 54 years	100.0	11.2	12.6	1.7	16.1	29.4	1.2	16.4	11.5
Employment status									
Employed	100.0	14.4	15.3	2.4	15.7	32.9	1.3	8.0	10.1
Full time	100.0	17.2	18.4	2.9	12.1	31.2	1.5	5.4	11.2
Part time	100.0	13.1	13.7	1.9	16.0	31.7	1.3	10.9	11.4
Unemployed	100.0	10.9	9.9	1.1	17.9	33.3	†	14.9	10.9
Not in labor force (see detail below)	100.0	9.3	10.1	1.2	18.3	27.5	1.4	22.8	9.4
Total	100.0	11.1	11.4	1.5	17.5	30.7	1.2	16.5	10.1
Detailed not in labor force									
Total individuals aged 25 to 54 years not in labor force	100.0	9.3	10.1	1.2	18.3	27.5	1.4	22.8	9.4
Disabled	100.0	7.4	7.8	†	19.4	24.9	1.6	29.2	9.1
Retired	100.0	12.2	8.2	†	14.9	29.0	1.1	21.6	12.5
Student	100.0	11.5	12.4	3.1	12.8	26.3	†	15.2	17.9
Homemaker	100.0	16.2	16.1	1.1	14.8	36.1	†	5.9	8.9
Sheltered/non-competitive employment	100.0	7.5	8.1	1.7	12.8	21.7	†	19.0	28.7
Other	100.0	10.7	12.3	1.7	18.1	30.2	1.3	17.5	8.3

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTES: Percentages may not sum to 100 percent because of rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.10a. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 25 to 54 years²	1,848,515	152,970	186,603	17,206	338,814	592,722	21,997	370,975	167,228
Employment status									
Employed	259,467	30,341	33,021	3,857	48,610	96,575	3,411	25,725	17,927
Full time	126,792	18,948	20,730	2,610	18,388	45,277	2,090	8,690	10,059
Part time	88,402	9,717	10,600	1,066	16,245	31,050	1,134	11,521	7,069
Unemployed	339,105	31,040	27,730	2,269	68,638	123,518	3,195	58,502	24,213
Not in labor force (see detail below)	469,077	37,902	38,609	3,445	94,993	136,256	6,628	122,783	28,461
Total	1,067,649	99,283	99,360	9,571	212,241	356,349	13,234	207,010	70,601
Detailed not in labor force									
Total individuals with SMI aged 25 to 54 years not in labor force	469,077	37,902	38,609	3,445	94,993	136,256	6,628	122,783	28,461
Disabled	251,326	16,915	16,724	1,344	52,181	64,528	4,061	81,204	14,369
Retired	2,783	309	181	10	468	850	23	680	262
Student	17,047	1,656	1,730	318	2,453	4,944	137	3,053	2,756
Homemaker	25,443	3,497	3,604	198	4,342	9,938	268	1,792	1,804
Sheltered/non-competitive employment	6,303	539	481	65	1,009	1,682	26	1,498	1,003
Other	166,175	14,986	15,889	1,510	34,540	54,314	2,113	34,556	8,267

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTES: *Employed* includes “full-time/part-time not differentiated,” which is not presented separately; total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.10b. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 25 to 54 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Employment status									
Employed	24.3	30.6	33.2	40.3	22.9	27.1	25.8	12.4	25.4
Full time	11.9	19.1	20.9	27.3	8.7	12.7	15.8	4.2	14.2
Part time	8.3	9.8	10.7	11.1	7.7	8.7	8.6	5.6	10.0
Unemployed	31.8	31.3	27.9	23.7	32.3	34.7	24.1	28.3	34.3
Not in labor force (see detail below)	43.9	38.2	38.9	36.0	44.8	38.2	50.1	59.3	40.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force									
Total individuals with SMI aged 25 to 54 years not in labor force	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Disabled	53.6	44.6	43.3	39.0	54.9	47.4	61.3	66.1	50.5
Retired	†	†	†	†	†	†	†	†	†
Student	3.6	4.4	4.5	9.2	2.6	3.6	2.1	2.5	9.7
Homemaker	5.4	9.2	9.3	5.7	4.6	7.3	4.0	1.5	6.3
Sheltered/non-competitive employment	1.3	1.4	1.2	1.9	1.1	1.2	†	1.2	3.5
Other	35.4	39.5	41.2	43.8	36.4	39.9	31.9	28.1	29.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTES: Percentages may not sum to 100 percent because of rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.10c. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2018

Employment status and ‘detailed not in labor force’	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years²	100.0	8.3	10.1	†	18.3	32.1	1.2	20.1	9.0
Employment status									
Employed	100.0	11.7	12.7	1.5	18.7	37.2	1.3	9.9	6.9
Full time	100.0	14.9	16.3	2.1	14.5	35.7	1.6	6.9	7.9
Part time	100.0	11.0	12.0	1.2	18.4	35.1	1.3	13.0	8.0
Unemployed	100.0	9.2	8.2	†	20.2	36.4	†	17.3	7.1
Not in labor force (see detail below)	100.0	8.1	8.2	†	20.3	29.0	1.4	26.2	6.1
Total	100.0	9.3	9.3	†	19.9	33.4	1.2	19.4	6.6
Detailed not in labor force									
Total individuals with SMI aged 25 to 54 years not in labor force	100.0	8.1	8.2	†	20.3	29.0	1.4	26.2	6.1
Disabled	100.0	6.7	6.7	†	20.8	25.7	1.6	32.3	5.7
Retired	100.0	11.1	6.5	†	16.8	30.5	†	24.4	9.4
Student	100.0	9.7	10.1	1.9	14.4	29.0	†	17.9	16.2
Homemaker	100.0	13.7	14.2	†	17.1	39.1	1.1	7.0	7.1
Sheltered/non-competitive employment	100.0	8.6	7.6	1.0	16.0	26.7	†	23.8	15.9
Other	100.0	9.0	9.6	†	20.8	32.7	1.3	20.8	5.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTES: Percentages may not sum to 100 percent because of rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.11a. Individuals aged 25 to 54 years, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018

<i>Timing of admission and mental health service setting</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 25 to 54 years	2,476,907	276,239	311,401	41,662	398,050	729,056	30,448	405,426	284,625
Timing of admission									
New individuals	1,271,179	172,666	181,656	21,660	173,495	395,846	14,750	144,101	167,005
Continuing individuals	1,205,728	103,573	129,745	20,002	224,555	333,210	15,698	261,325	117,620
Mental health service setting^{2,3}									
State psychiatric hospital	64,656	2,217	1,192	155	10,095	8,013	2,038	34,470	6,476
Community-based program	2,412,139	271,414	309,655	41,515	390,323	716,767	29,347	381,608	271,510
Residential treatment center	24,181	1,415	1,089	167	3,176	4,708	252	9,001	4,373
Other psychiatric inpatient	131,186	7,960	8,793	732	25,545	39,054	1,587	35,296	12,219
Institutions under the justice system	40,067	7,584	2,052	106	2,332	4,832	3,479	8,535	11,147

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.11b. Individuals aged 25 to 54 years, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018

Timing of admission and mental health service setting	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission									
New individuals	51.3	62.5	58.3	52.0	43.6	54.3	48.4	35.5	58.7
Continuing individuals	48.7	37.5	41.7	48.0	56.4	45.7	51.6	64.5	41.3
Mental health service setting^{2,3}									
State psychiatric hospital	2.6	†	†	†	2.5	1.1	6.7	8.5	2.3
Community-based program	97.4	98.3	99.4	99.6	98.1	98.3	96.4	94.1	95.4
Residential treatment center	†	†	†	†	†	†	†	2.2	1.5
Other psychiatric inpatient	5.3	2.9	2.8	1.8	6.4	5.4	5.2	8.7	4.3
Institutions under the justice system	1.6	2.7	†	†	†	†	11.4	2.1	3.9

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages for timing of admission may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.11c. Individuals aged 25 to 54 years, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018

Timing of admission and mental health service setting	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 25 to 54 years	100.0	11.2	12.6	1.7	16.1	29.4	1.2	16.4	11.5
Timing of admission									
New individuals	100.0	13.6	14.3	1.7	13.6	31.1	1.2	11.3	13.1
Continuing individuals	100.0	8.6	10.8	1.7	18.6	27.6	1.3	21.7	9.8
Mental health service setting^{2,3}									
State psychiatric hospital	100.0	3.4	1.8	†	15.6	12.4	3.2	53.3	10.0
Community-based program	100.0	11.3	12.8	1.7	16.2	29.7	1.2	15.8	11.3
Residential treatment center	100.0	5.9	4.5	†	13.1	19.5	1.0	37.2	18.1
Other psychiatric inpatient	100.0	6.1	6.7	†	19.5	29.8	1.2	26.9	9.3
Institutions under the justice system	100.0	18.9	5.1	†	5.8	12.1	8.7	21.3	27.8

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.12a. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years²	1,848,515	152,970	186,603	17,206	338,814	592,722	21,997	370,975	167,228
Timing of admission									
New individuals	903,036	94,498	110,554	9,490	143,894	314,359	9,186	126,443	94,612
Continuing individuals	945,479	58,472	76,049	7,716	194,920	278,363	12,811	244,532	72,616
Mental health service setting^{3,4}									
State psychiatric hospital	54,873	1,650	901	110	8,763	6,604	1,627	31,297	3,921
Community-based program	1,794,157	149,129	185,291	17,118	332,497	582,054	21,254	349,894	156,920
Residential treatment center	21,023	946	746	103	2,871	4,100	214	8,657	3,386
Other psychiatric inpatient	107,210	4,597	5,449	402	21,990	32,454	1,318	32,700	8,300
Institutions under the justice system	35,565	4,921	1,881	81	2,201	4,536	3,435	8,417	10,093

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.12b. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018

Timing of admission and mental health service setting	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 25 to 54 years²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission									
New individuals	48.9	61.8	59.2	55.2	42.5	53.0	41.8	34.1	56.6
Continuing individuals	51.1	38.2	40.8	44.8	57.5	47.0	58.2	65.9	43.4
Mental health service setting^{3,4}									
State psychiatric hospital	3.0	1.1	†	†	2.6	1.1	7.4	8.4	2.3
Community-based program	97.1	97.5	99.3	99.5	98.1	98.2	96.6	94.3	93.8
Residential treatment center	1.1	†	†	†	†	†	†	2.3	2.0
Other psychiatric inpatient	5.8	3.0	2.9	2.3	6.5	5.5	6.0	8.8	5.0
Institutions under the justice system	1.9	3.2	1.0	†	†	†	15.6	2.3	6.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages for timing of admission may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.12c. Individuals aged 25 to 54 years with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018

<i>Timing of admission and mental health service setting</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 25 to 54 years²	100.0	8.3	10.1	†	18.3	32.1	1.2	20.1	9.0
Timing of admission									
New individuals	100.0	10.5	12.2	1.1	15.9	34.8	1.0	14.0	10.5
Continuing individuals	100.0	6.2	8.0	†	20.6	29.4	1.4	25.9	7.7
Mental health service setting^{3,4}									
State psychiatric hospital	100.0	3.0	1.6	†	16.0	12.0	3.0	57.0	7.1
Community-based program	100.0	8.3	10.3	†	18.5	32.4	1.2	19.5	8.7
Residential treatment center	100.0	4.5	3.5	†	13.7	19.5	1.0	41.2	16.1
Other psychiatric inpatient	100.0	4.3	5.1	†	20.5	30.3	1.2	30.5	7.7
Institutions under the justice system	100.0	13.8	5.3	†	6.2	12.8	9.7	23.7	28.4

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 4.13. Individuals aged 25 to 54 years, by mental health service setting and SMI status: Number and percent distribution, 2018

<i>Mental health service setting</i>	<i>SMI status¹</i>								
	<i>Number of individuals</i>			<i>Column percent of individuals</i>			<i>Row percent distribution of individuals</i>		
	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>
Total individuals aged 25 to 54 years	2,654,952	1,973,914	681,038	100.0	100.0	100.0	100.0	74.3	25.7
Mental health service setting^{2,3}									
State psychiatric hospital	66,570	60,372	6,198	2.5	3.1	†	100.0	90.7	9.3
Community-based program	2,584,599	1,912,696	671,903	97.4	96.9	98.7	100.0	74.0	26.0
Residential treatment center	26,300	22,262	4,038	†	1.1	†	100.0	84.6	15.4
Other psychiatric inpatient	139,763	111,880	27,883	5.3	5.7	4.1	100.0	80.0	20.0
Institutions under the justice system	42,646	37,915	4,731	1.6	1.9	†	100.0	88.9	11.1

† Less than 1 percent of individuals.

SMI = serious emotional disturbance.

¹ The federal definition for SMI status can be found in Appendix D.

² Individuals could receive services in more than one mental health service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.1a. Individuals aged 55 years and older, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals aged 55 years and older	769,081	843,771	802,401	894,823	908,917	932,573
Gender						
Male	308,152	343,021	328,387	368,572	381,778	393,913
Female	460,273	499,360	472,697	523,535	523,113	537,368
Total	768,425	842,381	801,084	892,107	904,891	931,281
Age¹						
55 to 59 years	337,373	368,375	350,060	385,959	389,871	402,710
60 to 64 years	203,503	225,348	220,758	249,173	255,186	273,170
65 years and older	228,205	250,048	231,583	259,691	263,860	256,693
Total	769,081	843,771	802,401	894,823	908,917	932,573
Race						
American Indian or Alaska Native	5,883	7,066	7,502	8,336	8,065	8,849
Asian ²	13,120	15,292	15,049	17,264	17,564	19,385
Black or African American	128,520	144,630	137,884	156,840	163,511	175,479
Native Hawaiian or Other Pacific Islander	1,904	2,049	2,222	2,515	1,545	1,776
White	524,168	566,342	532,049	594,172	590,317	613,288
Other	35,715	41,532	37,444	43,717	50,868	54,359
Total	709,310	776,911	732,150	822,844	831,870	873,136
Ethnicity						
Not of Hispanic or Latino origin	539,090	584,680	542,769	608,005	709,514	738,800
Hispanic or Latino origin	56,716	65,860	65,796	74,605	83,462	82,984
Total	595,806	650,540	608,565	682,610	792,976	821,784
Specific Hispanic or Latino origin						
Mexican	1,501	2,266	2,786	2,523	2,960	2,952
Puerto Rican	3,709	10,283	11,018	9,763	12,830	5,007
Cuban	3,242	3,131	3,303	3,155	3,071	3,729
Other specific	5,005	5,742	5,698	6,253	6,920	7,332
Total	13,457	21,422	22,805	21,694	25,781	19,020

Continued. See notes at end of table.

Table 5.1a. Individuals aged 55 years and older, by gender, age, race, ethnicity, education, veteran status, and SMI status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Highest school grade completed						
0 to 8	53,662	61,936	64,324	66,708	55,504	53,792
9 to 11	48,342	52,752	56,755	60,452	58,711	62,033
12 (or GED)	141,712	165,836	181,050	196,706	176,720	198,585
More than 12	60,009	71,578	77,422	86,569	100,645	109,891
Total	303,725	352,102	379,551	410,435	391,580	424,301
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	18,028	17,795
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	303,590	313,206
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	321,618	331,001
SMI status³						
SMI	530,075	586,398	564,150	613,978	651,764	697,908
Not SMI	176,258	197,894	194,518	223,840	195,277	187,208
Total	706,333	784,292	758,668	837,818	847,041	885,116

n/c Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 5.1b. Individuals aged 55 years and older, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	47	45	47	48	48
Total individuals aged 55 years and older	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	40.1	40.7	41.0	41.3	42.2	42.3
Female	59.9	59.3	59.0	58.7	57.8	57.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age¹						
55 to 59 years	43.9	43.7	43.6	43.1	42.9	43.2
60 to 64 years	26.5	26.7	27.5	27.8	28.1	29.3
65 years and older	29.7	29.6	28.9	29.0	29.0	27.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	†	†	1.0	1.0	†	1.0
Asian ²	1.8	2.0	2.1	2.1	2.1	2.2
Black or African American	18.1	18.6	18.8	19.1	19.7	20.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	73.9	72.9	72.7	72.2	71.0	70.2
Other	5.0	5.3	5.1	5.3	6.1	6.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	90.5	89.9	89.2	89.1	89.5	89.9
Hispanic or Latino origin	9.5	10.1	10.8	10.9	10.5	10.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	11.2	10.6	12.2	11.6	11.5	15.5
Puerto Rican	27.6	48.0	48.3	45.0	49.8	26.3
Cuban	24.1	14.6	14.5	14.5	11.9	19.6
Other specific	37.2	26.8	25.0	28.8	26.8	38.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 5.1b. Individuals aged 55 years and older, by gender, age, race, ethnicity, education, veteran status, and SMI status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Highest school grade completed						
0 to 8	17.7	17.6	16.9	16.3	14.2	12.7
9 to 11	15.9	15.0	15.0	14.7	15.0	14.6
12 (or GED)	46.7	47.1	47.7	47.9	45.1	46.8
More than 12	19.8	20.3	20.4	21.1	25.7	25.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	5.6	5.4
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	94.4	94.6
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0
SMI status³						
SMI	75.0	74.8	74.4	73.3	76.9	78.8
Not SMI	25.0	25.2	25.6	26.7	23.1	21.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ Age is based on the individual's age at the midpoint of the reporting period.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

³ The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 5.2a. Individuals aged 55 years and older with SMI, by gender, age, race, ethnicity, education, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	44	47	45	46	46	47
Total individuals with SMI aged 55 years and older¹	530,075	586,398	564,150	613,978	651,764	697,908
Gender						
Male	209,242	237,256	229,413	251,369	270,700	290,740
Female	320,485	348,381	334,090	361,892	380,520	406,723
Total	529,727	585,637	563,503	613,261	651,220	697,463
Age²						
55 to 59 years	241,638	266,295	254,383	275,110	288,257	307,721
60 to 64 years	144,853	161,821	158,912	175,644	187,340	207,111
65 years and older	143,584	158,282	150,855	163,224	176,167	183,076
Total	530,075	586,398	564,150	613,978	651,764	697,908
Race						
American Indian or Alaska Native	3,531	4,528	4,628	5,141	5,619	6,081
Asian ³	9,861	11,889	11,825	12,294	13,134	14,603
Black or African American	95,387	105,161	105,151	114,705	123,068	136,346
Native Hawaiian or Other Pacific Islander	1,657	1,810	1,931	2,100	1,233	1,392
White	353,640	384,637	363,651	402,168	425,175	452,882
Other	26,207	31,998	29,378	32,523	40,886	44,828
Total	490,283	540,023	516,564	568,931	609,115	656,132
Ethnicity						
Not of Hispanic or Latino origin	385,235	416,542	398,994	437,087	505,022	542,229
Hispanic or Latino origin	43,603	52,485	51,350	58,076	67,196	69,953
Total	428,838	469,027	450,344	495,163	572,218	612,182
Specific Hispanic or Latino origin						
Mexican	1,110	1,774	2,222	1,952	2,359	2,357
Puerto Rican	3,097	7,780	8,380	7,529	9,388	4,164
Cuban	2,938	2,997	3,192	3,047	2,950	3,634
Other specific	4,206	4,800	4,872	5,268	5,745	6,281
Total	11,351	17,351	18,666	17,796	20,442	16,436

Continued. See notes at end of table.

**Table 5.2a. Individuals aged 55 years and older with SMI, by gender, age, race, ethnicity, education, and veteran status: Number, 2013–2018
(continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Highest school grade completed						
0 to 8	33,170	39,900	40,315	42,471	40,705	40,092
9 to 11	35,356	40,017	42,770	46,767	49,618	52,752
12 (or GED)	92,752	112,510	122,290	137,810	143,206	163,928
More than 12	44,029	56,984	60,847	70,359	83,727	92,064
Total	205,307	249,411	266,222	297,407	317,256	348,836
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	13,283	13,742
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	226,296	248,449
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	239,579	262,191

n/c Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ The federal definition for SMI status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 5.2b. Individuals aged 55 years and older with SMI, by gender, age, race, ethnicity, education, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	44	47	45	46	46	47
Total individuals with SMI aged 55 years and older¹	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	39.5	40.5	40.7	41.0	41.6	41.7
Female	60.5	59.5	59.3	59.0	58.4	58.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age²						
55 to 59 years	45.6	45.4	45.1	44.8	44.2	44.1
60 to 64 years	27.3	27.6	28.2	28.6	28.7	29.7
65 years and older	27.1	27.0	26.7	26.6	27.0	26.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ³	2.0	2.2	2.3	2.2	2.2	2.2
Black or African American	19.5	19.5	20.4	20.2	20.2	20.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	72.1	71.2	70.4	70.7	69.8	69.0
Other	5.3	5.9	5.7	5.7	6.7	6.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	89.8	88.8	88.6	88.3	88.3	88.6
Hispanic or Latino origin	10.2	11.2	11.4	11.7	11.7	11.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	9.8	10.2	11.9	11.0	11.5	14.3
Puerto Rican	27.3	44.8	44.9	42.3	45.9	25.3
Cuban	25.9	17.3	17.1	17.1	14.4	22.1
Other specific	37.1	27.7	26.1	29.6	28.1	38.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 5.2b. Individuals aged 55 years and older with SMI, by gender, age, race, ethnicity, education, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Highest school grade completed						
0 to 8	16.2	16.0	15.1	14.3	12.8	11.5
9 to 11	17.2	16.0	16.1	15.7	15.6	15.1
12 (or GED)	45.2	45.1	45.9	46.3	45.1	47.0
More than 12	21.4	22.8	22.9	23.7	26.4	26.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	5.5	5.2
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	94.5	94.8
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

GED = general equivalency diploma; SMI = serious mental illness.

¹ The federal definition for SMI status can be found in Appendix D.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 5.3a. Individuals aged 55 years and older, by mental health diagnosis, gender, race, and ethnicity: Number, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	827,060	59,957	76,046	6,966	115,042	291,333	7,301	194,943	75,472
Gender									
Male	341,045	23,132	26,196	3,492	39,917	102,726	3,382	103,781	38,419
Female	485,302	36,766	49,773	3,471	75,040	188,308	3,912	91,053	36,979
Total	826,347	59,898	75,969	6,963	114,957	291,034	7,294	194,834	75,398
Race									
American Indian or Alaska Native	8,154	959	954	87	886	2,747	86	1,356	1,079
Asian ²	16,348	1,780	1,269	139	1,009	6,889	81	4,175	1,006
Black or African American	153,387	9,452	7,684	1,289	16,082	48,456	1,219	58,305	10,900
Native Hawaiian or Other Pacific Islander	1,664	174	140	19	156	588	23	446	118
White	547,171	39,635	56,510	4,593	86,756	194,271	5,305	109,578	50,523
Other	50,080	3,117	3,801	443	5,281	20,733	274	11,372	5,059
Total	776,804	55,117	70,358	6,570	110,170	273,684	6,988	185,232	68,685
Ethnicity									
Not of Hispanic or Latino origin	657,403	48,428	58,864	5,895	96,531	227,566	6,214	157,530	56,375
Hispanic or Latino origin	73,389	4,714	6,381	396	7,948	31,328	362	15,290	6,970
Total	730,792	53,142	65,245	6,291	104,479	258,894	6,576	172,820	63,345
Specific Hispanic or Latino origin									
Mexican	2,860	329	264	10	328	1,094	18	596	221
Puerto Rican	4,327	264	316	19	599	1,855	18	966	290
Cuban	3,111	134	232	11	317	1,393	2	611	411
Other specific	6,438	581	628	49	795	2,575	27	1,173	610
Total	16,736	1,308	1,440	89	2,039	6,917	65	3,346	1,532

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.3b. Individuals aged 55 years and older, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	41.3	38.6	34.5	50.2	34.7	35.3	46.4	53.3	51.0
Female	58.7	61.4	65.5	49.8	65.3	64.7	53.6	46.7	49.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	1.0	1.7	1.4	1.3	†	1.0	1.2	†	1.6
Asian ²	2.1	3.2	1.8	2.1	†	2.5	1.2	2.3	1.5
Black or African American	19.7	17.1	10.9	19.6	14.6	17.7	17.4	31.5	15.9
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	70.4	71.9	80.3	69.9	78.7	71.0	75.9	59.2	73.6
Other	6.4	5.7	5.4	6.7	4.8	7.6	3.9	6.1	7.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity									
Not of Hispanic or Latino origin	90.0	91.1	90.2	93.7	92.4	87.9	94.5	91.2	89.0
Hispanic or Latino origin	10.0	8.9	9.8	6.3	7.6	12.1	5.5	8.8	11.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	17.1	25.2	18.3	11.2	16.1	15.8	27.7	17.8	14.4
Puerto Rican	25.9	20.2	21.9	21.3	29.4	26.8	27.7	28.9	18.9
Cuban	18.6	10.2	16.1	12.4	15.5	20.1	3.1	18.3	26.8
Other specific	38.5	44.4	43.6	55.1	39.0	37.2	41.5	35.1	39.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.3c. Individuals aged 55 years and older, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	100.0	7.2	9.2	†	13.9	35.2	†	23.6	9.1
Gender									
Male	100.0	6.8	7.7	1.0	11.7	30.1	†	30.4	11.3
Female	100.0	7.6	10.3	†	15.5	38.8	†	18.8	7.6
Total	100.0	7.2	9.2	†	13.9	35.2	†	23.6	9.1
Race									
American Indian or Alaska Native	100.0	11.8	11.7	1.1	10.9	33.7	1.1	16.6	13.2
Asian ²	100.0	10.9	7.8	†	6.2	42.1	†	25.5	6.2
Black or African American	100.0	6.2	5.0	†	10.5	31.6	†	38.0	7.1
Native Hawaiian or Other Pacific Islander	100.0	10.5	8.4	1.1	9.4	35.3	1.4	26.8	7.1
White	100.0	7.2	10.3	†	15.9	35.5	†	20.0	9.2
Other	100.0	6.2	7.6	†	10.5	41.4	†	22.7	10.1
Total	100.0	7.1	9.1	†	14.2	35.2	†	23.8	8.8
Ethnicity									
Not of Hispanic or Latino origin	100.0	7.4	9.0	†	14.7	34.6	†	24.0	8.6
Hispanic or Latino origin	100.0	6.4	8.7	†	10.8	42.7	†	20.8	9.5
Total	100.0	7.3	8.9	†	14.3	35.4	†	23.6	8.7
Specific Hispanic or Latino origin									
Mexican	100.0	11.5	9.2	†	11.5	38.3	†	20.8	7.7
Puerto Rican	100.0	6.1	7.3	†	13.8	42.9	†	22.3	6.7
Cuban	100.0	4.3	7.5	†	10.2	44.8	†	19.6	13.2
Other specific	100.0	9.0	9.8	†	12.3	40.0	†	18.2	9.5
Total	100.0	7.8	8.6	†	12.2	41.3	†	20.0	9.2

† Less than 1 percent of individuals.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.4a. Individuals aged 55 years and older with SMI, by mental health diagnosis, gender, race, and ethnicity: Number, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older²	655,241	33,694	47,785	3,504	100,014	242,891	4,881	177,878	44,594
Gender									
Male	270,098	12,533	16,248	1,779	34,536	85,421	2,234	94,997	22,350
Female	384,741	21,142	31,511	1,724	65,427	157,289	2,642	82,800	22,206
Total	654,839	33,675	47,759	3,503	99,963	242,710	4,876	177,797	44,556
Race									
American Indian or Alaska Native	5,845	551	573	31	748	2,135	56	1,201	550
Asian ³	14,046	1,223	893	99	939	6,066	41	4,009	776
Black or African American	128,247	4,851	4,803	673	14,386	42,150	955	53,796	6,633
Native Hawaiian or Other Pacific Islander	1,340	121	65	16	137	501	10	412	78
White	423,471	21,939	34,863	2,239	74,902	158,639	3,430	98,952	28,507
Other	43,218	2,051	2,724	257	4,827	18,758	201	10,707	3,693
Total	616,167	30,736	43,921	3,315	95,939	228,249	4,693	169,077	40,237
Ethnicity									
Not of Hispanic or Latino origin	510,809	25,843	35,074	2,902	83,372	185,258	4,137	142,724	31,499
Hispanic or Latino origin	64,071	3,198	4,643	228	7,391	28,666	261	14,577	5,107
Total	574,880	29,041	39,717	3,130	90,763	213,924	4,398	157,301	36,606
Specific Hispanic or Latino origin									
Mexican	2,320	173	145	7	302	1,013	9	560	111
Puerto Rican	3,824	193	223	12	574	1,725	12	941	144
Cuban	3,032	122	214	10	314	1,371	2	604	395
Other specific	5,655	378	469	24	752	2,443	17	1,124	448
Total	14,831	866	1,051	53	1,942	6,552	40	3,229	1,098

SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.4b. Individuals aged 55 years and older with SMI, by mental health diagnosis, gender, race, and ethnicity: Column percent distribution, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	41.2	37.2	34.0	50.8	34.5	35.2	45.8	53.4	50.2
Female	58.8	62.8	66.0	49.2	65.5	64.8	54.2	46.6	49.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	1.8	1.3	†	†	†	1.2	†	1.4
Asian ³	2.3	4.0	2.0	3.0	†	2.7	†	2.4	1.9
Black or African American	20.8	15.8	10.9	20.3	15.0	18.5	20.3	31.8	16.5
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	68.7	71.4	79.4	67.5	78.1	69.5	73.1	58.5	70.8
Other	7.0	6.7	6.2	7.8	5.0	8.2	4.3	6.3	9.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity									
Not of Hispanic or Latino origin	88.9	89.0	88.3	92.7	91.9	86.6	94.1	90.7	86.0
Hispanic or Latino origin	11.1	11.0	11.7	7.3	8.1	13.4	5.9	9.3	14.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	15.6	20.0	13.8	13.2	15.6	15.5	22.5	17.3	10.1
Puerto Rican	25.8	22.3	21.2	22.6	29.6	26.3	30.0	29.1	13.1
Cuban	20.4	14.1	20.4	18.9	16.2	20.9	5.0	18.7	36.0
Other specific	38.1	43.6	44.6	45.3	38.7	37.3	42.5	34.8	40.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.4c. Individuals aged 55 years and older with SMI, by mental health diagnosis, gender, race, and ethnicity: Row percent distribution, 2018

Gender, race, and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older²	100.0	5.1	7.3	†	15.3	37.1	†	27.1	6.8
Gender									
Male	100.0	4.6	6.0	†	12.8	31.6	†	35.2	8.3
Female	100.0	5.5	8.2	†	17.0	40.9	†	21.5	5.8
Total	100.0	5.1	7.3	†	15.3	37.1	†	27.2	6.8
Race									
American Indian or Alaska Native	100.0	9.4	9.8	†	12.8	36.5	†	20.5	9.4
Asian ³	100.0	8.7	6.4	†	6.7	43.2	†	28.5	5.5
Black or African American	100.0	3.8	3.7	†	11.2	32.9	†	41.9	5.2
Native Hawaiian or Other Pacific Islander	100.0	9.0	4.9	1.2	10.2	37.4	†	30.7	5.8
White	100.0	5.2	8.2	†	17.7	37.5	†	23.4	6.7
Other	100.0	4.7	6.3	†	11.2	43.4	†	24.8	8.5
Total	100.0	5.0	7.1	†	15.6	37.0	†	27.4	6.5
Ethnicity									
Not of Hispanic or Latino origin	100.0	5.1	6.9	†	16.3	36.3	†	27.9	6.2
Hispanic or Latino origin	100.0	5.0	7.2	†	11.5	44.7	†	22.8	8.0
Total	100.0	5.1	6.9	†	15.8	37.2	†	27.4	6.4
Specific Hispanic or Latino origin									
Mexican	100.0	7.5	6.3	†	13.0	43.7	†	24.1	4.8
Puerto Rican	100.0	5.0	5.8	†	15.0	45.1	†	24.6	3.8
Cuban	100.0	4.0	7.1	†	10.4	45.2	†	19.9	13.0
Other specific	100.0	6.7	8.3	†	13.3	43.2	†	19.9	7.9
Total	100.0	5.8	7.1	†	13.1	44.2	†	21.8	7.4

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.5a. Individuals aged 55 years and older, by mental health diagnosis, education, and marital status: Number, 2018

Education and marital status	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	827,060	59,957	76,046	6,966	115,042	291,333	7,301	194,943	75,472
Highest school grade completed									
0 to 8	50,495	4,750	5,212	520	4,884	18,011	337	11,473	5,308
9 to 11	58,036	3,331	4,075	369	7,569	20,238	518	18,270	3,666
12 (or GED)	186,183	12,961	15,805	1,179	28,212	66,767	1,464	46,800	12,995
More than 12	104,368	8,823	8,797	439	20,065	37,918	1,001	20,648	6,677
Total	399,082	29,865	33,889	2,507	60,730	142,934	3,320	97,191	28,646
Marital status									
Never married	185,741	9,393	11,964	1,454	23,534	52,360	1,023	68,245	17,768
Currently married	79,262	6,712	7,736	832	13,056	33,213	435	11,624	5,654
Separated	29,688	2,234	2,136	143	5,059	12,287	192	5,850	1,787
Divorced	128,360	9,140	9,511	774	24,820	49,016	834	25,695	8,570
Widowed	41,638	3,230	3,879	938	5,665	18,261	174	6,729	2,762
Total	464,689	30,709	35,226	4,141	72,134	165,137	2,658	118,143	36,541

GED = general equivalency diploma.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.5b. Individuals aged 55 years and older, by mental health diagnosis, education, and marital status: Column percent distribution, 2018

<i>Education and marital status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 55 years and older	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Highest school grade completed									
0 to 8	12.7	15.9	15.4	20.7	8.0	12.6	10.2	11.8	18.5
9 to 11	14.5	11.2	12.0	14.7	12.5	14.2	15.6	18.8	12.8
12 (or GED)	46.7	43.4	46.6	47.0	46.5	46.7	44.1	48.2	45.4
More than 12	26.2	29.5	26.0	17.5	33.0	26.5	30.2	21.2	23.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marital status									
Never married	40.0	30.6	34.0	35.1	32.6	31.7	38.5	57.8	48.6
Currently married	17.1	21.9	22.0	20.1	18.1	20.1	16.4	9.8	15.5
Separated	6.4	7.3	6.1	3.5	7.0	7.4	7.2	5.0	4.9
Divorced	27.6	29.8	27.0	18.7	34.4	29.7	31.4	21.7	23.5
Widowed	9.0	10.5	11.0	22.7	7.9	11.1	6.5	5.7	7.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

GED = general equivalency diploma.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.5c. Individuals aged 55 years and older, by mental health diagnosis, education, and marital status: Row percent distribution, 2018

<i>Education and marital status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 55 years and older	100.0	7.2	9.2	†	13.9	35.2	†	23.6	9.1
Highest school grade completed									
0 to 8	100.0	9.4	10.3	1.0	9.7	35.7	†	22.7	10.5
9 to 11	100.0	5.7	7.0	†	13.0	34.9	†	31.5	6.3
12 (or GED)	100.0	7.0	8.5	†	15.2	35.9	†	25.1	7.0
More than 12	100.0	8.5	8.4	†	19.2	36.3	†	19.8	6.4
Total	100.0	7.5	8.5	†	15.2	35.8	†	24.4	7.2
Marital status									
Never married	100.0	5.1	6.4	†	12.7	28.2	†	36.7	9.6
Currently married	100.0	8.5	9.8	1.0	16.5	41.9	†	14.7	7.1
Separated	100.0	7.5	7.2	†	17.0	41.4	†	19.7	6.0
Divorced	100.0	7.1	7.4	†	19.3	38.2	†	20.0	6.7
Widowed	100.0	7.8	9.3	2.3	13.6	43.9	†	16.2	6.6
Total	100.0	6.6	7.6	†	15.5	35.5	†	25.4	7.9

† Less than 1 percent of individuals.

GED = general equivalency diploma.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.6a. Individuals aged 55 years and older with SMI, by mental health diagnosis, education, and marital status: Number, 2018

<i>Education and marital status</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 55 years and older²	655,241	33,694	47,785	3,504	100,014	242,891	4,881	177,878	44,594
Highest school grade completed									
0 to 8	38,587	3,196	3,082	290	4,231	14,573	256	10,492	2,467
9 to 11	50,552	2,294	2,797	269	7,006	18,076	453	17,370	2,287
12 (or GED)	156,924	8,569	10,434	744	25,944	58,809	1,187	43,883	7,354
More than 12	89,042	5,774	5,984	300	18,662	33,907	822	19,499	4,094
Total	335,105	19,833	22,297	1,603	55,843	125,365	2,718	91,244	16,202
Marital status									
Never married	153,574	4,570	7,839	569	21,317	46,657	757	63,998	7,867
Currently married	62,165	3,312	4,930	371	11,469	28,084	304	10,646	3,049
Separated	25,595	1,232	1,582	61	4,716	11,295	138	5,571	1,000
Divorced	109,090	5,426	6,760	361	22,837	43,996	649	24,016	5,045
Widowed	32,415	1,647	2,308	434	4,962	15,437	111	6,183	1,333
Total	382,839	16,187	23,419	1,796	65,301	145,469	1,959	110,414	18,294

GED = general equivalency diploma; SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.6b. Individuals aged 55 years and older with SMI, by mental health diagnosis, education, and marital status: Column percent distribution, 2018

<i>Education and marital status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 55 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Highest school grade completed									
0 to 8	11.5	16.1	13.8	18.1	7.6	11.6	9.4	11.5	15.2
9 to 11	15.1	11.6	12.5	16.8	12.5	14.4	16.7	19.0	14.1
12 (or GED)	46.8	43.2	46.8	46.4	46.5	46.9	43.7	48.1	45.4
More than 12	26.6	29.1	26.8	18.7	33.4	27.0	30.2	21.4	25.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marital status									
Never married	40.1	28.2	33.5	31.7	32.6	32.1	38.6	58.0	43.0
Currently married	16.2	20.5	21.1	20.7	17.6	19.3	15.5	9.6	16.7
Separated	6.7	7.6	6.8	3.4	7.2	7.8	7.0	5.0	5.5
Divorced	28.5	33.5	28.9	20.1	35.0	30.2	33.1	21.8	27.6
Widowed	8.5	10.2	9.9	24.2	7.6	10.6	5.7	5.6	7.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

GED = general equivalency diploma; SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.6c. Individuals aged 55 years and older with SMI, by mental health diagnosis, education, and marital status: Row percent distribution, 2018

<i>Education and marital status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 55 years and older²	100.0	5.1	7.3	†	15.3	37.1	†	27.1	6.8
Highest school grade completed									
0 to 8	100.0	8.3	8.0	†	11.0	37.8	†	27.2	6.4
9 to 11	100.0	4.5	5.5	†	13.9	35.8	†	34.4	4.5
12 (or GED)	100.0	5.5	6.6	†	16.5	37.5	†	28.0	4.7
More than 12	100.0	6.5	6.7	†	21.0	38.1	†	21.9	4.6
Total	100.0	5.9	6.7	†	16.7	37.4	†	27.2	4.8
Marital status									
Never married	100.0	3.0	5.1	†	13.9	30.4	†	41.7	5.1
Currently married	100.0	5.3	7.9	†	18.4	45.2	†	17.1	4.9
Separated	100.0	4.8	6.2	†	18.4	44.1	†	21.8	3.9
Divorced	100.0	5.0	6.2	†	20.9	40.3	†	22.0	4.6
Widowed	100.0	5.1	7.1	1.3	15.3	47.6	†	19.1	4.1
Total	100.0	4.2	6.1	†	17.1	38.0	†	28.8	4.8

† Less than 1 percent of individuals.

GED = general equivalency diploma; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.7a. Individuals aged 55 years and older, by mental health diagnosis, living arrangement, SMI status, and veteran status: Number, 2018

<i>Living arrangement, SMI status, and veteran status</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 55 years and older	827,060	59,957	76,046	6,966	115,042	291,333	7,301	194,943	75,472
Living arrangement									
Homeless	23,247	1,966	1,174	77	3,111	8,205	228	5,529	2,957
Foster home/foster care	3,366	129	292	47	344	505	39	1,483	527
Residential care	38,346	886	1,330	477	3,800	5,442	288	20,772	5,351
Institutional setting	21,490	1,172	1,353	1,230	2,082	5,531	195	8,376	1,551
Jail/correctional facility	4,471	578	240	47	428	759	325	1,280	814
Private residence	408,711	29,695	35,472	1,671	64,167	153,552	3,431	89,082	31,641
Other	16,189	2,099	1,864	176	1,603	4,750	89	4,028	1,580
Total	515,820	36,525	41,725	3,725	75,535	178,744	4,595	130,550	44,421
SMI status²									
SMI	655,241	33,694	47,785	3,504	100,014	242,891	4,881	177,878	44,594
Not SMI	145,301	23,249	23,700	3,149	12,674	40,172	2,069	13,732	26,556
Total	800,542	56,943	71,485	6,653	112,688	283,063	6,950	191,610	71,150
Veteran status									
Veteran	15,875	1,728	1,025	159	2,120	4,924	100	4,471	1,348
Not a veteran	291,019	22,234	24,610	1,966	42,038	100,974	1,718	69,238	28,241
Total	306,894	23,962	25,635	2,125	44,158	105,898	1,818	73,709	29,589

SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.7b. Individuals aged 55 years and older, by mental health diagnosis, living arrangement, SMI status, and veteran status: Column percent distribution, 2018

<i>Living arrangement, SMI status, and veteran status</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 55 years and older	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement									
Homeless	4.5	5.4	2.8	2.1	4.1	4.6	5.0	4.2	6.7
Foster home/foster care	†	†	†	1.3	†	†	†	1.1	1.2
Residential care	7.4	2.4	3.2	12.8	5.0	3.0	6.3	15.9	12.0
Institutional setting	4.2	3.2	3.2	33.0	2.8	3.1	4.2	6.4	3.5
Jail/correctional facility	†	1.6	†	1.3	†	†	7.1	†	1.8
Private residence	79.2	81.3	85.0	44.9	85.0	85.9	74.7	68.2	71.2
Other	3.1	5.7	4.5	4.7	2.1	2.7	1.9	3.1	3.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SMI status²									
SMI	81.8	59.2	66.8	52.7	88.8	85.8	70.2	92.8	62.7
Not SMI	18.2	40.8	33.2	47.3	11.2	14.2	29.8	7.2	37.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status									
Veteran	5.2	7.2	4.0	7.5	4.8	4.6	5.5	6.1	4.6
Not a veteran	94.8	92.8	96.0	92.5	95.2	95.4	94.5	93.9	95.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.7c. Individuals aged 55 years and older, by mental health diagnosis, living arrangement, SMI status, and veteran status: Row percent distribution, 2018

Living arrangement, SMI status, and veteran status	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	100.0	7.2	9.2	†	13.9	35.2	†	23.6	9.1
Living arrangement									
Homeless	100.0	8.5	5.1	†	13.4	35.3	†	23.8	12.7
Foster home/foster care	100.0	3.8	8.7	1.4	10.2	15.0	1.2	44.1	15.7
Residential care	100.0	2.3	3.5	1.2	9.9	14.2	†	54.2	14.0
Institutional setting	100.0	5.5	6.3	5.7	9.7	25.7	†	39.0	7.2
Jail/correctional facility	100.0	12.9	5.4	1.1	9.6	17.0	7.3	28.6	18.2
Private residence	100.0	7.3	8.7	†	15.7	37.6	†	21.8	7.7
Other	100.0	13.0	11.5	1.1	9.9	29.3	†	24.9	9.8
Total	100.0	7.1	8.1	†	14.6	34.7	†	25.3	8.6
SMI status²									
SMI	100.0	5.1	7.3	†	15.3	37.1	†	27.1	6.8
Not SMI	100.0	16.0	16.3	2.2	8.7	27.6	1.4	9.5	18.3
Total	100.0	7.1	8.9	†	14.1	35.4	†	23.9	8.9
Veteran status									
Veteran	100.0	10.9	6.5	1.0	13.4	31.0	†	28.2	8.5
Not a veteran	100.0	7.6	8.5	†	14.4	34.7	†	23.8	9.7
Total	100.0	7.8	8.4	†	14.4	34.5	†	24.0	9.6

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.8a. Individuals aged 55 years and older with SMI, by mental health diagnosis, living arrangement, and veteran status: Number, 2018

<i>Living arrangement and veteran status</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 55 years and older²	655,241	33,694	47,785	3,504	100,014	242,891	4,881	177,878	44,594
Living arrangement									
Homeless	20,515	1,474	853	40	2,896	7,510	180	5,237	2,325
Foster home/foster care	2,154	70	123	15	239	343	22	1,202	140
Residential care	32,715	602	881	308	3,487	4,879	257	19,846	2,455
Institutional setting	15,768	630	619	785	1,674	3,661	145	7,354	900
Jail/correctional facility	3,655	295	183	32	383	667	309	1,197	589
Private residence	347,378	20,013	23,909	1,085	59,176	135,715	2,814	84,159	20,507
Other	13,274	1,037	1,532	63	1,492	4,358	72	3,696	1,024
Total	435,459	24,121	28,100	2,328	69,347	157,133	3,799	122,691	27,940
Veteran status									
Veteran	13,275	1,145	677	95	1,934	4,397	74	4,195	758
Not a veteran	240,522	14,893	16,259	1,271	38,156	89,770	1,330	64,397	14,446
Total	253,797	16,038	16,936	1,366	40,090	94,167	1,404	68,592	15,204

SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.8b. Individuals aged 55 years and older with SMI, by mental health diagnosis, living arrangement, and veteran status: Column percent distribution, 2018

<i>Living arrangement and veteran status</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 55 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement									
Homeless	4.7	6.1	3.0	1.7	4.2	4.8	4.7	4.3	8.3
Foster home/foster care	†	†	†	†	†	†	†	†	†
Residential care	7.5	2.5	3.1	13.2	5.0	3.1	6.8	16.2	8.8
Institutional setting	3.6	2.6	2.2	33.7	2.4	2.3	3.8	6.0	3.2
Jail/correctional facility	†	1.2	†	1.4	†	†	8.1	†	2.1
Private residence	79.8	83.0	85.1	46.6	85.3	86.4	74.1	68.6	73.4
Other	3.0	4.3	5.5	2.7	2.2	2.8	1.9	3.0	3.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status									
Veteran	5.2	7.1	4.0	7.0	4.8	4.7	5.3	6.1	5.0
Not a veteran	94.8	92.9	96.0	93.0	95.2	95.3	94.7	93.9	95.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.8c. Individuals aged 55 years and older with SMI, by mental health diagnosis, living arrangement, and veteran status: Row percent distribution, 2018

<i>Living arrangement and veteran status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 55 years and older²	100.0	5.1	7.3	†	15.3	37.1	†	27.1	6.8
Living arrangement									
Homeless	100.0	7.2	4.2	†	14.1	36.6	†	25.5	11.3
Foster home/foster care	100.0	3.2	5.7	†	11.1	15.9	1.0	55.8	6.5
Residential care	100.0	1.8	2.7	†	10.7	14.9	†	60.7	7.5
Institutional setting	100.0	4.0	3.9	5.0	10.6	23.2	†	46.6	5.7
Jail/correctional facility	100.0	8.1	5.0	†	10.5	18.2	8.5	32.7	16.1
Private residence	100.0	5.8	6.9	†	17.0	39.1	†	24.2	5.9
Other	100.0	7.8	11.5	†	11.2	32.8	†	27.8	7.7
Total	100.0	5.5	6.5	†	15.9	36.1	†	28.2	6.4
Veteran status									
Veteran	100.0	8.6	5.1	†	14.6	33.1	†	31.6	5.7
Not a veteran	100.0	6.2	6.8	†	15.9	37.3	†	26.8	6.0
Total	100.0	6.3	6.7	†	15.8	37.1	†	27.0	6.0

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.9a. Individuals aged 55 years and older, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 55 years and older	827,060	59,957	76,046	6,966	115,042	291,333	7,301	194,943	75,472
Employment status									
Employed	52,497	5,273	6,152	130	7,971	20,249	485	7,090	5,147
Full time	23,777	2,981	3,661	83	2,811	8,997	268	2,326	2,650
Part time	20,985	1,947	2,054	30	3,301	7,689	193	3,506	2,265
Unemployed	82,032	6,260	6,256	282	12,509	32,472	467	16,892	6,894
Not in labor force (see detail below)	311,620	20,978	25,577	2,298	46,925	106,753	3,023	82,858	23,208
Total	446,149	32,511	37,985	2,710	67,405	159,474	3,975	106,840	35,249
Detailed not in labor force									
Total individuals aged 55 years and older not in labor force	311,620	20,978	25,577	2,298	46,925	106,753	3,023	82,858	23,208
Disabled	190,519	11,055	14,258	1,047	30,251	62,077	2,204	56,884	12,743
Retired	41,347	3,500	3,982	808	5,592	16,939	259	6,669	3,598
Student	1,558	220	172	3	193	425	4	270	271
Homemaker	7,069	666	789	33	946	3,100	37	985	513
Sheltered/non-competitive employment	3,066	139	187	12	313	642	22	856	895
Other	68,061	5,398	6,189	395	9,630	23,570	497	17,194	5,188

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTES: *Employed* includes “full-time/part-time not differentiated,” which is not presented separately; total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.9b. Individuals aged 55 years and older, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/ dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 55 years and older	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Employment status									
Employed	11.8	16.2	16.2	4.8	11.8	12.7	12.2	6.6	14.6
Full time	5.3	9.2	9.6	3.1	4.2	5.6	6.7	2.2	7.5
Part time	4.7	6.0	5.4	1.1	4.9	4.8	4.9	3.3	6.4
Unemployed	18.4	19.3	16.5	10.4	18.6	20.4	11.7	15.8	19.6
Not in labor force (see detail below)	69.8	64.5	67.3	84.8	69.6	66.9	76.1	77.6	65.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force									
Total individuals aged 55 years and older not in labor force	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Disabled	61.1	52.7	55.7	45.6	64.5	58.2	72.9	68.7	54.9
Retired	13.3	16.7	15.6	35.2	11.9	15.9	8.6	8.0	15.5
Student	†	1.0	†	†	†	†	†	†	1.2
Homemaker	2.3	3.2	3.1	1.4	2.0	2.9	1.2	1.2	2.2
Sheltered/non-competitive employment	†	†	†	†	†	†	†	1.0	3.9
Other	21.8	25.7	24.2	17.2	20.5	22.1	16.4	20.8	22.4

† Less than 1 percent of individuals.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTES: Percentages may not sum to 100 percent because of rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.9c. Individuals aged 55 years and older, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 55 years and older	100.0	7.2	9.2	†	13.9	35.2	†	23.6	9.1
Employment status									
Employed	100.0	10.0	11.7	†	15.2	38.6	†	13.5	9.8
Full time	100.0	12.5	15.4	†	11.8	37.8	1.1	9.8	11.1
Part time	100.0	9.3	9.8	†	15.7	36.6	†	16.7	10.8
Unemployed	100.0	7.6	7.6	†	15.2	39.6	†	20.6	8.4
Not in labor force (see detail below)	100.0	6.7	8.2	†	15.1	34.3	†	26.6	7.4
Total	100.0	7.3	8.5	†	15.1	35.7	†	23.9	7.9
Detailed not in labor force									
Total individuals aged 55 years and older not in labor force	100.0	6.7	8.2	†	15.1	34.3	†	26.6	7.4
Disabled	100.0	5.8	7.5	†	15.9	32.6	1.2	29.9	6.7
Retired	100.0	8.5	9.6	2.0	13.5	41.0	†	16.1	8.7
Student	100.0	14.1	11.0	†	12.4	27.3	†	17.3	17.4
Homemaker	100.0	9.4	11.2	†	13.4	43.9	†	13.9	7.3
Sheltered/non-competitive employment	100.0	4.5	6.1	†	10.2	20.9	†	27.9	29.2
Other	100.0	7.9	9.1	†	14.1	34.6	†	25.3	7.6

† Less than 1 percent of individuals.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTES: Percentages may not sum to 100 percent because of rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.10a. Individuals aged 55 years and older with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 55 years and older²	655,241	33,694	47,785	3,504	100,014	242,891	4,881	177,878	44,594
Employment status									
Employed	40,365	3,021	3,729	45	7,092	17,011	363	6,359	2,745
Full time	16,349	1,642	2,095	25	2,295	6,884	183	1,899	1,326
Part time	17,145	1,237	1,432	15	2,987	6,792	156	3,232	1,294
Unemployed	69,378	4,310	4,305	141	11,508	28,985	342	15,814	3,973
Not in labor force (see detail below)	262,130	14,306	17,227	1,539	42,998	92,959	2,533	77,626	12,942
Total	371,873	21,637	25,261	1,725	61,598	138,955	3,238	99,799	19,660
Detailed not in labor force									
Total individuals with SMI aged 55 years and older not in labor force	262,130	14,306	17,227	1,539	42,998	92,959	2,533	77,626	12,942
Disabled	164,871	8,135	10,211	692	27,849	55,022	1,908	53,442	7,612
Retired	33,928	2,028	2,648	582	5,189	14,791	206	6,259	2,225
Student	1,189	119	126	2	183	397	3	252	107
Homemaker	5,846	450	546	21	870	2,634	33	929	363
Sheltered/non-competitive employment	2,116	112	102	8	281	546	17	785	265
Other	54,180	3,462	3,594	234	8,626	19,569	366	15,959	2,370

SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTES: *Employed* includes “full-time/part-time not differentiated,” which is not presented separately; total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.10b. Individuals aged 55 years and older with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2018

Employment status and ‘detailed not in labor force’	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Employment status									
Employed	10.9	14.0	14.8	2.6	11.5	12.2	11.2	6.4	14.0
Full time	4.4	7.6	8.3	1.4	3.7	5.0	5.7	1.9	6.7
Part time	4.6	5.7	5.7	†	4.8	4.9	4.8	3.2	6.6
Unemployed	18.7	19.9	17.0	8.2	18.7	20.9	10.6	15.8	20.2
Not in labor force (see detail below)	70.5	66.1	68.2	89.2	69.8	66.9	78.2	77.8	65.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force									
Total individuals with SMI aged 55 years and older not in labor force	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Disabled	62.9	56.9	59.3	45.0	64.8	59.2	75.3	68.8	58.8
Retired	12.9	14.2	15.4	37.8	12.1	15.9	8.1	8.1	17.2
Student	†	†	†	†	†	†	†	†	†
Homemaker	2.2	3.1	3.2	1.4	2.0	2.8	1.3	1.2	2.8
Sheltered/non-competitive employment	†	†	†	†	†	†	†	1.0	2.0
Other	20.7	24.2	20.9	15.2	20.1	21.1	14.4	20.6	18.3

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTES: Percentages may not sum to 100 percent because of rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.10c. Individuals aged 55 years and older with SMI, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2018

<i>Employment status and ‘detailed not in labor force’</i>		<i>Mental health diagnosis¹</i>								
		<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/ dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
						<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with SMI aged 55 years and older²		100.0	5.1	7.3	†	15.3	37.1	†	27.1	6.8
Employment status										
Employed		100.0	7.5	9.2	†	17.6	42.1	†	15.8	6.8
Full time		100.0	10.0	12.8	†	14.0	42.1	1.1	11.6	8.1
Part time		100.0	7.2	8.4	†	17.4	39.6	†	18.9	7.5
Unemployed		100.0	6.2	6.2	†	16.6	41.8	†	22.8	5.7
Not in labor force (see detail below)		100.0	5.5	6.6	†	16.4	35.5	†	29.6	4.9
	Total	100.0	5.8	6.8	†	16.6	37.4	†	26.8	5.3
Detailed not in labor force										
Total individuals with SMI aged 55 years and older not in labor force		100.0	5.5	6.6	†	16.4	35.5	†	29.6	4.9
Disabled		100.0	4.9	6.2	†	16.9	33.4	1.2	32.4	4.6
Retired		100.0	6.0	7.8	1.7	15.3	43.6	†	18.4	6.6
Student		100.0	10.0	10.6	†	15.4	33.4	†	21.2	9.0
Homemaker		100.0	7.7	9.3	†	14.9	45.1	†	15.9	6.2
Sheltered/non-competitive employment		100.0	5.3	4.8	†	13.3	25.8	†	37.1	12.5
Other		100.0	6.4	6.6	†	15.9	36.1	†	29.5	4.4

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

NOTES: Percentages may not sum to 100 percent because of rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.11a. Individuals aged 55 years and older, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018

<i>Timing of admission and mental health service setting</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>Delirium/ dementia</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals aged 55 years and older	827,060	59,957	76,046	6,966	115,042	291,333	7,301	194,943	75,472
Timing of admission									
New individuals	326,361	33,638	38,413	3,996	36,161	124,806	3,136	49,621	36,590
Continuing individuals	500,699	26,319	37,633	2,970	78,881	166,527	4,165	145,322	38,882
Mental health service setting^{2,3}									
State psychiatric hospital	19,823	362	271	969	2,315	1,883	434	11,761	1,828
Community-based program	808,552	59,321	75,655	6,120	113,267	288,481	7,044	185,926	72,738
Residential treatment center	8,977	279	345	67	991	1,718	63	4,574	940
Other psychiatric inpatient	32,647	1,490	1,941	485	4,970	11,035	252	10,198	2,276
Institutions under the justice system	5,255	700	201	38	326	724	353	1,570	1,343

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.11b. Individuals aged 55 years and older, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018

Timing of admission and mental health service setting	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission									
New individuals	39.5	56.1	50.5	57.4	31.4	42.8	43.0	25.5	48.5
Continuing individuals	60.5	43.9	49.5	42.6	68.6	57.2	57.0	74.5	51.5
Mental health service setting^{2,3}									
State psychiatric hospital	2.4	†	†	13.9	2.0	†	5.9	6.0	2.4
Community-based program	97.8	98.9	99.5	87.9	98.5	99.0	96.5	95.4	96.4
Residential treatment center	1.1	†	†	†	†	†	†	2.3	1.2
Other psychiatric inpatient	3.9	2.5	2.6	7.0	4.3	3.8	3.5	5.2	3.0
Institutions under the justice system	†	1.2	†	†	†	†	4.8	†	1.8

† Less than 1 percent of individuals.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages for timing of admission may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.11c. Individuals aged 55 years and older, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals aged 55 years and older	100.0	7.2	9.2	†	13.9	35.2	†	23.6	9.1
Timing of admission									
New individuals	100.0	10.3	11.8	1.2	11.1	38.2	†	15.2	11.2
Continuing individuals	100.0	5.3	7.5	†	15.8	33.3	†	29.0	7.8
Mental health service setting^{2,3}									
State psychiatric hospital	100.0	1.8	1.4	4.9	11.7	9.5	2.2	59.3	9.2
Community-based program	100.0	7.3	9.4	†	14.0	35.7	†	23.0	9.0
Residential treatment center	100.0	3.1	3.8	†	11.0	19.1	†	51.0	10.5
Other psychiatric inpatient	100.0	4.6	5.9	1.5	15.2	33.8	†	31.2	7.0
Institutions under the justice system	100.0	13.3	3.8	†	6.2	13.8	6.7	29.9	25.6

† Less than 1 percent of individuals.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.12a. Individuals aged 55 years and older with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018

Timing of admission and mental health service setting	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older²	655,241	33,694	47,785	3,504	100,014	242,891	4,881	177,878	44,594
Timing of admission									
New individuals	235,997	17,975	23,921	1,792	29,575	99,098	1,466	41,935	20,235
Continuing individuals	419,244	15,719	23,864	1,712	70,439	143,793	3,415	135,943	24,359
Mental health service setting^{3,4}									
State psychiatric hospital	17,411	271	208	726	2,020	1,573	340	10,860	1,413
Community-based program	639,335	33,202	47,467	2,961	98,532	240,397	4,704	169,705	42,367
Residential treatment center	8,248	199	294	54	936	1,588	56	4,384	737
Other psychiatric inpatient	28,201	1,039	1,451	270	4,433	9,652	209	9,670	1,477
Institutions under the justice system	4,818	475	186	27	317	703	351	1,557	1,202

SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.12b. Individuals aged 55 years and older with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018

Timing of admission and mental health service setting	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission									
New individuals	36.0	53.3	50.1	51.1	29.6	40.8	30.0	23.6	45.4
Continuing individuals	64.0	46.7	49.9	48.9	70.4	59.2	70.0	76.4	54.6
Mental health service setting^{3,4}									
State psychiatric hospital	2.7	†	†	20.7	2.0	†	7.0	6.1	3.2
Community-based program	97.6	98.5	99.3	84.5	98.5	99.0	96.4	95.4	95.0
Residential treatment center	1.3	†	†	1.5	†	†	1.1	2.5	1.7
Other psychiatric inpatient	4.3	3.1	3.0	7.7	4.4	4.0	4.3	5.4	3.3
Institutions under the justice system	†	1.4	†	†	†	†	7.2	†	2.7

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages for timing of admission may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.12c. Individuals aged 55 years and older with SMI, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	Delirium/dementia	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with SMI aged 55 years and older²	100.0	5.1	7.3	†	15.3	37.1	†	27.1	6.8
Timing of admission									
New individuals	100.0	7.6	10.1	†	12.5	42.0	†	17.8	8.6
Continuing individuals	100.0	3.7	5.7	†	16.8	34.3	†	32.4	5.8
Mental health service setting^{3,4}									
State psychiatric hospital	100.0	1.6	1.2	4.2	11.6	9.0	2.0	62.4	8.1
Community-based program	100.0	5.2	7.4	†	15.4	37.6	†	26.5	6.6
Residential treatment center	100.0	2.4	3.6	†	11.3	19.3	†	53.2	8.9
Other psychiatric inpatient	100.0	3.7	5.1	†	15.7	34.2	†	34.3	5.2
Institutions under the justice system	100.0	9.9	3.9	†	6.6	14.6	7.3	32.3	24.9

† Less than 1 percent of individuals.

SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² The federal definition for SMI status can be found in Appendix D.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 5.13. Individuals aged 55 years and older, by mental health service setting and SMI status: Number and percent distribution, 2018

<i>Mental health service setting</i>	<i>SMI status¹</i>								
	<i>Number of individuals</i>			<i>Column percent of individuals</i>			<i>Row percent distribution of individuals</i>		
	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>	<i>Total</i>	<i>SMI</i>	<i>Not SMI</i>
Total individuals aged 55 years and older	885,116	697,908	187,208	100.0	100.0	100.0	100.0	78.8	21.2
Mental health service setting^{2,3}									
State psychiatric hospital	21,324	19,338	1,986	2.4	2.8	1.1	100.0	90.7	9.3
Community-based program	864,212	679,922	184,290	97.6	97.4	98.4	100.0	78.7	21.3
Residential treatment center	9,503	8,627	876	1.1	1.2	†	100.0	90.8	9.2
Other psychiatric inpatient	35,016	29,506	5,510	4.0	4.2	2.9	100.0	84.3	15.7
Institutions under the justice system	5,595	5,145	450	†	†	†	100.0	92.0	8.0

† Less than 1 percent of individuals.

SMI = serious emotional disturbance.

¹ The federal definition for SMI status can be found in Appendix D.

² Individuals could receive services in more than one mental health service setting during the reporting period.

³ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.1a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by gender, age, race, ethnicity, education, veteran status, and SMI/SED status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Number of states reporting	45	46	44	44	45	45
Total individuals with co-occurring MH and SU disorders aged 12 years and older¹	775,830	773,536	755,227	687,825	881,434	1,026,726
Gender						
Male	430,791	429,567	419,031	378,379	466,825	543,815
Female	344,479	343,130	335,757	308,971	412,209	481,040
Total	775,270	772,697	754,788	687,350	879,034	1,024,855
Age²						
12 to 14 years	11,944	10,332	8,185	7,686	19,366	18,691
15 to 17 years	47,758	42,145	36,170	33,985	46,671	49,196
18 to 20 years	37,008	33,790	31,459	28,832	36,981	42,764
21 to 24 years	61,377	59,646	57,960	51,362	64,208	73,139
25 to 29 years	93,623	91,810	91,808	85,553	111,611	133,007
30 to 34 years	96,545	96,053	96,626	89,039	113,299	134,479
35 to 39 years	78,764	79,878	81,582	78,199	102,536	123,782
40 to 44 years	83,691	80,715	75,723	66,148	80,857	94,824
45 to 49 years	89,350	86,840	81,933	71,483	85,767	97,095
50 to 54 years	85,375	89,034	86,199	75,850	89,182	99,167
55 to 59 years	52,925	59,017	60,008	56,210	71,686	86,024
60 to 64 years	23,990	27,981	30,161	28,747	38,208	47,587
65 years and older	13,480	16,295	17,413	14,731	21,062	26,971
Total	775,830	773,536	755,227	687,825	881,434	1,026,726
Race						
American Indian or Alaska Native	12,109	13,544	14,259	13,746	15,471	18,997
Asian ³	5,512	6,135	6,277	7,165	7,546	9,411
Black or African American	140,440	142,174	134,457	121,258	163,377	195,090
Native Hawaiian or Other Pacific Islander	2,023	2,256	2,258	1,801	1,769	2,201
White	505,563	498,481	487,592	448,016	573,706	660,387
Other	56,334	58,690	59,032	55,766	72,700	88,857
Total	721,981	721,280	703,875	647,752	834,569	974,943
Ethnicity						
Not of Hispanic or Latino origin	580,684	577,112	567,933	539,587	723,990	830,134
Hispanic or Latino origin	93,684	96,111	92,994	92,313	99,291	109,036
Total	674,368	673,223	660,927	631,900	823,281	939,170
Specific Hispanic or Latino origin						
Mexican	3,505	4,755	5,558	2,370	4,243	4,399
Puerto Rican	3,613	7,360	6,850	4,966	6,282	6,609
Cuban	1,274	841	827	614	866	908
Other specific	5,833	9,034	7,263	5,352	6,711	6,989
Total	14,225	21,990	20,498	13,302	18,102	18,905

Continued. See notes at end of table.

Table 6.1a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by gender, age, race, ethnicity, education, veteran status, and SMI/SED status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Highest school grade completed						
0 to 8	46,123	44,569	43,895	34,511	47,128	58,473
9 to 11	100,296	105,126	98,296	80,782	105,721	135,664
12 (or GED)	161,028	198,179	196,289	160,469	207,535	282,872
More than 12	64,664	81,670	80,028	68,917	93,437	123,141
Total	372,111	429,544	418,508	344,679	453,821	600,150
Veteran status⁴						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	12,198	13,856
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	310,013	362,989
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	322,211	376,845
SMI/SED status⁵						
SMI	500,467	544,703	541,278	514,093	627,465	741,570
SED	42,398	40,170	35,809	34,789	52,513	55,627
Not SMI/SED	193,466	165,088	156,188	122,856	176,411	198,164
Total	736,331	749,961	733,275	671,738	856,389	995,361

n/c Not collected.

GED = general equivalency diploma; SED = serious emotional disturbance; SMI = serious mental illness.

¹ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

⁵ The federal definition for SMI and SED status can be found in Appendix D.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 6.1b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by gender, age, race, ethnicity, education, veteran status, and SMI/SED status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Number of states reporting	45	46	44	44	45	45
Total individuals with co-occurring MH and SU disorders aged 12 years and older¹	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	55.6	55.6	55.5	55.0	53.1	53.1
Female	44.4	44.4	44.5	45.0	46.9	46.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age²						
12 to 14 years	1.5	1.3	1.1	1.1	2.2	1.8
15 to 17 years	6.2	5.4	4.8	4.9	5.3	4.8
18 to 20 years	4.8	4.4	4.2	4.2	4.2	4.2
21 to 24 years	7.9	7.7	7.7	7.5	7.3	7.1
25 to 29 years	12.1	11.9	12.2	12.4	12.7	13.0
30 to 34 years	12.4	12.4	12.8	12.9	12.9	13.1
35 to 39 years	10.2	10.3	10.8	11.4	11.6	12.1
40 to 44 years	10.8	10.4	10.0	9.6	9.2	9.2
45 to 49 years	11.5	11.2	10.8	10.4	9.7	9.5
50 to 54 years	11.0	11.5	11.4	11.0	10.1	9.7
55 to 59 years	6.8	7.6	7.9	8.2	8.1	8.4
60 to 64 years	3.1	3.6	4.0	4.2	4.3	4.6
65 years and older	1.7	2.1	2.3	2.1	2.4	2.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Race						
American Indian or Alaska Native	1.7	1.9	2.0	2.1	1.9	1.9
Asian ³	†	†	†	1.1	†	†
Black or African American	19.5	19.7	19.1	18.7	19.6	20.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	70.0	69.1	69.3	69.2	68.7	67.7
Other	7.8	8.1	8.4	8.6	8.7	9.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	86.1	85.7	85.9	85.4	87.9	88.4
Hispanic or Latino origin	13.9	14.3	14.1	14.6	12.1	11.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	24.6	21.6	27.1	17.8	23.4	23.3
Puerto Rican	25.4	33.5	33.4	37.3	34.7	35.0
Cuban	9.0	3.8	4.0	4.6	4.8	4.8
Other specific	41.0	41.1	35.4	40.2	37.1	37.0
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 6.1b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by gender, age, race, ethnicity, education, veteran status, and SMI/SED status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Highest school grade completed						
0 to 8	12.4	10.4	10.5	10.0	10.4	9.7
9 to 11	27.0	24.5	23.5	23.4	23.3	22.6
12 (or GED)	43.3	46.1	46.9	46.6	45.7	47.1
More than 12	17.4	19.0	19.1	20.0	20.6	20.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	3.8	3.7
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	96.2	96.3
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0
SMI/SED status⁵						
SMI	68.0	72.6	73.8	76.5	73.3	74.5
SED	5.8	5.4	4.9	5.2	6.1	5.6
Not SMI/SED	26.3	22.0	21.3	18.3	20.6	19.9
Total	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

GED = general equivalency diploma; SED = serious emotional disturbance; SMI = serious mental illness.

¹ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

⁵ The federal definition for SMI and SED status can be found in Appendix D.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 6.2a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, gender, and age:
Number, 2018**

Gender and age	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	1,026,726	101,731	105,583	21,085	169,793	289,422	15,761	169,647	153,704
Gender									
Male	543,815	46,652	47,891	14,459	76,493	138,271	8,694	119,985	91,370
Female	481,040	54,778	57,489	6,563	92,931	150,648	7,043	49,490	62,098
Total	1,024,855	101,430	105,380	21,022	169,424	288,919	15,737	169,475	153,468
Age³									
12 to 14 years	18,691	2,937	1,368	2,964	432	3,882	14	115	6,979
15 to 17 years	49,196	6,546	4,239	4,751	2,171	12,883	98	783	17,725
18 to 20 years	42,764	4,962	4,440	2,056	5,122	12,416	578	3,683	9,507
21 to 24 years	73,139	7,965	8,333	1,719	11,245	19,767	1,402	10,815	11,893
25 to 29 years	133,007	14,853	16,768	2,645	21,978	34,948	2,478	19,883	19,454
30 to 34 years	134,479	15,090	17,190	2,351	23,405	35,025	2,546	20,979	17,893
35 to 39 years	123,782	13,301	14,436	1,660	23,336	32,364	2,270	20,707	15,708
40 to 44 years	94,824	9,459	9,730	982	18,593	26,114	1,618	16,733	11,595
45 to 49 years	97,095	8,814	8,872	777	19,025	28,897	1,500	17,362	11,848
50 to 54 years	99,167	7,908	8,162	543	18,432	31,482	1,379	19,530	11,731
55 to 59 years	86,024	5,801	6,662	397	14,698	28,145	1,033	19,358	9,930
60 to 64 years	47,587	2,742	3,548	161	7,461	15,586	541	11,993	5,555
65 years and older	26,971	1,353	1,835	79	3,895	7,913	304	7,706	3,886

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

³ Age is based on the individual's age at the midpoint of the reporting period.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.2b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, gender, and age: Column percent distribution, 2018

Gender and age	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	53.1	46.0	45.4	68.8	45.1	47.9	55.2	70.8	59.5
Female	46.9	54.0	54.6	31.2	54.9	52.1	44.8	29.2	40.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age³									
12 to 14 years	1.8	2.9	1.3	14.1	†	1.3	†	†	4.5
15 to 17 years	4.8	6.4	4.0	22.5	1.3	4.5	†	†	11.5
18 to 20 years	4.2	4.9	4.2	9.8	3.0	4.3	3.7	2.2	6.2
21 to 24 years	7.1	7.8	7.9	8.2	6.6	6.8	8.9	6.4	7.7
25 to 29 years	13.0	14.6	15.9	12.5	12.9	12.1	15.7	11.7	12.7
30 to 34 years	13.1	14.8	16.3	11.2	13.8	12.1	16.2	12.4	11.6
35 to 39 years	12.1	13.1	13.7	7.9	13.7	11.2	14.4	12.2	10.2
40 to 44 years	9.2	9.3	9.2	4.7	11.0	9.0	10.3	9.9	7.5
45 to 49 years	9.5	8.7	8.4	3.7	11.2	10.0	9.5	10.2	7.7
50 to 54 years	9.7	7.8	7.7	2.6	10.9	10.9	8.7	11.5	7.6
55 to 59 years	8.4	5.7	6.3	1.9	8.7	9.7	6.6	11.4	6.5
60 to 64 years	4.6	2.7	3.4	†	4.4	5.4	3.4	7.1	3.6
65 years and older	2.6	1.3	1.7	†	2.3	2.7	1.9	4.5	2.5

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

³ Age is based on the individual's age at the midpoint of the reporting period.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.2c. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, gender, and age: Row percent distribution, 2018

Gender and age	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	9.9	10.3	2.1	16.5	28.2	1.5	16.5	15.0
Gender									
Male	100.0	8.6	8.8	2.7	14.1	25.4	1.6	22.1	16.8
Female	100.0	11.4	12.0	1.4	19.3	31.3	1.5	10.3	12.9
Total	100.0	9.9	10.3	2.1	16.5	28.2	1.5	16.5	15.0
Age³									
12 to 14 years	100.0	15.7	7.3	15.9	2.3	20.8	†	†	37.3
15 to 17 years	100.0	13.3	8.6	9.7	4.4	26.2	†	1.6	36.0
18 to 20 years	100.0	11.6	10.4	4.8	12.0	29.0	1.4	8.6	22.2
21 to 25 years	100.0	10.9	11.4	2.4	15.4	27.0	1.9	14.8	16.3
26 to 29 years	100.0	11.2	12.6	2.0	16.5	26.3	1.9	14.9	14.6
30 to 34 years	100.0	11.2	12.8	1.7	17.4	26.0	1.9	15.6	13.3
35 to 39 years	100.0	10.7	11.7	1.3	18.9	26.1	1.8	16.7	12.7
40 to 44 years	100.0	10.0	10.3	1.0	19.6	27.5	1.7	17.6	12.2
45 to 49 years	100.0	9.1	9.1	†	19.6	29.8	1.5	17.9	12.2
50 to 54 years	100.0	8.0	8.2	†	18.6	31.7	1.4	19.7	11.8
55 to 59 years	100.0	6.7	7.7	†	17.1	32.7	1.2	22.5	11.5
60 to 64 years	100.0	5.8	7.5	†	15.7	32.8	1.1	25.2	11.7
65 years and older	100.0	5.0	6.8	†	14.4	29.3	1.1	28.6	14.4

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

³ Age is based on the individual's age at the midpoint of the reporting period.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.3a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, race, and ethnicity: Number, 2018

Race and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	1,026,726	101,731	105,583	21,085	169,793	289,422	15,761	169,647	153,704
Race									
American Indian or Alaska Native	18,997	3,248	2,908	350	1,850	5,516	266	2,295	2,564
Asian ³	9,411	842	713	118	1,120	2,090	98	2,583	1,847
Black or African American	195,090	16,363	9,703	3,311	29,176	52,935	3,365	55,654	24,583
Native Hawaiian or Other Pacific Islander	2,201	266	224	29	253	523	25	531	350
White	660,387	67,369	79,355	14,092	121,082	194,063	10,134	87,751	86,541
Other	88,857	7,122	5,762	1,855	10,659	21,403	1,080	14,513	26,463
Total	974,943	95,210	98,665	19,755	164,140	276,530	14,968	163,327	142,348
Ethnicity									
Not of Hispanic or Latino origin	830,134	83,030	86,206	17,173	144,087	239,181	13,121	138,068	109,268
Hispanic or Latino origin	109,036	9,350	8,567	1,863	14,697	27,988	1,437	17,592	27,542
Total	939,170	92,380	94,773	19,036	158,784	267,169	14,558	155,660	136,810
Specific Hispanic or Latino origin									
Mexican	4,399	650	438	129	510	1,444	48	559	621
Puerto Rican	6,609	677	472	121	1,041	2,109	80	1,386	723
Cuban	908	85	64	10	166	248	4	215	116
Other specific	6,989	797	601	146	1,161	2,147	83	1,221	833
Total	18,905	2,209	1,575	406	2,878	5,948	215	3,381	2,293

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

**Table 6.3b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, race, and ethnicity:
Column percent distribution, 2018**

Race and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	1.9	3.4	2.9	1.8	1.1	2.0	1.8	1.4	1.8
Asian ³	†	†	†	†	†	†	†	1.6	1.3
Black or African American	20.0	17.2	9.8	16.8	17.8	19.1	22.5	34.1	17.3
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	67.7	70.8	80.4	71.3	73.8	70.2	67.7	53.7	60.8
Other	9.1	7.5	5.8	9.4	6.5	7.7	7.2	8.9	18.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity									
Not of Hispanic or Latino origin	88.4	89.9	91.0	90.2	90.7	89.5	90.1	88.7	79.9
Hispanic or Latino origin	11.6	10.1	9.0	9.8	9.3	10.5	9.9	11.3	20.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	23.3	29.4	27.8	31.8	17.7	24.3	22.3	16.5	27.1
Puerto Rican	35.0	30.6	30.0	29.8	36.2	35.5	37.2	41.0	31.5
Cuban	4.8	3.8	4.1	2.5	5.8	4.2	1.9	6.4	5.1
Other specific	37.0	36.1	38.2	36.0	40.3	36.1	38.6	36.1	36.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

**Table 6.3c. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, race, and ethnicity:
Row percent distribution, 2018**

Race and ethnicity	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	9.9	10.3	2.1	16.5	28.2	1.5	16.5	15.0
Race									
American Indian or Alaska Native	100.0	17.1	15.3	1.8	9.7	29.0	1.4	12.1	13.5
Asian ³	100.0	8.9	7.6	1.3	11.9	22.2	1.0	27.4	19.6
Black or African American	100.0	8.4	5.0	1.7	15.0	27.1	1.7	28.5	12.6
Native Hawaiian or Other Pacific Islander	100.0	12.1	10.2	1.3	11.5	23.8	1.1	24.1	15.9
White	100.0	10.2	12.0	2.1	18.3	29.4	1.5	13.3	13.1
Other	100.0	8.0	6.5	2.1	12.0	24.1	1.2	16.3	29.8
Total	100.0	9.8	10.1	2.0	16.8	28.4	1.5	16.8	14.6
Ethnicity									
Not of Hispanic or Latino origin	100.0	10.0	10.4	2.1	17.4	28.8	1.6	16.6	13.2
Hispanic or Latino origin	100.0	8.6	7.9	1.7	13.5	25.7	1.3	16.1	25.3
Total	100.0	9.8	10.1	2.0	16.9	28.4	1.6	16.6	14.6
Specific Hispanic or Latino origin									
Mexican	100.0	14.8	10.0	2.9	11.6	32.8	1.1	12.7	14.1
Puerto Rican	100.0	10.2	7.1	1.8	15.8	31.9	1.2	21.0	10.9
Cuban	100.0	9.4	7.0	1.1	18.3	27.3	†	23.7	12.8
Other specific	100.0	11.4	8.6	2.1	16.6	30.7	1.2	17.5	11.9
Total	100.0	11.7	8.3	2.1	15.2	31.5	1.1	17.9	12.1

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.4a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis and substance use diagnosis: Number, 2018

Substance use diagnosis ¹	Mental health diagnosis ²								
	Total	Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ²
		Anxiety disorders	ADHD	Bipolar disorders	Depressive disorders				
Total individuals with co-occurring MH and SU disorders aged 12 years and older³	1,026,726	101,731	105,583	21,085	169,793	289,422	15,761	169,647	153,704
Alcoholic psychoses	12,859	1,192	1,363	145	1,889	4,094	206	1,933	2,037
Alcohol intoxication	8,348	968	1,047	123	959	2,800	107	919	1,425
Drug psychoses	32,510	3,020	3,422	733	4,765	7,917	761	6,314	5,578
Dependence									
Alcohol	146,426	13,999	14,997	1,541	25,595	48,251	3,039	21,718	17,286
Cocaine	28,855	2,643	1,763	230	6,579	8,546	583	6,066	2,445
Marijuana	89,375	9,253	7,406	2,641	16,212	22,359	1,911	15,868	13,725
Opioid	104,418	13,186	17,323	2,117	17,555	34,906	1,432	5,529	12,370
Other	159,691	17,115	18,390	2,732	26,129	37,762	2,583	32,435	22,545
Non-dependent substance abuse									
Alcohol	68,047	5,661	6,336	863	11,986	19,850	1,087	12,823	9,441
Cocaine	13,296	936	764	79	3,024	3,698	224	3,352	1,219
Marijuana	88,419	7,955	7,365	2,834	14,921	22,150	1,102	15,587	16,505
Opioid	12,769	1,379	1,810	178	2,228	4,294	170	1,054	1,656
Other	52,520	4,738	5,263	882	8,069	12,088	939	11,371	9,170
None reported	209,193	19,686	18,334	5,987	29,882	60,707	1,617	34,678	38,302

ADHD = attention deficit/hyperactivity disorder.

¹ Substance use diagnostic category definitions can be found in Appendix E.

² "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

³ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

NOTE: Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.4b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis and substance use diagnosis: Column percent distribution, 2018

Substance use diagnosis ¹	Mental health diagnosis ²								
	Total	Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ²
		Anxiety disorders	ADHD	Bipolar disorders	Depressive disorders				
Total individuals with co-occurring MH and SU disorders aged 12 years and older³	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Alcoholic psychoses	1.3	1.2	1.3	†	1.1	1.4	1.3	1.1	1.3
Alcohol intoxication	†	†	†	†	†	†	†	†	†
Drug psychoses	3.2	3.0	3.2	3.5	2.8	2.7	4.8	3.7	3.6
Substance dependence									
Alcohol	14.3	13.8	14.2	7.3	15.1	16.7	19.3	12.8	11.2
Cocaine	2.8	2.6	1.7	1.1	3.9	3.0	3.7	3.6	1.6
Marijuana	8.7	9.1	7.0	12.5	9.5	7.7	12.1	9.4	8.9
Opioid	10.2	13.0	16.4	10.0	10.3	12.1	9.1	3.3	8.0
Other	15.6	16.8	17.4	13.0	15.4	13.0	16.4	19.1	14.7
Non-dependent substance abuse									
Alcohol	6.6	5.6	6.0	4.1	7.1	6.9	6.9	7.6	6.1
Cocaine	1.3	†	†	†	1.8	1.3	1.4	2.0	†
Marijuana	8.6	7.8	7.0	13.4	8.8	7.7	7.0	9.2	10.7
Opioid	1.2	1.4	1.7	†	1.3	1.5	1.1	†	1.1
Other	5.1	4.7	5.0	4.2	4.8	4.2	6.0	6.7	6.0
None reported	20.4	19.4	17.4	28.4	17.6	21.0	10.3	20.4	24.9

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ Substance use diagnostic category definitions can be found in Appendix E.

² "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

³ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.4c. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis and substance use diagnosis: Row percent distribution, 2018

<i>Substance use diagnosis¹</i>	<i>Total</i>	<i>Mental health diagnosis²</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders²</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with co-occurring MH and SU disorders aged 12 years and older³	100.0	9.9	10.3	2.1	16.5	28.2	1.5	16.5	15.0
Alcoholic psychoses	100.0	9.3	10.6	1.1	14.7	31.8	1.6	15.0	15.8
Alcohol intoxication	100.0	11.6	12.5	1.5	11.5	33.5	1.3	11.0	17.1
Drug psychoses	100.0	9.3	10.5	2.3	14.7	24.4	2.3	19.4	17.2
Substance dependence									
Alcohol	100.0	9.6	10.2	1.1	17.5	33.0	2.1	14.8	11.8
Cocaine	100.0	9.2	6.1	†	22.8	29.6	2.0	21.0	8.5
Marijuana	100.0	10.4	8.3	3.0	18.1	25.0	2.1	17.8	15.4
Opioid	100.0	12.6	16.6	2.0	16.8	33.4	1.4	5.3	11.8
Other	100.0	10.7	11.5	1.7	16.4	23.6	1.6	20.3	14.1
Non-dependent substance abuse									
Alcohol	100.0	8.3	9.3	1.3	17.6	29.2	1.6	18.8	13.9
Cocaine	100.0	7.0	5.7	†	22.7	27.8	1.7	25.2	9.2
Marijuana	100.0	9.0	8.3	3.2	16.9	25.1	1.2	17.6	18.7
Opioid	100.0	10.8	14.2	1.4	17.4	33.6	1.3	8.3	13.0
Other	100.0	9.0	10.0	1.7	15.4	23.0	1.8	21.7	17.5
None reported	100.0	9.4	8.8	2.9	14.3	29.0	†	16.6	18.3

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ Substance use diagnostic category definitions can be found in Appendix E.

² “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

³ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.5a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, education, and marital status: Number, 2018

Education and marital status	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders			Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
		Anxiety disorders	ADHD	Bipolar disorders	Depressive disorders				
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	1,026,726	101,731	105,583	21,085	169,793	289,422	15,761	169,647	153,704
Highest school grade completed									
0 to 8	58,473	7,052	6,329	3,492	6,780	15,266	831	8,538	10,185
9 to 11	135,664	14,486	11,769	4,311	20,609	39,047	2,679	26,122	16,641
12 (or GED)	282,872	28,600	29,872	4,522	52,774	84,857	5,479	50,683	26,085
More than 12	123,141	12,899	13,538	1,599	26,674	38,771	2,275	18,270	9,115
Total	600,150	63,037	61,508	13,924	106,837	177,941	11,264	103,613	62,026
Marital status									
Never married	381,319	34,272	30,100	10,034	64,453	107,192	5,470	81,540	48,258
Currently married	65,820	7,815	6,786	648	13,581	22,264	1,063	7,590	6,073
Separated	32,333	3,566	2,479	185	7,424	11,187	420	4,530	2,542
Divorced	90,241	8,406	6,830	549	21,656	31,042	1,122	14,057	6,579
Widowed	13,408	1,137	998	65	2,579	5,016	128	2,045	1,440
Total	583,121	55,196	47,193	11,481	109,693	176,701	8,203	109,762	64,892

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.5b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, education, and marital status: Column percent distribution, 2018

Education and marital status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Highest school grade completed									
0 to 8	9.7	11.2	10.3	25.1	6.3	8.6	7.4	8.2	16.4
9 to 11	22.6	23.0	19.1	31.0	19.3	21.9	23.8	25.2	26.8
12 (or GED)	47.1	45.4	48.6	32.5	49.4	47.7	48.6	48.9	42.1
More than 12	20.5	20.5	22.0	11.5	25.0	21.8	20.2	17.6	14.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marital status									
Never married	65.4	62.1	63.8	87.4	58.8	60.7	66.7	74.3	74.4
Currently married	11.3	14.2	14.4	5.6	12.4	12.6	13.0	6.9	9.4
Separated	5.5	6.5	5.3	1.6	6.8	6.3	5.1	4.1	3.9
Divorced	15.5	15.2	14.5	4.8	19.7	17.6	13.7	12.8	10.1
Widowed	2.3	2.1	2.1	†	2.4	2.8	1.6	1.9	2.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.5c. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, education, and marital status: Row percent distribution, 2018

<i>Education and marital status</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	9.9	10.3	2.1	16.5	28.2	1.5	16.5	15.0
Highest school grade completed									
0 to 8	100.0	12.1	10.8	6.0	11.6	26.1	1.4	14.6	17.4
9 to 11	100.0	10.7	8.7	3.2	15.2	28.8	2.0	19.3	12.3
12 (or GED)	100.0	10.1	10.6	1.6	18.7	30.0	1.9	17.9	9.2
More than 12	100.0	10.5	11.0	1.3	21.7	31.5	1.8	14.8	7.4
Total	100.0	10.5	10.2	2.3	17.8	29.6	1.9	17.3	10.3
Marital status									
Never married	100.0	9.0	7.9	2.6	16.9	28.1	1.4	21.4	12.7
Currently married	100.0	11.9	10.3	†	20.6	33.8	1.6	11.5	9.2
Separated	100.0	11.0	7.7	†	23.0	34.6	1.3	14.0	7.9
Divorced	100.0	9.3	7.6	†	24.0	34.4	1.2	15.6	7.3
Widowed	100.0	8.5	7.4	†	19.2	37.4	†	15.3	10.7
Total	100.0	9.5	8.1	2.0	18.8	30.3	1.4	18.8	11.1

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; GED = general equivalency diploma.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.6a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, living arrangement, SMI/SED status, and veteran status: Number, 2018

Living arrangement, SMI/SED status, and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	1,026,726	101,731	105,583	21,085	169,793	289,422	15,761	169,647	153,704
Living arrangement									
Homeless	61,656	5,668	4,102	425	9,853	17,746	689	11,991	11,182
Foster home/foster care	4,450	651	292	197	372	729	64	1,228	917
Residential care	41,638	2,190	2,070	461	5,816	7,882	470	15,547	7,202
Institutional setting	13,573	849	916	149	1,721	2,661	481	4,625	2,171
Jail/correctional facility	30,065	5,467	2,283	445	2,862	4,998	2,914	4,425	6,671
Private residence	580,755	54,453	59,350	13,643	101,825	172,082	7,661	88,283	83,458
Other	24,857	3,936	3,608	436	3,199	6,762	212	3,694	3,010
Total	756,994	73,214	72,621	15,756	125,648	212,860	12,491	129,793	114,611
SMI/SED status³									
SMI	741,570	54,217	60,546	6,182	142,482	215,823	12,720	155,791	93,809
SED	55,627	7,637	4,438	6,304	2,167	14,037	82	776	20,186
Not SMI/SED	198,164	36,663	36,362	7,664	21,569	50,110	2,417	10,523	32,856
Total	995,361	98,517	101,346	20,150	166,218	279,970	15,219	167,090	146,851
Veteran status⁴									
Veteran	13,856	1,676	1,010	121	2,343	4,488	119	3,064	1,035
Not a veteran	362,989	36,336	33,318	4,758	65,162	108,440	4,270	66,176	44,529
Total	376,845	38,012	34,328	4,879	67,505	112,928	4,389	69,240	45,564

ADHD = attention deficit/hyperactivity disorder; SED = serious emotional disturbance; SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

³ The federal definitions for SMI and SED statuses can be found in Appendix D.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.6b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, living arrangement, SMI/SED status, and veteran status: Column percent distribution, 2018

Living arrangement, SMI/SED status, and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Living arrangement									
Homeless	8.1	7.7	5.6	2.7	7.8	8.3	5.5	9.2	9.8
Foster home/foster care	†	†	†	1.3	†	†	†	†	†
Residential care	5.5	3.0	2.9	2.9	4.6	3.7	3.8	12.0	6.3
Institutional setting	1.8	1.2	1.3	†	1.4	1.3	3.9	3.6	1.9
Jail/correctional facility	4.0	7.5	3.1	2.8	2.3	2.3	23.3	3.4	5.8
Private residence	76.7	74.4	81.7	86.6	81.0	80.8	61.3	68.0	72.8
Other	3.3	5.4	5.0	2.8	2.5	3.2	1.7	2.8	2.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SMI/SED status³									
SMI	74.5	55.0	59.7	30.7	85.7	77.1	83.6	93.2	63.9
SED	5.6	7.8	4.4	31.3	1.3	5.0	†	†	13.7
Not SMI/SED	19.9	37.2	35.9	38.0	13.0	17.9	15.9	6.3	22.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	3.7	4.4	2.9	2.5	3.5	4.0	2.7	4.4	2.3
Not a veteran	96.3	95.6	97.1	97.5	96.5	96.0	97.3	95.6	97.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SED = serious emotional disturbance; SMI = serious mental illness.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

³ The federal definitions for SMI and SED statuses can be found in Appendix D.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

Table 6.6c. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, living arrangement, SMI/SED status, and veteran status: Row percent distribution, 2018

Living arrangement, SMI/SED status, and veteran status	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	9.9	10.3	2.1	16.5	28.2	1.5	16.5	15.0
Living arrangement									
Homeless	100.0	9.2	6.7	†	16.0	28.8	1.1	19.4	18.1
Foster home/foster care	100.0	14.6	6.6	4.4	8.4	16.4	1.4	27.6	20.6
Residential care	100.0	5.3	5.0	1.1	14.0	18.9	1.1	37.3	17.3
Institutional setting	100.0	6.3	6.7	1.1	12.7	19.6	3.5	34.1	16.0
Jail/correctional facility	100.0	18.2	7.6	1.5	9.5	16.6	9.7	14.7	22.2
Private residence	100.0	9.4	10.2	2.3	17.5	29.6	1.3	15.2	14.4
Other	100.0	15.8	14.5	1.8	12.9	27.2	†	14.9	12.1
Total	100.0	9.7	9.6	2.1	16.6	28.1	1.7	17.1	15.1
SMI/SED status³									
SMI	100.0	7.3	8.2	†	19.2	29.1	1.7	21.0	12.7
SED	100.0	13.7	8.0	11.3	3.9	25.2	†	1.4	36.3
Not SMI/SED	100.0	18.5	18.3	3.9	10.9	25.3	1.2	5.3	16.6
Total	100.0	9.9	10.2	2.0	16.7	28.1	1.5	16.8	14.8
Veteran status⁴									
Veteran	100.0	12.1	7.3	†	16.9	32.4	†	22.1	7.6
Not a veteran	100.0	10.5	9.1	2.3	16.8	29.5	1.1	16.9	13.9
Total	100.0	10.5	9.0	2.2	16.8	29.6	1.1	17.1	13.6

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder; SED = serious emotional disturbance; SMI = serious mental illness.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

³ The federal definitions for SMI and SED statuses can be found in Appendix D.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

Table 6.7a. Individuals aged 18 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Number, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	958,839	92,248	99,976	13,370	167,190	272,657	15,649	168,749	129,000
Employment status									
Employed	127,514	15,345	17,450	2,838	22,493	41,258	2,233	12,039	13,858
Full time	65,689	9,599	11,328	1,865	8,890	20,355	1,466	4,756	7,430
Part time	44,057	4,941	5,133	830	7,475	13,709	669	5,309	5,991
Unemployed	181,163	16,730	16,254	2,190	34,229	61,159	2,043	26,924	21,634
Not in labor force (see detail below)	290,035	23,535	26,596	3,726	55,755	77,883	4,745	69,313	28,482
Total	598,712	55,610	60,300	8,754	112,477	180,300	9,021	108,276	63,974
Detailed not in labor force									
Disabled	160,331	10,346	11,191	1,138	32,438	40,499	2,773	47,819	14,127
Retired	7,106	508	505	24	1,099	2,534	43	1,273	1,120
Student	11,939	1,255	1,030	505	1,130	2,737	77	860	4,345
Homemaker	8,859	1,264	1,210	87	1,422	2,925	116	548	1,287
Sheltered/non-competitive employment	2,806	158	150	22	365	522	13	484	1,092
Other	98,994	10,004	12,510	1,950	19,301	28,666	1,723	18,329	6,511
Total	290,035	23,535	26,596	3,726	55,755	77,883	4,745	69,313	28,482

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

NOTES: *Employed* includes “full-time/part-time not differentiated,” which is not presented separately; total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.7b. Individuals aged 18 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Column percent distribution, 2018

Employment status and ‘detailed not in labor force’	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Employment status									
Employed	21.3	27.6	28.9	32.4	20.0	22.9	24.8	11.1	21.7
Full time	11.0	17.3	18.8	21.3	7.9	11.3	16.3	4.4	11.6
Part time	7.4	8.9	8.5	9.5	6.6	7.6	7.4	4.9	9.4
Unemployed	30.3	30.1	27.0	25.0	30.4	33.9	22.6	24.9	33.8
Not in labor force (see detail below)	48.4	42.3	44.1	42.6	49.6	43.2	52.6	64.0	44.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Detailed not in labor force									
Disabled	55.3	44.0	42.1	30.5	58.2	52.0	58.4	69.0	49.6
Retired	2.5	2.2	1.9	†	2.0	3.3	†	1.8	3.9
Student	4.1	5.3	3.9	13.6	2.0	3.5	1.6	1.2	15.3
Homemaker	3.1	5.4	4.5	2.3	2.6	3.8	2.4	†	4.5
Sheltered/non-competitive employment	†	†	†	†	†	†	†	†	3.8
Other	34.1	42.5	47.0	52.3	34.6	36.8	36.3	26.4	22.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

NOTES: Percentages may not sum to 100 percent because of rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.7c. Individuals aged 18 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, employment status, and ‘detailed not in labor force’: Row percent distribution, 2018

<i>Employment status and ‘detailed not in labor force’</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	9.6	10.4	1.4	17.4	28.4	1.6	17.6	13.5
Employment status									
Employed	100.0	12.0	13.7	2.2	17.6	32.4	1.8	9.4	10.9
Full time	100.0	14.6	17.2	2.8	13.5	31.0	2.2	7.2	11.3
Part time	100.0	11.2	11.7	1.9	17.0	31.1	1.5	12.1	13.6
Unemployed	100.0	9.2	9.0	1.2	18.9	33.8	1.1	14.9	11.9
Not in labor force (see detail below)	100.0	8.1	9.2	1.3	19.2	26.9	1.6	23.9	9.8
Total	100.0	9.3	10.1	1.5	18.8	30.1	1.5	18.1	10.7
Detailed not in labor force									
Disabled	100.0	6.5	7.0	†	20.2	25.3	1.7	29.8	8.8
Retired	100.0	7.1	7.1	†	15.5	35.7	†	17.9	15.8
Student	100.0	10.5	8.6	4.2	9.5	22.9	†	7.2	36.4
Homemaker	100.0	14.3	13.7	†	16.1	33.0	1.3	6.2	14.5
Sheltered/non-competitive employment	100.0	5.6	5.3	†	13.0	18.6	†	17.2	38.9
Other	100.0	10.1	12.6	2.0	19.5	29.0	1.7	18.5	6.6
Total	100.0	8.1	9.2	1.3	19.2	26.9	1.6	23.9	9.8

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

NOTES: Percentages may not sum to 100 percent because of rounding. *Employed* includes “full-time/part-time not differentiated,” which is not presented separately.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.8a. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, timing of admission, and mental health service setting: Number, 2018

<i>Timing of admission and mental health service setting</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>			<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
		<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Bipolar disorders</i>	<i>Depressive disorders</i>				
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	1,026,726	101,731	105,583	21,085	169,793	289,422	15,761	169,647	153,704
Timing of admission									
New individuals	492,985	59,702	56,050	9,990	72,617	155,466	6,435	54,319	78,406
Continuing individuals	533,741	42,029	49,533	11,095	97,176	133,956	9,326	115,328	75,298
Mental health service setting^{3,4}									
State psychiatric hospital	38,165	1,557	699	171	6,600	5,508	1,522	18,778	3,330
Community-based program	989,985	99,487	104,602	20,872	165,413	282,226	14,931	157,102	145,352
Residential treatment center	17,922	1,216	831	331	2,415	3,988	213	5,489	3,439
Other psychiatric inpatient	89,298	6,050	6,872	997	17,426	28,484	1,208	19,844	8,417
Institutions under the justice system	33,393	5,089	1,659	253	1,814	4,205	3,263	6,341	10,769

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.8b. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, timing of admission, and mental health service setting: Column percent, 2018

Timing of admission and mental health service setting	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Timing of admission									
New individuals	48.0	58.7	53.1	47.4	42.8	53.7	40.8	32.0	51.0
Continuing individuals	52.0	41.3	46.9	52.6	57.2	46.3	59.2	68.0	49.0
Mental health service setting^{3,4}									
State psychiatric hospital	3.7	1.5	†	†	3.9	1.9	9.7	11.1	2.2
Community-based program	96.4	97.8	99.1	99.0	97.4	97.5	94.7	92.6	94.6
Residential treatment center	1.7	1.2	†	1.6	1.4	1.4	1.4	3.2	2.2
Other psychiatric inpatient	8.7	5.9	6.5	4.7	10.3	9.8	7.7	11.7	5.5
Institutions under the justice system	3.3	5.0	1.6	1.2	1.1	1.5	20.7	3.7	7.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

³ Percentages may not sum to 100 percent, because individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages for timing of admission may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 6.8c. Individuals aged 12 years and older with co-occurring mental health and substance use disorders, by mental health diagnosis, timing of admission, and mental health service setting: Row percent distribution, 2018

Timing of admission and mental health service setting	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals with co-occurring MH and SU disorders aged 12 years and older²	100.0	9.9	10.3	2.1	16.5	28.2	1.5	16.5	15.0
Timing of admission									
New individuals	100.0	12.1	11.4	2.0	14.7	31.5	1.3	11.0	15.9
Continuing individuals	100.0	7.9	9.3	2.1	18.2	25.1	1.7	21.6	14.1
Mental health service setting^{3,4}									
State psychiatric hospital	100.0	4.1	1.8	†	17.3	14.4	4.0	49.2	8.7
Community-based program	100.0	10.0	10.6	2.1	16.7	28.5	1.5	15.9	14.7
Residential treatment center	100.0	6.8	4.6	1.8	13.5	22.3	1.2	30.6	19.2
Other psychiatric inpatient	100.0	6.8	7.7	1.1	19.5	31.9	1.4	22.2	9.4
Institutions under the justice system	100.0	15.2	5.0	†	5.4	12.6	9.8	19.0	32.2

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. This table focuses on the most frequently reported mental health diagnoses for this age group. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder.

³ Individuals could receive services in more than one service setting during the reporting period.

⁴ Mental health service setting category definitions can be found in Appendix C.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 7.1a. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2015: Number, 2015

Selected characteristics	Living arrangement							
	Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Number of states reporting	41	39	35	38	39	39	41	37
Total individuals¹	1,019,988	33,543	15,778	55,282	18,306	14,955	848,238	33,886
Gender								
Male	492,062	19,929	8,553	33,465	10,861	11,785	391,700	15,769
Female	527,835	13,604	7,225	21,814	7,444	3,167	456,466	18,115
Total	1,019,897	33,533	15,778	55,279	18,305	14,952	848,166	33,884
Age²								
0 to 11 years	133,491	436	5,017	1,169	135	28	123,952	2,754
12 to 14 years	63,475	160	2,024	1,469	182	361	57,905	1,374
15 to 17 years	40,853	157	1,450	1,549	218	727	35,673	1,079
18 to 20 years	28,147	707	679	1,189	412	606	23,214	1,340
21 to 25 years	48,794	2,090	688	2,483	942	1,463	38,897	2,231
26 to 29 years	76,055	3,480	715	3,990	1,523	2,316	60,597	3,434
30 to 34 years	83,280	3,957	665	4,525	1,677	2,153	66,877	3,426
35 to 39 years	78,738	3,712	596	4,123	1,464	1,749	64,206	2,888
40 to 44 years	81,048	3,904	607	4,526	1,441	1,485	66,106	2,979
45 to 49 years	93,896	4,868	670	5,888	1,583	1,549	75,911	3,427
50 to 54 years	108,423	5,068	834	7,664	2,168	1,258	87,772	3,659
55 to 59 years	89,000	3,160	720	7,280	2,177	745	72,189	2,729
60 to 64 years	53,310	1,329	543	5,026	1,713	313	42,884	1,502
65 years and older	41,245	483	567	4,379	2,661	178	31,922	1,055
Total	1,019,755	33,511	15,775	55,260	18,296	14,931	848,105	33,877
SED status³								
SED	184,369	590	7,167	3,626	408	945	167,459	4,174
Not SED	48,823	131	1,136	492	46	153	45,906	959
Total	233,192	721	8,303	4,118	454	1,098	213,365	5,133
SMI status³								
SMI	634,444	27,734	3,953	45,388	14,922	10,026	511,814	20,607
Not SMI	124,892	3,373	3,199	4,847	2,010	3,313	100,483	7,667
Total	759,336	31,107	7,152	50,235	16,932	13,339	612,297	28,274

Continued. See notes at end of table.

Table 7.1a. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2015: Number, 2015 (continued)

Selected characteristics	Living arrangement							
	Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Substance use diagnosis								
Total individuals aged 12 years and older with co-occurring MH and SU disorders	205,732	14,934	1,288	15,945	5,220	7,858	152,725	7,762
Alcoholic psychoses	1,489	223	6	131	42	37	944	106
Alcohol intoxication	596	109	1	37	4	5	410	30
Drug psychoses	4,218	578	18	326	113	194	2,763	226
Dependence								
Alcohol	29,820	2,227	140	2,273	687	1,556	21,939	998
Cocaine	5,861	539	12	436	84	400	4,217	173
Marijuana	12,140	765	82	660	198	750	9,362	323
Opioid	8,769	532	15	392	117	521	6,849	343
Other	35,858	3,271	204	3,290	1,043	1,987	24,858	1,205
Non-dependent abuse								
Alcohol	18,215	975	68	1,561	523	581	13,955	552
Cocaine	3,362	286	5	269	78	66	2,526	132
Marijuana	16,478	870	153	976	389	574	13,099	417
Opioid	1,573	112	8	73	32	51	1,232	65
Other	25,874	1,443	366	2,336	1,124	389	18,335	1,881
None reported	41,479	3,004	210	3,185	786	747	32,236	1,311

Continued. See notes at end of table.

Table 7.1a. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2015: Number, 2015 (continued)

<i>Selected characteristics</i>	<i>Living arrangement</i>							
	<i>Total</i>	<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Mental health diagnosis								
Trauma- and stressor-related disorders	89,687	2,746	3,291	2,327	563	1,665	74,673	4,422
Anxiety disorders	75,413	1,343	1,421	1,964	492	1,011	65,894	3,288
Attention deficit/hyperactivity disorder	72,910	286	1,457	896	108	155	68,605	1,403
Mood disorders	396,300	15,354	3,622	14,391	4,057	4,227	340,327	14,322
Bipolar disorders	169,460	7,002	1,672	7,711	2,014	2,029	143,307	5,725
Depressive disorders	226,840	8,352	1,950	6,680	2,043	2,198	197,020	8,597
Personality disorders	9,477	386	149	623	299	1,127	6,678	215
Schizophrenia and other psychotic disorders	207,122	8,665	2,147	26,984	9,055	3,655	149,549	7,067
Alcohol or substance-related disorders	8,038	1,186	36	422	230	747	5,169	248
Other mental disorders ⁴	73,443	722	2,876	3,411	1,118	1,050	62,669	1,597
No/deferred diagnosis	87,598	2,855	779	4,264	2,384	1,318	74,674	1,324

MH = mental health; SED = serious emotional disturbance; SMI = serious mental illness; SU = substance use.

¹ Of 1,800,068 individuals who were served in both the 2015 and 2018 reporting periods, the total number of individuals presented in this table consists only of those with known living arrangements in both reporting years.

² Age is based on the individual's age at the midpoint of the reporting period in 2015.

³ The federal definitions for SED and SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.1b. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2015: Column percent distribution, 2015

Selected characteristics	Total	Living arrangement						
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Number of states reporting	41	39	35	38	39	39	41	37
Total individuals¹	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender								
Male	48.2	59.4	54.2	60.5	59.3	78.8	46.2	46.5
Female	51.8	40.6	45.8	39.5	40.7	21.2	53.8	53.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²								
0 to 11 years	13.1	1.3	31.8	2.1	†	†	14.6	8.1
12 to 14 years	6.2	†	12.8	2.7	†	2.4	6.8	4.1
15 to 17 years	4.0	†	9.2	2.8	1.2	4.9	4.2	3.2
18 to 20 years	2.8	2.1	4.3	2.2	2.3	4.1	2.7	4.0
21 to 25 years	4.8	6.2	4.4	4.5	5.1	9.8	4.6	6.6
26 to 29 years	7.5	10.4	4.5	7.2	8.3	15.5	7.1	10.1
30 to 34 years	8.2	11.8	4.2	8.2	9.2	14.4	7.9	10.1
35 to 39 years	7.7	11.1	3.8	7.5	8.0	11.7	7.6	8.5
40 to 44 years	7.9	11.6	3.8	8.2	7.9	9.9	7.8	8.8
45 to 49 years	9.2	14.5	4.2	10.7	8.7	10.4	9.0	10.1
50 to 54 years	10.6	15.1	5.3	13.9	11.8	8.4	10.3	10.8
55 to 59 years	8.7	9.4	4.6	13.2	11.9	5.0	8.5	8.1
60 to 64 years	5.2	4.0	3.4	9.1	9.4	2.1	5.1	4.4
65 years and older	4.0	1.4	3.6	7.9	14.5	1.2	3.8	3.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SED status³								
SED	79.1	81.8	86.3	88.1	89.9	86.1	78.5	81.3
Not SED	20.9	18.2	13.7	11.9	10.1	13.9	21.5	18.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SMI status³								
SMI	83.6	89.2	55.3	90.4	88.1	75.2	83.6	72.9
Not SMI	16.4	10.8	44.7	9.6	11.9	24.8	16.4	27.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 7.1b. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2015: Column percent distribution, 2015 (continued)

Selected characteristics	Living arrangement							
	Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Substance use diagnosis								
Total individuals aged 12 years and older with co-occurring MH and SU disorders	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Alcoholic psychoses	†	1.5	†	†	†	†	†	1.4
Alcohol intoxication	†	†	†	†	†	†	†	†
Drug psychoses	2.1	3.9	1.4	2.0	2.2	2.5	1.8	2.9
Dependence								
Alcohol	14.5	14.9	10.9	14.3	13.2	19.8	14.4	12.9
Cocaine	2.8	3.6	†	2.7	1.6	5.1	2.8	2.2
Marijuana	5.9	5.1	6.4	4.1	3.8	9.5	6.1	4.2
Opioid	4.3	3.6	1.2	2.5	2.2	6.6	4.5	4.4
Other	17.4	21.9	15.8	20.6	20.0	25.3	16.3	15.5
Non-dependent abuse								
Alcohol	8.9	6.5	5.3	9.8	10.0	7.4	9.1	7.1
Cocaine	1.6	1.9	†	1.7	1.5	†	1.7	1.7
Marijuana	8.0	5.8	11.9	6.1	7.5	7.3	8.6	5.4
Opioid	†	†	†	†	†	†	†	†
Other	12.6	9.7	28.4	14.7	21.5	5.0	12.0	24.2
None reported	20.2	20.1	16.3	20.0	15.1	9.5	21.1	16.9

Continued. See notes at end of table.

Table 7.1b. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2015: Column percent distribution, 2015 (continued)

Selected characteristics	Total	Living arrangement						Other
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	
Mental health diagnosis								
Trauma- and stressor-related disorders	8.8	8.2	20.9	4.2	3.1	11.1	8.8	13.0
Anxiety disorders	7.4	4.0	9.0	3.6	2.7	6.8	7.8	9.7
Attention deficit/hyperactivity disorder	7.1	†	9.2	1.6	†	1.0	8.1	4.1
Mood disorders	38.9	45.8	23.0	26.0	22.2	28.3	40.1	42.3
Bipolar disorders	16.6	20.9	10.6	13.9	11.0	13.6	16.9	16.9
Depressive disorders	22.2	24.9	12.4	12.1	11.2	14.7	23.2	25.4
Personality disorders	†	1.2	†	1.1	1.6	7.5	†	†
Schizophrenia and other psychotic disorders	20.3	25.8	13.6	48.8	49.5	24.4	17.6	20.9
Alcohol or substance-related disorders	†	3.5	†	†	1.3	5.0	†	†
Other mental disorders ⁴	7.2	2.2	18.2	6.2	6.1	7.0	7.4	4.7
No/deferred diagnosis	8.6	8.5	4.9	7.7	13.0	8.8	8.8	3.9

† Less than 1 percent of individuals.

MH = mental health; SED = serious emotional disturbance; SMI = serious mental illness; SU = substance use.

¹ Of 1,800,068 individuals who were served in both the 2015 and 2018 reporting periods, the total number of individuals presented in this table consists only of those with known living arrangements in both reporting years.

² Age is based on the individual's age at the midpoint of the reporting period in 2015.

³ The federal definitions for SED and SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.1c. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2015: Row percent distribution, 2015

Selected characteristics	Total	Living arrangement						
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Number of states reporting	41	39	35	38	39	39	41	37
Total individuals¹	100.0	3.3	1.5	5.4	1.8	1.5	83.2	3.3
Gender								
Male	100.0	4.1	1.7	6.8	2.2	2.4	79.6	3.2
Female	100.0	2.6	1.4	4.1	1.4	†	86.5	3.4
Total	100.0	3.3	1.5	5.4	1.8	1.5	83.2	3.3
Age²								
0 to 11 years	100.0	†	3.8	†	†	†	92.9	2.1
12 to 14 years	100.0	†	3.2	2.3	†	†	91.2	2.2
15 to 17 years	100.0	†	3.5	3.8	†	1.8	87.3	2.6
18 to 20 years	100.0	2.5	2.4	4.2	1.5	2.2	82.5	4.8
21 to 25 years	100.0	4.3	1.4	5.1	1.9	3.0	79.7	4.6
26 to 29 years	100.0	4.6	†	5.2	2.0	3.0	79.7	4.5
30 to 34 years	100.0	4.8	†	5.4	2.0	2.6	80.3	4.1
35 to 39 years	100.0	4.7	†	5.2	1.9	2.2	81.5	3.7
40 to 44 years	100.0	4.8	†	5.6	1.8	1.8	81.6	3.7
45 to 49 years	100.0	5.2	†	6.3	1.7	1.6	80.8	3.6
50 to 54 years	100.0	4.7	†	7.1	2.0	1.2	81.0	3.4
55 to 59 years	100.0	3.6	†	8.2	2.4	†	81.1	3.1
60 to 64 years	100.0	2.5	1.0	9.4	3.2	†	80.4	2.8
65 years and older	100.0	1.2	1.4	10.6	6.5	†	77.4	2.6
Total	100.0	3.3	1.5	5.4	1.8	1.5	83.2	3.3
SED status³								
SED	100.0	†	3.9	2.0	†	†	90.8	2.3
Not SED	100.0	†	2.3	1.0	†	†	94.0	2.0
Total	100.0	†	3.6	1.8	†	†	91.5	2.2
SMI status³								
SMI	100.0	4.4	†	7.2	2.4	1.6	80.7	3.2
Not SMI	100.0	2.7	2.6	3.9	1.6	2.7	80.5	6.1
Total	100.0	4.1	†	6.6	2.2	1.8	80.6	3.7

Continued. See notes at end of table.

Table 7.1c. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2015: Row percent distribution, 2015 (continued)

Selected characteristics	Total	Living arrangement						Other
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	
Substance use diagnosis								
Total individuals aged 12 years and older with co-occurring MH and SU disorders	100.0	7.3	†	7.8	2.5	3.8	74.2	3.8
Alcoholic psychoses	100.0	15.0	†	8.8	2.8	2.5	63.4	7.1
Alcohol intoxication	100.0	18.3	†	6.2	†	†	68.8	5.0
Drug psychoses	100.0	13.7	†	7.7	2.7	4.6	65.5	5.4
Dependence								
Alcohol	100.0	7.5	†	7.6	2.3	5.2	73.6	3.3
Cocaine	100.0	9.2	†	7.4	1.4	6.8	72.0	3.0
Marijuana	100.0	6.3	†	5.4	1.6	6.2	77.1	2.7
Opioid	100.0	6.1	†	4.5	1.3	5.9	78.1	3.9
Other	100.0	9.1	†	9.2	2.9	5.5	69.3	3.4
Non-dependent abuse								
Alcohol	100.0	5.4	†	8.6	2.9	3.2	76.6	3.0
Cocaine	100.0	8.5	†	8.0	2.3	2.0	75.1	3.9
Marijuana	100.0	5.3	†	5.9	2.4	3.5	79.5	2.5
Opioid	100.0	7.1	†	4.6	2.0	3.2	78.3	4.1
Other	100.0	5.6	1.4	9.0	4.3	1.5	70.9	7.3
None reported	100.0	7.2	†	7.7	1.9	1.8	77.7	3.2

Continued. See notes at end of table.

Table 7.1c. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2015: Row percent distribution, 2015 (continued)

Selected characteristics	Total	Living arrangement						
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Mental health diagnosis								
Trauma- and stressor-related disorders	100.0	3.1	3.7	2.6	†	1.9	83.3	4.9
Anxiety disorders	100.0	1.8	1.9	2.6	†	1.3	87.4	4.4
Attention deficit/hyperactivity disorder	100.0	†	2.0	1.2	†	†	94.1	1.9
Mood disorders	100.0	3.9	†	3.6	1.0	1.1	85.9	3.6
Bipolar disorders	100.0	4.1	†	4.6	1.2	1.2	84.6	3.4
Depressive disorders	100.0	3.7	†	2.9	†	†	86.9	3.8
Personality disorders	100.0	4.1	1.6	6.6	3.2	11.9	70.5	2.3
Schizophrenia and other psychotic disorders	100.0	4.2	1.0	13.0	4.4	1.8	72.2	3.4
Alcohol or substance-related disorders	100.0	14.8	†	5.3	2.9	9.3	64.3	3.1
Other mental disorders ⁴	100.0	†	3.9	4.6	1.5	1.4	85.3	2.2
No/deferred diagnosis	100.0	3.3	†	4.9	2.7	1.5	85.2	1.5

† Less than 1 percent of individuals.

MH = mental health; SED = serious emotional disturbance; SMI = serious mental illness; SU = substance use.

¹ Of 1,800,068 individuals who were served in both the 2015 and 2018 reporting periods, the total number of individuals presented in this table consists only of those with known living arrangements in both reporting years.

² Age is based on the individual's age at the midpoint of the reporting period in 2015.

³ The federal definitions for SED and SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.2a. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2018: Number, 2018

Selected characteristics	Total	Living arrangement						
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Number of states reporting	41	39	31	39	39	38	40	38
Total individuals¹	1,019,988	37,218	14,567	59,746	20,221	15,587	851,340	21,309
Gender								
Male	492,062	21,792	7,937	36,390	11,573	12,193	391,321	10,856
Female	527,835	15,419	6,628	23,351	8,648	3,391	459,948	10,450
Total	1,019,897	37,211	14,565	59,741	20,221	15,584	851,269	21,306
Age²								
0 to 11 years	74,372	277	2,761	766	82	16	70,053	417
12 to 14 years	59,548	200	1,765	1,159	131	184	55,640	469
15 to 17 years	63,394	266	2,200	2,126	254	1,169	56,544	835
18 to 20 years	40,668	829	1,060	1,398	272	885	35,111	1,113
21 to 25 years	38,983	1,678	584	1,902	634	961	32,211	1,013
26 to 29 years	67,097	3,556	731	3,872	1,375	2,087	53,790	1,686
30 to 34 years	79,539	4,436	639	4,587	1,582	2,235	64,136	1,924
35 to 39 years	83,629	4,564	636	4,743	1,664	1,938	68,053	2,031
40 to 44 years	75,883	4,070	540	4,315	1,365	1,502	62,363	1,728
45 to 49 years	87,011	4,735	567	5,194	1,586	1,481	71,484	1,964
50 to 54 years	101,413	5,242	704	7,021	1,906	1,330	83,083	2,127
55 to 59 years	103,923	4,123	839	8,193	2,373	963	85,062	2,370
60 to 64 years	75,944	2,192	658	6,862	2,354	509	61,529	1,840
65 years and older	68,351	997	881	7,583	4,637	309	52,157	1,787
Total	1,019,755	37,165	14,565	59,721	20,215	15,569	851,216	21,304
SED status³								
SED	169,490	637	6,220	3,752	407	1,275	155,673	1,526
Not SED	23,506	53	405	174	43	50	22,644	137
Total	192,996	690	6,625	3,926	450	1,325	178,317	1,663
SMI status³								
SMI	704,797	32,387	4,934	50,313	16,832	11,799	571,589	16,943
Not SMI	98,102	2,276	2,773	4,350	2,154	1,613	82,815	2,121
Total	802,899	34,663	7,707	54,663	18,986	13,412	654,404	19,064

Continued. See notes at end of table.

Table 7.2a. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2018: Number, 2018 (continued)

<i>Selected characteristics</i>	<i>Living arrangement</i>							
	<i>Total</i>	<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Substance use diagnosis								
Total individuals aged 12 years and older with co-occurring MH and SU disorders	254,688	18,236	2,031	19,749	5,720	8,542	192,997	7,413
Alcoholic psychoses	3,005	213	22	272	114	54	2,252	78
Alcohol intoxication	1,431	225	14	87	30	56	959	60
Drug psychoses	7,137	801	44	467	318	246	5,096	165
Dependence								
Alcohol	38,643	2,825	214	3,034	916	1,887	28,696	1,071
Cocaine	7,731	694	12	653	100	369	5,693	210
Marijuana	19,557	1,081	173	1,296	279	1,227	15,067	434
Opioid	20,167	1,393	52	1,088	212	825	15,743	854
Other	46,441	3,609	661	4,188	1,179	1,360	33,695	1,749
Non-dependent abuse								
Alcohol	19,268	944	121	1,610	477	515	15,188	413
Cocaine	3,825	322	10	329	77	87	2,929	71
Marijuana	20,928	1,015	227	1,284	345	604	17,009	444
Opioid	2,451	135	13	151	54	73	1,943	82
Other	18,002	1,334	306	1,625	731	403	13,111	492
None reported	46,102	3,645	162	3,665	888	836	35,616	1,290

Continued. See notes at end of table.

Table 7.2a. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2018: Number, 2018 (continued)

<i>Selected characteristics</i>	<i>Living arrangement</i>							
	<i>Total</i>	<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Mental health diagnosis								
Trauma- and stressor-related disorders	79,994	3,004	2,732	2,284	607	1,554	67,825	1,988
Anxiety disorders	86,598	1,898	1,498	2,450	716	786	77,369	1,881
Attention deficit/hyperactivity disorder	79,087	358	1,630	1,238	147	335	74,686	693
Mood disorders	374,848	13,996	2,926	13,524	4,316	3,299	329,369	7,418
Bipolar disorders	150,056	5,957	983	6,549	1,940	1,406	130,261	2,960
Depressive disorders	224,792	8,039	1,943	6,975	2,376	1,893	199,108	4,458
Personality disorders	11,208	417	155	616	334	1,492	8,039	155
Schizophrenia and other psychotic disorders	225,209	10,370	2,287	29,612	10,250	3,659	162,652	6,379
Alcohol or substance-related disorders	20,502	3,008	84	1,553	433	744	14,100	580
Other mental disorders ⁴	93,256	1,686	2,950	5,365	1,289	2,474	77,857	1,635
No/deferred diagnosis	49,286	2,481	305	3,104	2,129	1,244	39,443	580

MH = mental health; SED = serious emotional disturbance; SMI = serious mental illness; SU = substance use.

¹ Of 1,800,068 individuals who were served in both the 2015 and 2018 reporting periods, the total number of individuals presented in this table consists only of those with known living arrangements in both reporting years.

² Age is based on the individual's age at the midpoint of the reporting period in 2018.

³ The federal definitions for SED and SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.2b. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2018: Column percent distribution, 2018

Selected characteristics	Total	Living arrangement						
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Number of states reporting	41	39	31	39	39	38	40	38
Total individuals¹	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender								
Male	48.2	58.6	54.5	60.9	57.2	78.2	46.0	51.0
Female	51.8	41.4	45.5	39.1	42.8	21.8	54.0	49.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²								
0 to 11 years	7.3	†	19.0	1.3	†	†	8.2	2.0
12 to 14 years	5.8	†	12.1	1.9	†	1.2	6.5	2.2
15 to 17 years	6.2	†	15.1	3.6	1.3	7.5	6.6	3.9
18 to 20 years	4.0	2.2	7.3	2.3	1.3	5.7	4.1	5.2
21 to 25 years	3.8	4.5	4.0	3.2	3.1	6.2	3.8	4.8
26 to 29 years	6.6	9.6	5.0	6.5	6.8	13.4	6.3	7.9
30 to 34 years	7.8	11.9	4.4	7.7	7.8	14.4	7.5	9.0
35 to 39 years	8.2	12.3	4.4	7.9	8.2	12.4	8.0	9.5
40 to 44 years	7.4	11.0	3.7	7.2	6.8	9.6	7.3	8.1
45 to 49 years	8.5	12.7	3.9	8.7	7.8	9.5	8.4	9.2
50 to 54 years	9.9	14.1	4.8	11.8	9.4	8.5	9.8	10.0
55 to 59 years	10.2	11.1	5.8	13.7	11.7	6.2	10.0	11.1
60 to 64 years	7.4	5.9	4.5	11.5	11.6	3.3	7.2	8.6
65 years and older	6.7	2.7	6.0	12.7	22.9	2.0	6.1	8.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SED status³								
SED	87.8	92.3	93.9	95.6	90.4	96.2	87.3	91.8
Not SED	12.2	7.7	6.1	4.4	9.6	3.8	12.7	8.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SMI status³								
SMI	87.8	93.4	64.0	92.0	88.7	88.0	87.3	88.9
Not SMI	12.2	6.6	36.0	8.0	11.3	12.0	12.7	11.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 7.2b. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2018: Column percent distribution, 2018 (continued)

Selected characteristics	Living arrangement							
	Total	Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Substance use diagnosis								
Total individuals aged 12 years and older with co-occurring MH and SU disorders	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Alcoholic psychoses	1.2	1.2	1.1	1.4	2.0	†	1.2	1.1
Alcohol intoxication	†	1.2	†	†	†	†	†	†
Drug psychoses	2.8	4.4	2.2	2.4	5.6	2.9	2.6	2.2
Dependence								
Alcohol	15.2	15.5	10.5	15.4	16.0	22.1	14.9	14.4
Cocaine	3.0	3.8	†	3.3	1.7	4.3	2.9	2.8
Marijuana	7.7	5.9	8.5	6.6	4.9	14.4	7.8	5.9
Opioid	7.9	7.6	2.6	5.5	3.7	9.7	8.2	11.5
Other	18.2	19.8	32.5	21.2	20.6	15.9	17.5	23.6
Non-dependent abuse								
Alcohol	7.6	5.2	6.0	8.2	8.3	6.0	7.9	5.6
Cocaine	1.5	1.8	†	1.7	1.3	1.0	1.5	†
Marijuana	8.2	5.6	11.2	6.5	6.0	7.1	8.8	6.0
Opioid	†	†	†	†	†	†	1.0	1.1
Other	7.1	7.3	15.1	8.2	12.8	4.7	6.8	6.6
None reported	18.1	20.0	8.0	18.6	15.5	9.8	18.5	17.4

Continued. See notes at end of table.

Table 7.2b. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2018: Column percent distribution, 2018 (continued)

<i>Selected characteristics</i>	<i>Living arrangement</i>							
	<i>Total</i>	<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Mental health diagnosis								
Trauma- and stressor-related disorders	7.8	8.1	18.8	3.8	3.0	10.0	8.0	9.3
Anxiety disorders	8.5	5.1	10.3	4.1	3.5	5.0	9.1	8.8
Attention deficit/hyperactivity disorder	7.8	†	11.2	2.1	†	2.1	8.8	3.3
Mood disorders	36.8	37.6	20.1	22.6	21.3	21.2	38.7	34.8
Bipolar disorders	14.7	16.0	6.7	11.0	9.6	9.0	15.3	13.9
Depressive disorders	22.0	21.6	13.3	11.7	11.8	12.1	23.4	20.9
Personality disorders	1.1	1.1	1.1	1.0	1.7	9.6	†	†
Schizophrenia and other psychotic disorders	22.1	27.9	15.7	49.6	50.7	23.5	19.1	29.9
Alcohol or substance-related disorders	2.0	8.1	†	2.6	2.1	4.8	1.7	2.7
Other mental disorders ⁴	9.1	4.5	20.3	9.0	6.4	15.9	9.1	7.7
No/deferred diagnosis	4.8	6.7	2.1	5.2	10.5	8.0	4.6	2.7

† Less than 1 percent of individuals.

MH = mental health; SED = serious emotional disturbance; SMI = serious mental illness; SU = substance use.

¹ Of 1,800,068 individuals who were served in both the 2015 and 2018 reporting periods, the total number of individuals presented in this table consists only of those with known living arrangements in both reporting years.

² Age is based on the individual's age at the midpoint of the reporting period in 2018.

³ The federal definitions for SED and SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.2c. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2018: Row percent distribution, 2018

Selected characteristics	Total	Living arrangement						
		Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Number of states reporting	41	39	31	39	39	38	40	38
Total individuals¹	100.0	3.6	1.4	5.9	2.0	1.5	83.5	2.1
Gender								
Male	100.0	4.4	1.6	7.4	2.4	2.5	79.5	2.2
Female	100.0	2.9	1.3	4.4	1.6	†	87.1	2.0
Total	100.0	3.6	1.4	5.9	2.0	1.5	83.5	2.1
Age²								
0 to 11 years	100.0	†	3.7	1.0	†	†	94.2	†
12 to 15 years	100.0	†	3.0	1.9	†	†	93.4	†
15 to 17 years	100.0	†	3.5	3.4	†	1.8	89.2	1.3
18 to 20 years	100.0	2.0	2.6	3.4	†	2.2	86.3	2.7
21 to 25 years	100.0	4.3	1.5	4.9	1.6	2.5	82.6	2.6
26 to 29 years	100.0	5.3	1.1	5.8	2.0	3.1	80.2	2.5
30 to 34 years	100.0	5.6	†	5.8	2.0	2.8	80.6	2.4
35 to 39 years	100.0	5.5	†	5.7	2.0	2.3	81.4	2.4
40 to 44 years	100.0	5.4	†	5.7	1.8	2.0	82.2	2.3
45 to 49 years	100.0	5.4	†	6.0	1.8	1.7	82.2	2.3
50 to 54 years	100.0	5.2	†	6.9	1.9	1.3	81.9	2.1
55 to 59 years	100.0	4.0	†	7.9	2.3	†	81.9	2.3
60 to 64 years	100.0	2.9	†	9.0	3.1	†	81.0	2.4
65 years and older	100.0	1.5	1.3	11.1	6.8	†	76.3	2.6
Total	100.0	3.6	1.4	5.9	2.0	1.5	83.5	2.1
SED status³								
SED	100.0	†	3.7	2.2	†	†	91.8	†
Not SED	100.0	†	1.7	†	†	†	96.3	†
Total	100.0	†	3.4	2.0	†	†	92.4	†
SMI status³								
SMI	100.0	4.6	†	7.1	2.4	1.7	81.1	2.4
Not SMI	100.0	2.3	2.8	4.4	2.2	1.6	84.4	2.2
Total	100.0	4.3	†	6.8	2.4	1.7	81.5	2.4

Continued. See notes at end of table.

Table 7.2c. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2018: Row percent distribution, 2018 (continued)

<i>Selected characteristics</i>	<i>Total</i>	<i>Living arrangement</i>						<i>Other</i>
		<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	
Substance use diagnosis								
Total individuals aged 12 and older with co-occurring MH and SU disorders	100.0	7.2	†	7.8	2.2	3.4	75.8	2.9
Alcoholic psychoses	100.0	7.1	†	9.1	3.8	1.8	74.9	2.6
Alcohol intoxication	100.0	15.7	†	6.1	2.1	3.9	67.0	4.2
Drug psychoses	100.0	11.2	†	6.5	4.5	3.4	71.4	2.3
Dependence								
Alcohol	100.0	7.3	†	7.9	2.4	4.9	74.3	2.8
Cocaine	100.0	9.0	†	8.4	1.3	4.8	73.6	2.7
Marijuana	100.0	5.5	†	6.6	1.4	6.3	77.0	2.2
Opioid	100.0	6.9	†	5.4	1.1	4.1	78.1	4.2
Other	100.0	7.8	1.4	9.0	2.5	2.9	72.6	3.8
Non-dependent abuse								
Alcohol	100.0	4.9	†	8.4	2.5	2.7	78.8	2.1
Cocaine	100.0	8.4	†	8.6	2.0	2.3	76.6	1.9
Marijuana	100.0	4.8	1.1	6.1	1.6	2.9	81.3	2.1
Opioid	100.0	5.5	†	6.2	2.2	3.0	79.3	3.3
Other	100.0	7.4	1.7	9.0	4.1	2.2	72.8	2.7
None reported	100.0	7.9	†	7.9	1.9	1.8	77.3	2.8

Continued. See notes at end of table.

Table 7.2c. Individuals served in both 2015 and 2018, by living arrangement and selected characteristics in 2018: Row percent distribution, 2018 (continued)

<i>Selected characteristics</i>	<i>Living arrangement</i>							
	<i>Total</i>	<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Mental health diagnosis								
Trauma- and stressor-related disorders	100.0	3.8	3.4	2.9	†	1.9	84.8	2.5
Anxiety disorders	100.0	2.2	1.7	2.8	†	†	89.3	2.2
Attention deficit/hyperactivity disorder	100.0	†	2.1	1.6	†	†	94.4	†
Mood disorders	100.0	3.7	†	3.6	1.2	†	87.9	2.0
Bipolar disorders	100.0	4.0	†	4.4	1.3	†	86.8	2.0
Depressive disorders	100.0	3.6	†	3.1	1.1	†	88.6	2.0
Personality disorders	100.0	3.7	1.4	5.5	3.0	13.3	71.7	1.4
Schizophrenia and other psychotic disorders	100.0	4.6	1.0	13.1	4.6	1.6	72.2	2.8
Alcohol or substance-related disorders	100.0	14.7	†	7.6	2.1	3.6	68.8	2.8
Other mental disorders ⁴	100.0	1.8	3.2	5.8	1.4	2.7	83.5	1.8
No/deferred diagnosis	100.0	5.0	†	6.3	4.3	2.5	80.0	1.2

† Less than 1 percent of individuals.

MH = mental health; SED = serious emotional disturbance; SMI = serious mental illness; SU = substance use.

¹ Of 1,800,068 individuals who were served in both the 2015 and 2018 reporting periods, the total number of individuals presented in this table consists only of those with known living arrangements in both reporting years.

² Age is based on the individual's age at the midpoint of the reporting period in 2018.

³ The federal definitions for SED and SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.3a. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2015: Number, 2015

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Number of states reporting	41	40	40	40
Total individuals aged 18 years and older¹	636,139	116,827	127,628	391,684
Gender				
Male	277,694	47,877	59,178	170,639
Female	358,403	68,939	68,440	221,024
Total	636,097	116,816	127,618	391,663
Age²				
18 to 20 years	23,355	3,861	4,345	15,149
21 to 25 years	39,652	9,565	9,993	20,094
26 to 29 years	61,508	15,876	15,408	30,224
30 to 34 years	67,001	16,510	16,169	34,322
35 to 39 years	64,025	14,189	14,918	34,918
40 to 44 years	65,740	13,025	14,365	38,350
45 to 49 years	75,869	13,208	15,551	47,110
50 to 54 years	88,296	13,299	16,242	58,755
55 to 59 years	72,750	9,851	11,667	51,232
60 to 64 years	43,793	5,153	5,901	32,739
65 years and older	34,150	2,290	3,069	28,791
SMI status³				
SMI	516,101	84,493	104,550	327,058
Not SMI	100,396	29,479	18,163	52,754
Total	616,497	113,972	122,713	379,812

Continued. See notes at end of table.

Table 7.3a. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2015: Number, 2015 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Substance use diagnosis				
Total individuals aged 18 years and older with co-occurring MH and SU disorders	153,599	27,911	34,202	91,486
Alcoholic psychoses	1,019	149	298	572
Alcohol intoxication	390	81	116	193
Drug psychoses	2,760	369	943	1,448
Dependence				
Alcohol	23,536	4,761	5,423	13,352
Cocaine	4,621	694	1,022	2,905
Marijuana	8,750	1,791	2,331	4,628
Opioid	6,584	1,211	1,882	3,491
Other	23,421	4,299	5,645	13,477
Non-dependent abuse				
Alcohol	14,240	2,843	2,876	8,521
Cocaine	2,809	364	636	1,809
Marijuana	11,543	2,139	2,766	6,638
Opioid	1,200	220	321	659
Other	20,035	3,932	3,857	12,246
None reported	32,691	5,058	6,086	21,547

Continued. See notes at end of table.

Table 7.3a. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2015: Number, 2015 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Mental health diagnosis				
Trauma- and stressor-related disorders	41,888	10,105	8,580	23,203
Anxiety disorders	44,725	12,680	8,061	23,984
Attention deficit/hyperactivity disorder	9,810	3,492	1,250	5,068
Mood disorders	289,126	55,918	63,921	169,287
Bipolar disorders	122,155	22,517	27,959	71,679
Depressive disorders	166,971	33,401	35,962	97,608
Personality disorders	7,694	1,751	1,076	4,867
Schizophrenia and other psychotic disorders	155,150	16,908	28,845	109,397
Alcohol or substance-related disorders	3,789	671	1,494	1,624
Other mental disorders ⁴	20,641	4,643	2,792	13,206
No/deferred diagnosis	63,316	10,659	11,609	41,048

MH = mental health; SMI = serious mental illness; SU = substance use.

¹ Of 1,800,068 individuals who were served in both the 2015 and 2018 reporting periods, the total number of individuals presented in this table consists only of those with known employment status in both reporting years. The total number of individuals aged 18 years and older in Table 7.3a is lower than that shown in Table 7.4a because of the progression in age of individuals who were served in 2015, but who did not become 18 years or older until 2018.

² Age is based on the individual's age at the midpoint of the reporting period in 2015.

³ The federal definition for SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.3b. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2015: Column percent distribution, 2018

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Number of states reporting	41	40	40	40
Total individuals aged 18 years and older¹	100.0	100.0	100.0	100.0
Gender				
Male	43.7	41.0	46.4	43.6
Female	56.3	59.0	53.6	56.4
Total	100.0	100.0	100.0	100.0
Age²				
18 to 20 years	3.7	3.3	3.4	3.9
21 to 25 years	6.2	8.2	7.8	5.1
26 to 29 years	9.7	13.6	12.1	7.7
30 to 34 years	10.5	14.1	12.7	8.8
35 to 39 years	10.1	12.1	11.7	8.9
40 to 44 years	10.3	11.1	11.3	9.8
45 to 49 years	11.9	11.3	12.2	12.0
50 to 54 years	13.9	11.4	12.7	15.0
55 to 59 years	11.4	8.4	9.1	13.1
60 to 64 years	6.9	4.4	4.6	8.4
65 years and older	5.4	2.0	2.4	7.4
SMI status³				
SMI	83.7	74.1	85.2	86.1
Not SMI	16.3	25.9	14.8	13.9
Total	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 7.3b. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2015: Column percent distribution, 2018 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Substance use diagnosis				
Total individuals aged 18 years and older with co-occurring MH and SU disorders	100.0	100.0	100.0	100.0
Alcoholic psychoses	†	†	†	†
Alcohol intoxication	†	†	†	†
Drug psychoses	1.8	1.3	2.8	1.6
Dependence				
Alcohol	15.3	17.1	15.9	14.6
Cocaine	3.0	2.5	3.0	3.2
Marijuana	5.7	6.4	6.8	5.1
Opioid	4.3	4.3	5.5	3.8
Other	15.2	15.4	16.5	14.7
Non-dependent abuse				
Alcohol	9.3	10.2	8.4	9.3
Cocaine	1.8	1.3	1.9	2.0
Marijuana	7.5	7.7	8.1	7.3
Opioid	†	†	†	†
Other	13.0	14.1	11.3	13.4
None reported	21.3	18.1	17.8	23.6

Continued. See notes at end of table.

Table 7.3b. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2015: Column percent distribution, 2018 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Mental health diagnosis				
Trauma- and stressor-related disorders	6.6	8.6	6.7	5.9
Anxiety disorders	7.0	10.9	6.3	6.1
Attention deficit/hyperactivity disorder	1.5	3.0	†	1.3
Mood disorders	45.5	47.9	50.1	43.2
Bipolar disorders	19.2	19.3	21.9	18.3
Depressive disorders	26.2	28.6	28.2	24.9
Personality disorders	1.2	1.5	†	1.2
Schizophrenia and other psychotic disorders	24.4	14.5	22.6	27.9
Alcohol or substance-related disorders	†	†	1.2	†
Other mental disorders ⁴	3.2	4.0	2.2	3.4
No/deferred diagnosis	10.0	9.1	9.1	10.5

† Less than 1 percent of individuals.

MH = mental health; SMI = serious mental illness; SU = substance use.

¹ Of 1,800,068 individuals who were served in both the 2015 and 2018 reporting periods, the total number of individuals presented in this table consists only of those with known employment status in both reporting years. The total number of individuals aged 18 years and older in Table 7.3a is lower than that shown in Table 7.4a because of the progression in age of individuals who were served in 2015, but who did not become 18 years or older until 2018.

² Age is based on the individual's age at the midpoint of the reporting period in 2015.

³ The federal definition for SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.3c. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2015: Row percent distribution, 2018

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Number of states reporting	41	40	40	40
Total individuals aged 18 years and older¹	100.0	18.4	20.1	61.6
Gender				
Male	100.0	17.2	21.3	61.4
Female	100.0	19.2	19.1	61.7
Total	100.0	18.4	20.1	61.6
Age²				
18 to 20 years	100.0	16.5	18.6	64.9
21 to 25 years	100.0	24.1	25.2	50.7
26 to 29 years	100.0	25.8	25.1	49.1
30 to 34 years	100.0	24.6	24.1	51.2
35 to 39 years	100.0	22.2	23.3	54.5
40 to 44 years	100.0	19.8	21.9	58.3
45 to 49 years	100.0	17.4	20.5	62.1
50 to 54 years	100.0	15.1	18.4	66.5
55 to 59 years	100.0	13.5	16.0	70.4
60 to 64 years	100.0	11.8	13.5	74.8
65 years and older	100.0	6.7	9.0	84.3
SMI status³				
SMI	100.0	16.4	20.3	63.4
Not SMI	100.0	29.4	18.1	52.5
Total	100.0	18.5	19.9	61.6

Continued. See notes at end of table.

Table 7.3c. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2015: Row percent distribution, 2018 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Substance use diagnosis				
Total individuals aged 18 years and older with co-occurring MH and SU disorders	100.0	18.2	22.3	59.6
Alcoholic psychoses	100.0	14.6	29.2	56.1
Alcohol intoxication	100.0	20.8	29.7	49.5
Drug psychoses	100.0	13.4	34.2	52.5
Dependence				
Alcohol	100.0	20.2	23.0	56.7
Cocaine	100.0	15.0	22.1	62.9
Marijuana	100.0	20.5	26.6	52.9
Opioid	100.0	18.4	28.6	53.0
Other	100.0	18.4	24.1	57.5
Non-dependent abuse				
Alcohol	100.0	20.0	20.2	59.8
Cocaine	100.0	13.0	22.6	64.4
Marijuana	100.0	18.5	24.0	57.5
Opioid	100.0	18.3	26.8	54.9
Other	100.0	19.6	19.3	61.1
None reported	100.0	15.5	18.6	65.9

Continued. See notes at end of table.

Table 7.3c. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2015: Row percent distribution, 2018 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Mental health diagnosis				
Trauma- and stressor-related disorders	100.0	24.1	20.5	55.4
Anxiety disorders	100.0	28.4	18.0	53.6
Attention deficit/hyperactivity disorder	100.0	35.6	12.7	51.7
Mood disorders	100.0	19.3	22.1	58.6
Bipolar disorders	100.0	18.4	22.9	58.7
Depressive disorders	100.0	20.0	21.5	58.5
Personality disorders	100.0	22.8	14.0	63.3
Schizophrenia and other psychotic disorders	100.0	10.9	18.6	70.5
Alcohol or substance-related disorders	100.0	17.7	39.4	42.9
Other mental disorders ⁴	100.0	22.5	13.5	64.0
No/deferred diagnosis	100.0	16.8	18.3	64.8

MH = mental health; SMI = serious mental illness; SU = substance use.

¹ Of 1,800,068 individuals who were served in both the 2015 and 2018 reporting periods, the total number of individuals presented in this table consists only of those with known employment status in both reporting years. The total number of individuals aged 18 years and older in Table 7.3a is lower than that shown in Table 7.4a because of the progression in age of individuals who were served in 2015, but who did not become 18 years or older until 2018.

² Age is based on the individual's age at the midpoint of the reporting period in 2015.

³ The federal definition for SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.4a. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2018: Number, 2018

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Number of states reporting	41	40	40	40
Total individuals aged 18 years and older¹	653,998	125,126	123,565	405,307
Gender				
Male	286,269	51,059	57,397	177,813
Female	367,685	74,057	66,160	227,468
Total	653,954	125,116	123,557	405,281
Age²				
18 to 20 years	17,950	3,248	2,912	11,790
21 to 25 years	32,202	8,431	7,548	16,223
26 to 29 years	54,453	14,987	13,320	26,146
30 to 34 years	64,058	17,566	15,353	31,139
35 to 39 years	67,684	16,894	15,582	35,208
40 to 44 years	61,773	13,885	13,206	34,682
45 to 49 years	70,347	13,661	14,108	42,578
50 to 54 years	82,085	13,270	14,808	54,007
55 to 59 years	84,990	11,949	13,402	59,639
60 to 64 years	61,857	7,308	8,232	46,317
65 years and older	56,599	3,927	5,094	47,578
SMI status³				
SMI	558,096	99,010	106,367	352,719
Not SMI	81,836	23,903	13,390	44,543
Total	639,932	122,913	119,757	397,262

Continued. See notes at end of table.

Table 7.4a. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2018: Number, 2018 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Substance use diagnosis				
Total individuals aged 18 years and older with co-occurring MH and SU disorders	192,609	37,177	39,950	115,482
Alcoholic psychoses	2,417	446	463	1,508
Alcohol intoxication	920	185	255	480
Drug psychoses	5,052	845	1,452	2,755
Dependence				
Alcohol	30,942	6,509	5,755	18,678
Cocaine	6,391	863	1,169	4,359
Marijuana	13,235	2,841	2,832	7,562
Opioid	12,413	2,404	3,299	6,710
Other	34,578	7,131	5,992	21,455
Non-dependent abuse				
Alcohol	15,889	3,417	2,665	9,807
Cocaine	3,364	421	619	2,324
Marijuana	14,802	3,045	2,807	8,950
Opioid	1,833	324	430	1,079
Other	14,304	3,236	2,082	8,986
None reported	36,469	5,510	10,130	20,829

Continued. See notes at end of table.

Table 7.4a. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2018: Number, 2018 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Mental health diagnosis				
Trauma- and stressor-related disorders	42,150	10,805	8,496	22,849
Anxiety disorders	53,768	15,424	9,009	29,335
Attention deficit/hyperactivity disorder	10,587	3,673	1,556	5,358
Mood disorders	287,444	58,773	56,135	172,536
Bipolar disorders	120,754	23,955	24,237	72,562
Depressive disorders	166,690	34,818	31,898	99,974
Personality disorders	8,861	2,041	1,197	5,623
Schizophrenia and other psychotic disorders	171,664	19,081	29,071	123,512
Alcohol or substance-related disorders	8,382	1,763	3,498	3,121
Other mental disorders ⁴	35,638	7,986	5,387	22,265
No/deferred diagnosis	35,504	5,580	9,216	20,708

MH = mental health; SMI = serious mental illness; SU = substance use.

¹ Of 1,800,068 individuals who were served in both the 2015 and 2018 reporting periods, the total number of individuals presented in this table consists only of those with known employment status in both reporting years. The total number of individuals aged 18 years and older in Table 7.3a is lower than that shown in Table 7.4a because of the progression in age of individuals who were served in 2015, but who did not become 18 years or older until 2018.

² Age is based on the individual's age at the midpoint of the reporting period in 2018.

³ The federal definition for SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.4b. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2018: Column percent distribution, 2018

Selected characteristics	Employment status			
	Total	Employed	Unemployed	Not in labor force
Number of states reporting	41	40	40	40
Total individuals aged 18 years and older¹	100.0	100.0	100.0	100.0
Gender				
Male	43.8	40.8	46.5	43.9
Female	56.2	59.2	53.5	56.1
Total	100.0	100.0	100.0	100.0
Age²				
18 to 20 years	2.7	2.6	2.4	2.9
21 to 25 years	4.9	6.7	6.1	4.0
26 to 29 years	8.3	12.0	10.8	6.5
30 to 34 years	9.8	14.0	12.4	7.7
35 to 39 years	10.3	13.5	12.6	8.7
40 to 44 years	9.4	11.1	10.7	8.6
45 to 49 years	10.8	10.9	11.4	10.5
50 to 54 years	12.6	10.6	12.0	13.3
55 to 59 years	13.0	9.5	10.8	14.7
60 to 64 years	9.5	5.8	6.7	11.4
65 years and older	8.7	3.1	4.1	11.7
SMI status³				
SMI	87.2	80.6	88.8	88.8
Not SMI	12.8	19.4	11.2	11.2
Total	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 7.4b. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2018: Column percent distribution, 2018 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Substance use diagnosis				
Total individuals aged 18 years and older with co-occurring MH and SU disorders	100.0	100.0	100.0	100.0
Alcoholic psychoses	1.3	1.2	1.2	1.3
Alcohol intoxication	†	†	†	†
Drug psychoses	2.6	2.3	3.6	2.4
Dependence				
Alcohol	16.1	17.5	14.4	16.2
Cocaine	3.3	2.3	2.9	3.8
Marijuana	6.9	7.6	7.1	6.5
Opioid	6.4	6.5	8.3	5.8
Other	18.0	19.2	15.0	18.6
Non-dependent abuse				
Alcohol	8.2	9.2	6.7	8.5
Cocaine	1.7	1.1	1.5	2.0
Marijuana	7.7	8.2	7.0	7.8
Opioid	†	†	1.1	†
Other	7.4	8.7	5.2	7.8
None reported	18.9	14.8	25.4	18.0

Continued. See notes at end of table.

Table 7.4b. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2018: Column percent distribution, 2018 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Mental health diagnosis				
Trauma- and stressor-related disorders	6.4	8.6	6.9	5.6
Anxiety disorders	8.2	12.3	7.3	7.2
Attention deficit/hyperactivity disorder	1.6	2.9	1.3	1.3
Mood disorders	44.0	47.0	45.4	42.6
Bipolar disorders	18.5	19.1	19.6	17.9
Depressive disorders	25.5	27.8	25.8	24.7
Personality disorders	1.4	1.6	†	1.4
Schizophrenia and other psychotic disorders	26.2	15.2	23.5	30.5
Alcohol or substance-related disorders	1.3	1.4	2.8	†
Other mental disorders ⁴	5.4	6.4	4.4	5.5
No/deferred diagnosis	5.4	4.5	7.5	5.1

† Less than 1 percent of individuals.

MH = mental health; SMI = serious mental illness; SU = substance use.

¹ Of 1,800,068 individuals who were served in both the 2015 and 2018 reporting periods, the total number of individuals presented in this table consists only of those with known employment status in both reporting years. The total number of individuals aged 18 years and older in Table 7.3a is lower than that shown in Table 7.4a because of the progression in age of individuals who were served in 2015, but who did not become 18 years or older until 2018.

² Age is based on the individual's age at the midpoint of the reporting period in 2018.

³ The federal definition for SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.4c. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2018: Row percent distribution, 2018

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Number of states reporting	41	40	40	40
Total individuals aged 18 years and older¹	100.0	19.1	18.9	62.0
Gender				
Male	100.0	17.8	20.1	62.1
Female	100.0	20.1	18.0	61.9
Total	100.0	19.1	18.9	62.0
Age²				
18 to 20 years	100.0	18.1	16.2	65.7
21 to 25 years	100.0	26.2	23.4	50.4
26 to 29 years	100.0	27.5	24.5	48.0
30 to 34 years	100.0	27.4	24.0	48.6
35 to 39 years	100.0	25.0	23.0	52.0
40 to 44 years	100.0	22.5	21.4	56.1
45 to 49 years	100.0	19.4	20.1	60.5
50 to 54 years	100.0	16.2	18.0	65.8
55 to 59 years	100.0	14.1	15.8	70.2
60 to 64 years	100.0	11.8	13.3	74.9
65 years and older	100.0	6.9	9.0	84.1
SMI status³				
SMI	100.0	17.7	19.1	63.2
Not SMI	100.0	29.2	16.4	54.4
Total	100.0	19.2	18.7	62.1

Continued. See notes at end of table.

Table 7.4c. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2018: Row percent distribution, 2018 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Substance use diagnosis				
Total individuals aged 18 years and older with co-occurring MH and SU disorders	100.0	19.3	20.7	60.0
Alcoholic psychoses	100.0	18.5	19.2	62.4
Alcohol intoxication	100.0	20.1	27.7	52.2
Drug psychoses	100.0	16.7	28.7	54.5
Dependence				
Alcohol	100.0	21.0	18.6	60.4
Cocaine	100.0	13.5	18.3	68.2
Marijuana	100.0	21.5	21.4	57.1
Opioid	100.0	19.4	26.6	54.1
Other	100.0	20.6	17.3	62.0
Non-dependent abuse				
Alcohol	100.0	21.5	16.8	61.7
Cocaine	100.0	12.5	18.4	69.1
Marijuana	100.0	20.6	19.0	60.5
Opioid	100.0	17.7	23.5	58.9
Other	100.0	22.6	14.6	62.8
None reported	100.0	15.1	27.8	57.1

Continued. See notes at end of table.

Table 7.4c. Individuals served in both 2015 and 2018, by employment status and selected characteristics in 2018: Row percent distribution, 2018 (continued)

<i>Selected characteristics</i>	<i>Employment status</i>			
	<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Mental health diagnosis				
Trauma- and stressor-related disorders	100.0	25.6	20.2	54.2
Anxiety disorders	100.0	28.7	16.8	54.6
Attention deficit/hyperactivity disorder	100.0	34.7	14.7	50.6
Mood disorders	100.0	20.4	19.5	60.0
Bipolar disorders	100.0	19.8	20.1	60.1
Depressive disorders	100.0	20.9	19.1	60.0
Personality disorders	100.0	23.0	13.5	63.5
Schizophrenia and other psychotic disorders	100.0	11.1	16.9	71.9
Alcohol or substance-related disorders	100.0	21.0	41.7	37.2
Other mental disorders ⁴	100.0	22.4	15.1	62.5
No/deferred diagnosis	100.0	15.7	26.0	58.3

MH = mental health; SMI = serious mental illness; SU = substance use.

¹ Of 1,800,068 individuals who were served in both the 2015 and 2018 reporting periods, the total number of individuals presented in this table consists only of those with known employment status in both reporting years. The total number of individuals aged 18 years and older in Table 7.3a is lower than that shown in Table 7.4a because of the progression in age of individuals who were served in 2015, but who did not become 18 years or older until 2018.

² Age is based on the individual's age at the midpoint of the reporting period in 2018.

³ The federal definition for SMI status can be found in Appendix D.

⁴ "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol or substance-related disorders. See Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.5a. Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and living arrangement in 2015 and 2018: Number

<i>Living arrangement in 2015</i>	<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Living arrangement in 2018</i>						
			<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Homeless	Total	29,960	12,736	46	1,649	566	590	13,232	1,141
	Bipolar disorders	6,933	2,903	7	326	78	113	3,263	243
	Depressive disorders	8,269	3,280	8	303	74	115	4,136	353
	Schizophrenia and other psychotic disorders	8,652	3,943	21	770	328	230	3,021	339
	Alcohol or substance-related disorders	1,179	611	--	67	28	28	427	18
Foster home/care	Total	6,995	113	4,641	404	123	17	1,530	167
	Bipolar disorders	922	20	561	44	21	5	246	25
	Depressive disorders	1,269	15	773	68	12	1	373	27
	Schizophrenia and other psychotic disorders	2,098	49	1,287	183	59	6	448	66
	Alcohol or substance-related disorders	15	--	12	--	--	1	2	--
Residential care	Total	46,973	1,549	492	30,873	1,429	295	11,163	1,172
	Bipolar disorders	6,857	357	59	3,782	146	46	2,262	205
	Depressive disorders	6,052	305	63	3,266	121	36	2,107	154
	Schizophrenia and other psychotic disorders	26,861	679	244	19,157	1,014	163	4,959	645
	Alcohol or substance-related disorders	392	57	--	191	5	10	119	10
Institutional settings	Total	15,464	456	105	1,751	8,530	289	3,923	410
	Bipolar disorders	1,892	77	9	209	860	43	649	45
	Depressive disorders	1,969	63	12	125	1,057	21	636	55
	Schizophrenia and other psychotic disorders	9,017	258	60	1,195	5,149	181	1,931	243
	Alcohol or substance-related disorders	227	10	--	12	113	5	86	1

Continued. See notes at end of table.

Table 7.5a. Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and living arrangement in 2015 and 2018: Number (continued)

Living arrangement in 2015	Diagnostic group in 2015 ¹	Total	Living arrangement in 2018						
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Jail/correctional facility	Total	12,591	783	11	679	355	7,522	3,025	216
	Bipolar disorders	1,892	166	2	108	48	846	696	26
	Depressive disorders	2,072	88	3	65	23	1,314	546	33
	Schizophrenia and other psychotic disorders	3,624	306	2	350	235	1,711	935	85
	Alcohol or substance-related disorders	593	72	--	39	9	275	183	15
Private residence	Total	572,990	15,127	1,028	13,033	5,636	3,425	527,162	7,579
	Bipolar disorders	126,771	3,997	140	2,601	1,011	801	116,630	1,591
	Depressive disorders	173,936	3,991	198	2,383	1,101	715	163,427	2,121
	Schizophrenia and other psychotic disorders	147,864	3,878	360	5,928	2,551	1,022	132,006	2,119
	Alcohol or substance-related disorders	4,755	472	2	124	52	114	3,861	130
Other	Total	27,438	1,957	262	1,729	589	271	16,132	6,498
	Bipolar disorders	5,264	422	27	317	74	52	3,371	1,001
	Depressive disorders	7,922	598	26	322	78	63	5,341	1,494
	Schizophrenia and other psychotic disorders	7,017	462	146	757	335	85	2,716	2,516
	Alcohol or substance-related disorders	234	35	--	9	5	7	76	102

-- Quantity is zero.

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.5b. Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and living arrangement in 2015 and 2018: Row percent distribution

<i>Living arrangement in 2015</i>	<i>Diagnostic group in 2015¹</i>	<i>Total</i>	<i>Living arrangement in 2018</i>							
			<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>	
Homeless	Total	100.0	42.5	†	5.5	1.9	2.0	44.2	3.8	
	Bipolar disorders	100.0	41.9	†	4.7	1.1	1.6	47.1	3.5	
	Depressive disorders	100.0	39.7	†	3.7	†	1.4	50.0	4.3	
	Schizophrenia and other psychotic disorders	100.0	45.6	†	8.9	3.8	2.7	34.9	3.9	
	Alcohol or substance-related disorders	100.0	51.8	--	5.7	2.4	2.4	36.2	1.5	
	Foster home/care	Total	100.0	1.6	66.3	5.8	1.8	†	21.9	2.4
Foster home/care	Bipolar disorders	100.0	2.2	60.8	4.8	2.3	†	26.7	2.7	
	Depressive disorders	100.0	1.2	60.9	5.4	†	†	29.4	2.1	
	Schizophrenia and other psychotic disorders	100.0	2.3	61.3	8.7	2.8	†	21.4	3.1	
	Alcohol or substance-related disorders	100.0	--	80.0	--	--	6.7	13.3	--	
	Residential care	Total	100.0	3.3	1.0	65.7	3.0	†	23.8	2.5
	Residential care	Bipolar disorders	100.0	5.2	†	55.2	2.1	†	33.0	3.0
Depressive disorders		100.0	5.0	1.0	54.0	2.0	†	34.8	2.5	
Schizophrenia and other psychotic disorders		100.0	2.5	†	71.3	3.8	†	18.5	2.4	
Alcohol or substance-related disorders		100.0	14.5	--	48.7	1.3	2.6	30.4	2.6	
Institutional settings		Total	100.0	2.9	†	11.3	55.2	1.9	25.4	2.7
Institutional settings		Bipolar disorders	100.0	4.1	†	11.0	45.5	2.3	34.3	2.4
	Depressive disorders	100.0	3.2	†	6.3	53.7	1.1	32.3	2.8	
	Schizophrenia and other psychotic disorders	100.0	2.9	†	13.3	57.1	2.0	21.4	2.7	
	Alcohol or substance-related disorders	100.0	4.4	--	5.3	49.8	2.2	37.9	†	

Continued. See notes at end of table.

Table 7.5b. Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and living arrangement in 2015 and 2018: Row percent distribution (continued)

Living arrangement in 2015	Diagnostic group in 2015 ¹	Total	Living arrangement in 2018						
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Jail/correctional facility	Total	100.0	6.2	†	5.4	2.8	59.7	24.0	1.7
	Bipolar disorders	100.0	8.8	†	5.7	2.5	44.7	36.8	1.4
	Depressive disorders	100.0	4.2	†	3.1	1.1	63.4	26.4	1.6
	Schizophrenia and other psychotic disorders	100.0	8.4	†	9.7	6.5	47.2	25.8	2.3
	Alcohol or substance-related disorders	100.0	12.1	--	6.6	1.5	46.4	30.9	2.5
Private residence	Total	100.0	2.6	†	2.3	†	†	92.0	1.3
	Bipolar disorders	100.0	3.2	†	2.1	†	†	92.0	1.3
	Depressive disorders	100.0	2.3	†	1.4	†	†	94.0	1.2
	Schizophrenia and other psychotic disorders	100.0	2.6	†	4.0	1.7	†	89.3	1.4
	Alcohol or substance-related disorders	100.0	9.9	†	2.6	1.1	2.4	81.2	2.7
Other	Total	100.0	7.1	†	6.3	2.1	†	58.8	23.7
	Bipolar disorders	100.0	8.0	†	6.0	1.4	†	64.0	19.0
	Depressive disorders	100.0	7.5	†	4.1	†	†	67.4	18.9
	Schizophrenia and other psychotic disorders	100.0	6.6	2.1	10.8	4.8	1.2	38.7	35.9
	Alcohol or substance-related disorders	100.0	15.0	--	3.8	2.1	3.0	32.5	43.6

-- Quantity is zero; † Less than 1 percent individuals.

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.6a. Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and employment status in 2015 and 2018: Number

<i>Employment status in 2015</i>	<i>Diagnostic group in 2015¹</i>	<i>Employment status in 2018</i>			
		<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Employed	Total	106,168	70,180	13,287	22,701
	Bipolar disorders	22,517	14,068	3,436	5,013
	Depressive disorders	33,401	22,150	4,433	6,818
	Schizophrenia and other psychotic disorders	16,908	10,647	1,793	4,468
	Alcohol or substance-related disorders	671	397	179	95
Unemployed	Total	116,019	15,614	65,083	35,322
	Bipolar disorders	27,959	3,984	15,800	8,175
	Depressive disorders	35,962	5,403	19,855	10,704
	Schizophrenia and other psychotic disorders	28,845	2,218	16,408	10,219
	Alcohol or substance-related disorders	1,494	239	1,013	242
Not in labor force	Total	350,636	25,136	31,458	294,042
	Bipolar disorders	71,679	5,507	7,296	58,876
	Depressive disorders	97,608	7,396	8,861	81,351
	Schizophrenia and other psychotic disorders	109,397	4,606	8,970	95,821
	Alcohol or substance-related disorders	1,624	142	270	1,212

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.6b. Individuals aged 18 years and older served in both 2015 and 2018, by mental health diagnosis in 2015 and employment status in 2015 and 2018: Row percent distribution

<i>Employment status in 2015</i>	<i>Diagnostic group in 2015¹</i>	<i>Employment status in 2018</i>			
		<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Employed	Total	100.0	66.1	12.5	21.4
	Bipolar disorders	100.0	62.5	15.3	22.3
	Depressive disorders	100.0	66.3	13.3	20.4
	Schizophrenia and other psychotic disorders	100.0	63.0	10.6	26.4
	Alcohol or substance-related disorders	100.0	59.2	26.7	14.2
Unemployed	Total	100.0	13.5	56.1	30.4
	Bipolar disorders	100.0	14.2	56.5	29.2
	Depressive disorders	100.0	15.0	55.2	29.8
	Schizophrenia and other psychotic disorders	100.0	7.7	56.9	35.4
	Alcohol or substance-related disorders	100.0	16.0	67.8	16.2
Not in labor force	Total	100.0	7.2	9.0	83.9
	Bipolar disorders	100.0	7.7	10.2	82.1
	Depressive disorders	100.0	7.6	9.1	83.3
	Schizophrenia and other psychotic disorders	100.0	4.2	8.2	87.6
	Alcohol or substance-related disorders	100.0	8.7	16.6	74.6

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2015 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.7. Individuals aged 18 years and older served in consecutive reporting periods, by mental health diagnosis and living arrangement: Number, 2013–2018

Living arrangement	Diagnostic group ¹	Reporting periods				
		2013	2013–2015	2013–2016	2013–2017	2013–2018
Homeless	Total	97,844	8,378	3,038	1,329	859
	Bipolar disorders	24,308	2,026	613	214	126
	Depressive disorders	28,878	2,284	701	294	184
	Schizophrenia and other psychotic disorders	21,712	3,198	1,454	776	528
	Alcohol or substance-related disorders	6,377	284	128	10	2
Foster home/care	Total	9,550	1,555	331	179	108
	Bipolar disorders	1,762	221	53	27	16
	Depressive disorders	2,309	393	58	26	15
	Schizophrenia and other psychotic disorders	1,911	363	145	94	59
	Alcohol or substance-related disorders	212	108	4	--	--
Residential care	Total	95,595	29,902	17,469	12,464	10,000
	Bipolar disorders	16,331	3,352	1,531	869	652
	Depressive disorders	15,886	2,906	1,272	708	499
	Schizophrenia and other psychotic disorders	47,233	20,814	13,517	10,298	8,408
	Alcohol or substance-related disorders	1,349	162	77	2	--
Institutional settings	Total	41,695	7,790	4,059	2,442	1,764
	Bipolar disorders	5,893	751	288	139	90
	Depressive disorders	7,646	814	306	161	102
	Schizophrenia and other psychotic disorders	18,590	5,327	3,122	1,995	1,472
	Alcohol or substance-related disorders	1,668	25	10	2	2

Continued. See notes at end of table.

Table 7.7. Individuals aged 18 years and older served in consecutive reporting periods, by mental health diagnosis and living arrangement: Number, 2013–2018 (continued)

<i>Living arrangement</i>	<i>Diagnostic group¹</i>	<i>Reporting periods</i>				
		<i>2013</i>	<i>2013–2015</i>	<i>2013–2016</i>	<i>2013–2017</i>	<i>2013–2018</i>
Jail/correctional facility	Total	49,468	5,772	2,729	1,480	1,126
	Bipolar disorders	9,532	775	268	119	92
	Depressive disorders	8,746	1,086	581	180	130
	Schizophrenia and other psychotic disorders	9,131	1,803	1,014	707	553
	Alcohol or substance-related disorders	3,496	207	81	2	--
Private residence	Total	1,770,915	411,463	206,418	133,971	102,259
	Bipolar disorders	421,100	104,343	50,040	30,932	22,845
	Depressive disorders	601,964	130,798	60,797	37,609	27,332
	Schizophrenia and other psychotic disorders	321,843	125,041	77,845	56,784	46,033
	Alcohol or substance-related disorders	30,848	2,182	990	57	21
Other	Total	62,459	6,243	2,505	1,234	842
	Bipolar disorders	13,192	1,236	411	145	87
	Depressive disorders	18,927	1,544	481	202	138
	Schizophrenia and other psychotic disorders	15,251	2,609	1,379	822	576
	Alcohol or substance-related disorders	1,478	121	70	2	2

-- Quantity is zero.

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 7.8a. Individuals aged 18 years and older served in both 2013 and 2015, by mental health diagnosis in 2013 and living arrangement in 2013 and 2015: Number

<i>Living arrangement in 2013</i>	<i>Diagnostic group in 2013¹</i>	<i>Total</i>	<i>Living arrangement in 2015</i>						
			<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Homeless	Total	34,245	16,360	29	1,857	544	509	13,359	1,587
	Bipolar disorders	8,949	4,085	7	380	110	128	3,857	382
	Depressive disorders	9,529	4,473	5	371	79	77	4,054	470
	Schizophrenia and other psychotic disorders	9,718	4,897	11	862	298	201	3,066	383
	Alcohol or substance-related disorders	1,536	852	--	92	16	28	491	57
Foster home/care	Total	4,699	68	2,239	220	43	9	1,937	183
	Bipolar disorders	858	16	340	42	8	3	415	34
	Depressive disorders	1,059	6	513	26	5	1	473	35
	Schizophrenia and other psychotic disorders	1,380	24	480	89	17	4	719	47
	Alcohol or substance-related disorders	123	--	119	--	--	--	3	1
Residential care	Total	54,414	1,161	200	39,019	1,686	298	10,538	1,512
	Bipolar disorders	8,235	269	31	5,066	222	60	2,326	261
	Depressive disorders	6,822	253	29	4,221	126	45	1,905	243
	Schizophrenia and other psychotic disorders	32,057	497	84	24,539	1,146	155	4,888	748
	Alcohol or substance-related disorders	457	29	1	303	7	11	100	6
Institutional settings	Total	17,847	416	39	1,667	10,745	275	3,865	840
	Bipolar disorders	2,321	71	6	191	1,211	36	702	104
	Depressive disorders	2,080	50	1	116	1,227	14	609	63
	Schizophrenia and other psychotic disorders	10,286	238	26	1,183	6,501	180	1,547	611
	Alcohol or substance-related disorders	607	12	--	6	210	7	370	2

Continued. See notes at end of table.

Table 7.8a. Individuals aged 18 years and older served in both 2013 and 2015, by mental health diagnosis in 2013 and living arrangement in 2013 and 2015: Number (continued)

Living arrangement in 2013	Diagnostic group in 2013 ¹	Total	Living arrangement in 2015						
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Jail/correctional facility	Total	16,224	727	6	583	450	10,899	3,277	282
	Bipolar disorders	3,268	165	1	110	54	2,050	838	50
	Depressive disorders	2,716	103	1	70	27	1,855	613	47
	Schizophrenia and other psychotic disorders	4,273	244	2	273	296	2,361	983	114
	Alcohol or substance-related disorders	822	67	--	44	3	527	164	17
Private residence	Total	703,846	12,395	732	11,684	5,312	3,267	656,640	13,816
	Bipolar disorders	179,131	3,619	135	2,451	1,029	913	167,267	3,717
	Depressive disorders	217,101	3,328	183	1,969	818	596	205,792	4,415
	Schizophrenia and other psychotic disorders	185,048	3,210	205	5,808	2,824	1,061	169,432	2,508
	Alcohol or substance-related disorders	6,410	355	9	116	68	95	5,679	88
Other	Total	23,062	1,153	124	970	491	179	10,600	9,545
	Bipolar disorders	5,056	296	24	192	61	40	2,497	1,946
	Depressive disorders	6,248	320	22	124	79	36	3,264	2,403
	Schizophrenia and other psychotic disorders	7,465	321	53	528	307	66	2,787	3,403
	Alcohol or substance-related disorders	374	32	1	15	3	8	115	200

-- Quantity is zero.

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2015 reporting periods. Data as of 07.08.2019.

Table 7.8b. Individuals aged 18 years and older served in both 2013 and 2015, by mental health diagnosis in 2013 and living arrangement in 2013 and 2015: Row percent distribution

Living arrangement in 2013	Diagnostic group in 2013 ¹	Total	Living arrangement in 2015						
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Homeless	Total	100.0	47.8	†	5.4	1.6	1.5	39.0	4.6
	Bipolar disorders	100.0	45.6	†	4.2	1.2	1.4	43.1	4.3
	Depressive disorders	100.0	46.9	†	3.9	†	†	42.5	4.9
	Schizophrenia and other psychotic disorders	100.0	50.4	†	8.9	3.1	2.1	31.5	3.9
	Alcohol or substance-related disorders	100.0	55.5	--	6.0	1.0	1.8	32.0	3.7
Foster home/care	Total	100.0	1.4	47.6	4.7	†	†	41.2	3.9
	Bipolar disorders	100.0	1.9	39.6	4.9	†	†	48.4	4.0
	Depressive disorders	100.0	†	48.4	2.5	†	†	44.7	3.3
	Schizophrenia and other psychotic disorders	100.0	1.7	34.8	6.4	1.2	†	52.1	3.4
	Alcohol or substance-related disorders	100.0	--	96.7	--	--	--	2.4	†
Residential care	Total	100.0	2.1	†	71.7	3.1	†	19.4	2.8
	Bipolar disorders	100.0	3.3	†	61.5	2.7	†	28.2	3.2
	Depressive disorders	100.0	3.7	†	61.9	1.8	†	27.9	3.6
	Schizophrenia and other psychotic disorders	100.0	1.6	†	76.5	3.6	†	15.2	2.3
	Alcohol or substance-related disorders	100.0	6.3	†	66.3	1.5	2.4	21.9	1.3
Institutional settings	Total	100.0	2.3	†	9.3	60.2	1.5	21.7	4.7
	Bipolar disorders	100.0	3.1	†	8.2	52.2	1.6	30.2	4.5
	Depressive disorders	100.0	2.4	†	5.6	59.0	†	29.3	3.0
	Schizophrenia and other psychotic disorders	100.0	2.3	†	11.5	63.2	1.7	15.0	5.9
	Alcohol or substance-related disorders	100.0	2.0	--	†	34.6	1.2	61.0	†

Continued. See notes at end of table.

Table 7.8b. Individuals aged 18 years and older served in both 2013 and 2015, by mental health diagnosis in 2013 and living arrangement in 2013 and 2015: Row percent distribution (continued)

Living arrangement in 2013	Diagnostic group in 2013 ¹	Total	Living arrangement in 2015						
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Jail/correctional facility	Total	100.0	4.5	†	3.6	2.8	67.2	20.2	1.7
	Bipolar disorders	100.0	5.0	†	3.4	1.7	62.7	25.6	1.5
	Depressive disorders	100.0	3.8	†	2.6	†	68.3	22.6	1.7
	Schizophrenia and other psychotic disorders	100.0	5.7	†	6.4	6.9	55.3	23.0	2.7
	Alcohol or substance-related disorders	100.0	8.2	--	5.4	†	64.1	20.0	2.1
Private residence	Total	100.0	1.8	†	1.7	†	†	93.3	2.0
	Bipolar disorders	100.0	2.0	†	1.4	†	†	93.4	2.1
	Depressive disorders	100.0	1.5	†	†	†	†	94.8	2.0
	Schizophrenia and other psychotic disorders	100.0	1.7	†	3.1	1.5	†	91.6	1.4
	Alcohol or substance-related disorders	100.0	5.5	†	1.8	1.1	1.5	88.6	1.4
Other	Total	100.0	5.0	†	4.2	2.1	†	46.0	#
	Bipolar disorders	100.0	5.9	†	3.8	1.2	†	49.4	#
	Depressive disorders	100.0	5.1	†	2.0	1.3	†	52.2	#
	Schizophrenia and other psychotic disorders	100.0	4.3	†	7.1	4.1	†	37.3	#
	Alcohol or substance-related disorders	100.0	8.6	†	4.0	†	2.1	30.7	#

-- Quantity is zero; † Less than 1 percent individuals.

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2015 reporting periods. Data as of 07.08.2019.

Table 7.9a. Individuals aged 18 years and older served in both 2013 and 2016, by mental health diagnosis in 2013 and living arrangement in 2013 and 2016: Number

<i>Living arrangement in 2013</i>	<i>Diagnostic group in 2013¹</i>	<i>Total</i>	<i>Living arrangement in 2016</i>						
			<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Homeless	Total	25,704	9,997	22	1,722	474	399	11,753	1,337
	Bipolar disorders	6,530	2,413	2	331	91	98	3,242	353
	Depressive disorders	6,757	2,449	3	328	56	62	3,435	424
	Schizophrenia and other psychotic disorders	7,858	3,299	10	824	281	158	2,984	302
	Alcohol or substance-related disorders	1,222	601	1	89	15	21	457	38
Foster home/care	Total	3,849	52	1,276	211	45	14	2,033	218
	Bipolar disorders	662	16	161	35	8	1	407	34
	Depressive disorders	868	12	310	26	3	3	479	35
	Schizophrenia and other psychotic disorders	1,236	10	227	97	22	9	781	90
	Alcohol or substance-related disorders	119	--	117	--	--	--	2	--
Residential care	Total	45,330	1,104	168	29,713	1,435	273	10,920	1,717
	Bipolar disorders	6,573	270	20	3,499	173	56	2,212	343
	Depressive disorders	5,342	193	19	2,932	119	36	1,802	241
	Schizophrenia and other psychotic disorders	27,643	492	87	19,521	1,016	146	5,540	841
	Alcohol or substance-related disorders	381	27	1	246	7	7	87	6
Institutional settings	Total	13,519	358	25	1,586	7,370	223	3,233	724
	Bipolar disorders	1,683	60	3	159	718	29	610	104
	Depressive disorders	1,467	37	2	112	790	13	469	44
	Schizophrenia and other psychotic disorders	8,064	220	15	1,132	4,701	139	1,324	533
	Alcohol or substance-related disorders	492	9	--	13	128	7	335	--

Continued. See notes at end of table.

Table 7.9a. Individuals aged 18 years and older served in both 2013 and 2016, by mental health diagnosis in 2013 and living arrangement in 2013 and 2016: Number (continued)

Living arrangement in 2013	Diagnostic group in 2013 ¹	Total	Living arrangement in 2016						
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Jail/correctional facility	Total	12,692	681	7	639	382	7,486	3,236	261
	Bipolar disorders	2,496	171	--	125	45	1,298	810	47
	Depressive disorders	2,066	82	--	78	20	1,286	553	47
	Schizophrenia and other psychotic disorders	3,682	222	6	301	264	1,753	1,041	95
	Alcohol or substance-related disorders	672	57	--	42	2	376	180	15
Private residence	Total	540,873	11,969	460	12,678	4,708	2,791	489,971	18,296
	Bipolar disorders	136,106	3,391	73	2,490	868	786	123,368	5,130
	Depressive disorders	158,719	3,121	112	2,024	754	488	146,322	5,898
	Schizophrenia and other psychotic disorders	157,896	3,364	168	6,660	2,565	887	141,023	3,229
	Alcohol or substance-related disorders	5,160	345	5	124	50	86	4,465	85
Other	Total	16,721	1,039	134	1,038	480	140	8,715	5,175
	Bipolar disorders	3,499	255	26	199	68	43	1,894	1,014
	Depressive disorders	4,288	283	23	140	65	25	2,602	1,150
	Schizophrenia and other psychotic disorders	6,105	305	68	597	308	44	2,579	2,204
	Alcohol or substance-related disorders	306	30	--	12	3	6	98	157

-- Quantity is zero.

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2016 reporting periods. Data as of 07.08.2019.

Table 7.9b. Individuals aged 18 years and older served in both 2013 and 2016, by mental health diagnosis in 2013 and living arrangement in 2013 and 2016: Row percent distribution

Living arrangement in 2013	Diagnostic group in 2013 ¹	Total	Living arrangement in 2016						
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Homeless	Total	100.0	38.9	†	6.7	1.8	1.6	45.7	5.2
	Bipolar disorders	100.0	37.0	†	5.1	1.4	1.5	49.6	5.4
	Depressive disorders	100.0	36.2	†	4.9	†	†	50.8	6.3
	Schizophrenia and other psychotic disorders	100.0	42.0	†	10.5	3.6	2.0	38.0	3.8
	Alcohol or substance-related disorders	100.0	49.2	†	7.3	1.2	1.7	37.4	3.1
Foster home/care	Total	100.0	1.4	33.2	5.5	1.2	†	52.8	5.7
	Bipolar disorders	100.0	2.4	24.3	5.3	1.2	†	61.5	5.1
	Depressive disorders	100.0	1.4	35.7	3.0	†	†	55.2	4.0
	Schizophrenia and other psychotic disorders	100.0	†	18.4	7.8	1.8	†	63.2	7.3
	Alcohol or substance-related disorders	100.0	--	98.3	--	--	--	1.7	--
Residential care	Total	100.0	2.4	†	65.5	3.2	†	24.1	3.8
	Bipolar disorders	100.0	4.1	†	53.2	2.6	†	33.7	5.2
	Depressive disorders	100.0	3.6	†	54.9	2.2	†	33.7	4.5
	Schizophrenia and other psychotic disorders	100.0	1.8	†	70.6	3.7	†	20.0	3.0
	Alcohol or substance-related disorders	100.0	7.1	†	64.6	1.8	1.8	22.8	1.6
Institutional settings	Total	100.0	2.6	†	11.7	54.5	1.6	23.9	5.4
	Bipolar disorders	100.0	3.6	†	9.4	42.7	1.7	36.2	6.2
	Depressive disorders	100.0	2.5	†	7.6	53.9	†	32.0	3.0
	Schizophrenia and other psychotic disorders	100.0	2.7	†	14.0	58.3	1.7	16.4	6.6
	Alcohol or substance-related disorders	100.0	1.8	--	2.6	26.0	1.4	68.1	--

Continued. See notes at end of table.

Table 7.9b. Individuals aged 18 years and older served in both 2013 and 2016, by mental health diagnosis in 2013 and living arrangement in 2013 and 2016: Row percent distribution (continued)

Living arrangement in 2013	Diagnostic group in 2013 ¹	Total	Living arrangement in 2016						
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Jail/correctional facility	Total	100.0	5.4	†	5.0	3.0	59.0	25.5	2.1
	Bipolar disorders	100.0	6.9	--	5.0	1.8	52.0	32.5	1.9
	Depressive disorders	100.0	4.0	--	3.8	†	62.2	26.8	2.3
	Schizophrenia and other psychotic disorders	100.0	6.0	†	8.2	7.2	47.6	28.3	2.6
	Alcohol or substance-related disorders	100.0	8.5	--	6.3	†	56.0	26.8	2.2
Private residence	Total	100.0	2.2	†	2.3	†	†	90.6	3.4
	Bipolar disorders	100.0	2.5	†	1.8	†	†	90.6	3.8
	Depressive disorders	100.0	2.0	†	1.3	†	†	92.2	3.7
	Schizophrenia and other psychotic disorders	100.0	2.1	†	4.2	1.6	†	89.3	2.0
	Alcohol or substance-related disorders	100.0	6.7	†	2.4	†	1.7	86.5	1.6
Other	Total	100.0	6.2	†	6.2	2.9	†	52.1	#
	Bipolar disorders	100.0	7.3	†	5.7	1.9	1.2	54.1	#
	Depressive disorders	100.0	6.6	†	3.3	1.5	†	60.7	#
	Schizophrenia and other psychotic disorders	100.0	5.0	1.1	9.8	5.0	†	42.2	#
	Alcohol or substance-related disorders	100.0	9.8	--	3.9	†	2.0	32.0	#

-- Quantity is zero; † Less than 1 percent individuals.

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2016 reporting periods. Data as of 07.08.2019.

Table 7.10a. Individuals aged 18 years and older served in both 2013 and 2017, by mental health diagnosis in 2013 and living arrangement in 2013 and 2017: Number

<i>Living arrangement in 2013</i>	<i>Diagnostic group in 2013¹</i>	<i>Total</i>	<i>Living arrangement in 2017</i>						
			<i>Homeless</i>	<i>Foster home/care</i>	<i>Residential care</i>	<i>Institutional setting</i>	<i>Jail/correctional facility</i>	<i>Private residence</i>	<i>Other</i>
Homeless	Total	21,894	7,695	25	1,563	434	393	10,541	1,243
	Bipolar disorders	5,551	1,841	3	321	91	93	2,867	335
	Depressive disorders	5,754	1,879	2	247	54	67	3,125	380
	Schizophrenia and other psychotic disorders	6,835	2,633	15	793	251	152	2,695	296
	Alcohol or substance-related disorders	911	420	1	60	14	15	372	29
Foster home/care	Total	2,924	58	573	235	47	21	1,775	215
	Bipolar disorders	592	20	107	37	7	5	371	45
	Depressive disorders	586	11	98	34	2	2	400	39
	Schizophrenia and other psychotic disorders	1,138	10	196	114	31	9	704	74
	Alcohol or substance-related disorders	9	1	5	1	--	1	1	--
Residential care	Total	40,894	1,080	185	26,155	1,445	265	10,112	1,652
	Bipolar disorders	5,759	254	19	2,897	151	52	2,075	311
	Depressive disorders	4,518	165	18	2,310	119	23	1,645	238
	Schizophrenia and other psychotic disorders	25,596	539	94	17,842	1,051	147	5,099	824
	Alcohol or substance-related disorders	188	23	--	89	6	9	61	--
Institutional settings	Total	11,675	353	31	1,739	5,784	200	2,859	709
	Bipolar disorders	1,433	68	6	172	580	30	496	81
	Depressive disorders	1,085	37	3	110	520	10	375	30
	Schizophrenia and other psychotic disorders	7,252	193	20	1,280	3,849	122	1,252	536
	Alcohol or substance-related disorders	401	11	--	11	94	1	284	--

Continued. See notes at end of table.

Table 7.10a. Individuals aged 18 years and older served in both 2013 and 2017, by mental health diagnosis in 2013 and living arrangement in 2013 and 2017: Number (continued)

Living arrangement in 2013	Diagnostic group in 2013 ¹	Total	Living arrangement in 2017						
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Jail/correctional facility	Total	10,999	719	5	698	380	5,986	2,981	230
	Bipolar disorders	2,114	160	1	122	47	1,031	714	39
	Depressive disorders	1,754	103	--	68	21	1,010	521	31
	Schizophrenia and other psychotic disorders	3,325	235	2	359	252	1,440	945	92
	Alcohol or substance-related disorders	493	61	--	47	5	207	159	14
Private residence	Total	450,921	12,033	434	12,952	4,870	2,806	401,384	16,442
	Bipolar disorders	113,463	3,411	86	2,474	932	798	101,299	4,463
	Depressive disorders	129,437	2,968	100	1,935	733	541	118,011	5,149
	Schizophrenia and other psychotic disorders	137,792	3,583	158	6,971	2,725	878	120,265	3,212
	Alcohol or substance-related disorders	2,973	312	4	95	36	70	2,374	82
Other	Total	13,767	1,001	77	922	408	156	7,579	3,624
	Bipolar disorders	2,841	240	14	162	45	34	1,651	695
	Depressive disorders	3,403	254	14	105	49	26	2,197	758
	Schizophrenia and other psychotic disorders	5,278	308	38	559	280	65	2,326	1,702
	Alcohol or substance-related disorders	160	24	--	5	4	6	88	33

-- Quantity is zero.

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2017 reporting periods. Data as of 07.08.2019.

Table 7.10b. Individuals aged 18 years and older served in both 2013 and 2017, by mental health diagnosis in 2013 and living arrangement in 2013 and 2017: Row percent distribution

Living arrangement in 2013	Diagnostic group in 2013 ¹	Total	Living arrangement in 2017						
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Homeless	Total	100.0	35.1	†	7.1	2.0	1.8	48.1	5.7
	Bipolar disorders	100.0	33.2	†	5.8	1.6	1.7	51.6	6.0
	Depressive disorders	100.0	32.7	†	4.3	†	1.2	54.3	6.6
	Schizophrenia and other psychotic disorders	100.0	38.5	†	11.6	3.7	2.2	39.4	4.3
	Alcohol or substance-related disorders	100.0	46.1	†	6.6	1.5	1.6	40.8	3.2
Foster home/care	Total	100.0	2.0	19.6	8.0	1.6	†	60.7	7.4
	Bipolar disorders	100.0	3.4	18.1	6.3	1.2	†	62.7	7.6
	Depressive disorders	100.0	1.9	16.7	5.8	†	†	68.3	6.7
	Schizophrenia and other psychotic disorders	100.0	†	17.2	10.0	2.7	†	61.9	6.5
	Alcohol or substance-related disorders	100.0	11.1	55.6	11.1	--	11.1	11.1	--
Residential care	Total	100.0	2.6	†	64.0	3.5	†	24.7	4.0
	Bipolar disorders	100.0	4.4	†	50.3	2.6	†	36.0	5.4
	Depressive disorders	100.0	3.7	†	51.1	2.6	†	36.4	5.3
	Schizophrenia and other psychotic disorders	100.0	2.1	†	69.7	4.1	†	19.9	3.2
	Alcohol or substance-related disorders	100.0	12.2	--	47.3	3.2	4.8	32.4	--
Institutional settings	Total	100.0	3.0	†	14.9	49.5	1.7	24.5	6.1
	Bipolar disorders	100.0	4.7	†	12.0	40.5	2.1	34.6	5.7
	Depressive disorders	100.0	3.4	†	10.1	47.9	†	34.6	2.8
	Schizophrenia and other psychotic disorders	100.0	2.7	†	17.7	53.1	1.7	17.3	7.4
	Alcohol or substance-related disorders	100.0	2.7	--	2.7	23.4	†	70.8	--

Continued. See notes at end of table.

Table 7.10b. Individuals aged 18 years and older served in both 2013 and 2017, by mental health diagnosis in 2013 and living arrangement in 2013 and 2017: Row percent distribution (continued)

Living arrangement in 2013	Diagnostic group in 2013 ¹	Total	Living arrangement in 2017						
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Jail/correctional facility	Total	100.0	6.5	†	6.3	3.5	54.4	27.1	2.1
	Bipolar disorders	100.0	7.6	†	5.8	2.2	48.8	33.8	1.8
	Depressive disorders	100.0	5.9	--	3.9	1.2	57.6	29.7	1.8
	Schizophrenia and other psychotic disorders	100.0	7.1	†	10.8	7.6	43.3	28.4	2.8
	Alcohol or substance-related disorders	100.0	12.4	--	9.5	1.0	42.0	32.3	2.8
Private residence	Total	100.0	2.7	†	2.9	1.1	†	89.0	3.6
	Bipolar disorders	100.0	3.0	†	2.2	†	†	89.3	3.9
	Depressive disorders	100.0	2.3	†	1.5	†	†	91.2	4.0
	Schizophrenia and other psychotic disorders	100.0	2.6	†	5.1	2.0	†	87.3	2.3
	Alcohol or substance-related disorders	100.0	10.5	†	3.2	1.2	2.4	79.9	2.8
Other	Total	100.0	7.3	†	6.7	3.0	1.1	55.1	#
	Bipolar disorders	100.0	8.4	†	5.7	1.6	1.2	58.1	#
	Depressive disorders	100.0	7.5	†	3.1	1.4	†	64.6	#
	Schizophrenia and other psychotic disorders	100.0	5.8	†	10.6	5.3	1.2	44.1	#
	Alcohol or substance-related disorders	100.0	15.0	--	3.1	2.5	3.8	55.0	#

-- Quantity is zero; † Less than 1 percent individuals.

¹ "Diagnostic group" refers to the first-mentioned mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2017 reporting periods. Data as of 07.08.2019.

Table 7.11a. Individuals aged 18 years and older served in both 2013 and 2018, by mental health diagnosis in 2013 and living arrangement in 2013 and 2018: Number

Living arrangement in 2013	Diagnostic group in 2013 ¹	Total	Living arrangement in 2018						
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Homeless	Total	19,822	7,043	18	1,439	390	390	9,833	709
	Bipolar disorders	4,963	1,671	--	303	66	82	2,687	154
	Depressive disorders	5,186	1,759	2	228	43	63	2,887	204
	Schizophrenia and other psychotic disorders	6,328	2,398	8	738	241	177	2,530	236
	Alcohol or substance-related disorders	813	373	1	45	13	12	348	21
Foster home/care	Total	2,689	73	472	254	51	22	1,702	115
	Bipolar disorders	519	16	86	39	10	2	348	18
	Depressive disorders	547	12	88	32	1	5	387	22
	Schizophrenia and other psychotic disorders	1,055	30	141	115	33	7	683	46
	Alcohol or substance-related disorders	12	1	8	1	--	--	--	2
Residential care	Total	35,710	1,104	170	22,153	1,344	245	9,889	805
	Bipolar disorders	5,110	234	23	2,544	143	43	2,005	118
	Depressive disorders	4,035	203	13	2,005	96	22	1,595	101
	Schizophrenia and other psychotic disorders	22,212	538	92	14,928	985	143	5,012	514
	Alcohol or substance-related disorders	182	25	--	78	7	7	63	2
Institutional settings	Total	9,668	352	27	1,480	4,629	174	2,724	282
	Bipolar disorders	1,148	63	6	144	428	27	452	28
	Depressive disorders	876	36	3	88	383	12	334	20
	Schizophrenia and other psychotic disorders	6,028	210	16	1,094	3,124	104	1,276	204
	Alcohol or substance-related disorders	379	8	--	13	101	1	256	--

Continued. See notes at end of table.

Table 7.11a. Individuals aged 18 years and older served in both 2013 and 2018, by mental health diagnosis in 2013 and living arrangement in 2013 and 2018: Number (continued)

Living arrangement in 2013	Diagnostic group in 2013 ¹	Total	Living arrangement in 2018						
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other
Jail/correctional facility	Total	10,103	724	5	589	333	5,396	2,836	220
	Bipolar disorders	1,929	136	1	108	40	936	669	39
	Depressive disorders	1,605	107	--	47	14	917	489	31
	Schizophrenia and other psychotic disorders	2,978	241	2	311	221	1,229	899	75
	Alcohol or substance-related disorders	489	58	--	35	10	211	158	17
Private residence	Total	407,134	12,361	321	12,898	4,921	2,780	367,940	5,913
	Bipolar disorders	101,533	3,477	58	2,594	903	733	92,390	1,378
	Depressive disorders	114,933	3,044	59	1,939	771	542	107,160	1,418
	Schizophrenia and other psychotic disorders	128,222	3,682	118	6,813	2,773	901	112,065	1,870
	Alcohol or substance-related disorders	2,707	297	3	80	34	55	2,150	88
Other	Total	12,137	1,002	56	891	412	129	6,957	2,690
	Bipolar disorders	2,490	239	5	154	44	26	1,556	466
	Depressive disorders	2,929	243	9	106	39	21	1,968	543
	Schizophrenia and other psychotic disorders	4,745	301	29	545	297	52	2,176	1,345
	Alcohol or substance-related disorders	128	18	--	5	3	3	73	26

-- Quantity is zero.

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.11b. Individuals aged 18 years and older served in both 2013 and 2018, by mental health diagnosis in 2013 and living arrangement in 2013 and 2018: Row percent distribution

Living arrangement in 2013	Diagnostic group in 2013 ¹	Total	Living arrangement in 2018							
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other	
Homeless	Total	100.0	35.5	†	7.3	2.0	2.0	49.6	3.6	
	Bipolar disorders	100.0	33.7	--	6.1	1.3	1.7	54.1	3.1	
	Depressive disorders	100.0	33.9	†	4.4	†	1.2	55.7	3.9	
	Schizophrenia and other psychotic disorders	100.0	37.9	†	11.7	3.8	2.8	40.0	3.7	
	Alcohol or substance-related disorders	100.0	45.9	†	5.5	1.6	1.5	42.8	2.6	
	Foster home/care	Total	100.0	2.7	17.6	9.4	1.9	†	63.3	4.3
Foster home/care	Bipolar disorders	100.0	3.1	16.6	7.5	1.9	†	67.1	3.5	
	Depressive disorders	100.0	2.2	16.1	5.9	†	†	70.7	4.0	
	Schizophrenia and other psychotic disorders	100.0	2.8	13.4	10.9	3.1	†	64.7	4.4	
	Alcohol or substance-related disorders	100.0	8.3	66.7	8.3	--	--	--	#	
	Residential care	Total	100.0	3.1	†	62.0	3.8	†	27.7	2.3
	Residential care	Bipolar disorders	100.0	4.6	†	49.8	2.8	†	39.2	2.3
Depressive disorders		100.0	5.0	†	49.7	2.4	†	39.5	2.5	
Schizophrenia and other psychotic disorders		100.0	2.4	†	67.2	4.4	†	22.6	2.3	
Alcohol or substance-related disorders		100.0	13.7	--	42.9	3.8	3.8	34.6	1.1	
Institutional settings		Total	100.0	3.6	†	15.3	47.9	1.8	28.2	2.9
Institutional settings		Bipolar disorders	100.0	5.5	†	12.5	37.3	2.4	39.4	2.4
	Depressive disorders	100.0	4.1	†	10.0	43.7	1.4	38.1	2.3	
	Schizophrenia and other psychotic disorders	100.0	3.5	†	18.1	51.8	1.7	21.2	3.4	
	Alcohol or substance-related disorders	100.0	2.1	--	3.4	26.6	†	67.5	--	

Continued. See notes at end of table.

Table 7.11b. Individuals aged 18 years and older served in both 2013 and 2018, by mental health diagnosis in 2013 and living arrangement in 2013 and 2018: Row percent distribution (continued)

Living arrangement in 2013	Diagnostic group in 2013 ¹	Total	Living arrangement in 2018							
			Homeless	Foster home/care	Residential care	Institutional setting	Jail/correctional facility	Private residence	Other	
Jail/correctional facility	Total	100.0	7.2	†	5.8	3.3	53.4	28.1	2.2	
	Bipolar disorders	100.0	7.1	†	5.6	2.1	48.5	34.7	2.0	
	Depressive disorders	100.0	6.7	--	2.9	†	57.1	30.5	1.9	
	Schizophrenia and other psychotic disorders	100.0	8.1	†	10.4	7.4	41.3	30.2	2.5	
	Alcohol or substance-related disorders	100.0	11.9	--	7.2	2.0	43.1	32.3	3.5	
	Private residence	Total	100.0	3.0	†	3.2	1.2	†	90.4	1.5
Private residence	Bipolar disorders	100.0	3.4	†	2.6	†	†	91.0	1.4	
	Depressive disorders	100.0	2.6	†	1.7	†	†	93.2	1.2	
	Schizophrenia and other psychotic disorders	100.0	2.9	†	5.3	2.2	†	87.4	1.5	
	Alcohol or substance-related disorders	100.0	11.0	†	3.0	1.3	2.0	79.4	3.3	
	Other	Total	100.0	8.3	†	7.3	3.4	1.1	57.3	#
	Other	Bipolar disorders	100.0	9.6	†	6.2	1.8	1.0	62.5	#
Depressive disorders		100.0	8.3	†	3.6	1.3	†	67.2	#	
Schizophrenia and other psychotic disorders		100.0	6.3	†	11.5	6.3	1.1	45.9	#	
Alcohol or substance-related disorders		100.0	14.1	--	3.9	2.3	2.3	57.0	#	

-- Quantity is zero; † Less than 1 percent individuals.

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.12. Individuals aged 18 years and older served in consecutive reporting periods, by mental health diagnosis and employment status: Number, 2013–2018

<i>Living arrangement</i>	<i>Diagnostic group¹</i>	<i>Reporting periods</i>				
		<i>2013</i>	<i>2013–2015</i>	<i>2013–2016</i>	<i>2013–2017</i>	<i>2013–2018</i>
Employed	Total	300,146	37,398	13,756	8,148	5,653
	Bipolar disorders	69,561	10,484	3,926	2,240	1,477
	Depressive disorders	110,003	13,010	4,295	2,405	1,618
	Schizophrenia and other psychotic disorders	28,958	7,566	3,836	2,624	1,969
	Alcohol or substance-related disorders	5,232	118	14	2	1
Unemployed	Total	573,679	76,941	17,196	9,383	6,242
	Bipolar disorders	143,816	20,478	4,374	2,272	1,439
	Depressive disorders	202,815	26,240	5,284	2,620	1,632
	Schizophrenia and other psychotic disorders	86,536	19,499	5,861	3,788	2,751
	Alcohol or substance-related disorders	12,716	294	75	12	5
Not in labor force	Total	885,266	215,832	114,606	78,757	58,258
	Bipolar disorders	206,881	48,809	24,046	15,624	10,887
	Depressive disorders	274,479	59,499	29,744	19,188	13,566
	Schizophrenia and other psychotic disorders	221,172	83,755	52,165	39,179	30,427
	Alcohol or substance-related disorders	8,395	241	80	8	3

¹ “Diagnostic group” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. “Total” includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 7.13a. Individuals aged 18 years and older served in both 2013 and 2015, by mental health diagnosis in 2013 and employment status in 2013 and 2015: Number

<i>Employment status in 2013</i>	<i>Diagnostic group in 2013¹</i>	<i>Employment status in 2015</i>			
		<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Employed	Total	94,229	64,501	12,712	17,016
	Bipolar disorders	26,385	17,643	3,707	5,035
	Depressive disorders	31,663	22,002	4,325	5,336
	Schizophrenia and other psychotic disorders	16,543	10,892	1,954	3,697
	Alcohol or substance-related disorders	470	318	87	65
Unemployed	Total	187,696	16,190	131,262	40,244
	Bipolar disorders	50,820	4,931	34,963	10,926
	Depressive disorders	60,778	5,672	42,908	12,198
	Schizophrenia and other psychotic disorders	40,144	2,199	27,387	10,558
	Alcohol or substance-related disorders	1,558	162	1,166	230
Not in labor force	Total	384,547	18,880	33,672	331,995
	Bipolar disorders	93,140	5,887	8,730	78,523
	Depressive disorders	105,512	5,644	8,854	91,014
	Schizophrenia and other psychotic disorders	127,966	4,179	10,795	112,992
	Alcohol or substance-related disorders	1,522	55	167	1,300

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2015 reporting periods. Data as of 07.08.2019.

Table 7.13b. Individuals aged 18 years and older served in both 2013 and 2015, by mental health diagnosis in 2013 and employment status in 2013 and 2015: Row percent distribution

<i>Employment status in 2013</i>	<i>Diagnostic group in 2013¹</i>	<i>Employment status in 2015</i>			
		<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Employed	Total	100.0	68.5	13.5	18.1
	Bipolar disorders	100.0	66.9	14.0	19.1
	Depressive disorders	100.0	69.5	13.7	16.9
	Schizophrenia and other psychotic disorders	100.0	65.8	11.8	22.3
	Alcohol or substance-related disorders	100.0	67.7	18.5	13.8
Unemployed	Total	100.0	8.6	69.9	21.4
	Bipolar disorders	100.0	9.7	68.8	21.5
	Depressive disorders	100.0	9.3	70.6	20.1
	Schizophrenia and other psychotic disorders	100.0	5.5	68.2	26.3
	Alcohol or substance-related disorders	100.0	10.4	74.8	14.8
Not in labor force	Total	100.0	4.9	8.8	86.3
	Bipolar disorders	100.0	6.3	9.4	84.3
	Depressive disorders	100.0	5.3	8.4	86.3
	Schizophrenia and other psychotic disorders	100.0	3.3	8.4	88.3
	Alcohol or substance-related disorders	100.0	3.6	11.0	85.4

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2015 reporting periods. Data as of 07.08.2019.

Table 7.14a. Individuals aged 18 years and older served in both 2013 and 2016, by mental health diagnosis in 2013 and employment status in 2013 and 2016: Number

<i>Employment status in 2013</i>	<i>Diagnostic group in 2013¹</i>	<i>Total</i>	<i>Employment status in 2016</i>		
			<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Employed	Total	65,433	39,616	9,652	16,165
	Bipolar disorders	18,289	10,781	2,903	4,605
	Depressive disorders	21,026	12,916	3,200	4,910
	Schizophrenia and other psychotic disorders	13,470	8,038	1,500	3,932
	Alcohol or substance-related disorders	350	221	76	53
Unemployed	Total	107,917	13,221	58,166	36,530
	Bipolar disorders	29,039	3,964	15,462	9,613
	Depressive disorders	33,041	4,449	17,703	10,889
	Schizophrenia and other psychotic disorders	27,052	2,052	14,567	10,433
	Alcohol or substance-related disorders	1,171	144	827	200
Not in labor force	Total	304,111	17,595	25,424	261,092
	Bipolar disorders	72,739	5,314	6,763	60,662
	Depressive disorders	79,978	5,119	6,803	68,056
	Schizophrenia and other psychotic disorders	108,133	4,267	7,571	96,295
	Alcohol or substance-related disorders	1,194	50	146	998

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2016 reporting periods. Data as of 07.08.2019.

Table 7.14b. Individuals aged 18 years and older served in both 2013 and 2016, by mental health diagnosis in 2013 and employment status in 2013 and 2016: Row percent distribution

<i>Employment status in 2013</i>	<i>Diagnostic group in 2013¹</i>	<i>Employment status in 2016</i>			
		<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Employed	Total	100.0	60.5	14.8	24.7
	Bipolar disorders	100.0	58.9	15.9	25.2
	Depressive disorders	100.0	61.4	15.2	23.4
	Schizophrenia and other psychotic disorders	100.0	59.7	11.1	29.2
	Alcohol or substance-related disorders	100.0	63.1	21.7	15.1
Unemployed	Total	100.0	12.3	53.9	33.9
	Bipolar disorders	100.0	13.7	53.2	33.1
	Depressive disorders	100.0	13.5	53.6	33.0
	Schizophrenia and other psychotic disorders	100.0	7.6	53.8	38.6
	Alcohol or substance-related disorders	100.0	12.3	70.6	17.1
Not in labor force	Total	100.0	5.8	8.4	85.9
	Bipolar disorders	100.0	7.3	9.3	83.4
	Depressive disorders	100.0	6.4	8.5	85.1
	Schizophrenia and other psychotic disorders	100.0	3.9	7.0	89.1
	Alcohol or substance-related disorders	100.0	4.2	12.2	83.6

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2016 reporting periods. Data as of 07.08.2019.

Table 7.15a. Individuals aged 18 years and older served in both 2013 and 2017, by mental health diagnosis in 2013 and employment status in 2013 and 2017: Number

<i>Employment status in 2013</i>	<i>Diagnostic group in 2013¹</i>	<i>Employment status in 2017</i>			
		<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Employed	Total	55,130	30,801	8,918	15,411
	Bipolar disorders	15,408	8,368	2,636	4,404
	Depressive disorders	17,363	9,861	2,860	4,642
	Schizophrenia and other psychotic disorders	11,828	6,520	1,449	3,859
	Alcohol or substance-related disorders	215	103	66	46
Unemployed	Total	91,617	11,676	45,707	34,234
	Bipolar disorders	24,641	3,519	12,151	8,971
	Depressive disorders	27,402	3,934	13,578	9,890
	Schizophrenia and other psychotic disorders	23,971	1,827	12,018	10,126
	Alcohol or substance-related disorders	941	129	620	192
Not in labor force	Total	259,074	16,624	24,639	217,811
	Bipolar disorders	61,817	4,949	6,530	50,338
	Depressive disorders	66,322	4,750	6,679	54,893
	Schizophrenia and other psychotic disorders	94,340	4,080	7,219	83,041
	Alcohol or substance-related disorders	862	42	157	663

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2017 reporting periods. Data as of 07.08.2019.

Table 7.15b. Individuals aged 18 years and older served in both 2013 and 2017, by mental health diagnosis in 2013 and employment status in 2013 and 2017: Row percent distribution

<i>Employment status in 2013</i>	<i>Diagnostic group in 2013¹</i>	<i>Employment status in 2017</i>			
		<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Employed	Total	100.0	55.9	16.2	28.0
	Bipolar disorders	100.0	54.3	17.1	28.6
	Depressive disorders	100.0	56.8	16.5	26.7
	Schizophrenia and other psychotic disorders	100.0	55.1	12.3	32.6
	Alcohol or substance-related disorders	100.0	47.9	30.7	21.4
Unemployed	Total	100.0	12.7	49.9	37.4
	Bipolar disorders	100.0	14.3	49.3	36.4
	Depressive disorders	100.0	14.4	49.6	36.1
	Schizophrenia and other psychotic disorders	100.0	7.6	50.1	42.2
	Alcohol or substance-related disorders	100.0	13.7	65.9	20.4
Not in labor force	Total	100.0	6.4	9.5	84.1
	Bipolar disorders	100.0	8.0	10.6	81.4
	Depressive disorders	100.0	7.2	10.1	82.8
	Schizophrenia and other psychotic disorders	100.0	4.3	7.7	88.0
	Alcohol or substance-related disorders	100.0	4.9	18.2	76.9

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual.

"Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2017 reporting periods. Data as of 07.08.2019.

Table 7.16a. Individuals aged 18 years and older served in both 2013 and 2018, by mental health diagnosis in 2013 and employment status in 2013 and 2018: Number

<i>Employment status in 2013</i>	<i>Diagnostic group in 2013¹</i>	<i>Employment status in 2018</i>			
		<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Employed	Total	50,492	27,089	8,954	14,449
	Bipolar disorders	14,100	7,338	2,633	4,129
	Depressive disorders	15,726	8,571	2,813	4,342
	Schizophrenia and other psychotic disorders	11,097	5,886	1,582	3,629
	Alcohol or substance-related disorders	231	107	77	47
Unemployed	Total	83,248	11,024	41,014	31,210
	Bipolar disorders	22,378	3,360	10,809	8,209
	Depressive disorders	24,373	3,640	11,772	8,961
	Schizophrenia and other psychotic disorders	22,326	1,768	11,255	9,303
	Alcohol or substance-related disorders	895	130	572	193
Not in labor force	Total	241,301	15,651	27,996	197,654
	Bipolar disorders	56,766	4,703	7,031	45,032
	Depressive disorders	60,431	4,408	7,108	48,915
	Schizophrenia and other psychotic disorders	90,235	3,724	9,412	77,099
	Alcohol or substance-related disorders	928	47	154	727

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2018 reporting periods. Data as of 07.08.2019.

Table 7.16b. Individuals aged 18 years and older served in both 2013 and 2018, by mental health diagnosis in 2013 and employment status in 2013 and 2018: Row percent distribution

<i>Employment status in 2013</i>	<i>Diagnostic group in 2013¹</i>	<i>Employment status in 2018</i>			
		<i>Total</i>	<i>Employed</i>	<i>Unemployed</i>	<i>Not in labor force</i>
Employed	Total	100.0	53.7	17.7	28.6
	Bipolar disorders	100.0	52.0	18.7	29.3
	Depressive disorders	100.0	54.5	17.9	27.6
	Schizophrenia and other psychotic disorders	100.0	53.0	14.3	32.7
	Alcohol or substance-related disorders	100.0	46.3	33.3	20.3
Unemployed	Total	100.0	13.2	49.3	37.5
	Bipolar disorders	100.0	15.0	48.3	36.7
	Depressive disorders	100.0	14.9	48.3	36.8
	Schizophrenia and other psychotic disorders	100.0	7.9	50.4	41.7
	Alcohol or substance-related disorders	100.0	14.5	63.9	21.6
Not in labor force	Total	100.0	6.5	11.6	81.9
	Bipolar disorders	100.0	8.3	12.4	79.3
	Depressive disorders	100.0	7.3	11.8	80.9
	Schizophrenia and other psychotic disorders	100.0	4.1	10.4	85.4
	Alcohol or substance-related disorders	100.0	5.1	16.6	78.3

¹ "Diagnostic group" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Total" includes all mental health diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013 and 2018 reporting periods. Data as of 07.08.2019.

Table 8.1. Individuals, by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2018

<i>Census region, Census division, and state or jurisdiction</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders²</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total	5,494,012	793,054	654,028	461,444	601,023	1,410,277	45,746	660,521	867,919
Census region³									
Northeast	680,681	106,028	70,454	71,608	72,218	168,294	7,425	76,430	108,224
Midwest	1,499,950	226,145	209,559	138,659	154,566	376,047	17,945	150,902	226,127
South	1,622,266	187,464	143,139	150,543	234,027	468,261	9,994	222,488	206,350
West	1,688,168	273,304	230,766	100,431	140,028	396,941	10,363	209,919	326,416
Census division³									
New England	136,773	24,497	13,220	9,518	13,515	31,334	1,125	18,893	24,671
Middle Atlantic	543,908	81,531	57,234	62,090	58,703	136,960	6,300	57,537	83,553
East North Central	940,886	127,390	99,449	80,151	108,921	240,523	11,572	104,768	168,112
West North Central	559,064	98,755	110,110	58,508	45,645	135,524	6,373	46,134	58,015
South Atlantic	776,905	105,152	80,758	71,951	103,678	211,769	6,903	106,124	90,570
East South Central	406,543	45,334	37,755	43,578	43,801	105,401	2,157	45,634	82,883
West South Central	438,818	36,978	24,626	35,014	86,548	151,091	934	70,730	32,897
Mountain	614,391	96,741	94,375	42,948	62,366	139,486	4,541	54,083	119,851
Pacific	1,073,777	176,563	136,391	57,483	77,662	257,455	5,822	155,836	206,565
State or jurisdiction									
Alabama	101,599	8,158	5,900	13,669	11,831	27,591	437	17,752	16,261
Alaska	‡	‡	‡	‡	‡	‡	‡	‡	‡
Arizona	169,861	12,659	29,586	20,587	28,143	42,320	1,319	21,951	13,296
Arkansas	59,565	10,359	7,699	1,075	6,044	18,416	418	6,758	8,796
California	660,567	75,912	54,673	28,971	49,619	160,164	1,057	121,780	168,391
Colorado	148,846	36,755	18,175	7,276	15,560	39,107	974	12,703	18,296
Connecticut	77,860	13,690	8,023	6,462	7,914	18,344	506	10,517	12,404
Delaware	8,800	612	205	254	1,490	1,925	33	1,683	2,598
District of Columbia	35,828	2,880	1,263	2,399	6,155	12,846	38	6,670	3,577
Florida	185,688	26,900	14,027	13,081	26,120	53,057	515	32,965	19,023
Georgia	‡	‡	‡	‡	‡	‡	‡	‡	‡
Hawaii	9,097	2,401	275	395	1,032	1,557	15	2,719	703
Idaho	7,666	291	383	218	1,536	2,271	154	1,985	828
Illinois	66,235	8,809	5,655	5,127	10,132	18,504	241	10,409	7,358

Continued. See notes at end of table.

**Table 8.1. Individuals, by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2018
(continued)**

Census region, Census division, and state or jurisdiction	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ²
					Bipolar disorders	Depressive disorders			
Indiana	137,363	11,718	13,401	14,142	13,813	47,509	1,044	14,899	20,837
Iowa	174,011	29,171	35,102	25,669	11,833	43,078	790	10,473	17,895
Kansas	‡	‡	‡	‡	‡	‡	‡	‡	‡
Kentucky	155,989	22,995	17,814	13,405	11,021	35,319	941	10,367	44,127
Louisiana	23,000	1,127	1,729	1,993	3,559	8,596	78	4,631	1,287
Maine	‡	‡	‡	‡	‡	‡	‡	‡	‡
Maryland	211,849	33,567	29,107	28,739	26,331	56,339	498	16,323	20,945
Massachusetts	9,248	712	827	318	1,656	1,861	31	3,322	521
Michigan	224,567	18,469	13,515	12,034	28,423	53,417	1,302	30,743	66,664
Minnesota	269,422	59,657	61,843	23,686	15,009	58,737	4,578	17,509	28,403
Mississippi	84,918	7,577	6,237	13,776	8,855	20,203	300	10,365	17,605
Missouri	73,128	3,028	9,068	6,764	12,909	22,356	797	13,187	5,019
Montana	61,189	9,788	13,719	4,220	3,759	14,071	550	3,370	11,712
Nebraska	20,011	2,462	1,760	721	3,479	5,700	116	2,264	3,509
Nevada	13,250	1,681	1,589	367	1,008	2,022	437	2,003	4,143
New Hampshire	‡	‡	‡	‡	‡	‡	‡	‡	‡
New Jersey	‡	‡	‡	‡	‡	‡	‡	‡	‡
New Mexico	149,498	25,568	20,189	7,559	6,705	20,739	230	5,557	62,951
New York ⁴	56,509	7,579	3,271	1,870	5,166	7,603	4,586	19,316	7,118
North Carolina	90,175	11,858	8,114	2,439	14,038	29,505	1,110	13,345	9,766
North Dakota	7,182	505	776	443	925	1,853	87	1,259	1,334
Ohio	449,140	81,609	56,369	46,037	50,839	109,867	2,835	39,583	62,001
Oklahoma	87,512	14,598	9,960	3,580	10,429	32,048	141	9,690	7,066
Oregon	134,637	42,073	25,243	7,672	8,850	25,810	691	11,182	13,116
Pennsylvania	487,399	73,952	53,963	60,220	53,537	129,357	1,714	38,221	76,435
Puerto Rico	2,669	105	101	193	180	718	19	670	683

Continued. See notes at end of table.

**Table 8.1. Individuals, by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2018
(continued)**

<i>Census region, Census division, and state or jurisdiction</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders²</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Rhode Island	20,637	2,447	1,671	1,223	2,860	7,445	233	3,414	1,344
South Carolina	84,322	10,564	6,100	10,116	10,409	20,051	872	15,733	10,477
South Dakota	15,310	3,932	1,561	1,225	1,490	3,800	5	1,442	1,855
Tennessee ⁴	64,037	6,604	7,804	2,728	12,094	22,288	479	7,150	4,890
Texas	268,741	10,894	5,238	28,366	66,516	92,031	297	49,651	15,748
Utah	48,319	7,028	8,436	1,859	4,168	14,276	561	5,575	6,416
Vermont	29,028	7,648	2,699	1,515	1,085	3,684	355	1,640	10,402
Virginia	117,261	15,114	14,530	11,621	13,368	25,589	3,583	17,143	16,313
Washington	269,476	56,177	56,200	20,445	18,161	69,924	4,059	20,155	24,355
West Virginia	42,982	3,657	7,412	3,302	5,767	12,457	254	2,262	7,871
Wisconsin	63,581	6,785	10,509	2,811	5,714	11,226	6,150	9,134	11,252
Wyoming	15,762	2,971	2,298	862	1,487	4,680	316	939	2,209
Other jurisdictions ⁵	278	8	9	10	4	16	‡	112	119

‡ No data reported.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

² "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes substance use diagnoses. See Appendix E.

³ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

⁴ New York and Tennessee report only a subset of the total individuals served through their SMHAs.

⁵ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 8.2. Individuals per 100,000 population (2018 estimate), by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2018

<i>Census region, Census division, and state or jurisdiction</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders²</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total	1,823	263	217	153	199	468	15	219	288
Census region³									
Northeast	1,529	238	158	161	162	378	17	172	243
Midwest	2,294	346	320	212	236	575	27	231	346
South	1,420	164	125	132	205	410	9	195	181
West	2,185	354	299	130	181	514	13	272	423
Census division³									
New England	1,125	201	109	78	111	258	9	155	203
Middle Atlantic	1,681	252	177	192	181	423	19	178	258
East North Central	2,005	271	212	171	232	512	25	223	358
West North Central	3,028	535	596	317	247	734	35	250	314
South Atlantic	1,418	192	147	131	189	386	13	194	165
East South Central	2,127	237	198	228	229	551	11	239	434
West South Central	1,088	92	61	87	215	375	2	175	82
Mountain	2,502	394	384	175	254	568	19	220	488
Pacific	2,037	335	259	109	147	488	11	296	392
State or jurisdiction									
Alabama	2,079	167	121	280	242	564	9	363	333
Alaska	‡	‡	‡	‡	‡	‡	‡	‡	‡
Arizona	2,369	177	413	287	392	590	18	306	185
Arkansas	1,976	344	255	36	201	611	14	224	292
California	1,670	192	138	73	125	405	3	308	426
Colorado	2,613	645	319	128	273	687	17	223	321
Connecticut	2,179	383	225	181	222	513	14	294	347
Delaware	910	63	21	26	154	199	3	174	269
District of Columbia	5,100	410	180	342	876	1,829	5	950	509
Florida	872	126	66	61	123	249	2	155	89
Georgia	‡	‡	‡	‡	‡	‡	‡	‡	‡
Hawaii	640	169	19	28	73	110	1	191	49
Idaho	437	17	22	12	88	129	9	113	47
Illinois	520	69	44	40	80	145	2	82	58

Continued. See notes at end of table.

Table 8.2. Individuals per 100,000 population (2018 estimate), by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2018 (continued)

Census region, Census division, and state or jurisdiction	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ²
					Bipolar disorders	Depressive disorders			
Indiana	2,053	175	200	211	206	710	16	223	311
Iowa	5,513	924	1,112	813	375	1,365	25	332	567
Kansas	‡	‡	‡	‡	‡	‡	‡	‡	‡
Kentucky	3,491	515	399	300	247	790	21	232	988
Louisiana	494	24	37	43	76	184	2	99	28
Maine	‡	‡	‡	‡	‡	‡	‡	‡	‡
Maryland	3,506	556	482	476	436	932	8	270	347
Massachusetts	134	10	12	5	24	27	†	48	8
Michigan	2,247	185	135	120	284	534	13	308	667
Minnesota	4,802	1,063	1,102	422	267	1,047	82	312	506
Mississippi	2,843	254	209	461	297	676	10	347	589
Missouri	1,194	49	148	110	211	365	13	215	82
Montana	5,760	921	1,291	397	354	1,325	52	317	1,103
Nebraska	1,037	128	91	37	180	295	6	117	182
Nevada	437	55	52	12	33	67	14	66	137
New Hampshire	‡	‡	‡	‡	‡	‡	‡	‡	‡
New Jersey	‡	‡	‡	‡	‡	‡	‡	‡	‡
New Mexico	7,134	1,220	963	361	320	990	11	265	3,004
New York ⁴	289	39	17	10	26	39	23	99	36
North Carolina	868	114	78	23	135	284	11	129	94
North Dakota	945	66	102	58	122	244	11	166	176
Ohio	3,842	698	482	394	435	940	24	339	530
Oklahoma	2,219	370	253	91	264	813	4	246	179
Oregon	3,213	1,004	602	183	211	616	16	267	313
Pennsylvania	3,806	577	421	470	418	1,010	13	298	597
Puerto Rico	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>

Continued. See notes at end of table.

Table 8.2. Individuals per 100,000 population (2018 estimate), by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2018 (continued)

Census region, Census division, and state or jurisdiction	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ²
					Bipolar disorders	Depressive disorders			
Rhode Island	1,952	231	158	116	271	704	22	323	127
South Carolina	1,659	208	120	199	205	394	17	309	206
South Dakota	1,735	446	177	139	169	431	†	163	210
Tennessee ⁴	946	98	115	40	179	329	7	106	72
Texas	936	38	18	99	232	321	1	173	55
Utah	1,529	222	267	59	132	452	18	176	203
Vermont	4,635	1,221	431	242	173	588	57	262	1,661
Virginia	1,377	177	171	136	157	300	42	201	192
Washington	3,576	745	746	271	241	928	54	267	323
West Virginia	2,380	203	410	183	319	690	14	125	436
Wisconsin	1,094	117	181	48	98	193	106	157	194
Wyoming	2,728	514	398	149	257	810	55	163	382
Other jurisdictions ⁵	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>

‡ No data reported; † Less than 1 per 100,000 population; *n/a* Population detail required for adjustment was not available.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. See Appendix E.

³ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

⁴ New York and Tennessee report only a subset of the total individuals served through their SMHAs.

⁵ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: States that submitted insufficient data for inclusion in this report, as well as their reference populations, are excluded from calculations presented in this table.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2018-alldata6: Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2018.

Table 8.3. Individuals per 100,000 population, by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2018, adjusted to 2010 population estimate for age, gender, and race

<i>Census region, Census division, and state or jurisdiction</i>	<i>Mental health diagnosis¹</i>								
	<i>Total</i>	<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders²</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Total	1,721	246	204	151	195	443	14	201	267
Census region³									
Northeast	1,535	247	162	178	162	371	16	158	242
Midwest	2,156	314	290	196	228	544	26	227	331
South	1,356	159	125	126	202	395	9	169	173
West	2,039	329	274	129	180	472	13	265	376
Census division³									
New England	1,038	200	103	81	104	238	9	149	153
Middle Atlantic	1,722	264	185	213	185	422	18	160	274
East North Central	1,929	258	201	168	223	491	24	214	351
West North Central	2,747	458	518	261	242	695	32	268	273
South Atlantic	1,336	187	150	129	181	361	13	157	158
East South Central	2,112	248	211	220	237	546	12	202	436
West South Central	1,050	84	57	81	216	369	2	168	74
Mountain	2,313	352	349	170	254	532	18	233	405
Pacific	1,845	312	232	106	142	427	11	271	344
State or jurisdiction									
Alabama	1,937	166	126	263	250	537	9	270	316
Alaska	‡	‡	‡	‡	‡	‡	‡	‡	‡
Arizona	2,597	194	437	321	433	634	20	356	203
Arkansas	1,657	284	215	29	186	512	12	193	226
California	1,396	159	104	67	115	321	2	266	362
Colorado	2,353	573	272	120	250	606	16	225	291
Connecticut	2,136	383	219	186	216	490	14	281	348
Delaware	880	63	22	26	152	192	3	147	275
District of Columbia	1,694	135	67	126	296	563	2	333	171
Florida	863	129	67	67	122	243	2	139	93
Georgia	‡	‡	‡	‡	‡	‡	‡	‡	‡
Hawaii	572	131	18	19	94	103	†	179	27
Idaho	457	18	26	11	82	127	8	139	45
Illinois	495	66	43	39	77	139	2	76	54

Continued. See notes at end of table.

Table 8.3. Individuals per 100,000 population, by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2018, adjusted to 2010 population estimate for age, gender, and race (continued)

Census region, Census division, and state or jurisdiction	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ²
					Bipolar disorders	Depressive disorders			
Indiana	2,100	176	201	216	212	722	16	240	318
Iowa	4,791	768	965	632	348	1,237	23	340	478
Kansas	‡	‡	‡	‡	‡	‡	‡	‡	‡
Kentucky	3,517	520	393	314	246	776	22	242	1,005
Louisiana	432	22	38	31	75	175	2	67	22
Maine	‡	‡	‡	‡	‡	‡	‡	‡	‡
Maryland	3,175	515	504	438	404	819	9	194	293
Massachusetts	106	8	8	3	19	18	†	44	6
Michigan	2,251	190	138	124	287	534	13	294	671
Minnesota	4,508	975	989	354	270	1,044	78	348	449
Mississippi	2,466	250	225	348	309	594	9	222	508
Missouri	1,189	49	147	111	211	361	13	217	81
Montana	5,941	930	1,281	440	377	1,280	54	369	1,211
Nebraska	1,101	129	90	36	189	310	6	144	199
Nevada	375	49	40	9	30	57	12	58	119
New Hampshire	‡	‡	‡	‡	‡	‡	‡	‡	‡
New Jersey	‡	‡	‡	‡	‡	‡	‡	‡	‡
New Mexico	6,180	1,119	880	310	295	881	12	241	2,443
New York ⁴	258	35	17	10	25	36	20	81	35
North Carolina	843	114	82	22	136	281	11	106	92
North Dakota	890	63	93	54	124	229	10	170	146
Ohio	3,475	638	436	371	387	837	21	298	488
Oklahoma	2,361	377	264	93	284	873	4	280	187
Oregon	3,225	1,007	586	191	209	605	16	307	304
Pennsylvania	4,027	624	441	521	434	1,050	14	299	644
Puerto Rico	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a

Continued. See notes at end of table.

Table 8.3. Individuals per 100,000 population, by mental health diagnosis, Census region, Census division, and state or jurisdiction: Number, 2018, adjusted to 2010 population estimate for age, gender, and race (continued)

Census region, Census division, and state or jurisdiction	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ²
					Bipolar disorders	Depressive disorders			
Rhode Island	1,753	209	141	105	247	621	20	295	116
South Carolina	1,492	201	126	170	204	370	18	217	187
South Dakota	1,948	528	186	149	189	474	†	192	230
Tennessee ⁴	1,036	115	133	48	196	350	10	100	83
Texas	939	37	18	94	236	326	1	174	53
Utah	1,544	211	249	52	138	464	21	225	183
Vermont	3,775	1,371	425	303	150	528	52	264	681
Virginia	1,341	178	177	139	156	296	41	168	185
Washington	3,796	812	769	291	255	982	56	297	335
West Virginia	2,544	223	423	215	341	703	15	140	483
Wisconsin	1,115	123	175	49	99	189	133	169	177
Wyoming	2,720	514	358	145	247	807	58	202	389
Other jurisdictions ⁵	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>	<i>n/a</i>

‡ No data reported; † Less than 1 per 100,000 population; *n/a* Population detail required for adjustment was not available.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

² "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. See Appendix E.

³ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

⁴ New York and Tennessee report only a subset of the total individuals served through their SMHAs.

⁵ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: States that submitted insufficient data for inclusion in this report, as well as their reference populations, are excluded from calculations presented in this table.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019; Population: U.S. Census Bureau, Population Estimates Program, SC-EST2018-alldata6: Annual State Resident Population Estimates for 6 Race Groups (5 Race Alone Groups and Two or More Races) by Age, Sex, and Hispanic Origin: April 1, 2010 to July 1, 2018.

Table 8.4. Individuals with primary diagnosis of trauma- and stressor-related disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Total individuals with primary diagnosis of trauma- and stressor-related disorders	645,568	660,241	637,555	664,355	734,330	793,054
Census region¹						
Northeast	134,983	119,514	64,637	88,726	105,106	106,028
Midwest	132,067	174,299	180,344	164,715	189,178	226,145
South	159,658	150,914	164,523	159,707	178,987	187,464
West	218,853	215,507	228,041	251,192	260,950	273,304
Census division¹						
New England	40,714	24,921	19,819	20,182	29,443	24,497
Middle Atlantic	94,269	94,593	44,818	68,544	75,663	81,531
East North Central	89,849	95,862	98,512	90,253	120,603	127,390
West North Central	42,218	78,437	81,832	74,462	68,575	98,755
South Atlantic	94,050	88,473	98,002	85,520	103,815	105,152
East South Central	42,045	39,348	42,574	42,822	40,215	45,334
West South Central	23,563	23,093	23,947	31,365	34,957	36,978
Mountain	86,277	85,790	86,072	96,252	99,751	96,741
Pacific	132,576	129,717	141,969	154,940	161,199	176,563
State or jurisdiction						
Alabama	6,252	5,816	7,441	7,155	7,554	8,158
Alaska	‡	‡	‡	‡	‡	‡
Arizona	28,555	24,135	23,991	27,951	11,691	12,659
Arkansas	7,906	7,656	8,591	10,346	11,029	10,359
California	77,098	71,965	72,517	68,428	66,218	75,912
Colorado	19,974	22,576	19,246	21,484	32,917	36,755
Connecticut	12,751	13,467	8,930	9,590	13,849	13,690
Delaware	937	1,125	1,052	682	938	612
District of Columbia	‡	1,212	1,642	1,736	2,651	2,880
Florida	26,189	18,589	21,424	24,090	31,330	26,900
Georgia	‡	‡	‡	‡	‡	‡
Hawaii	2,001	2,320	2,211	1,603	1,947	2,401
Idaho	237	92	53	249	331	291
Illinois	15,368	15,659	14,715	1,807	8,831	8,809

Continued. See notes at end of table.

Table 8.4. Individuals with primary diagnosis of trauma- and stressor-related disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Indiana	11,882	11,914	12,312	10,419	11,186	11,718
Iowa	17,196	19,429	22,973	14,460	6,670	29,171
Kansas	11,635	6,155	‡	‡	‡	‡
Kentucky	20,730	20,240	20,797	21,711	21,059	22,995
Louisiana	1,131	1,365	1,374	967	1,500	1,127
Maine	15,961	‡	‡	‡	3,886	‡
Maryland	23,683	25,052	27,598	28,087	31,136	33,567
Massachusetts	2,523	2,659	2,833	1,425	1,776	712
Michigan	‡	‡	‡	‡	12,110	18,469
Minnesota	‡	39,250	44,738	51,213	52,900	59,657
Mississippi	7,584	5,024	6,776	6,158	5,742	7,577
Missouri	6,209	6,154	6,491	1,846	2,510	3,028
Montana	6,128	6,625	7,543	5,906	8,789	9,788
Nebraska	2,371	2,603	2,487	2,260	2,358	2,462
Nevada	2,197	2,060	2,245	1,505	1,725	1,681
New Hampshire	‡	‡	‡	‡	‡	‡
New Jersey	41,080	40,255	38,639	3,442	‡	‡
New Mexico	21,370	21,759	23,462	29,202	33,836	25,568
New York ²	5,450	5,880	6,179	6,945	7,316	7,579
North Carolina	25,936	24,040	26,502	10,791	15,541	11,858
North Dakota	1,479	1,380	1,361	868	595	505
Ohio	51,257	56,696	63,277	69,878	81,945	81,609
Oklahoma	7,770	8,438	8,231	11,446	12,600	14,598
Oregon	25,323	24,297	30,886	42,457	41,374	42,073
Pennsylvania	47,739	48,458	‡	58,157	68,347	73,952
Puerto Rico	‡	2	5	5	88	105

Continued. See notes at end of table.

Table 8.4. Individuals with primary diagnosis of trauma- and stressor-related disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Rhode Island	3,224	2,974	2,537	2,895	3,183	2,447
South Carolina	6,864	7,381	7,964	8,336	9,105	10,564
South Dakota	3,328	3,466	3,782	3,815	3,542	3,932
Tennessee ²	7,479	8,268	7,560	7,798	5,860	6,604
Texas	6,756	5,634	5,751	8,606	9,828	10,894
Utah	4,699	5,383	6,171	6,428	7,163	7,028
Vermont	6,255	5,821	5,519	6,272	6,749	7,648
Virginia	10,441	11,074	11,820	11,798	13,114	15,114
Washington	28,154	31,135	36,355	42,452	51,660	56,177
West Virginia	‡	‡	‡	‡	‡	3,657
Wisconsin	11,342	11,593	8,208	8,149	6,531	6,785
Wyoming	3,117	3,160	3,361	3,527	3,299	2,971
Other jurisdictions ³	7	5	5	10	21	8

‡ No data reported.

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

² New York and Tennessee report only a subset of the total individuals served through their SMHAs.

³ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 8.5. Individuals with primary diagnosis of anxiety disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Total individuals with primary diagnosis of anxiety disorders	376,408	409,035	427,333	493,641	579,070	654,028
Census region¹						
Northeast	72,870	63,863	32,599	55,329	67,685	70,454
Midwest	71,902	106,562	117,137	126,898	172,250	209,559
South	114,648	108,995	123,910	131,935	131,647	143,139
West	116,983	129,613	153,682	179,464	207,430	230,766
Census division¹						
New England	23,719	12,013	9,648	11,814	16,393	13,220
Middle Atlantic	49,151	51,850	22,951	43,515	51,292	57,234
East North Central	47,967	55,038	61,095	60,690	94,889	99,449
West North Central	23,935	51,524	56,042	66,208	77,361	110,110
South Atlantic	61,891	61,140	73,230	64,215	71,941	80,758
East South Central	34,214	32,207	34,234	45,913	34,786	37,755
West South Central	18,543	15,648	16,446	21,807	24,920	24,626
Mountain	48,116	48,321	57,069	67,352	84,082	94,375
Pacific	68,867	81,292	96,613	112,112	123,348	136,391
State or jurisdiction						
Alabama	4,532	4,572	5,188	5,683	5,782	5,900
Alaska	‡	‡	‡	‡	‡	‡
Arizona	19,483	15,762	16,751	22,735	24,187	29,586
Arkansas	6,006	6,133	6,724	7,938	9,121	7,699
California	39,751	49,420	53,060	51,854	44,488	54,673
Colorado	8,259	10,218	13,739	13,318	17,711	18,175
Connecticut	7,499	6,910	5,123	6,782	7,852	8,023
Delaware	374	357	233	127	216	205
District of Columbia	‡	497	592	585	1,147	1,263
Florida	12,925	9,593	11,283	12,984	12,942	14,027
Georgia	‡	‡	‡	‡	‡	‡
Hawaii	249	245	205	239	226	275
Idaho	80	29	15	357	400	383
Illinois	8,102	8,930	9,513	1,397	5,786	5,655

Continued. See notes at end of table.

Table 8.5. Individuals with primary diagnosis of anxiety disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Indiana	8,754	9,046	9,756	11,258	12,857	13,401
Iowa	7,863	9,388	11,739	10,573	5,592	35,102
Kansas	7,951	5,137	‡	‡	‡	‡
Kentucky	14,064	14,433	15,495	19,823	17,650	17,814
Louisiana	1,650	1,630	1,812	1,506	1,896	1,729
Maine	11,191	‡	‡	‡	2,881	‡
Maryland	13,115	15,541	20,474	23,610	27,138	29,107
Massachusetts	851	1,035	1,162	662	971	827
Michigan	‡	‡	‡	‡	9,657	13,515
Minnesota	‡	28,571	34,963	43,596	58,572	61,843
Mississippi	5,562	3,306	5,460	12,746	4,384	6,237
Missouri	4,540	4,546	5,258	8,101	9,169	9,068
Montana	4,176	4,613	5,267	4,408	8,963	13,719
Nebraska	1,062	1,150	1,203	1,533	1,741	1,760
Nevada	1,670	1,668	1,604	1,571	1,619	1,589
New Hampshire	‡	‡	‡	‡	‡	‡
New Jersey	19,535	19,695	19,274	1,179	‡	‡
New Mexico	6,725	7,480	10,884	15,280	20,191	20,189
New York ²	2,234	2,413	3,677	4,536	3,409	3,271
North Carolina	20,420	18,684	22,955	8,663	11,305	8,114
North Dakota	1,572	1,678	1,705	1,189	922	776
Ohio	23,972	29,221	36,066	40,529	57,230	56,369
Oklahoma	8,088	5,104	4,913	8,082	9,392	9,960
Oregon	9,678	8,060	13,826	18,366	27,102	25,243
Pennsylvania	27,382	29,742	‡	37,800	47,883	53,963
Puerto Rico	‡	1	3	3	49	101

Continued. See notes at end of table.

Table 8.5. Individuals with primary diagnosis of anxiety disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Rhode Island	2,031	1,907	1,437	2,039	2,182	1,671
South Carolina	4,160	4,411	4,955	5,336	5,650	6,100
South Dakota	947	1,054	1,174	1,216	1,365	1,561
Tennessee ²	10,056	9,896	8,091	7,661	6,970	7,804
Texas	2,799	2,781	2,997	4,281	4,511	5,238
Utah	6,030	6,662	6,693	7,406	8,549	8,436
Vermont	2,147	2,161	1,926	2,331	2,507	2,699
Virginia	10,897	12,057	12,738	12,910	13,543	14,530
Washington	19,189	23,567	29,522	41,653	51,532	56,200
West Virginia	‡	‡	‡	‡	‡	7,412
Wisconsin	7,139	7,841	5,760	7,506	9,359	10,509
Wyoming	1,693	1,889	2,116	2,277	2,462	2,298
Other jurisdictions ³	5	1	2	12	9	9

‡ No data reported.

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

² New York and Tennessee report only a subset of the total individuals served through their SMHAs.

³ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 8.6. Individuals with primary diagnosis of attention deficit/hyperactivity disorder, by Census region, Census division, and state or jurisdiction: Number, 2013–2018

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Total individuals with primary diagnosis of attention deficit/hyperactivity disorder	442,351	420,745	351,786	406,362	429,499	461,444
Census region¹						
Northeast	76,619	75,149	18,040	70,822	74,985	71,608
Midwest	99,389	116,749	114,966	106,276	120,435	138,659
South	165,982	143,982	145,159	144,532	145,282	150,543
West	100,352	84,830	73,561	84,661	88,528	100,431
Census division¹						
New England	13,254	6,782	5,300	5,186	10,082	9,518
Middle Atlantic	63,365	68,367	12,740	65,636	64,903	62,090
East North Central	65,618	67,789	69,688	62,101	81,224	80,151
West North Central	33,771	48,960	45,278	44,175	39,211	58,508
South Atlantic	84,459	72,175	73,900	64,669	69,810	71,951
East South Central	48,704	40,581	47,008	45,807	39,765	43,578
West South Central	32,819	31,226	24,251	34,056	35,707	35,014
Mountain	39,538	35,474	34,098	36,407	40,767	42,948
Pacific	60,814	49,356	39,463	48,254	47,761	57,483
State or jurisdiction						
Alabama	12,546	12,227	13,299	12,736	13,059	13,669
Alaska	‡	‡	‡	‡	‡	‡
Arizona	20,780	16,395	15,086	17,925	18,596	20,587
Arkansas	5,313	3,670	3,103	2,704	2,480	1,075
California	38,514	23,568	22,677	21,051	27,333	28,971
Colorado	6,286	6,852	6,491	5,653	7,393	7,276
Connecticut	2,888	3,149	2,134	1,968	6,327	6,462
Delaware	570	487	209	230	367	254
District of Columbia	‡	1,255	1,473	1,586	1,945	2,399
Florida	22,589	12,187	12,731	12,562	13,859	13,081
Georgia	‡	‡	‡	‡	‡	‡
Hawaii	362	339	361	382	336	395
Idaho	103	64	26	174	182	218
Illinois	11,749	11,367	10,858	983	5,799	5,127

Continued. See notes at end of table.

Table 8.6. Individuals with primary diagnosis of attention deficit/hyperactivity disorder, by Census region, Census division, and state or jurisdiction: Number, 2013–2018 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Indiana	13,894	14,060	14,549	16,154	14,908	14,142
Iowa	14,531	14,573	15,119	11,309	4,883	25,669
Kansas	10,816	6,004	‡	‡	‡	‡
Kentucky	17,061	16,176	15,424	14,469	13,618	13,405
Louisiana	3,432	3,522	3,282	2,045	2,188	1,993
Maine	6,652	‡	‡	‡	354	‡
Maryland	21,017	22,585	24,326	26,093	27,773	28,739
Massachusetts	768	850	909	245	400	318
Michigan	‡	‡	‡	‡	9,150	12,034
Minnesota	‡	20,054	21,940	24,463	25,165	23,686
Mississippi	13,292	7,061	13,535	13,398	10,599	13,776
Missouri	5,657	5,503	5,511	5,898	6,835	6,764
Montana	2,593	2,689	2,659	2,093	3,364	4,220
Nebraska	641	728	590	614	621	721
Nevada	466	442	478	427	425	367
New Hampshire	‡	‡	‡	‡	‡	‡
New Jersey	11,668	11,559	11,016	3,287	‡	‡
New Mexico	7,062	6,632	6,819	7,191	7,835	7,559
New York ²	1,486	1,557	1,724	2,074	2,004	1,870
North Carolina	20,459	15,393	14,810	2,601	4,227	2,439
North Dakota	893	841	832	640	497	443
Ohio	36,461	38,717	41,678	42,247	48,854	46,037
Oklahoma	2,512	2,655	2,302	3,506	3,604	3,580
Oregon	7,524	10,203	258	8,900	‡	7,672
Pennsylvania	50,211	55,251	‡	60,275	62,899	60,220
Puerto Rico	‡	31	58	57	262	193

Continued. See notes at end of table.

Table 8.6. Individuals with primary diagnosis of attention deficit/hyperactivity disorder, by Census region, Census division, and state or jurisdiction: Number, 2013–2018 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Rhode Island	1,470	1,325	994	1,419	1,461	1,223
South Carolina	9,814	9,830	10,122	10,242	10,216	10,116
South Dakota	1,233	1,257	1,286	1,251	1,210	1,225
Tennessee ²	5,805	5,117	4,750	5,204	2,489	2,728
Texas	21,562	21,379	15,564	25,801	27,435	28,366
Utah	1,262	1,320	1,455	1,952	2,016	1,859
Vermont	1,476	1,458	1,263	1,554	1,540	1,515
Virginia	10,010	10,438	10,229	11,355	11,423	11,621
Washington	14,414	15,246	16,167	17,921	20,092	20,445
West Virginia	‡	‡	‡	‡	‡	3,302
Wisconsin	3,514	3,645	2,603	2,717	2,513	2,811
Wyoming	986	1,080	1,084	992	956	862
Other jurisdictions ³	9	4	2	14	7	10

‡ No data reported.

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

² New York and Tennessee report only a subset of the total individuals served through their SMHAs.

³ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 8.7. Individuals with primary diagnosis of bipolar disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Total individuals with primary diagnosis of bipolar disorders	854,076	880,161	758,549	663,415	607,293	601,023
Census region¹						
Northeast	122,567	131,966	57,879	99,714	93,464	72,218
Midwest	159,424	177,303	165,296	125,876	142,698	154,566
South	342,927	339,810	312,193	247,676	236,170	234,027
West	229,150	231,076	223,172	190,128	134,853	140,028
Census division¹						
New England	37,509	27,805	24,428	16,232	18,935	13,515
Middle Atlantic	85,058	104,161	33,451	83,482	74,529	58,703
East North Central	110,296	116,640	114,928	77,556	102,309	108,921
West North Central	49,128	60,663	50,368	48,320	40,389	45,645
South Atlantic	157,951	154,383	148,375	102,102	103,886	103,678
East South Central	77,330	74,470	76,516	54,481	42,462	43,801
West South Central	107,646	110,957	87,302	91,093	89,822	86,548
Mountain	98,952	92,380	80,327	63,655	62,738	62,366
Pacific	130,198	138,696	142,845	126,473	72,115	77,662
State or jurisdiction						
Alabama	16,224	15,817	16,897	12,250	12,089	11,831
Alaska	‡	‡	‡	‡	‡	‡
Arizona	46,004	38,731	36,936	31,853	26,306	28,143
Arkansas	12,317	12,356	12,520	7,074	7,029	6,044
California	100,515	110,464	109,058	95,803	42,971	49,619
Colorado	20,990	22,503	13,568	11,926	15,826	15,560
Connecticut	14,137	14,902	12,126	10,418	8,659	7,914
Delaware	1,719	1,875	1,913	1,199	1,253	1,490
District of Columbia	‡	3,719	4,272	4,307	5,117	6,155
Florida	52,256	38,727	39,892	32,613	25,588	26,120
Georgia	‡	‡	‡	‡	‡	‡
Hawaii	1,746	1,658	1,280	1,022	1,136	1,032
Idaho	980	523	389	1,507	1,540	1,536
Illinois	29,321	30,002	27,885	2,341	11,339	10,132

Continued. See notes at end of table.

Table 8.7. Individuals with primary diagnosis of bipolar disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Indiana	18,558	18,516	18,821	13,291	14,437	13,813
Iowa	7,397	8,405	9,268	13,588	8,110	11,833
Kansas	14,603	9,719	‡	‡	‡	‡
Kentucky	26,098	25,722	26,046	18,919	11,029	11,021
Louisiana	9,275	10,245	9,176	4,519	4,233	3,559
Maine	9,903	‡	‡	‡	3,829	‡
Maryland	31,847	33,179	29,887	23,719	24,846	26,331
Massachusetts	6,482	6,894	7,383	1,543	2,062	1,656
Michigan	‡	‡	‡	‡	22,555	28,423
Minnesota	‡	16,082	15,949	14,357	11,982	15,009
Mississippi	11,876	6,142	11,155	9,840	6,692	8,855
Missouri	18,144	17,578	17,094	14,021	14,105	12,909
Montana	4,836	4,750	4,722	3,529	3,692	3,759
Nebraska	5,545	5,399	4,641	3,541	3,659	3,479
Nevada	7,091	6,758	6,226	1,128	1,126	1,008
New Hampshire	‡	‡	‡	‡	‡	‡
New Jersey	27,199	26,632	25,784	487	‡	‡
New Mexico	9,607	9,782	9,170	8,172	8,656	6,705
New York ²	9,519	9,523	7,667	6,327	5,005	5,166
North Carolina	42,350	46,274	41,700	16,653	23,957	14,038
North Dakota	1,654	1,660	1,534	1,241	1,057	925
Ohio	47,037	52,876	57,789	52,505	48,193	50,839
Oklahoma	12,688	15,956	13,087	12,296	11,417	10,429
Oregon	8,004	5,299	9,757	9,926	9,182	8,850
Pennsylvania	48,340	68,006	‡	76,668	69,524	53,537
Puerto Rico	‡	‡	3	3	93	180

Continued. See notes at end of table.

Table 8.7. Individuals with primary diagnosis of bipolar disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Rhode Island	5,270	4,455	3,513	3,223	3,305	2,860
South Carolina	12,133	12,239	12,204	10,534	10,037	10,409
South Dakota	1,785	1,820	1,882	1,572	1,476	1,490
Tennessee ²	23,132	26,789	22,418	13,472	12,652	12,094
Texas	73,366	72,400	52,519	67,204	67,143	66,516
Utah	7,340	7,170	7,139	3,631	4,048	4,168
Vermont	1,717	1,554	1,406	1,048	1,080	1,085
Virginia	17,646	18,370	18,507	13,077	13,088	13,368
Washington	19,933	21,275	22,750	19,722	18,826	18,161
West Virginia	‡	‡	‡	‡	‡	5,767
Wisconsin	15,380	15,246	10,433	9,419	5,785	5,714
Wyoming	2,104	2,163	2,177	1,909	1,544	1,487
Other jurisdictions ³	8	6	6	18	15	4

‡ No data reported.

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

² New York and Tennessee report only a subset of the total individuals served through their SMHAs.

³ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 8.8. Individuals with primary diagnosis of primary depressive disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Total individuals with primary diagnosis of depressive disorders	1,267,441	1,327,354	1,255,030	1,257,761	1,315,673	1,410,277
Census region¹						
Northeast	213,712	212,492	101,106	166,424	183,083	168,294
Midwest	249,411	310,222	305,749	282,273	314,844	376,047
South	451,602	447,288	460,152	442,732	453,340	468,261
West	352,694	357,332	387,999	366,297	363,998	396,941
Census division¹						
New England	50,944	35,030	30,748	30,945	39,976	31,334
Middle Atlantic	162,768	177,462	70,358	135,479	143,107	136,960
East North Central	182,318	195,696	196,289	157,652	213,221	240,523
West North Central	67,093	114,526	109,460	124,621	101,623	135,524
South Atlantic	211,234	207,054	228,600	191,189	203,394	211,769
East South Central	117,038	110,968	119,513	110,199	99,751	105,401
West South Central	123,330	129,266	112,039	141,344	150,195	151,091
Mountain	117,701	111,399	117,687	116,870	135,030	139,486
Pacific	234,993	245,933	270,312	249,427	228,968	257,455
State or jurisdiction						
Alabama	27,650	26,308	28,054	28,567	27,550	27,591
Alaska	‡	‡	‡	‡	‡	‡
Arizona	42,082	32,908	33,191	38,082	37,269	42,320
Arkansas	16,526	16,260	17,547	20,324	20,363	18,416
California	164,907	169,118	171,277	148,454	128,608	160,164
Colorado	25,751	28,983	31,406	29,202	38,811	39,107
Connecticut	18,905	20,203	16,904	17,090	18,326	18,344
Delaware	1,723	1,884	1,637	1,241	1,447	1,925
District of Columbia	‡	9,945	11,014	10,699	10,148	12,846
Florida	69,401	49,238	53,391	55,125	50,328	53,057
Georgia	‡	‡	‡	‡	‡	‡
Hawaii	2,914	2,887	2,113	2,106	1,869	1,557
Idaho	875	394	262	1,982	2,206	2,271
Illinois	44,252	43,297	40,118	2,809	19,959	18,504

Continued. See notes at end of table.

Table 8.8. Individuals with primary diagnosis of primary depressive disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Indiana	41,804	43,831	46,369	42,521	46,902	47,509
Iowa	13,909	16,762	20,918	36,476	19,808	43,078
Kansas	18,666	12,202	‡	‡	‡	‡
Kentucky	39,189	38,837	39,373	37,459	34,359	35,319
Louisiana	14,970	16,406	14,395	7,409	9,097	8,596
Maine	16,265	‡	‡	‡	6,220	‡
Maryland	38,820	42,894	50,617	48,561	52,952	56,339
Massachusetts	4,172	4,929	5,077	2,054	2,916	1,861
Michigan	‡	‡	‡	‡	38,198	53,417
Minnesota	‡	50,858	54,562	56,208	48,280	58,737
Mississippi	22,156	11,807	21,650	18,724	14,937	20,203
Missouri	21,255	21,222	20,944	20,353	22,256	22,356
Montana	9,593	9,906	10,367	7,375	10,845	14,071
Nebraska	6,195	6,334	5,450	5,296	5,659	5,700
Nevada	7,442	7,446	6,709	2,060	2,185	2,022
New Hampshire	‡	‡	‡	‡	‡	‡
New Jersey	62,772	61,212	60,830	1,652	‡	‡
New Mexico	15,857	15,480	18,636	21,178	25,067	20,739
New York ²	8,687	8,795	9,528	9,681	7,431	7,603
North Carolina	60,067	61,116	67,649	32,080	44,421	29,505
North Dakota	3,674	3,628	3,729	2,821	2,144	1,853
Ohio	72,123	84,345	95,157	98,206	97,177	109,867
Oklahoma	20,665	24,389	20,980	29,046	31,174	32,048
Oregon	17,247	16,212	25,331	25,472	26,745	25,810
Pennsylvania	91,309	107,455	‡	124,146	135,676	129,357
Puerto Rico	‡	5	14	14	388	718

Continued. See notes at end of table.

Table 8.8. Individuals with primary diagnosis of primary depressive disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Rhode Island	7,094	5,894	5,282	8,499	9,112	7,445
South Carolina	18,479	19,115	20,400	20,039	19,630	20,051
South Dakota	3,394	3,520	3,857	3,467	3,476	3,800
Tennessee ²	28,043	34,016	30,436	25,449	22,905	22,288
Texas	71,169	72,211	59,117	84,565	89,561	92,031
Utah	11,317	11,303	12,017	11,868	13,891	14,276
Vermont	4,508	4,004	3,485	3,302	3,402	3,684
Virginia	22,744	22,862	23,892	23,444	24,468	25,589
Washington	49,925	57,716	71,591	73,395	71,746	69,924
West Virginia	‡	‡	‡	‡	‡	12,457
Wisconsin	24,139	24,223	14,645	14,116	10,985	11,226
Wyoming	4,784	4,979	5,099	5,123	4,756	4,680
Other jurisdictions ³	22	15	10	21	20	16

‡ No data reported.

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

² New York and Tennessee report only a subset of the total individuals served through their SMHAs.

³ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 8.9. Individuals with primary diagnosis of primary schizophrenia and other psychotic disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Total individuals with primary diagnosis of schizophrenia and other psychotic disorders	630,500	656,286	593,824	576,500	626,384	660,521
Census region¹						
Northeast	99,540	109,978	74,026	76,457	80,001	76,430
Midwest	110,085	125,378	110,748	98,271	136,698	150,902
South	227,088	221,255	210,420	211,638	218,030	222,488
West	193,763	199,649	198,608	190,028	191,279	209,919
Census division¹						
New England	29,304	25,349	24,519	19,087	21,339	18,893
Middle Atlantic	70,236	84,629	49,507	57,370	58,662	57,537
East North Central	77,553	79,968	71,580	57,242	98,038	104,768
West North Central	32,532	45,410	39,168	41,029	38,660	46,134
South Atlantic	108,672	103,657	106,327	93,694	101,698	106,124
East South Central	46,823	44,413	49,929	45,820	42,707	45,634
West South Central	71,593	73,185	54,164	72,124	73,625	70,730
Mountain	50,404	51,287	47,025	46,035	52,328	54,083
Pacific	143,359	148,362	151,583	143,993	138,951	155,836
State or jurisdiction						
Alabama	17,127	17,275	17,529	17,315	17,546	17,752
Alaska	‡	‡	‡	‡	‡	‡
Arizona	19,382	18,483	18,155	18,789	19,727	21,951
Arkansas	8,332	8,390	8,307	8,418	8,526	6,758
California	112,502	119,375	118,726	109,781	104,008	121,780
Colorado	11,470	12,460	9,587	10,379	12,911	12,703
Connecticut	11,360	11,226	10,790	10,869	10,984	10,517
Delaware	1,571	1,825	1,833	1,226	1,184	1,683
District of Columbia	‡	4,896	5,332	5,183	5,752	6,670
Florida	41,224	29,048	30,180	29,169	29,624	32,965
Georgia	‡	‡	‡	‡	‡	‡
Hawaii	3,595	3,500	2,872	3,139	2,866	2,719
Idaho	671	423	272	1,817	1,961	1,985
Illinois	20,986	21,067	17,832	3,304	11,241	10,409

Continued. See notes at end of table.

Table 8.9. Individuals with primary diagnosis of primary schizophrenia and other psychotic disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Indiana	14,486	14,735	14,704	14,599	15,442	14,899
Iowa	4,586	4,965	5,229	8,045	4,796	10,473
Kansas	7,724	5,804	‡	‡	‡	‡
Kentucky	11,370	11,396	11,300	10,440	10,171	10,367
Louisiana	8,355	8,933	7,956	4,972	4,962	4,631
Maine	3,563	‡	‡	‡	1,301	‡
Maryland	13,870	14,479	15,041	15,451	15,869	16,323
Massachusetts	10,360	10,842	10,934	2,877	3,721	3,322
Michigan	‡	‡	‡	‡	25,726	30,743
Minnesota	‡	14,810	14,955	14,932	15,314	17,509
Mississippi	10,682	4,175	11,920	10,439	7,565	10,365
Missouri	13,745	13,471	13,419	13,138	13,563	13,187
Montana	3,116	3,128	3,091	2,407	2,804	3,370
Nebraska	3,275	3,145	2,199	2,002	2,230	2,264
Nevada	5,232	5,253	5,510	1,967	2,042	2,003
New Hampshire	‡	‡	‡	‡	‡	‡
New Jersey	30,622	30,106	29,744	112	‡	‡
New Mexico	4,787	5,669	4,566	5,139	6,610	5,557
New York ²	20,381	19,950	19,763	19,372	19,398	19,316
North Carolina	21,329	22,349	22,853	11,250	17,546	13,345
North Dakota	1,523	1,576	1,578	1,447	1,349	1,259
Ohio	26,975	29,170	28,773	29,290	37,068	39,583
Oklahoma	7,789	8,545	7,179	9,486	9,803	9,690
Oregon	9,076	6,921	10,314	11,282	11,794	11,182
Pennsylvania	19,233	34,573	‡	37,886	39,264	38,221
Puerto Rico	‡	2	8	7	289	670

Continued. See notes at end of table.

Table 8.9. Individuals with primary diagnosis of primary schizophrenia and other psychotic disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Rhode Island	2,376	2,031	1,656	3,757	3,730	3,414
South Carolina	15,026	15,231	15,051	14,787	14,778	15,733
South Dakota	1,679	1,639	1,788	1,465	1,408	1,442
Tennessee ²	7,644	11,567	9,180	7,626	7,425	7,150
Texas	47,117	47,317	30,722	49,248	50,334	49,651
Utah	4,950	4,930	4,883	4,574	5,336	5,575
Vermont	1,645	1,250	1,139	1,584	1,603	1,640
Virginia	15,652	15,829	16,037	16,628	16,945	17,143
Washington	18,186	18,566	19,671	19,791	20,283	20,155
West Virginia	‡	‡	‡	‡	‡	2,262
Wisconsin	15,106	14,996	10,271	10,049	8,561	9,134
Wyoming	796	941	961	963	937	939
Other jurisdictions ³	24	24	14	99	87	112

‡ No data reported.

¹ The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

² New York and Tennessee report only a subset of the total individuals served through their SMHAs.

³ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 8.10. Individuals with co-occurring mental health and substance use disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Total individuals with co-occurring mental health and substance use disorders¹	782,501	777,946	758,082	690,705	905,849	1,051,340
Census region²						
Northeast	104,773	113,018	104,243	85,971	79,863	97,988
Midwest	98,289	147,307	131,733	137,383	338,939	414,162
South	315,075	265,273	254,516	215,041	242,483	278,160
West	264,351	252,284	267,446	252,094	243,320	259,850
Census division²						
New England	45,555	56,382	52,255	36,151	34,630	40,327
Middle Atlantic	59,218	56,636	51,988	49,820	45,233	57,661
East North Central	51,695	53,204	42,125	42,627	246,006	277,646
West North Central	46,594	94,103	89,608	94,756	92,933	136,516
South Atlantic	165,863	133,025	141,468	99,404	120,543	147,113
East South Central	54,339	60,985	57,569	48,149	50,292	53,258
West South Central	94,873	71,263	55,479	67,488	71,648	77,789
Mountain	97,164	81,135	71,700	60,340	72,775	81,718
Pacific	167,187	171,149	195,746	191,754	170,545	178,132
State or jurisdiction						
Alabama	10,380	10,213	11,798	12,107	14,437	14,695
Alaska	‡	‡	‡	‡	‡	‡
Arizona	40,113	35,761	35,777	38,513	38,188	44,102
Arkansas	656	636	853	1,441	1,310	957
California	136,083	140,256	140,503	126,167	99,491	108,370
Colorado	22,046	20,852	14,886	1,527	383	376
Connecticut	6,355	40,981	36,944	26,826	24,740	25,035
Delaware	511	419	357	3,840	3,593	5,590
District of Columbia	‡	2,669	3,095	3,919	2,609	4,142
Florida	62,522	26,212	28,594	26,949	26,394	26,637
Georgia	‡	‡	‡	‡	‡	‡
Hawaii	2,402	2,887	2,232	2,276	2,785	3,045
Idaho	2,023	892	525	2,238	2,365	2,385
Illinois	18,659	19,135	17,033	1,594	5,492	5,101

Continued. See notes at end of table.

Table 8.10. Individuals with co-occurring mental health and substance use disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Indiana	19,968	18,321	18,243	21,488	23,079	22,827
Iowa	688	984	1,346	1,202	755	18,636
Kansas	10,944	8,606	‡	‡	‡	‡
Kentucky	23,392	23,621	25,226	20,393	22,326	23,286
Louisiana	8,529	203	968	1,356	3,188	3,127
Maine	22,981	‡	‡	‡	13	‡
Maryland	13,037	14,633	15,475	18,038	40,084	52,541
Massachusetts	6,679	7,118	7,559	2,413	2,948	9,204
Michigan	‡	‡	‡	‡	97,518	163,272
Minnesota	‡	36,682	40,789	44,995	46,771	80,278
Mississippi	4,045	‡	‡	‡	‡	‡
Missouri	15,599	28,797	30,058	33,034	33,522	31,182
Montana	127	149	211	3,942	9,466	18,488
Nebraska	10,658	10,380	9,170	8,338	5,495	87
Nevada	2,016	1,958	1,950	673	651	587
New Hampshire	‡	‡	‡	‡	‡	‡
New Jersey	32,408	30,706	29,387	4	‡	‡
New Mexico	14,538	8,271	7,378	10,822	15,331	9,197
New York ³	22,531	22,440	22,601	25,668	26,146	26,029
North Carolina	43,372	41,627	46,741	‡	‡	‡
North Dakota	5,699	5,644	5,430	4,327	3,585	3,415
Ohio	3,210	6,466	1,823	14,890	117,466	83,419
Oklahoma	19,155	8,424	7,016	8,672	8,700	9,493
Oregon	12,219	9,938	32,187	41,306	45,929	43,303
Pennsylvania	4,279	3,490	‡	24,148	19,087	31,632
Puerto Rico	‡	52	132	129	1,207	1,149

Continued. See notes at end of table.

Table 8.10. Individuals with co-occurring mental health and substance use disorders, by Census region, Census division, and state or jurisdiction: Number, 2013–2018 (continued)

<i>Census region, Census division, and state or jurisdiction</i>	2013	2014	2015	2016	2017	2018
Rhode Island	5,618	4,829	4,573	6,912	6,929	6,088
South Carolina	13,457	13,709	11,405	10,963	10,908	12,911
South Dakota	3,006	3,010	2,815	2,860	2,805	2,918
Tennessee ³	16,522	27,151	20,545	15,649	13,529	15,277
Texas	66,533	62,000	46,642	56,019	58,450	64,212
Utah	5,389	4,944	5,008	632	4,666	4,875
Vermont	3,922	3,454	3,179	‡	‡	‡
Virginia	32,964	33,756	35,801	35,695	36,955	40,198
Washington	16,483	18,068	20,824	22,005	22,340	23,414
West Virginia	‡	‡	‡	‡	‡	5,094
Wisconsin	9,858	9,282	5,026	4,655	2,451	3,027
Wyoming	10,912	8,308	5,965	1,993	1,725	1,708
Other jurisdictions ⁴	13	12	12	87	37	31

‡ No data reported.

¹ Total individuals with co-occurring mental health and substance use disorders who had at least one valid mental health diagnosis other than an alcohol- or substance-related disorder. Mental health diagnostic category definitions can be found in Appendix E.

² The Commonwealth of the Northern Mariana Islands, the Republic of Palau, and the Commonwealth of Puerto Rico are not included in any Census region or division.

³ New York and Tennessee report only a subset of the total individuals served through their SMHAs.

⁴ Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

NOTE: This table refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for the individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.1a. Alabama individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	97,362	94,474	101,582	104,320	103,959	102,424
Mental health diagnosis¹						
Trauma- and stressor-related disorders	6,252	5,816	7,441	7,155	7,554	8,158
Anxiety disorders	4,532	4,572	5,188	5,683	5,782	5,900
Attention deficit/hyperactivity disorder	12,546	12,227	13,299	12,736	13,059	13,669
Mood disorders	43,874	42,125	44,951	40,817	39,639	39,422
Bipolar disorders	16,224	15,817	16,897	12,250	12,089	11,831
Depressive disorders	27,650	26,308	28,054	28,567	27,550	27,591
Personality disorders	45	41	44	305	440	437
Schizophrenia and other psychotic disorders	17,127	17,275	17,529	17,315	17,546	17,752
Alcohol or substance-related disorders	1,763	1,426	2,471	14	6	7
Other mental disorders ²	9,084	10,601	10,382	10,375	14,187	16,254
Total	95,223	94,083	101,305	94,400	98,213	101,599
Gender						
Male	45,219	44,115	47,601	48,854	48,896	48,528
Female	52,001	50,101	53,748	55,338	55,045	53,888
Total	97,220	94,216	101,349	104,192	103,941	102,416
Age³						
0 to 11 years	13,456	13,441	15,025	15,238	15,662	16,028
12 to 14 years	7,054	7,023	7,519	7,664	7,907	8,417
15 to 17 years	7,659	7,774	8,478	8,535	8,801	8,728
18 to 20 years	3,989	3,642	3,999	4,232	4,393	4,459
21 to 24 years	4,918	4,591	5,052	5,043	4,752	4,554
25 to 29 years	6,673	6,303	7,022	7,464	7,417	7,162
30 to 34 years	7,341	6,809	7,356	7,541	7,305	7,143
35 to 39 years	6,698	6,629	7,245	7,836	7,673	7,528
40 to 44 years	7,318	6,799	7,120	7,153	6,901	6,579
45 to 49 years	7,672	7,043	7,386	7,492	7,233	6,867
50 to 54 years	8,144	7,749	7,958	8,110	7,708	7,110
55 to 59 years	6,580	6,726	7,008	7,211	7,149	6,951
60 to 64 years	4,117	4,251	4,518	4,770	5,012	5,116
65 years and older	5,431	5,328	5,669	5,839	5,726	5,774
Total	97,050	94,108	101,355	104,128	103,639	102,416

Continued. See notes at end of table.

Table 9.1a. Alabama individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	273	276	306	290	293	294
Asian ⁴	163	166	175	190	220	211
Black or African American	35,274	33,907	36,644	38,864	39,054	38,046
Native Hawaiian or Other Pacific Islander	26	26	28	36	51	58
White	57,103	55,232	59,412	59,892	59,477	56,220
Other	3,116	3,101	3,864	4,357	4,848	6,267
Total	95,955	92,708	100,429	103,629	103,943	101,096
Ethnicity						
Not of Hispanic or Latino origin	95,363	92,363	99,175	101,321	101,138	99,687
Hispanic or Latino origin	1,499	1,502	1,870	2,662	2,767	2,729
Total	96,862	93,865	101,045	103,983	103,905	102,416
Specific Hispanic or Latino origin						
Mexican	629	685	738	1,303	1,151	1,098
Puerto Rican	175	171	213	223	266	368
Cuban	42	32	36	35	32	46
Other specific	620	599	862	1,076	1,294	1,200
Total	1,466	1,487	1,849	2,637	2,743	2,712
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	1,209	1,000
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	62,342	54,256
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	63,551	55,256

n/c Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.1b. Alabama individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	6.6	6.2	7.3	7.6	7.7	8.0
Anxiety disorders	4.8	4.9	5.1	6.0	5.9	5.8
Attention deficit/hyperactivity disorder	13.2	13.0	13.1	13.5	13.3	13.5
Mood disorders	46.1	44.8	44.4	43.2	40.4	38.8
Bipolar disorders	17.0	16.8	16.7	13.0	12.3	11.6
Depressive disorders	29.0	28.0	27.7	30.3	28.1	27.2
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	18.0	18.4	17.3	18.3	17.9	17.5
Alcohol or substance-related disorders	1.9	1.5	2.4	†	†	†
Other mental disorders ²	9.5	11.3	10.2	11.0	14.4	16.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	46.5	46.8	47.0	46.9	47.0	47.4
Female	53.5	53.2	53.0	53.1	53.0	52.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	13.9	14.3	14.8	14.6	15.1	15.6
12 to 14 years	7.3	7.5	7.4	7.4	7.6	8.2
15 to 17 years	7.9	8.3	8.4	8.2	8.5	8.5
18 to 20 years	4.1	3.9	3.9	4.1	4.2	4.4
21 to 24 years	5.1	4.9	5.0	4.8	4.6	4.4
25 to 29 years	6.9	6.7	6.9	7.2	7.2	7.0
30 to 34 years	7.6	7.2	7.3	7.2	7.0	7.0
35 to 39 years	6.9	7.0	7.1	7.5	7.4	7.4
40 to 44 years	7.5	7.2	7.0	6.9	6.7	6.4
45 to 49 years	7.9	7.5	7.3	7.2	7.0	6.7
50 to 54 years	8.4	8.2	7.9	7.8	7.4	6.9
55 to 59 years	6.8	7.1	6.9	6.9	6.9	6.8
60 to 64 years	4.2	4.5	4.5	4.6	4.8	5.0
65 years and older	5.6	5.7	5.6	5.6	5.5	5.6
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.1b. Alabama individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	36.8	36.6	36.5	37.5	37.6	37.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	59.5	59.6	59.2	57.8	57.2	55.6
Other	3.2	3.3	3.8	4.2	4.7	6.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	98.5	98.4	98.1	97.4	97.3	97.3
Hispanic or Latino origin	1.5	1.6	1.9	2.6	2.7	2.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	42.9	46.1	39.9	49.4	42.0	40.5
Puerto Rican	11.9	11.5	11.5	8.5	9.7	13.6
Cuban	2.9	2.2	1.9	1.3	1.2	1.7
Other specific	42.3	40.3	46.6	40.8	47.2	44.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	1.9	1.8
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	98.1	98.2
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual’s age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.2a. Arizona individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	191,093	158,173	155,122	187,972	196,212	224,773
Mental health diagnosis¹						
Trauma- and stressor-related disorders	28,555	24,135	23,991	27,951	11,691	12,659
Anxiety disorders	19,483	15,762	16,751	22,735	24,187	29,586
Attention deficit/hyperactivity disorder	20,780	16,395	15,086	17,925	18,596	20,587
Mood disorders	88,086	71,639	70,127	69,935	63,575	70,463
Bipolar disorders	46,004	38,731	36,936	31,853	26,306	28,143
Depressive disorders	42,082	32,908	33,191	38,082	37,269	42,320
Personality disorders	566	395	375	769	1,001	1,319
Schizophrenia and other psychotic disorders	19,382	18,483	18,155	18,789	19,727	21,951
Alcohol or substance-related disorders	2	4	6	2,367	1	1
Other mental disorders ²	14,201	11,334	10,609	15,352	11,639	13,295
Total	191,055	158,147	155,100	175,823	150,417	169,861
Gender						
Male	89,088	74,268	73,212	88,531	93,785	108,015
Female	102,005	83,905	81,910	99,441	102,417	116,758
Total	191,093	158,173	155,122	187,972	196,202	224,773
Age³						
0 to 11 years	34,762	29,853	26,843	37,380	42,808	47,293
12 to 14 years	15,428	13,276	12,685	14,955	16,084	18,761
15 to 17 years	14,703	12,708	12,090	14,602	15,576	18,143
18 to 20 years	9,297	6,334	5,719	7,949	7,767	9,330
21 to 24 years	9,914	7,263	7,493	9,182	10,021	11,770
25 to 29 years	14,208	11,143	11,155	13,946	15,361	18,437
30 to 34 years	14,831	11,789	11,598	14,021	14,865	17,631
35 to 39 years	13,217	10,765	10,871	13,119	13,841	16,014
40 to 44 years	13,619	10,977	10,635	11,838	11,810	13,318
45 to 49 years	13,413	10,882	11,002	12,125	12,059	13,521
50 to 54 years	13,968	12,009	12,250	13,178	12,477	13,712
55 to 59 years	11,232	9,930	10,458	12,026	11,606	13,028
60 to 64 years	6,915	6,310	6,902	7,869	7,732	8,942
65 years and older	5,586	4,933	5,419	5,782	4,204	4,869
Total	191,093	158,172	155,120	187,972	196,211	224,769

Continued. See notes at end of table.

Table 9.2a. Arizona individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	6,342	6,225	5,473	7,246	6,161	6,615
Asian ⁴	1,234	1,106	1,054	1,333	1,355	1,647
Black or African American	13,039	11,448	10,650	13,073	14,193	17,455
Native Hawaiian or Other Pacific Islander	460	398	385	430	470	587
White	147,091	132,079	128,930	157,567	165,711	193,386
Other	2,718	2,302	2,272	3,351	4,593	4,557
Total	170,884	153,558	148,764	183,000	192,483	224,247
Ethnicity						
Not of Hispanic or Latino origin	118,344	107,469	104,917	132,345	135,954	155,405
Hispanic or Latino origin	52,495	45,988	43,732	50,420	56,460	68,767
Total	170,839	153,457	148,649	182,765	192,414	224,172
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2,200	175
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	106,671	8,465
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	108,871	8,640

‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.2b. Arizona individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	14.9	15.3	15.5	15.9	7.8	7.5
Anxiety disorders	10.2	10.0	10.8	12.9	16.1	17.4
Attention deficit/hyperactivity disorder	10.9	10.4	9.7	10.2	12.4	12.1
Mood disorders	46.1	45.3	45.2	39.8	42.3	41.5
Bipolar disorders	24.1	24.5	23.8	18.1	17.5	16.6
Depressive disorders	22.0	20.8	21.4	21.7	24.8	24.9
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	10.1	11.7	11.7	10.7	13.1	12.9
Alcohol or substance-related disorders	†	†	†	1.3	†	†
Other mental disorders ²	7.4	7.2	6.8	8.7	7.7	7.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	46.6	47.0	47.2	47.1	47.8	48.1
Female	53.4	53.0	52.8	52.9	52.2	51.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	18.2	18.9	17.3	19.9	21.8	21.0
12 to 14 years	8.1	8.4	8.2	8.0	8.2	8.3
15 to 17 years	7.7	8.0	7.8	7.8	7.9	8.1
18 to 20 years	4.9	4.0	3.7	4.2	4.0	4.2
21 to 24 years	5.2	4.6	4.8	4.9	5.1	5.2
25 to 29 years	7.4	7.0	7.2	7.4	7.8	8.2
30 to 34 years	7.8	7.5	7.5	7.5	7.6	7.8
35 to 39 years	6.9	6.8	7.0	7.0	7.1	7.1
40 to 44 years	7.1	6.9	6.9	6.3	6.0	5.9
45 to 49 years	7.0	6.9	7.1	6.5	6.1	6.0
50 to 54 years	7.3	7.6	7.9	7.0	6.4	6.1
55 to 59 years	5.9	6.3	6.7	6.4	5.9	5.8
60 to 64 years	3.6	4.0	4.4	4.2	3.9	4.0
65 years and older	2.9	3.1	3.5	3.1	2.1	2.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.2b. Arizona individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	3.7	4.1	3.7	4.0	3.2	2.9
Asian ⁴	†	†	†	†	†	†
Black or African American	7.6	7.5	7.2	7.1	7.4	7.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	86.1	86.0	86.7	86.1	86.1	86.2
Other	1.6	1.5	1.5	1.8	2.4	2.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	69.3	70.0	70.6	72.4	70.7	69.3
Hispanic or Latino origin	30.7	30.0	29.4	27.6	29.3	30.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2.0	2.0
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	98.0	98.0
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected;

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual’s age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.3a. Arkansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	69,963	67,641	69,398	74,458	74,985	74,888
Mental health diagnosis¹						
Trauma- and stressor-related disorders	7,906	7,656	8,591	10,346	11,029	10,359
Anxiety disorders	6,006	6,133	6,724	7,938	9,121	7,699
Attention deficit/hyperactivity disorder	5,313	3,670	3,103	2,704	2,480	1,075
Mood disorders	28,843	28,616	30,067	27,398	27,392	24,460
Bipolar disorders	12,317	12,356	12,520	7,074	7,029	6,044
Depressive disorders	16,526	16,260	17,547	20,324	20,363	18,416
Personality disorders	158	157	140	245	454	418
Schizophrenia and other psychotic disorders	8,332	8,390	8,307	8,418	8,526	6,758
Alcohol or substance-related disorders	--	3	4	--	--	--
Other mental disorders ²	10,050	9,588	8,619	10,620	10,104	8,796
Total	66,608	64,213	65,555	67,669	69,106	59,565
Gender						
Male	33,315	32,121	32,721	35,054	35,106	35,099
Female	36,431	35,357	36,622	39,372	39,791	39,690
Total	69,746	67,478	69,343	74,426	74,897	74,789
Age³						
0 to 11 years	12,204	11,132	10,602	10,902	6,300	10,585
12 to 14 years	4,985	4,768	4,644	4,852	3,271	5,227
15 to 17 years	5,164	4,840	5,059	5,545	3,557	5,896
18 to 20 years	3,244	2,944	3,194	3,697	2,492	4,246
21 to 24 years	3,973	3,788	3,873	4,090	2,245	3,958
25 to 29 years	5,441	5,370	5,663	6,187	3,600	6,062
30 to 34 years	5,870	5,726	5,981	6,435	4,008	6,165
35 to 39 years	5,200	5,321	5,764	6,359	4,201	6,283
40 to 44 years	5,187	5,132	5,220	5,577	3,683	5,377
45 to 49 years	5,232	4,897	5,004	5,360	3,695	5,148
50 to 54 years	5,192	5,274	5,306	5,468	3,937	4,983
55 to 59 years	3,847	3,898	4,235	4,496	3,498	4,711
60 to 64 years	2,261	2,367	2,534	2,825	2,231	3,116
65 years and older	2,163	2,184	2,319	2,664	28,267	3,131
Total	69,963	67,641	69,398	74,457	74,985	74,888

Continued. See notes at end of table.

Table 9.3a. Arkansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	214	249	210	250	235	269
Asian ⁴	165	161	157	204	199	220
Black or African American	13,507	12,927	13,864	14,572	14,992	14,746
Native Hawaiian or Other Pacific Islander	79	59	64	66	60	64
White	43,633	41,626	43,964	47,003	49,329	47,259
Other	781	690	717	808	874	846
Total	58,379	55,712	58,976	62,903	65,689	63,404
Ethnicity						
Not of Hispanic or Latino origin	56,370	52,812	56,808	63,207	65,827	64,741
Hispanic or Latino origin	2,184	2,238	2,216	2,435	2,693	2,938
Total	58,554	55,050	59,024	65,642	68,520	67,679
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.3b. Arkansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11.9	11.9	13.1	15.3	16.0	17.4
Anxiety disorders	9.0	9.6	10.3	11.7	13.2	12.9
Attention deficit/hyperactivity disorder	8.0	5.7	4.7	4.0	3.6	1.8
Mood disorders	43.3	44.6	45.9	40.5	39.6	41.1
Bipolar disorders	18.5	19.2	19.1	10.5	10.2	10.1
Depressive disorders	24.8	25.3	26.8	30.0	29.5	30.9
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	12.5	13.1	12.7	12.4	12.3	11.3
Alcohol or substance-related disorders	--	†	†	--	--	--
Other mental disorders ²	15.1	14.9	13.1	15.7	14.6	14.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.8	47.6	47.2	47.1	46.9	46.9
Female	52.2	52.4	52.8	52.9	53.1	53.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	17.4	16.5	15.3	14.6	8.4	14.1
12 to 14 years	7.1	7.0	6.7	6.5	4.4	7.0
15 to 17 years	7.4	7.2	7.3	7.4	4.7	7.9
18 to 20 years	4.6	4.4	4.6	5.0	3.3	5.7
21 to 24 years	5.7	5.6	5.6	5.5	3.0	5.3
25 to 29 years	7.8	7.9	8.2	8.3	4.8	8.1
30 to 34 years	8.4	8.5	8.6	8.6	5.3	8.2
35 to 39 years	7.4	7.9	8.3	8.5	5.6	8.4
40 to 44 years	7.4	7.6	7.5	7.5	4.9	7.2
45 to 49 years	7.5	7.2	7.2	7.2	4.9	6.9
50 to 54 years	7.4	7.8	7.6	7.3	5.3	6.7
55 to 59 years	5.5	5.8	6.1	6.0	4.7	6.3
60 to 64 years	3.2	3.5	3.7	3.8	3.0	4.2
65 years and older	3.1	3.2	3.3	3.6	37.7	4.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.3b. Arkansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	23.1	23.2	23.5	23.2	22.8	23.3
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	74.7	74.7	74.5	74.7	75.1	74.5
Other	1.3	1.2	1.2	1.3	1.3	1.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	96.3	95.9	96.2	96.3	96.1	95.7
Hispanic or Latino origin	3.7	4.1	3.8	3.7	3.9	4.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.4a. California individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	672,620	694,961	693,694	646,599	631,998	693,510
Mental health diagnosis¹						
Trauma- and stressor-related disorders	77,098	71,965	72,517	68,428	66,218	75,912
Anxiety disorders	39,751	49,420	53,060	51,854	44,488	54,673
Attention deficit/hyperactivity disorder	38,514	23,568	22,677	21,051	27,333	28,971
Mood disorders	265,422	279,582	280,335	244,257	171,579	209,783
Bipolar disorders	100,515	110,464	109,058	95,803	42,971	49,619
Depressive disorders	164,907	169,118	171,277	148,454	128,608	160,164
Personality disorders	1,056	3,297	3,030	2,835	883	1,057
Schizophrenia and other psychotic disorders	112,502	119,375	118,726	109,781	104,008	121,780
Alcohol or substance-related disorders	23,449	37,515	36,617	33,542	56,819	61,899
Other mental disorders ²	108,270	101,942	100,227	108,118	99,843	106,492
Total	666,062	686,664	687,189	639,866	571,171	660,567
Gender						
Male	355,050	364,722	365,541	346,739	335,840	362,635
Female	316,556	329,278	327,252	298,711	292,958	330,051
Total	671,606	694,000	692,793	645,450	628,798	692,686
Age³						
0 to 11 years	112,079	116,576	116,812	111,267	109,217	118,892
12 to 14 years	51,068	53,392	54,607	51,705	50,559	56,145
15 to 17 years	68,820	67,614	66,475	63,620	61,247	65,959
18 to 20 years	35,799	35,127	34,038	31,673	30,801	32,072
21 to 24 years	36,199	37,515	37,129	35,491	34,597	37,161
25 to 29 years	47,589	49,608	49,297	47,169	48,244	54,210
30 to 34 years	45,864	49,397	50,269	47,326	46,503	50,586
35 to 39 years	40,118	42,444	42,981	40,974	41,602	46,679
40 to 44 years	46,224	45,906	43,721	38,185	35,592	38,733
45 to 49 years	51,190	51,281	48,987	42,946	40,110	42,918
50 to 54 years	51,429	53,188	53,100	46,880	43,332	46,916
55 to 59 years	39,312	42,247	43,335	39,709	38,129	44,260
60 to 64 years	25,124	26,968	28,268	26,447	25,858	30,435
65 years and older	20,008	21,555	21,980	20,544	21,528	26,170
Total	670,823	692,818	690,999	643,936	627,319	691,136

Continued. See notes at end of table.

Table 9.4a. California individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	4,295	4,705	4,823	4,738	4,606	5,258
Asian ⁴	23,609	24,791	24,983	23,433	23,806	25,927
Black or African American	75,962	85,850	85,893	81,094	80,916	94,337
Native Hawaiian or Other Pacific Islander	1,612	1,711	1,903	2,088	1,912	2,209
White	198,210	209,191	212,155	210,846	199,695	211,795
Other	189,519	215,100	222,726	214,947	222,294	242,141
Total	493,207	541,348	552,483	537,146	533,229	581,667
Ethnicity						
Not of Hispanic or Latino origin	276,671	294,213	294,482	280,717	271,981	313,028
Hispanic or Latino origin	201,176	227,661	235,812	225,941	228,755	248,124
Total	477,847	521,874	530,294	506,658	500,736	561,152
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	478	431
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	6,384	6,244
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	6,862	6,675

‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.4b. California individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11.6	10.5	10.6	10.7	11.6	11.5
Anxiety disorders	6.0	7.2	7.7	8.1	7.8	8.3
Attention deficit/hyperactivity disorder	5.8	3.4	3.3	3.3	4.8	4.4
Mood disorders	39.8	40.7	40.8	38.2	30.0	31.8
Bipolar disorders	15.1	16.1	15.9	15.0	7.5	7.5
Depressive disorders	24.8	24.6	24.9	23.2	22.5	24.2
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	16.9	17.4	17.3	17.2	18.2	18.4
Alcohol or substance-related disorders	3.5	5.5	5.3	5.2	9.9	9.4
Other mental disorders ²	16.3	14.8	14.6	16.9	17.5	16.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	52.9	52.6	52.8	53.7	53.4	52.4
Female	47.1	47.4	47.2	46.3	46.6	47.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	16.7	16.8	16.9	17.3	17.4	17.2
12 to 14 years	7.6	7.7	7.9	8.0	8.1	8.1
15 to 17 years	10.3	9.8	9.6	9.9	9.8	9.5
18 to 20 years	5.3	5.1	4.9	4.9	4.9	4.6
21 to 24 years	5.4	5.4	5.4	5.5	5.5	5.4
25 to 29 years	7.1	7.2	7.1	7.3	7.7	7.8
30 to 34 years	6.8	7.1	7.3	7.3	7.4	7.3
35 to 39 years	6.0	6.1	6.2	6.4	6.6	6.8
40 to 44 years	6.9	6.6	6.3	5.9	5.7	5.6
45 to 49 years	7.6	7.4	7.1	6.7	6.4	6.2
50 to 54 years	7.7	7.7	7.7	7.3	6.9	6.8
55 to 59 years	5.9	6.1	6.3	6.2	6.1	6.4
60 to 64 years	3.7	3.9	4.1	4.1	4.1	4.4
65 years and older	3.0	3.1	3.2	3.2	3.4	3.8
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.4b. California individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	4.8	4.6	4.5	4.4	4.5	4.5
Black or African American	15.4	15.9	15.5	15.1	15.2	16.2
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	40.2	38.6	38.4	39.3	37.5	36.4
Other	38.4	39.7	40.3	40.0	41.7	41.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	57.9	56.4	55.5	55.4	54.3	55.8
Hispanic or Latino origin	42.1	43.6	44.5	44.6	45.7	44.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	7.0	6.5
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	93.0	93.5
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual’s age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.5a. Colorado individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	103,284	114,725	130,555	124,029	147,260	150,558
Mental health diagnosis¹						
Trauma- and stressor-related disorders	19,974	22,576	19,246	21,484	32,917	36,755
Anxiety disorders	8,259	10,218	13,739	13,318	17,711	18,175
Attention deficit/hyperactivity disorder	6,286	6,852	6,491	5,653	7,393	7,276
Mood disorders	46,741	51,486	44,974	41,128	54,637	54,667
Bipolar disorders	20,990	22,503	13,568	11,926	15,826	15,560
Depressive disorders	25,751	28,983	31,406	29,202	38,811	39,107
Personality disorders	273	234	565	719	955	974
Schizophrenia and other psychotic disorders	11,470	12,460	9,587	10,379	12,911	12,703
Alcohol or substance-related disorders	1,638	2,262	4,259	4,307	6,091	6,471
Other mental disorders ²	7,507	7,462	9,102	12,872	12,741	11,825
Total	102,148	113,550	107,963	109,860	145,356	148,846
Gender						
Male	47,170	52,985	60,369	57,160	67,351	69,161
Female	56,114	61,740	70,186	66,869	79,909	81,396
Total	103,284	114,725	130,555	124,029	147,260	150,557
Age³						
0 to 11 years	16,473	16,813	16,681	14,237	18,623	19,138
12 to 14 years	9,050	9,921	10,578	9,295	12,020	12,180
15 to 17 years	10,085	10,777	11,511	10,456	13,547	13,684
18 to 20 years	4,664	5,047	5,780	5,560	6,678	7,203
21 to 24 years	5,677	6,523	7,904	7,649	8,586	8,663
25 to 29 years	8,443	9,733	11,993	12,007	13,984	14,518
30 to 34 years	8,991	10,254	12,398	12,002	13,912	13,936
35 to 39 years	7,349	8,609	10,349	10,289	12,587	13,006
40 to 44 years	7,287	8,124	9,374	8,810	9,620	9,961
45 to 49 years	6,807	7,668	8,817	8,575	9,361	9,366
50 to 54 years	6,885	7,937	9,248	8,680	9,397	8,951
55 to 59 years	5,153	6,066	7,332	7,361	8,350	8,407
60 to 64 years	3,157	3,708	4,428	4,746	5,493	5,833
65 years and older	3,258	3,544	4,159	4,361	5,102	5,711
Total	103,279	114,724	130,552	124,028	147,260	150,557

Continued. See notes at end of table.

Table 9.5a. Colorado individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	1,766	2,761	2,951	2,628	3,282	3,387
Asian ⁴	921	1,181	1,210	1,194	1,491	1,544
Black or African American	7,521	8,538	8,636	7,749	9,092	8,865
Native Hawaiian or Other Pacific Islander	240	323	344	341	422	436
White	69,538	90,781	100,677	96,934	112,430	103,483
Other	23,194	3,080	4,464	4,592	6,374	15,380
Total	103,180	106,664	118,282	113,438	133,091	133,095
Ethnicity						
Not of Hispanic or Latino origin	75,617	83,377	95,926	89,697	101,194	101,999
Hispanic or Latino origin	27,653	28,682	30,088	28,493	36,664	37,882
Total	103,270	112,059	126,014	118,190	137,858	139,881
Specific Hispanic or Latino origin						
Mexican	‡	13,130	21,739	18,313	23,380	23,892
Puerto Rican	‡	194	414	669	949	885
Cuban	‡	64	128	306	474	400
Other specific	‡	15,142	7,692	9,114	11,748	12,576
Total	‡	28,530	29,973	28,402	36,551	37,753
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	3,125	3,178
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	99,661	101,924
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	102,786	105,102

‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.5b. Colorado individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	19.6	19.9	17.8	19.6	22.6	24.7
Anxiety disorders	8.1	9.0	12.7	12.1	12.2	12.2
Attention deficit/hyperactivity disorder	6.2	6.0	6.0	5.1	5.1	4.9
Mood disorders	45.8	45.3	41.7	37.4	37.6	36.7
Bipolar disorders	20.5	19.8	12.6	10.9	10.9	10.5
Depressive disorders	25.2	25.5	29.1	26.6	26.7	26.3
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	11.2	11.0	8.9	9.4	8.9	8.5
Alcohol or substance-related disorders	1.6	2.0	3.9	3.9	4.2	4.3
Other mental disorders ²	7.3	6.6	8.4	11.7	8.8	7.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	45.7	46.2	46.2	46.1	45.7	45.9
Female	54.3	53.8	53.8	53.9	54.3	54.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	16.0	14.7	12.8	11.5	12.6	12.7
12 to 14 years	8.8	8.6	8.1	7.5	8.2	8.1
15 to 17 years	9.8	9.4	8.8	8.4	9.2	9.1
18 to 20 years	4.5	4.4	4.4	4.5	4.5	4.8
21 to 24 years	5.5	5.7	6.1	6.2	5.8	5.8
25 to 29 years	8.2	8.5	9.2	9.7	9.5	9.6
30 to 34 years	8.7	8.9	9.5	9.7	9.4	9.3
35 to 39 years	7.1	7.5	7.9	8.3	8.5	8.6
40 to 44 years	7.1	7.1	7.2	7.1	6.5	6.6
45 to 49 years	6.6	6.7	6.8	6.9	6.4	6.2
50 to 54 years	6.7	6.9	7.1	7.0	6.4	5.9
55 to 59 years	5.0	5.3	5.6	5.9	5.7	5.6
60 to 64 years	3.1	3.2	3.4	3.8	3.7	3.9
65 years and older	3.2	3.1	3.2	3.5	3.5	3.8
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.5b. Colorado individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	1.7	2.6	2.5	2.3	2.5	2.5
Asian ⁴	†	1.1	1.0	1.1	1.1	1.2
Black or African American	7.3	8.0	7.3	6.8	6.8	6.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	67.4	85.1	85.1	85.5	84.5	77.8
Other	22.5	2.9	3.8	4.0	4.8	11.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	73.2	74.4	76.1	75.9	73.4	72.9
Hispanic or Latino origin	26.8	25.6	23.9	24.1	26.6	27.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	46.0	72.5	64.5	64.0	63.3
Puerto Rican	‡	†	1.4	2.4	2.6	2.3
Cuban	‡	†	†	1.1	1.3	1.1
Other specific	‡	53.1	25.7	32.1	32.1	33.3
Total	‡	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	3.0	3.0
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	97.0	97.0
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.6a. Connecticut individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	89,639	91,525	90,524	94,541	97,130	93,947
Mental health diagnosis¹						
Trauma- and stressor-related disorders	12,751	13,467	8,930	9,590	13,849	13,690
Anxiety disorders	7,499	6,910	5,123	6,782	7,852	8,023
Attention deficit/hyperactivity disorder	2,888	3,149	2,134	1,968	6,327	6,462
Mood disorders	33,042	35,105	29,030	27,508	26,985	26,258
Bipolar disorders	14,137	14,902	12,126	10,418	8,659	7,914
Depressive disorders	18,905	20,203	16,904	17,090	18,326	18,344
Personality disorders	72	512	401	431	461	506
Schizophrenia and other psychotic disorders	11,360	11,226	10,790	10,869	10,984	10,517
Alcohol or substance-related disorders	--	--	425	144	1,668	1,539
Other mental disorders ²	7,361	8,722	5,736	6,697	10,390	10,865
Total	74,973	79,091	62,569	63,989	78,516	77,860
Gender						
Male	46,377	46,981	46,134	48,903	50,586	49,096
Female	43,224	44,519	44,375	45,579	46,492	44,781
Total	89,601	91,500	90,509	94,482	97,078	93,877
Age³						
0 to 11 years	13,816	14,462	14,686	15,291	16,377	16,606
12 to 14 years	8,413	8,941	8,819	8,697	9,119	9,304
15 to 17 years	8,996	9,493	9,663	9,748	9,885	9,828
18 to 20 years	3,815	4,040	3,908	4,133	4,182	4,216
21 to 24 years	4,132	4,145	4,067	4,226	4,240	3,926
25 to 29 years	5,435	5,353	5,198	5,616	5,817	5,528
30 to 34 years	5,370	5,543	5,311	5,592	5,825	5,545
35 to 39 years	4,713	4,854	4,734	5,101	5,367	5,184
40 to 44 years	5,906	5,464	5,023	5,015	4,923	4,441
45 to 49 years	7,563	7,092	6,499	6,633	6,314	5,530
50 to 54 years	7,815	7,608	7,427	7,717	7,624	6,888
55 to 59 years	6,199	6,427	6,545	7,096	7,237	6,786
60 to 64 years	3,694	4,079	4,363	4,840	5,135	5,030
65 years and older	3,768	4,024	4,256	4,832	5,082	5,132
Total	89,635	91,525	90,499	94,537	97,127	93,944

Continued. See notes at end of table.

Table 9.6a. Connecticut individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	451	491	480	582	579	448
Asian ⁴	251	709	758	832	892	931
Black or African American	15,546	16,002	15,539	16,956	17,693	16,920
Native Hawaiian or Other Pacific Islander	195	204	229	225	228	235
White	54,460	55,994	54,748	57,439	58,806	55,337
Other	15,278	15,045	14,726	14,432	13,977	14,584
Total	86,181	88,445	86,480	90,466	92,175	88,455
Ethnicity						
Not of Hispanic or Latino origin	62,327	64,242	62,983	65,714	66,955	63,629
Hispanic or Latino origin	22,458	23,303	23,280	24,551	25,878	25,282
Total	84,785	87,545	86,263	90,265	92,833	88,911
Specific Hispanic or Latino origin						
Mexican	614	608	794	818	574	385
Puerto Rican	11,609	12,340	11,633	12,025	10,151	8,363
Cuban	155	156	147	162	172	113
Other specific	4,847	4,243	4,444	5,107	5,093	4,947
Total	17,225	17,347	17,018	18,112	15,990	13,808
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2,292	2,109
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	48,872	46,267
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	51,164	48,376

-- Quantity is zero; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.6b. Connecticut individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	17.0	17.0	14.3	15.0	17.6	17.6
Anxiety disorders	10.0	8.7	8.2	10.6	10.0	10.3
Attention deficit/hyperactivity disorder	3.9	4.0	3.4	3.1	8.1	8.3
Mood disorders	44.1	44.4	46.4	43.0	34.4	33.7
Bipolar disorders	18.9	18.8	19.4	16.3	11.0	10.2
Depressive disorders	25.2	25.5	27.0	26.7	23.3	23.6
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	15.2	14.2	17.2	17.0	14.0	13.5
Alcohol or substance-related disorders	--	--	†	†	2.1	2.0
Other mental disorders ²	9.8	11.0	9.2	10.5	13.2	14.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	51.8	51.3	51.0	51.8	52.1	52.3
Female	48.2	48.7	49.0	48.2	47.9	47.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	15.4	15.8	16.2	16.2	16.9	17.7
12 to 14 years	9.4	9.8	9.7	9.2	9.4	9.9
15 to 17 years	10.0	10.4	10.7	10.3	10.2	10.5
18 to 20 years	4.3	4.4	4.3	4.4	4.3	4.5
21 to 24 years	4.6	4.5	4.5	4.5	4.4	4.2
25 to 29 years	6.1	5.8	5.7	5.9	6.0	5.9
30 to 34 years	6.0	6.1	5.9	5.9	6.0	5.9
35 to 39 years	5.3	5.3	5.2	5.4	5.5	5.5
40 to 44 years	6.6	6.0	5.6	5.3	5.1	4.7
45 to 49 years	8.4	7.7	7.2	7.0	6.5	5.9
50 to 54 years	8.7	8.3	8.2	8.2	7.8	7.3
55 to 59 years	6.9	7.0	7.2	7.5	7.5	7.2
60 to 64 years	4.1	4.5	4.8	5.1	5.3	5.4
65 years and older	4.2	4.4	4.7	5.1	5.2	5.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.6b. Connecticut individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	1.1
Black or African American	18.0	18.1	18.0	18.7	19.2	19.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	63.2	63.3	63.3	63.5	63.8	62.6
Other	17.7	17.0	17.0	16.0	15.2	16.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	73.5	73.4	73.0	72.8	72.1	71.6
Hispanic or Latino origin	26.5	26.6	27.0	27.2	27.9	28.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	3.6	3.5	4.7	4.5	3.6	2.8
Puerto Rican	67.4	71.1	68.4	66.4	63.5	60.6
Cuban	†	†	†	†	1.1	†
Other specific	28.1	24.5	26.1	28.2	31.9	35.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	4.5	4.4
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	95.5	95.6
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.7a. Delaware individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	9,127	10,065	9,680	7,456	9,578	9,889
Mental health diagnosis¹						
Trauma- and stressor-related disorders	937	1,125	1,052	682	938	612
Anxiety disorders	374	357	233	127	216	205
Attention deficit/hyperactivity disorder	570	487	209	230	367	254
Mood disorders	3,442	3,759	3,550	2,440	2,700	3,415
Bipolar disorders	1,719	1,875	1,913	1,199	1,253	1,490
Depressive disorders	1,723	1,884	1,637	1,241	1,447	1,925
Personality disorders	7	13	7	22	21	33
Schizophrenia and other psychotic disorders	1,571	1,825	1,833	1,226	1,184	1,683
Alcohol or substance-related disorders	447	834	1,021	1,298	1,658	2,210
Other mental disorders ²	697	692	685	447	537	388
Total	8,045	9,092	8,590	6,472	7,621	8,800
Gender						
Male	4,807	5,372	5,252	4,017	5,042	5,334
Female	4,273	4,671	4,408	3,370	4,343	4,399
Total	9,080	10,043	9,660	7,387	9,385	9,733
Age³						
0 to 11 years	1,133	1,098	878	414	880	311
12 to 14 years	870	943	855	416	835	400
15 to 17 years	1,200	1,184	1,127	488	841	464
18 to 20 years	222	239	252	334	393	477
21 to 24 years	340	445	479	600	651	786
25 to 29 years	525	629	728	852	965	1,167
30 to 34 years	577	705	699	716	855	1,028
35 to 39 years	482	580	619	637	788	1,063
40 to 44 years	529	625	577	522	617	795
45 to 49 years	669	742	700	571	636	820
50 to 54 years	837	936	863	651	673	797
55 to 59 years	706	780	762	572	612	767
60 to 64 years	472	546	551	302	361	496
65 years and older	565	612	590	377	467	516
Total	9,127	10,064	9,680	7,452	9,574	9,887

Continued. See notes at end of table.

Table 9.7a. Delaware individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	38	45	36	46	68	73
Asian ⁴	56	60	70	55	59	60
Black or African American	2,997	3,341	3,238	2,286	3,044	3,144
Native Hawaiian or Other Pacific Islander	77	61	56	27	38	53
White	5,599	6,173	5,987	4,754	5,991	6,111
Other	73	84	68	84	94	136
Total	8,840	9,764	9,455	7,252	9,294	9,577
Ethnicity						
Not of Hispanic or Latino origin	7,669	8,498	8,464	6,689	8,422	8,651
Hispanic or Latino origin	701	817	718	399	610	560
Total	8,370	9,315	9,182	7,088	9,032	9,211
Specific Hispanic or Latino origin						
Mexican	52	95	90	51	28	66
Puerto Rican	93	104	87	68	79	118
Cuban	9	9	13	1	3	5
Other specific	94	149	116	81	110	144
Total	248	357	306	201	220	333
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	193	291
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	4,248	6,172
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	4,441	6,463

n/c Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.7b. Delaware individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11.6	12.4	12.2	10.5	12.3	7.0
Anxiety disorders	4.6	3.9	2.7	2.0	2.8	2.3
Attention deficit/hyperactivity disorder	7.1	5.4	2.4	3.6	4.8	2.9
Mood disorders	42.8	41.3	41.3	37.7	35.4	38.8
Bipolar disorders	21.4	20.6	22.3	18.5	16.4	16.9
Depressive disorders	21.4	20.7	19.1	19.2	19.0	21.9
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	19.5	20.1	21.3	18.9	15.5	19.1
Alcohol or substance-related disorders	5.6	9.2	11.9	20.1	21.8	25.1
Other mental disorders ²	8.7	7.6	8.0	6.9	7.0	4.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	52.9	53.5	54.4	54.4	53.7	54.8
Female	47.1	46.5	45.6	45.6	46.3	45.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	12.4	10.9	9.1	5.6	9.2	3.1
12 to 14 years	9.5	9.4	8.8	5.6	8.7	4.0
15 to 17 years	13.1	11.8	11.6	6.5	8.8	4.7
18 to 20 years	2.4	2.4	2.6	4.5	4.1	4.8
21 to 24 years	3.7	4.4	4.9	8.1	6.8	7.9
25 to 29 years	5.8	6.3	7.5	11.4	10.1	11.8
30 to 34 years	6.3	7.0	7.2	9.6	8.9	10.4
35 to 39 years	5.3	5.8	6.4	8.5	8.2	10.8
40 to 44 years	5.8	6.2	6.0	7.0	6.4	8.0
45 to 49 years	7.3	7.4	7.2	7.7	6.6	8.3
50 to 54 years	9.2	9.3	8.9	8.7	7.0	8.1
55 to 59 years	7.7	7.8	7.9	7.7	6.4	7.8
60 to 64 years	5.2	5.4	5.7	4.1	3.8	5.0
65 years and older	6.2	6.1	6.1	5.1	4.9	5.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.7b. Delaware individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	33.9	34.2	34.2	31.5	32.8	32.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	63.3	63.2	63.3	65.6	64.5	63.8
Other	†	†	†	1.2	1.0	1.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	91.6	91.2	92.2	94.4	93.2	93.9
Hispanic or Latino origin	8.4	8.8	7.8	5.6	6.8	6.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	21.0	26.6	29.4	25.4	12.7	19.8
Puerto Rican	37.5	29.1	28.4	33.8	35.9	35.4
Cuban	3.6	2.5	4.2	†	1.4	1.5
Other specific	37.9	41.7	37.9	40.3	50.0	43.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	4.3	4.5
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	95.7	95.5
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual’s age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.8a. District of Columbia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	‡	25,101	28,531	35,028	36,668	45,560
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	1,212	1,642	1,736	2,651	2,880
Anxiety disorders	‡	497	592	585	1,147	1,263
Attention deficit/hyperactivity disorder	‡	1,255	1,473	1,586	1,945	2,399
Mood disorders	‡	13,664	15,286	15,006	15,265	19,001
Bipolar disorders	‡	3,719	4,272	4,307	5,117	6,155
Depressive disorders	‡	9,945	11,014	10,699	10,148	12,846
Personality disorders	‡	30	32	31	30	38
Schizophrenia and other psychotic disorders	‡	4,896	5,332	5,183	5,752	6,670
Alcohol or substance-related disorders	‡	637	1,029	925	457	607
Other mental disorders ²	‡	735	1,034	1,145	2,376	2,970
Total	‡	22,926	26,420	26,197	29,623	35,828
Gender						
Male	‡	13,117	14,986	18,306	18,923	23,509
Female	‡	11,939	13,466	16,629	17,626	21,925
Total	‡	25,056	28,452	34,935	36,549	45,434
Age³						
0 to 11 years	‡	1,843	1,997	2,580	2,335	2,744
12 to 14 years	‡	1,319	1,460	1,566	1,378	1,708
15 to 17 years	‡	1,354	1,543	1,899	1,797	2,158
18 to 20 years	‡	991	1,166	1,583	1,534	2,033
21 to 24 years	‡	1,377	1,647	2,062	2,188	2,634
25 to 29 years	‡	1,823	2,295	3,156	3,439	4,296
30 to 34 years	‡	1,960	2,233	2,867	3,095	3,876
35 to 39 years	‡	1,644	2,044	2,639	2,936	3,725
40 to 44 years	‡	1,768	1,832	2,177	2,314	3,016
45 to 49 years	‡	2,734	2,801	2,960	3,000	3,385
50 to 54 years	‡	3,191	3,523	4,006	4,026	4,684
55 to 59 years	‡	2,589	2,950	3,554	4,009	4,977
60 to 64 years	‡	1,516	1,833	2,339	2,714	3,573
65 years and older	‡	991	1,207	1,640	1,903	2,751
Total	‡	25,100	28,531	35,028	36,668	45,560

Continued. See notes at end of table.

**Table 9.8a. District of Columbia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018 (continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	‡	37	59	90	77	114
Asian ⁴	‡	84	97	128	116	156
Black or African American	‡	22,306	25,417	31,262	32,907	40,747
Native Hawaiian or Other Pacific Islander	‡	2	3	2	2	2
White	‡	991	1,122	1,368	1,314	1,641
Other	‡	22	29	59	88	115
Total	‡	23,442	26,727	32,909	34,504	42,775
Ethnicity						
Not of Hispanic or Latino origin	‡	1,719	2,570	9,834	12,968	17,370
Hispanic or Latino origin	‡	314	377	686	749	974
Total	‡	2,033	2,947	10,520	13,717	18,344
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.8b. District of Columbia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	‡	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	5.3	6.2	6.6	8.9	8.0
Anxiety disorders	‡	2.2	2.2	2.2	3.9	3.5
Attention deficit/hyperactivity disorder	‡	5.5	5.6	6.1	6.6	6.7
Mood disorders	‡	59.6	57.9	57.3	51.5	53.0
Bipolar disorders	‡	16.2	16.2	16.4	17.3	17.2
Depressive disorders	‡	43.4	41.7	40.8	34.3	35.9
Personality disorders	‡	†	†	†	†	†
Schizophrenia and other psychotic disorders	‡	21.4	20.2	19.8	19.4	18.6
Alcohol or substance-related disorders	‡	2.8	3.9	3.5	1.5	1.7
Other mental disorders ²	‡	3.2	3.9	4.4	8.0	8.3
Total	‡	100.0	100.0	100.0	100.0	100.0
Gender						
Male	‡	52.4	52.7	52.4	51.8	51.7
Female	‡	47.6	47.3	47.6	48.2	48.3
Total	‡	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	‡	7.3	7.0	7.4	6.4	6.0
12 to 14 years	‡	5.3	5.1	4.5	3.8	3.7
15 to 17 years	‡	5.4	5.4	5.4	4.9	4.7
18 to 20 years	‡	3.9	4.1	4.5	4.2	4.5
21 to 24 years	‡	5.5	5.8	5.9	6.0	5.8
25 to 29 years	‡	7.3	8.0	9.0	9.4	9.4
30 to 34 years	‡	7.8	7.8	8.2	8.4	8.5
35 to 39 years	‡	6.5	7.2	7.5	8.0	8.2
40 to 44 years	‡	7.0	6.4	6.2	6.3	6.6
45 to 49 years	‡	10.9	9.8	8.5	8.2	7.4
50 to 54 years	‡	12.7	12.3	11.4	11.0	10.3
55 to 59 years	‡	10.3	10.3	10.1	10.9	10.9
60 to 64 years	‡	6.0	6.4	6.7	7.4	7.8
65 years and older	‡	3.9	4.2	4.7	5.2	6.0
Total	‡	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.8b. District of Columbia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018 (continued)**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	‡	†	†	†	†	†
Asian ⁴	‡	†	†	†	†	†
Black or African American	‡	95.2	95.1	95.0	95.4	95.3
Native Hawaiian or Other Pacific Islander	‡	†	†	†	†	†
White	‡	4.2	4.2	4.2	3.8	3.8
Other	‡	†	†	†	†	†
Total	‡	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	‡	84.6	87.2	93.5	94.5	94.7
Hispanic or Latino origin	‡	15.4	12.8	6.5	5.5	5.3
Total	‡	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.9a. Florida individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	260,065	233,512	230,652	214,257	213,758	218,368
Mental health diagnosis¹						
Trauma- and stressor-related disorders	26,189	18,589	21,424	24,090	31,330	26,900
Anxiety disorders	12,925	9,593	11,283	12,984	12,942	14,027
Attention deficit/hyperactivity disorder	22,589	12,187	12,731	12,562	13,859	13,081
Mood disorders	121,657	87,965	93,283	87,738	75,916	79,177
Bipolar disorders	52,256	38,727	39,892	32,613	25,588	26,120
Depressive disorders	69,401	49,238	53,391	55,125	50,328	53,057
Personality disorders	462	250	242	485	491	515
Schizophrenia and other psychotic disorders	41,224	29,048	30,180	29,169	29,624	32,965
Alcohol or substance-related disorders	1,220	817	788	1,337	813	1,169
Other mental disorders ²	11,644	6,696	7,473	10,083	16,013	17,854
Total	237,910	165,145	177,404	178,448	180,988	185,688
Gender						
Male	127,163	113,779	112,591	105,358	106,001	109,510
Female	132,902	119,732	118,014	108,868	107,756	108,857
Total	260,065	233,511	230,605	214,226	213,757	218,367
Age³						
0 to 11 years	25,374	16,314	16,552	15,196	14,618	13,904
12 to 14 years	16,080	10,980	10,509	9,933	9,931	10,382
15 to 17 years	17,881	14,971	14,859	13,356	12,858	12,642
18 to 20 years	11,374	10,892	10,428	9,731	9,889	10,003
21 to 24 years	16,620	14,590	13,821	12,698	12,662	13,017
25 to 29 years	22,115	20,447	20,901	20,033	20,631	20,762
30 to 34 years	22,292	21,195	21,155	20,058	20,420	21,321
35 to 39 years	19,599	18,479	18,848	18,387	19,075	20,053
40 to 44 years	21,059	18,690	17,843	16,207	16,164	16,544
45 to 49 years	23,583	20,650	19,703	17,978	17,458	17,272
50 to 54 years	24,600	24,018	22,835	20,511	19,202	18,572
55 to 59 years	18,916	19,544	19,617	18,453	18,478	18,763
60 to 64 years	11,204	12,394	12,726	12,139	12,333	13,170
65 years and older	9,366	10,345	10,851	9,574	10,038	11,963
Total	260,063	233,509	230,648	214,254	213,757	218,368

Continued. See notes at end of table.

Table 9.9a. Florida individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	1,214	846	909	821	725	790
Asian ⁴	1,321	1,298	1,287	1,421	1,348	1,283
Black or African American	60,604	50,644	51,231	47,285	46,889	47,903
Native Hawaiian or Other Pacific Islander	491	480	446	414	383	417
White	174,951	163,602	158,827	145,982	144,107	146,370
Other	21,451	16,600	17,020	14,770	13,524	14,189
Total	260,032	233,470	229,720	210,693	206,976	210,952
Ethnicity						
Not of Hispanic or Latino origin	219,539	198,963	195,131	179,464	177,696	178,651
Hispanic or Latino origin	40,472	34,472	35,444	34,741	36,022	39,685
Total	260,011	233,435	230,575	214,205	213,718	218,336
Specific Hispanic or Latino origin						
Mexican	3,120	3,113	3,035	2,696	2,878	3,061
Puerto Rican	8,950	6,984	6,990	7,236	8,027	8,143
Cuban	7,986	7,202	7,289	6,777	6,275	7,280
Other specific	17,288	13,702	14,470	14,440	14,628	16,379
Total	37,344	31,001	31,784	31,149	31,808	34,863
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	9,233	7,199
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	144,614	151,279
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	153,847	158,478

n/c Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.9b. Florida individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11.0	11.3	12.1	13.5	17.3	14.5
Anxiety disorders	5.4	5.8	6.4	7.3	7.2	7.6
Attention deficit/hyperactivity disorder	9.5	7.4	7.2	7.0	7.7	7.0
Mood disorders	51.1	53.3	52.6	49.2	41.9	42.6
Bipolar disorders	22.0	23.5	22.5	18.3	14.1	14.1
Depressive disorders	29.2	29.8	30.1	30.9	27.8	28.6
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	17.3	17.6	17.0	16.3	16.4	17.8
Alcohol or substance-related disorders	†	†	†	†	†	†
Other mental disorders ²	4.9	4.1	4.2	5.7	8.8	9.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	48.9	48.7	48.8	49.2	49.6	50.1
Female	51.1	51.3	51.2	50.8	50.4	49.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	9.8	7.0	7.2	7.1	6.8	6.4
12 to 14 years	6.2	4.7	4.6	4.6	4.6	4.8
15 to 17 years	6.9	6.4	6.4	6.2	6.0	5.8
18 to 20 years	4.4	4.7	4.5	4.5	4.6	4.6
21 to 24 years	6.4	6.2	6.0	5.9	5.9	6.0
25 to 29 years	8.5	8.8	9.1	9.4	9.7	9.5
30 to 34 years	8.6	9.1	9.2	9.4	9.6	9.8
35 to 39 years	7.5	7.9	8.2	8.6	8.9	9.2
40 to 44 years	8.1	8.0	7.7	7.6	7.6	7.6
45 to 49 years	9.1	8.8	8.5	8.4	8.2	7.9
50 to 54 years	9.5	10.3	9.9	9.6	9.0	8.5
55 to 59 years	7.3	8.4	8.5	8.6	8.6	8.6
60 to 64 years	4.3	5.3	5.5	5.7	5.8	6.0
65 years and older	3.6	4.4	4.7	4.5	4.7	5.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.9b. Florida individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	23.3	21.7	22.3	22.4	22.7	22.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	67.3	70.1	69.1	69.3	69.6	69.4
Other	8.2	7.1	7.4	7.0	6.5	6.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	84.4	85.2	84.6	83.8	83.1	81.8
Hispanic or Latino origin	15.6	14.8	15.4	16.2	16.9	18.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	8.4	10.0	9.5	8.7	9.0	8.8
Puerto Rican	24.0	22.5	22.0	23.2	25.2	23.4
Cuban	21.4	23.2	22.9	21.8	19.7	20.9
Other specific	46.3	44.2	45.5	46.4	46.0	47.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	6.0	4.5
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	94.0	95.5
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.10a. Hawaii individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	12,809	12,594	10,645	10,117	9,915	10,006
Mental health diagnosis¹						
Trauma- and stressor-related disorders	2,001	2,320	2,211	1,603	1,947	2,401
Anxiety disorders	249	245	205	239	226	275
Attention deficit/hyperactivity disorder	362	339	361	382	336	395
Mood disorders	4,660	4,545	3,393	3,128	3,005	2,589
Bipolar disorders	1,746	1,658	1,280	1,022	1,136	1,032
Depressive disorders	2,914	2,887	2,113	2,106	1,869	1,557
Personality disorders	94	26	34	34	24	15
Schizophrenia and other psychotic disorders	3,595	3,500	2,872	3,139	2,866	2,719
Alcohol or substance-related disorders	159	138	101	35	22	19
Other mental disorders ²	847	775	703	834	673	684
Total	11,967	11,888	9,880	9,394	9,099	9,097
Gender						
Male	7,022	6,946	6,044	5,838	5,706	5,820
Female	5,608	5,437	4,393	4,255	4,206	4,183
Total	12,630	12,383	10,437	10,093	9,912	10,003
Age³						
0 to 11 years	537	589	714	807	819	793
12 to 14 years	511	525	561	547	580	594
15 to 17 years	884	900	890	891	825	831
18 to 20 years	412	393	386	398	338	319
21 to 24 years	448	425	386	303	323	334
25 to 29 years	743	686	591	547	557	599
30 to 34 years	887	811	704	665	668	680
35 to 39 years	854	800	678	628	616	682
40 to 44 years	1,057	1,019	758	656	625	636
45 to 49 years	1,363	1,317	960	837	737	762
50 to 54 years	1,688	1,616	1,121	1,068	965	895
55 to 59 years	1,457	1,478	1,163	1,132	1,084	1,039
60 to 64 years	990	995	837	779	818	869
65 years and older	957	1,035	894	855	952	964
Total	12,788	12,589	10,643	10,113	9,907	9,997

Continued. See notes at end of table.

Table 9.10a. Hawaii individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	77	68	50	64	56	68
Asian ⁴	1,899	1,784	1,508	1,498	1,457	1,408
Black or African American	210	204	153	153	176	214
Native Hawaiian or Other Pacific Islander	1,531	1,326	962	994	950	994
White	3,067	2,643	2,018	2,071	2,088	2,130
Other	2,371	2,840	2,352	2,069	1,804	1,553
Total	9,155	8,865	7,043	6,849	6,531	6,367
Ethnicity						
Not of Hispanic or Latino origin	8,231	7,904	6,518	7,973	5,738	5,637
Hispanic or Latino origin	878	930	809	831	690	551
Total	9,109	8,834	7,327	8,804	6,428	6,188
Specific Hispanic or Latino origin						
Mexican	27	29	33	40	39	31
Puerto Rican	123	131	124	150	134	132
Cuban	8	8	5	6	7	4
Other specific	86	104	123	151	142	115
Total	244	272	285	347	322	282
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.10b. Hawaii individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	16.7	19.5	22.4	17.1	21.4	26.4
Anxiety disorders	2.1	2.1	2.1	2.5	2.5	3.0
Attention deficit/hyperactivity disorder	3.0	2.9	3.7	4.1	3.7	4.3
Mood disorders	38.9	38.2	34.3	33.3	33.0	28.5
Bipolar disorders	14.6	13.9	13.0	10.9	12.5	11.3
Depressive disorders	24.4	24.3	21.4	22.4	20.5	17.1
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	30.0	29.4	29.1	33.4	31.5	29.9
Alcohol or substance-related disorders	1.3	1.2	1.0	†	†	†
Other mental disorders ²	7.1	6.5	7.1	8.9	7.4	7.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	55.6	56.1	57.9	57.8	57.6	58.2
Female	44.4	43.9	42.1	42.2	42.4	41.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	4.2	4.7	6.7	8.0	8.3	7.9
12 to 14 years	4.0	4.2	5.3	5.4	5.9	5.9
15 to 17 years	6.9	7.1	8.4	8.8	8.3	8.3
18 to 20 years	3.2	3.1	3.6	3.9	3.4	3.2
21 to 24 years	3.5	3.4	3.6	3.0	3.3	3.3
25 to 29 years	5.8	5.4	5.6	5.4	5.6	6.0
30 to 34 years	6.9	6.4	6.6	6.6	6.7	6.8
35 to 39 years	6.7	6.4	6.4	6.2	6.2	6.8
40 to 44 years	8.3	8.1	7.1	6.5	6.3	6.4
45 to 49 years	10.7	10.5	9.0	8.3	7.4	7.6
50 to 54 years	13.2	12.8	10.5	10.6	9.7	9.0
55 to 59 years	11.4	11.7	10.9	11.2	10.9	10.4
60 to 64 years	7.7	7.9	7.9	7.7	8.3	8.7
65 years and older	7.5	8.2	8.4	8.5	9.6	9.6
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.10b. Hawaii individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	1.1
Asian ⁴	20.7	20.1	21.4	21.9	22.3	22.1
Black or African American	2.3	2.3	2.2	2.2	2.7	3.4
Native Hawaiian or Other Pacific Islander	16.7	15.0	13.7	14.5	14.5	15.6
White	33.5	29.8	28.7	30.2	32.0	33.5
Other	25.9	32.0	33.4	30.2	27.6	24.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	90.4	89.5	89.0	90.6	89.3	91.1
Hispanic or Latino origin	9.6	10.5	11.0	9.4	10.7	8.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	11.1	10.7	11.6	11.5	12.1	11.0
Puerto Rican	50.4	48.2	43.5	43.2	41.6	46.8
Cuban	3.3	2.9	1.8	1.7	2.2	1.4
Other specific	35.2	38.2	43.2	43.5	44.1	40.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.11a. Idaho individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	11,880	14,107	14,425	15,320	15,424	16,718
Mental health diagnosis¹						
Trauma- and stressor-related disorders	237	92	53	249	331	291
Anxiety disorders	80	29	15	357	400	383
Attention deficit/hyperactivity disorder	103	64	26	174	182	218
Mood disorders	1,855	917	651	3,489	3,746	3,807
Bipolar disorders	980	523	389	1,507	1,540	1,536
Depressive disorders	875	394	262	1,982	2,206	2,271
Personality disorders	1,777	1,050	941	151	148	154
Schizophrenia and other psychotic disorders	671	423	272	1,817	1,961	1,985
Alcohol or substance-related disorders	490	172	96	3	2	1
Other mental disorders ²	3,096	1,719	1,222	1,037	890	827
Total	8,309	4,466	3,276	7,277	7,660	7,666
Gender						
Male	6,631	8,486	8,581	9,125	9,191	9,925
Female	5,246	5,614	5,838	6,191	6,224	6,782
Total	11,877	14,100	14,419	15,316	15,415	16,707
Age³						
0 to 11 years	251	221	176	230	355	729
12 to 14 years	615	572	546	516	485	704
15 to 17 years	1,081	1,052	1,005	974	1,011	1,059
18 to 20 years	779	1,016	1,094	1,094	1,060	1,138
21 to 24 years	1,072	1,409	1,492	1,596	1,548	1,559
25 to 29 years	1,304	1,778	1,872	1,989	2,058	2,187
30 to 34 years	1,373	1,757	1,730	1,915	1,906	1,980
35 to 39 years	1,108	1,397	1,544	1,720	1,703	1,834
40 to 44 years	1,048	1,160	1,162	1,243	1,266	1,341
45 to 49 years	964	1,146	1,114	1,195	1,234	1,224
50 to 54 years	961	1,072	1,066	1,141	998	984
55 to 59 years	604	698	774	800	821	918
60 to 64 years	352	415	442	459	473	512
65 years and older	353	402	399	421	479	521
Total	11,865	14,095	14,416	15,293	15,397	16,690

Continued. See notes at end of table.

Table 9.11a. Idaho individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	336	393	332	402	364	386
Asian ⁴	65	87	74	71	81	93
Black or African American	185	250	274	271	293	291
Native Hawaiian or Other Pacific Islander	34	37	33	43	40	54
White	9,599	11,385	11,361	11,787	11,768	12,648
Other	509	865	1,206	1,380	1,435	1,754
Total	10,728	13,017	13,280	13,954	13,981	15,226
Ethnicity						
Not of Hispanic or Latino origin	9,418	11,321	11,460	11,899	11,888	12,793
Hispanic or Latino origin	1,011	1,287	1,409	1,572	1,562	1,750
Total	10,429	12,608	12,869	13,471	13,450	14,543
Specific Hispanic or Latino origin						
Mexican	274	303	288	254	205	209
Puerto Rican	18	25	22	25	21	23
Cuban	5	6	5	9	8	5
Other specific	394	365	333	221	152	160
Total	691	699	648	509	386	397
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	494	471
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	10,262	10,723
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	10,756	11,194

n/c Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.11b. Idaho individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	2.9	2.1	1.6	3.4	4.3	3.8
Anxiety disorders	†	†	†	4.9	5.2	5.0
Attention deficit/hyperactivity disorder	1.2	1.4	†	2.4	2.4	2.8
Mood disorders	22.3	20.5	19.9	47.9	48.9	49.7
Bipolar disorders	11.8	11.7	11.9	20.7	20.1	20.0
Depressive disorders	10.5	8.8	8.0	27.2	28.8	29.6
Personality disorders	21.4	23.5	28.7	2.1	1.9	2.0
Schizophrenia and other psychotic disorders	8.1	9.5	8.3	25.0	25.6	25.9
Alcohol or substance-related disorders	5.9	3.9	2.9	†	†	†
Other mental disorders ²	37.3	38.5	37.3	14.3	11.6	10.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	55.8	60.2	59.5	59.6	59.6	59.4
Female	44.2	39.8	40.5	40.4	40.4	40.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	2.1	1.6	1.2	1.5	2.3	4.4
12 to 14 years	5.2	4.1	3.8	3.4	3.1	4.2
15 to 17 years	9.1	7.5	7.0	6.4	6.6	6.3
18 to 20 years	6.6	7.2	7.6	7.2	6.9	6.8
21 to 24 years	9.0	10.0	10.3	10.4	10.1	9.3
25 to 29 years	11.0	12.6	13.0	13.0	13.4	13.1
30 to 34 years	11.6	12.5	12.0	12.5	12.4	11.9
35 to 39 years	9.3	9.9	10.7	11.2	11.1	11.0
40 to 44 years	8.8	8.2	8.1	8.1	8.2	8.0
45 to 49 years	8.1	8.1	7.7	7.8	8.0	7.3
50 to 54 years	8.1	7.6	7.4	7.5	6.5	5.9
55 to 59 years	5.1	5.0	5.4	5.2	5.3	5.5
60 to 64 years	3.0	2.9	3.1	3.0	3.1	3.1
65 years and older	3.0	2.9	2.8	2.8	3.1	3.1
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.11b. Idaho individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	3.1	3.0	2.5	2.9	2.6	2.5
Asian ⁴	†	†	†	†	†	†
Black or African American	1.7	1.9	2.1	1.9	2.1	1.9
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	89.5	87.5	85.5	84.5	84.2	83.1
Other	4.7	6.6	9.1	9.9	10.3	11.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	90.3	89.8	89.1	88.3	88.4	88.0
Hispanic or Latino origin	9.7	10.2	10.9	11.7	11.6	12.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	39.7	43.3	44.4	49.9	53.1	52.6
Puerto Rican	2.6	3.6	3.4	4.9	5.4	5.8
Cuban	†	†	†	1.8	2.1	1.3
Other specific	57.0	52.2	51.4	43.4	39.4	40.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	4.6	4.2
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	95.4	95.8
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.12a. Illinois individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	140,482	139,699	130,002	78,703	71,195	66,656
Mental health diagnosis¹						
Trauma- and stressor-related disorders	15,368	15,659	14,715	1,807	8,831	8,809
Anxiety disorders	8,102	8,930	9,513	1,397	5,786	5,655
Attention deficit/hyperactivity disorder	11,749	11,367	10,858	983	5,799	5,127
Mood disorders	73,573	73,299	68,003	5,150	31,298	28,636
Bipolar disorders	29,321	30,002	27,885	2,341	11,339	10,132
Depressive disorders	44,252	43,297	40,118	2,809	19,959	18,504
Personality disorders	237	204	198	34	282	241
Schizophrenia and other psychotic disorders	20,986	21,067	17,832	3,304	11,241	10,409
Alcohol or substance-related disorders	7	6	1	268	293	326
Other mental disorders ²	10,435	9,147	8,876	4,375	7,268	7,032
Total	140,457	139,679	129,996	17,318	70,798	66,235
Gender						
Male	66,502	66,491	62,453	38,763	34,141	32,109
Female	73,979	73,207	67,547	39,938	37,054	34,547
Total	140,481	139,698	130,000	78,701	71,195	66,656
Age³						
0 to 11 years	17,012	16,738	15,205	9,084	8,204	7,745
12 to 14 years	10,018	10,124	9,458	5,513	4,872	4,733
15 to 17 years	10,066	10,380	10,197	5,883	5,276	5,037
18 to 20 years	6,679	6,565	6,649	3,813	3,498	3,214
21 to 24 years	9,002	8,603	8,402	4,731	4,370	3,971
25 to 29 years	11,984	11,606	10,904	6,634	6,188	5,876
30 to 34 years	12,279	11,955	10,972	6,415	5,868	5,480
35 to 39 years	10,728	10,706	10,120	6,113	5,726	5,356
40 to 44 years	10,939	10,754	9,730	5,749	5,080	4,600
45 to 49 years	11,438	11,051	9,624	5,976	5,089	4,730
50 to 54 years	12,188	11,906	10,658	6,597	5,634	4,952
55 to 59 years	8,911	9,409	8,396	5,728	5,207	4,965
60 to 64 years	5,016	5,299	5,019	3,471	3,217	3,205
65 years and older	4,221	4,602	4,666	2,993	2,965	2,792
Total	140,481	139,698	130,000	78,700	71,194	66,656

Continued. See notes at end of table.

Table 9.12a. Illinois individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	439	374	333	232	197	213
Asian ⁴	1,729	1,735	1,655	926	775	678
Black or African American	30,619	30,314	25,584	15,463	12,413	11,909
Native Hawaiian or Other Pacific Islander	163	155	128	61	60	50
White	95,786	96,105	90,659	55,367	51,775	48,400
Other	1,086	1,095	1,330	1,006	996	918
Total	129,822	129,778	119,689	73,055	66,216	62,168
Ethnicity						
Not of Hispanic or Latino origin	112,760	114,886	107,171	67,382	62,039	57,962
Hispanic or Latino origin	17,242	18,568	17,271	8,093	5,701	5,256
Total	130,002	133,454	124,442	75,475	67,740	63,218
Specific Hispanic or Latino origin						
Mexican	11,236	11,760	11,320	4,887	3,279	2,979
Puerto Rican	2,743	2,810	2,329	1,244	929	822
Cuban	205	200	189	83	70	55
Other specific	3,036	3,254	3,130	1,496	1,207	1,249
Total	17,220	18,024	16,968	7,710	5,485	5,105
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	1,266	1,158
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	48,004	44,146
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	49,270	45,304

n/c Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.12b. Illinois individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	10.9	11.2	11.3	10.4	12.5	13.3
Anxiety disorders	5.8	6.4	7.3	8.1	8.2	8.5
Attention deficit/hyperactivity disorder	8.4	8.1	8.4	5.7	8.2	7.7
Mood disorders	52.4	52.5	52.3	29.7	44.2	43.2
Bipolar disorders	20.9	21.5	21.5	13.5	16.0	15.3
Depressive disorders	31.5	31.0	30.9	16.2	28.2	27.9
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	14.9	15.1	13.7	19.1	15.9	15.7
Alcohol or substance-related disorders	†	†	†	1.5	†	†
Other mental disorders ²	7.4	6.5	6.8	25.3	10.3	10.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.3	47.6	48.0	49.3	48.0	48.2
Female	52.7	52.4	52.0	50.7	52.0	51.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	12.1	12.0	11.7	11.5	11.5	11.6
12 to 14 years	7.1	7.2	7.3	7.0	6.8	7.1
15 to 17 years	7.2	7.4	7.8	7.5	7.4	7.6
18 to 20 years	4.8	4.7	5.1	4.8	4.9	4.8
21 to 24 years	6.4	6.2	6.5	6.0	6.1	6.0
25 to 29 years	8.5	8.3	8.4	8.4	8.7	8.8
30 to 34 years	8.7	8.6	8.4	8.2	8.2	8.2
35 to 39 years	7.6	7.7	7.8	7.8	8.0	8.0
40 to 44 years	7.8	7.7	7.5	7.3	7.1	6.9
45 to 49 years	8.1	7.9	7.4	7.6	7.1	7.1
50 to 54 years	8.7	8.5	8.2	8.4	7.9	7.4
55 to 59 years	6.3	6.7	6.5	7.3	7.3	7.4
60 to 64 years	3.6	3.8	3.9	4.4	4.5	4.8
65 years and older	3.0	3.3	3.6	3.8	4.2	4.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.12b. Illinois individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	1.3	1.3	1.4	1.3	1.2	1.1
Black or African American	23.6	23.4	21.4	21.2	18.7	19.2
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	73.8	74.1	75.7	75.8	78.2	77.9
Other	†	†	1.1	1.4	1.5	1.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	86.7	86.1	86.1	89.3	91.6	91.7
Hispanic or Latino origin	13.3	13.9	13.9	10.7	8.4	8.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	65.2	65.2	66.7	63.4	59.8	58.4
Puerto Rican	15.9	15.6	13.7	16.1	16.9	16.1
Cuban	1.2	1.1	1.1	1.1	1.3	1.1
Other specific	17.6	18.1	18.4	19.4	22.0	24.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2.6	2.6
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	97.4	97.4
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected;

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.13a. Indiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	126,032	129,176	134,129	135,393	137,649	137,388
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11,882	11,914	12,312	10,419	11,186	11,718
Anxiety disorders	8,754	9,046	9,756	11,258	12,857	13,401
Attention deficit/hyperactivity disorder	13,894	14,060	14,549	16,154	14,908	14,142
Mood disorders	60,362	62,347	65,190	55,812	61,339	61,322
Bipolar disorders	18,558	18,516	18,821	13,291	14,437	13,813
Depressive disorders	41,804	43,831	46,369	42,521	46,902	47,509
Personality disorders	469	473	410	937	931	1,044
Schizophrenia and other psychotic disorders	14,486	14,735	14,704	14,599	15,442	14,899
Alcohol or substance-related disorders	--	--	--	3	1	--
Other mental disorders ²	15,526	15,802	16,379	19,082	20,944	20,837
Total	125,373	128,377	133,300	128,264	137,608	137,363
Gender						
Male	60,160	61,093	63,174	63,926	64,846	65,064
Female	65,872	68,083	70,955	71,467	72,803	72,324
Total	126,032	129,176	134,129	135,393	137,649	137,388
Age³						
0 to 11 years	25,782	26,880	28,517	28,618	29,159	29,661
12 to 14 years	12,494	12,528	13,095	12,841	13,048	13,682
15 to 17 years	11,429	11,557	11,967	12,588	12,451	12,419
18 to 20 years	5,342	5,350	5,331	5,365	5,744	5,885
21 to 24 years	6,492	6,401	6,595	6,470	6,648	6,659
25 to 29 years	8,612	8,504	8,775	8,928	9,369	9,447
30 to 34 years	9,068	9,556	9,251	9,073	9,191	8,842
35 to 39 years	8,269	8,448	8,759	8,809	9,144	8,869
40 to 44 years	8,361	8,299	8,398	8,305	8,147	7,932
45 to 49 years	8,492	8,337	8,421	8,390	8,327	8,043
50 to 54 years	8,459	8,832	9,184	9,256	8,893	8,313
55 to 59 years	6,341	6,934	7,376	7,704	7,967	7,857
60 to 64 years	3,515	3,902	4,431	4,762	5,104	5,218
65 years and older	3,375	3,648	4,029	4,284	4,457	4,561
Total	126,031	129,176	134,129	135,393	137,649	137,388

Continued. See notes at end of table.

Table 9.13a. Indiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	669	644	676	649	579	506
Asian ⁴	320	381	438	434	520	527
Black or African American	18,472	18,835	19,388	19,528	20,143	20,176
Native Hawaiian or Other Pacific Islander	98	101	104	103	101	127
White	96,339	99,333	103,495	104,597	105,912	106,710
Other	10,134	9,880	10,023	10,079	10,389	9,341
Total	126,032	129,174	134,124	135,390	137,644	137,387
Ethnicity						
Not of Hispanic or Latino origin	112,802	116,429	121,103	122,499	124,863	125,528
Hispanic or Latino origin	7,272	7,457	8,132	8,423	8,723	9,613
Total	120,074	123,886	129,235	130,922	133,586	135,141
Specific Hispanic or Latino origin						
Mexican	2,482	2,770	3,144	3,231	3,258	3,410
Puerto Rican	527	587	603	614	614	739
Cuban	41	45	52	55	52	69
Other specific	4,188	4,018	4,287	4,470	4,764	5,363
Total	7,238	7,420	8,086	8,370	8,688	9,581
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2,681	2,628
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	627	660
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	3,308	3,288

-- Quantity is zero; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.13b. Indiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	9.5	9.3	9.2	8.1	8.1	8.5
Anxiety disorders	7.0	7.0	7.3	8.8	9.3	9.8
Attention deficit/hyperactivity disorder	11.1	11.0	10.9	12.6	10.8	10.3
Mood disorders	48.1	48.6	48.9	43.5	44.6	44.6
Bipolar disorders	14.8	14.4	14.1	10.4	10.5	10.1
Depressive disorders	33.3	34.1	34.8	33.2	34.1	34.6
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	11.6	11.5	11.0	11.4	11.2	10.8
Alcohol or substance-related disorders	--	--	--	†	†	--
Other mental disorders ²	12.4	12.3	12.3	14.9	15.2	15.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.7	47.3	47.1	47.2	47.1	47.4
Female	52.3	52.7	52.9	52.8	52.9	52.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	20.5	20.8	21.3	21.1	21.2	21.6
12 to 14 years	9.9	9.7	9.8	9.5	9.5	10.0
15 to 17 years	9.1	8.9	8.9	9.3	9.0	9.0
18 to 20 years	4.2	4.1	4.0	4.0	4.2	4.3
21 to 24 years	5.2	5.0	4.9	4.8	4.8	4.8
25 to 29 years	6.8	6.6	6.5	6.6	6.8	6.9
30 to 34 years	7.2	7.4	6.9	6.7	6.7	6.4
35 to 39 years	6.6	6.5	6.5	6.5	6.6	6.5
40 to 44 years	6.6	6.4	6.3	6.1	5.9	5.8
45 to 49 years	6.7	6.5	6.3	6.2	6.0	5.9
50 to 54 years	6.7	6.8	6.8	6.8	6.5	6.1
55 to 59 years	5.0	5.4	5.5	5.7	5.8	5.7
60 to 64 years	2.8	3.0	3.3	3.5	3.7	3.8
65 years and older	2.7	2.8	3.0	3.2	3.2	3.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.13b. Indiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	14.7	14.6	14.5	14.4	14.6	14.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	76.4	76.9	77.2	77.3	76.9	77.7
Other	8.0	7.6	7.5	7.4	7.5	6.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	93.9	94.0	93.7	93.6	93.5	92.9
Hispanic or Latino origin	6.1	6.0	6.3	6.4	6.5	7.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	34.3	37.3	38.9	38.6	37.5	35.6
Puerto Rican	7.3	7.9	7.5	7.3	7.1	7.7
Cuban	†	†	†	†	†	†
Other specific	57.9	54.2	53.0	53.4	54.8	56.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	81.0	79.9
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	19.0	20.1
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.14a. Iowa individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	106,151	111,351	116,443	115,628	101,763	189,347
Mental health diagnosis¹						
Trauma- and stressor-related disorders	17,196	19,429	22,973	14,460	6,670	29,171
Anxiety disorders	7,863	9,388	11,739	10,573	5,592	35,102
Attention deficit/hyperactivity disorder	14,531	14,573	15,119	11,309	4,883	25,669
Mood disorders	21,306	25,167	30,186	50,064	27,918	54,911
Bipolar disorders	7,397	8,405	9,268	13,588	8,110	11,833
Depressive disorders	13,909	16,762	20,918	36,476	19,808	43,078
Personality disorders	169	217	286	408	207	790
Schizophrenia and other psychotic disorders	4,586	4,965	5,229	8,045	4,796	10,473
Alcohol or substance-related disorders	271	578	966	941	430	--
Other mental disorders ²	10,519	10,416	10,194	9,996	5,304	17,895
Total	76,441	84,733	96,692	105,796	55,800	174,011
Gender						
Male	50,428	52,676	55,013	51,623	43,115	84,183
Female	55,713	58,675	61,430	64,004	58,460	105,025
Total	106,141	111,351	116,443	115,627	101,575	189,208
Age³						
0 to 11 years	21,624	24,655	24,236	15,664	7,496	33,234
12 to 14 years	9,736	9,954	10,109	7,672	5,159	14,783
15 to 17 years	9,932	9,518	9,900	8,688	7,097	14,450
18 to 20 years	6,475	5,262	5,592	5,563	5,274	9,946
21 to 24 years	6,434	6,625	7,195	7,150	6,785	11,400
25 to 29 years	8,158	8,822	9,828	10,285	9,734	17,156
30 to 34 years	8,224	8,739	9,669	10,087	9,589	15,526
35 to 39 years	6,779	7,258	8,264	9,263	9,007	14,450
40 to 44 years	5,930	6,437	6,935	7,828	7,427	11,280
45 to 49 years	5,791	6,269	6,596	7,618	7,510	10,686
50 to 54 years	6,171	6,757	7,174	8,333	8,043	10,668
55 to 59 years	4,931	5,223	5,579	7,299	7,547	10,498
60 to 64 years	3,067	3,141	3,115	4,770	5,029	7,085
65 years and older	2,898	2,690	2,249	5,407	6,062	8,181
Total	106,150	111,350	116,441	115,627	101,759	189,343

Continued. See notes at end of table.

Table 9.14a. Iowa individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	892	1,051	1,413	1,342	1,249	2,524
Asian ⁴	397	435	512	489	458	923
Black or African American	6,566	6,825	7,248	6,734	5,877	10,404
Native Hawaiian or Other Pacific Islander	60	198	427	382	292	743
White	69,689	74,642	78,122	78,216	68,567	121,633
Other	2,957	3,122	3,573	3,146	2,420	6,233
Total	80,561	86,273	91,295	90,309	78,863	142,460
Ethnicity						
Not of Hispanic or Latino origin	430	291	409	461	483	487
Hispanic or Latino origin	36	41	38	39	43	54
Total	466	332	447	500	526	541
Specific Hispanic or Latino origin						
Mexican	‡	‡	4	3	1	‡
Puerto Rican	‡	‡	--	--	--	‡
Cuban	‡	‡	--	--	--	‡
Other specific	‡	‡	--	--	--	‡
Total	‡	‡	4	3	1	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.14b. Iowa individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	22.5	22.9	23.8	13.7	12.0	16.8
Anxiety disorders	10.3	11.1	12.1	10.0	10.0	20.2
Attention deficit/hyperactivity disorder	19.0	17.2	15.6	10.7	8.8	14.8
Mood disorders	27.9	29.7	31.2	47.3	50.0	31.6
Bipolar disorders	9.7	9.9	9.6	12.8	14.5	6.8
Depressive disorders	18.2	19.8	21.6	34.5	35.5	24.8
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	6.0	5.9	5.4	7.6	8.6	6.0
Alcohol or substance-related disorders	†	†	†	†	†	--
Other mental disorders ²	13.8	12.3	10.5	9.4	9.5	10.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.5	47.3	47.2	44.6	42.4	44.5
Female	52.5	52.7	52.8	55.4	57.6	55.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	20.4	22.1	20.8	13.5	7.4	17.6
12 to 14 years	9.2	8.9	8.7	6.6	5.1	7.8
15 to 17 years	9.4	8.5	8.5	7.5	7.0	7.6
18 to 20 years	6.1	4.7	4.8	4.8	5.2	5.3
21 to 24 years	6.1	5.9	6.2	6.2	6.7	6.0
25 to 29 years	7.7	7.9	8.4	8.9	9.6	9.1
30 to 34 years	7.7	7.8	8.3	8.7	9.4	8.2
35 to 39 years	6.4	6.5	7.1	8.0	8.9	7.6
40 to 44 years	5.6	5.8	6.0	6.8	7.3	6.0
45 to 49 years	5.5	5.6	5.7	6.6	7.4	5.6
50 to 54 years	5.8	6.1	6.2	7.2	7.9	5.6
55 to 59 years	4.6	4.7	4.8	6.3	7.4	5.5
60 to 64 years	2.9	2.8	2.7	4.1	4.9	3.7
65 years and older	2.7	2.4	1.9	4.7	6.0	4.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.14b. Iowa individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	1.1	1.2	1.5	1.5	1.6	1.8
Asian ⁴	†	†	†	†	†	†
Black or African American	8.2	7.9	7.9	7.5	7.5	7.3
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	86.5	86.5	85.6	86.6	86.9	85.4
Other	3.7	3.6	3.9	3.5	3.1	4.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	92.3	87.7	91.5	92.2	91.8	90.0
Hispanic or Latino origin	7.7	12.3	8.5	7.8	8.2	10.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	100.0	100.0	100.0	‡
Puerto Rican	‡	‡	--	--	--	‡
Cuban	‡	‡	--	--	--	‡
Other specific	‡	‡	--	--	--	‡
Total	‡	‡	100.0	100.0	100.0	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.15a. Kansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	84,636	80,991	‡	‡	‡	‡
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11,635	6,155	‡	‡	‡	‡
Anxiety disorders	7,951	5,137	‡	‡	‡	‡
Attention deficit/hyperactivity disorder	10,816	6,004	‡	‡	‡	‡
Mood disorders	33,269	21,921	‡	‡	‡	‡
Bipolar disorders	14,603	9,719	‡	‡	‡	‡
Depressive disorders	18,666	12,202	‡	‡	‡	‡
Personality disorders	427	223	‡	‡	‡	‡
Schizophrenia and other psychotic disorders	7,724	5,804	‡	‡	‡	‡
Alcohol or substance-related disorders	--	--	‡	‡	‡	‡
Other mental disorders ²	11,228	6,269	‡	‡	‡	‡
Total	83,050	51,513	‡	‡	‡	‡
Gender						
Male	40,991	38,524	‡	‡	‡	‡
Female	43,213	42,023	‡	‡	‡	‡
Total	84,204	80,547	‡	‡	‡	‡
Age³						
0 to 11 years	18,793	19,844	‡	‡	‡	‡
12 to 14 years	8,288	8,307	‡	‡	‡	‡
15 to 17 years	8,503	8,525	‡	‡	‡	‡
18 to 20 years	4,308	3,731	‡	‡	‡	‡
21 to 24 years	4,826	4,266	‡	‡	‡	‡
25 to 29 years	6,057	5,387	‡	‡	‡	‡
30 to 34 years	6,042	5,510	‡	‡	‡	‡
35 to 39 years	4,802	4,502	‡	‡	‡	‡
40 to 44 years	4,585	3,917	‡	‡	‡	‡
45 to 49 years	4,755	3,955	‡	‡	‡	‡
50 to 54 years	5,078	4,388	‡	‡	‡	‡
55 to 59 years	3,810	3,397	‡	‡	‡	‡
60 to 64 years	2,080	2,019	‡	‡	‡	‡
65 years and older	2,282	2,806	‡	‡	‡	‡
Total	84,209	80,554	‡	‡	‡	‡

Continued. See notes at end of table.

Table 9.15a. Kansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	2,159	1,945	‡	‡	‡	‡
Asian ⁴	584	649	‡	‡	‡	‡
Black or African American	8,946	8,455	‡	‡	‡	‡
Native Hawaiian or Other Pacific Islander	122	98	‡	‡	‡	‡
White	64,648	63,129	‡	‡	‡	‡
Other	2,703	2,280	‡	‡	‡	‡
Total	79,162	76,556	‡	‡	‡	‡
Ethnicity						
Not of Hispanic or Latino origin	77,780	73,364	‡	‡	‡	‡
Hispanic or Latino origin	5,206	6,011	‡	‡	‡	‡
Total	82,986	79,375	‡	‡	‡	‡
Specific Hispanic or Latino origin						
Mexican	42	44	‡	‡	‡	‡
Puerto Rican	9	6	‡	‡	‡	‡
Cuban	3	--	‡	‡	‡	‡
Other specific	--	--	‡	‡	‡	‡
Total	54	50	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.15b. Kansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	‡	‡	‡	‡
Mental health diagnosis¹						
Trauma- and stressor-related disorders	14.0	11.9	‡	‡	‡	‡
Anxiety disorders	9.6	10.0	‡	‡	‡	‡
Attention deficit/hyperactivity disorder	13.0	11.7	‡	‡	‡	‡
Mood disorders	40.1	42.6	‡	‡	‡	‡
Bipolar disorders	17.6	18.9	‡	‡	‡	‡
Depressive disorders	22.5	23.7	‡	‡	‡	‡
Personality disorders	†	†	‡	‡	‡	‡
Schizophrenia and other psychotic disorders	9.3	11.3	‡	‡	‡	‡
Alcohol or substance-related disorders	--	--	‡	‡	‡	‡
Other mental disorders ²	13.5	12.2	‡	‡	‡	‡
Total	100.0	100.0	‡	‡	‡	‡
Gender						
Male	48.7	47.8	‡	‡	‡	‡
Female	51.3	52.2	‡	‡	‡	‡
Total	100.0	100.0	‡	‡	‡	‡
Age³						
0 to 11 years	22.3	24.6	‡	‡	‡	‡
12 to 14 years	9.8	10.3	‡	‡	‡	‡
15 to 17 years	10.1	10.6	‡	‡	‡	‡
18 to 20 years	5.1	4.6	‡	‡	‡	‡
21 to 24 years	5.7	5.3	‡	‡	‡	‡
25 to 29 years	7.2	6.7	‡	‡	‡	‡
30 to 34 years	7.2	6.8	‡	‡	‡	‡
35 to 39 years	5.7	5.6	‡	‡	‡	‡
40 to 44 years	5.4	4.9	‡	‡	‡	‡
45 to 49 years	5.6	4.9	‡	‡	‡	‡
50 to 54 years	6.0	5.4	‡	‡	‡	‡
55 to 59 years	4.5	4.2	‡	‡	‡	‡
60 to 64 years	2.5	2.5	‡	‡	‡	‡
65 years and older	2.7	3.5	‡	‡	‡	‡
Total	100.0	100.0	‡	‡	‡	‡

Continued. See notes at end of table.

Table 9.15b. Kansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	2.7	2.5	‡	‡	‡	‡
Asian ⁴	†	†	‡	‡	‡	‡
Black or African American	11.3	11.0	‡	‡	‡	‡
Native Hawaiian or Other Pacific Islander	†	†	‡	‡	‡	‡
White	81.7	82.5	‡	‡	‡	‡
Other	3.4	3.0	‡	‡	‡	‡
Total	100.0	100.0	‡	‡	‡	‡
Ethnicity						
Not of Hispanic or Latino origin	93.7	92.4	‡	‡	‡	‡
Hispanic or Latino origin	6.3	7.6	‡	‡	‡	‡
Total	100.0	100.0	‡	‡	‡	‡
Specific Hispanic or Latino origin						
Mexican	77.8	88.0	‡	‡	‡	‡
Puerto Rican	16.7	12.0	‡	‡	‡	‡
Cuban	5.6	--	‡	‡	‡	‡
Other specific	--	--	‡	‡	‡	‡
Total	100.0	100.0	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.16a. Kentucky individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	161,441	158,086	160,096	155,014	155,192	157,732
Mental health diagnosis¹						
Trauma- and stressor-related disorders	20,730	20,240	20,797	21,711	21,059	22,995
Anxiety disorders	14,064	14,433	15,495	19,823	17,650	17,814
Attention deficit/hyperactivity disorder	17,061	16,176	15,424	14,469	13,618	13,405
Mood disorders	65,287	64,559	65,419	56,378	45,388	46,340
Bipolar disorders	26,098	25,722	26,046	18,919	11,029	11,021
Depressive disorders	39,189	38,837	39,373	37,459	34,359	35,319
Personality disorders	568	528	605	1,003	915	941
Schizophrenia and other psychotic disorders	11,370	11,396	11,300	10,440	10,171	10,367
Alcohol or substance-related disorders	4,742	4,852	4,832	5,848	15,326	16,205
Other mental disorders ²	23,232	21,622	21,776	18,724	29,103	27,922
Total	157,054	153,806	155,648	148,396	153,230	155,989
Gender						
Male	78,199	76,245	77,168	74,017	73,957	75,775
Female	83,162	81,816	82,905	80,947	81,210	81,913
Total	161,361	158,061	160,073	154,964	155,167	157,688
Age³						
0 to 11 years	33,422	31,502	29,922	26,021	25,881	26,645
12 to 14 years	14,176	13,875	13,525	12,491	12,460	12,711
15 to 17 years	13,245	13,082	13,071	12,718	12,824	12,933
18 to 20 years	6,790	6,473	6,735	6,730	6,842	7,336
21 to 24 years	9,161	8,976	9,206	9,182	9,014	9,139
25 to 29 years	11,463	11,519	12,016	12,539	12,824	13,277
30 to 34 years	12,186	11,986	12,341	12,032	12,101	12,247
35 to 39 years	10,501	10,763	11,503	11,798	11,811	11,753
40 to 44 years	10,833	10,505	10,664	10,308	9,913	9,976
45 to 49 years	11,241	10,762	10,716	10,544	10,286	9,883
50 to 54 years	10,722	10,752	11,048	10,867	10,613	10,035
55 to 59 years	8,020	8,156	8,506	8,652	8,580	8,654
60 to 64 years	4,816	4,892	5,148	5,332	5,603	5,730
65 years and older	4,859	4,832	5,686	5,750	6,405	7,371
Total	161,435	158,075	160,087	154,964	155,157	157,690

Continued. See notes at end of table.

Table 9.16a. Kentucky individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	309	307	311	311	335	304
Asian ⁴	189	240	256	376	419	425
Black or African American	16,945	16,532	16,570	15,622	15,555	15,047
Native Hawaiian or Other Pacific Islander	130	83	103	112	135	151
White	134,281	130,428	132,093	126,185	125,398	125,431
Other	9,554	10,365	10,685	10,123	10,544	13,186
Total	161,408	157,955	160,018	152,729	152,386	154,544
Ethnicity						
Not of Hispanic or Latino origin	155,736	152,240	154,189	148,371	148,170	150,202
Hispanic or Latino origin	2,932	3,008	3,154	3,323	3,480	3,452
Total	158,668	155,248	157,343	151,694	151,650	153,654
Specific Hispanic or Latino origin						
Mexican	1,418	1,377	1,395	1,344	1,400	1,393
Puerto Rican	504	531	551	499	494	506
Cuban	239	241	263	266	286	287
Other specific	756	832	920	1,182	1,264	1,241
Total	2,917	2,981	3,129	3,291	3,444	3,427
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2,353	1,156
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	97,031	59,377
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	99,384	60,533

n/c Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.16b. Kentucky individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	13.2	13.2	13.4	14.6	13.7	14.7
Anxiety disorders	9.0	9.4	10.0	13.4	11.5	11.4
Attention deficit/hyperactivity disorder	10.9	10.5	9.9	9.8	8.9	8.6
Mood disorders	41.6	42.0	42.0	38.0	29.6	29.7
Bipolar disorders	16.6	16.7	16.7	12.7	7.2	7.1
Depressive disorders	25.0	25.3	25.3	25.2	22.4	22.6
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	7.2	7.4	7.3	7.0	6.6	6.6
Alcohol or substance-related disorders	3.0	3.2	3.1	3.9	10.0	10.4
Other mental disorders ²	14.8	14.1	14.0	12.6	19.0	17.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	48.5	48.2	48.2	47.8	47.7	48.1
Female	51.5	51.8	51.8	52.2	52.3	51.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	20.7	19.9	18.7	16.8	16.7	16.9
12 to 14 years	8.8	8.8	8.4	8.1	8.0	8.1
15 to 17 years	8.2	8.3	8.2	8.2	8.3	8.2
18 to 20 years	4.2	4.1	4.2	4.3	4.4	4.7
21 to 24 years	5.7	5.7	5.8	5.9	5.8	5.8
25 to 29 years	7.1	7.3	7.5	8.1	8.3	8.4
30 to 34 years	7.5	7.6	7.7	7.8	7.8	7.8
35 to 39 years	6.5	6.8	7.2	7.6	7.6	7.5
40 to 44 years	6.7	6.6	6.7	6.7	6.4	6.3
45 to 49 years	7.0	6.8	6.7	6.8	6.6	6.3
50 to 54 years	6.6	6.8	6.9	7.0	6.8	6.4
55 to 59 years	5.0	5.2	5.3	5.6	5.5	5.5
60 to 64 years	3.0	3.1	3.2	3.4	3.6	3.6
65 years and older	3.0	3.1	3.6	3.7	4.1	4.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.16b. Kentucky individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	10.5	10.5	10.4	10.2	10.2	9.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	83.2	82.6	82.5	82.6	82.3	81.2
Other	5.9	6.6	6.7	6.6	6.9	8.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	98.2	98.1	98.0	97.8	97.7	97.8
Hispanic or Latino origin	1.8	1.9	2.0	2.2	2.3	2.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	48.6	46.2	44.6	40.8	40.7	40.6
Puerto Rican	17.3	17.8	17.6	15.2	14.3	14.8
Cuban	8.2	8.1	8.4	8.1	8.3	8.4
Other specific	25.9	27.9	29.4	35.9	36.7	36.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2.4	1.9
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	97.6	98.1
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.17a. Louisiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	51,153	45,373	43,831	34,508	49,302	39,413
Mental health diagnosis¹						
Trauma- and stressor-related disorders	1,131	1,365	1,374	967	1,500	1,127
Anxiety disorders	1,650	1,630	1,812	1,506	1,896	1,729
Attention deficit/hyperactivity disorder	3,432	3,522	3,282	2,045	2,188	1,993
Mood disorders	24,245	26,651	23,571	11,928	13,330	12,155
Bipolar disorders	9,275	10,245	9,176	4,519	4,233	3,559
Depressive disorders	14,970	16,406	14,395	7,409	9,097	8,596
Personality disorders	117	178	295	158	91	78
Schizophrenia and other psychotic disorders	8,355	8,933	7,956	4,972	4,962	4,631
Alcohol or substance-related disorders	--	--	--	--	--	--
Other mental disorders ²	1,881	1,935	2,693	1,990	1,590	1,287
Total	40,811	44,214	40,983	23,566	25,557	23,000
Gender						
Male	22,251	19,433	19,128	15,482	22,354	17,869
Female	27,938	24,882	23,927	18,746	26,641	21,438
Total	50,189	44,315	43,055	34,228	48,995	39,307
Age³						
0 to 11 years	4,011	3,219	2,997	2,603	3,208	2,320
12 to 14 years	2,340	2,087	1,946	1,547	2,142	1,551
15 to 17 years	2,359	1,954	1,973	1,435	2,232	1,644
18 to 20 years	1,887	1,669	1,581	1,214	1,855	1,379
21 to 24 years	2,935	2,757	2,542	2,031	2,974	2,195
25 to 29 years	4,605	4,036	3,877	3,076	4,606	3,664
30 to 34 years	4,660	4,239	4,107	3,304	4,948	3,953
35 to 39 years	4,128	3,792	3,754	3,115	4,745	3,925
40 to 44 years	4,300	3,687	3,493	2,718	3,868	3,226
45 to 49 years	5,329	4,607	4,109	2,998	4,171	3,346
50 to 54 years	5,904	5,153	4,924	3,697	4,920	4,025
55 to 59 years	4,611	4,171	4,240	3,359	4,665	3,860
60 to 64 years	2,437	2,370	2,522	2,020	2,900	2,566
65 years and older	1,521	1,632	1,766	1,390	2,067	1,759
Total	51,027	45,373	43,831	34,507	49,301	39,413

Continued. See notes at end of table.

Table 9.17a. Louisiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	268	272	246	221	368	350
Asian ⁴	154	130	149	121	140	120
Black or African American	21,093	18,491	19,156	15,056	20,907	16,553
Native Hawaiian or Other Pacific Islander	19	42	66	29	57	43
White	24,096	19,922	18,975	15,303	22,602	18,066
Other	1,058	631	562	415	658	1,916
Total	46,688	39,488	39,154	31,145	44,732	37,048
Ethnicity						
Not of Hispanic or Latino origin	43,395	36,465	36,374	30,471	44,419	36,270
Hispanic or Latino origin	927	661	789	839	1,245	1,033
Total	44,322	37,126	37,163	31,310	45,664	37,303
Specific Hispanic or Latino origin						
Mexican	113	14	18	38	68	43
Puerto Rican	57	11	14	19	36	26
Cuban	48	10	13	16	19	13
Other specific	42	14	17	32	55	44
Total	260	49	62	105	178	126
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	445	459
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	25,864	28,168
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	26,309	28,627

-- Quantity is zero; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.17b. Louisiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	2.8	3.1	3.4	4.1	5.9	4.9
Anxiety disorders	4.0	3.7	4.4	6.4	7.4	7.5
Attention deficit/hyperactivity disorder	8.4	8.0	8.0	8.7	8.6	8.7
Mood disorders	59.4	60.3	57.5	50.6	52.2	52.8
Bipolar disorders	22.7	23.2	22.4	19.2	16.6	15.5
Depressive disorders	36.7	37.1	35.1	31.4	35.6	37.4
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	20.5	20.2	19.4	21.1	19.4	20.1
Alcohol or substance-related disorders	--	--	--	--	--	--
Other mental disorders ²	4.6	4.4	6.6	8.4	6.2	5.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	44.3	43.9	44.4	45.2	45.6	45.5
Female	55.7	56.1	55.6	54.8	54.4	54.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	7.9	7.1	6.8	7.5	6.5	5.9
12 to 14 years	4.6	4.6	4.4	4.5	4.3	3.9
15 to 17 years	4.6	4.3	4.5	4.2	4.5	4.2
18 to 20 years	3.7	3.7	3.6	3.5	3.8	3.5
21 to 24 years	5.8	6.1	5.8	5.9	6.0	5.6
25 to 29 years	9.0	8.9	8.8	8.9	9.3	9.3
30 to 34 years	9.1	9.3	9.4	9.6	10.0	10.0
35 to 39 years	8.1	8.4	8.6	9.0	9.6	10.0
40 to 44 years	8.4	8.1	8.0	7.9	7.8	8.2
45 to 49 years	10.4	10.2	9.4	8.7	8.5	8.5
50 to 54 years	11.6	11.4	11.2	10.7	10.0	10.2
55 to 59 years	9.0	9.2	9.7	9.7	9.5	9.8
60 to 64 years	4.8	5.2	5.8	5.9	5.9	6.5
65 years and older	3.0	3.6	4.0	4.0	4.2	4.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.17b. Louisiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	45.2	46.8	48.9	48.3	46.7	44.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	51.6	50.5	48.5	49.1	50.5	48.8
Other	2.3	1.6	1.4	1.3	1.5	5.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	97.9	98.2	97.9	97.3	97.3	97.2
Hispanic or Latino origin	2.1	1.8	2.1	2.7	2.7	2.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	43.5	28.6	29.0	36.2	38.2	34.1
Puerto Rican	21.9	22.4	22.6	18.1	20.2	20.6
Cuban	18.5	20.4	21.0	15.2	10.7	10.3
Other specific	16.2	28.6	27.4	30.5	30.9	34.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	1.7	1.6
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	98.3	98.4
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.18a. Maine individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	71,236	‡	‡	‡	49,502	‡
Mental health diagnosis¹						
Trauma- and stressor-related disorders	15,961	‡	‡	‡	3,886	‡
Anxiety disorders	11,191	‡	‡	‡	2,881	‡
Attention deficit/hyperactivity disorder	6,652	‡	‡	‡	354	‡
Mood disorders	26,168	‡	‡	‡	10,049	‡
Bipolar disorders	9,903	‡	‡	‡	3,829	‡
Depressive disorders	16,265	‡	‡	‡	6,220	‡
Personality disorders	186	‡	‡	‡	16	‡
Schizophrenia and other psychotic disorders	3,563	‡	‡	‡	1,301	‡
Alcohol or substance-related disorders	1,039	‡	‡	‡	133	‡
Other mental disorders ²	6,012	‡	‡	‡	235	‡
Total	70,772	‡	‡	‡	18,855	‡
Gender						
Male	30,155	‡	‡	‡	17,394	‡
Female	41,081	‡	‡	‡	25,677	‡
Total	71,236	‡	‡	‡	43,071	‡
Age³						
0 to 11 years	12,514	‡	‡	‡	1	‡
12 to 14 years	5,325	‡	‡	‡	--	‡
15 to 17 years	5,207	‡	‡	‡	--	‡
18 to 20 years	3,580	‡	‡	‡	384	‡
21 to 24 years	3,529	‡	‡	‡	2,213	‡
25 to 29 years	5,866	‡	‡	‡	4,819	‡
30 to 34 years	6,411	‡	‡	‡	5,649	‡
35 to 39 years	5,585	‡	‡	‡	5,799	‡
40 to 44 years	5,249	‡	‡	‡	5,184	‡
45 to 49 years	5,082	‡	‡	‡	5,209	‡
50 to 54 years	4,547	‡	‡	‡	5,453	‡
55 to 59 years	3,419	‡	‡	‡	5,283	‡
60 to 64 years	2,073	‡	‡	‡	4,016	‡
65 years and older	2,848	‡	‡	‡	5,488	‡
Total	71,235	‡	‡	‡	49,498	‡

Continued. See notes at end of table.

Table 9.18a. Maine individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	831	‡	‡	‡	18	‡
Asian ⁴	210	‡	‡	‡	2	‡
Black or African American	1,417	‡	‡	‡	8	‡
Native Hawaiian or Other Pacific Islander	31	‡	‡	‡	--	‡
White	61,311	‡	‡	‡	1,074	‡
Other	1,541	‡	‡	‡	10	‡
Total	65,341	‡	‡	‡	1,112	‡
Ethnicity						
Not of Hispanic or Latino origin	65,336	‡	‡	‡	29	‡
Hispanic or Latino origin	147	‡	‡	‡	17	‡
Total	65,483	‡	‡	‡	46	‡
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.18b. Maine individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	‡	‡	‡	100.0	‡
Mental health diagnosis¹						
Trauma- and stressor-related disorders	22.6	‡	‡	‡	20.6	‡
Anxiety disorders	15.8	‡	‡	‡	15.3	‡
Attention deficit/hyperactivity disorder	9.4	‡	‡	‡	1.9	‡
Mood disorders	37.0	‡	‡	‡	53.3	‡
Bipolar disorders	14.0	‡	‡	‡	20.3	‡
Depressive disorders	23.0	‡	‡	‡	33.0	‡
Personality disorders	†	‡	‡	‡	†	‡
Schizophrenia and other psychotic disorders	5.0	‡	‡	‡	6.9	‡
Alcohol or substance-related disorders	1.5	‡	‡	‡	†	‡
Other mental disorders ²	8.5	‡	‡	‡	1.2	‡
Total	100.0	‡	‡	‡	100.0	‡
Gender						
Male	42.3	‡	‡	‡	40.4	‡
Female	57.7	‡	‡	‡	59.6	‡
Total	100.0	‡	‡	‡	100.0	‡
Age³						
0 to 11 years	17.6	‡	‡	‡	†	‡
12 to 14 years	7.5	‡	‡	‡	--	‡
15 to 17 years	7.3	‡	‡	‡	--	‡
18 to 20 years	5.0	‡	‡	‡	†	‡
21 to 24 years	5.0	‡	‡	‡	4.5	‡
25 to 29 years	8.2	‡	‡	‡	9.7	‡
30 to 34 years	9.0	‡	‡	‡	11.4	‡
35 to 39 years	7.8	‡	‡	‡	11.7	‡
40 to 44 years	7.4	‡	‡	‡	10.5	‡
45 to 49 years	7.1	‡	‡	‡	10.5	‡
50 to 54 years	6.4	‡	‡	‡	11.0	‡
55 to 59 years	4.8	‡	‡	‡	10.7	‡
60 to 64 years	2.9	‡	‡	‡	8.1	‡
65 years and older	4.0	‡	‡	‡	11.1	‡
Total	100.0	‡	‡	‡	100.0	‡

Continued. See notes at end of table.

Table 9.18b. Maine individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	1.3	‡	‡	‡	1.6	‡
Asian ⁴	†	‡	‡	‡	†	‡
Black or African American	2.2	‡	‡	‡	†	‡
Native Hawaiian or Other Pacific Islander	†	‡	‡	‡	--	‡
White	93.8	‡	‡	‡	96.6	‡
Other	2.4	‡	‡	‡	†	‡
Total	100.0	‡	‡	‡	100.0	‡
Ethnicity						
Not of Hispanic or Latino origin	99.8	‡	‡	‡	63.0	‡
Hispanic or Latino origin	†	‡	‡	‡	37.0	‡
Total	100.0	‡	‡	‡	100.0	‡
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.19a. Maryland individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	153,768	165,346	185,306	202,972	201,718	213,605
Mental health diagnosis¹						
Trauma- and stressor-related disorders	23,683	25,052	27,598	28,087	31,136	33,567
Anxiety disorders	13,115	15,541	20,474	23,610	27,138	29,107
Attention deficit/hyperactivity disorder	21,017	22,585	24,326	26,093	27,773	28,739
Mood disorders	70,667	76,073	80,504	72,280	77,798	82,670
Bipolar disorders	31,847	33,179	29,887	23,719	24,846	26,331
Depressive disorders	38,820	42,894	50,617	48,561	52,952	56,339
Personality disorders	266	325	410	334	383	498
Schizophrenia and other psychotic disorders	13,870	14,479	15,041	15,451	15,869	16,323
Alcohol or substance-related disorders	69	128	130	24	50	--
Other mental disorders ²	10,815	10,798	11,791	18,979	20,124	20,945
Total	153,502	164,981	180,274	184,858	200,271	211,849
Gender						
Male	71,351	76,717	85,362	91,881	92,962	98,289
Female	82,414	88,620	99,938	111,078	108,740	115,299
Total	153,765	165,337	185,300	202,959	201,702	213,588
Age³						
0 to 11 years	28,621	29,796	31,157	32,416	33,966	37,514
12 to 14 years	14,358	15,014	15,697	15,978	16,483	17,841
15 to 17 years	13,191	13,734	15,039	15,845	16,258	16,690
18 to 20 years	8,507	8,155	8,710	9,570	9,814	10,549
21 to 24 years	8,578	9,353	10,750	12,188	11,500	12,056
25 to 29 years	12,817	14,281	17,355	20,000	19,583	20,957
30 to 34 years	12,312	13,905	16,288	18,164	17,948	19,213
35 to 39 years	9,660	10,728	12,823	14,663	14,928	15,901
40 to 44 years	10,327	10,895	11,655	12,243	11,532	12,035
45 to 49 years	11,459	12,027	13,183	14,201	13,033	13,050
50 to 54 years	11,011	12,273	14,123	15,675	14,492	14,463
55 to 59 years	7,298	8,642	10,429	12,414	12,055	12,464
60 to 64 years	3,889	4,688	5,937	7,126	7,285	7,688
65 years and older	1,738	1,855	2,158	2,488	2,841	3,180
Total	153,766	165,346	185,304	202,971	201,718	213,601

Continued. See notes at end of table.

Table 9.19a. Maryland individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	537	580	728	848	825	948
Asian ⁴	1,400	1,715	2,050	2,451	2,505	2,866
Black or African American	74,929	79,178	86,724	93,644	92,425	98,376
Native Hawaiian or Other Pacific Islander	206	260	327	424	435	583
White	71,883	78,126	87,824	95,425	94,896	100,151
Other	2,287	2,486	3,088	3,578	4,013	4,472
Total	151,242	162,345	180,741	196,370	195,099	207,396
Ethnicity						
Not of Hispanic or Latino origin	132,713	141,882	157,034	166,808	169,942	180,626
Hispanic or Latino origin	12,824	14,138	16,336	18,259	19,971	22,705
Total	145,537	156,020	173,370	185,067	189,913	203,331
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	3,862	4,071
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	109,313	129,200
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	113,175	133,271

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.19b. Maryland individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	15.4	15.2	15.3	15.2	15.5	15.8
Anxiety disorders	8.5	9.4	11.4	12.8	13.6	13.7
Attention deficit/hyperactivity disorder	13.7	13.7	13.5	14.1	13.9	13.6
Mood disorders	46.0	46.1	44.7	39.1	38.8	39.0
Bipolar disorders	20.7	20.1	16.6	12.8	12.4	12.4
Depressive disorders	25.3	26.0	28.1	26.3	26.4	26.6
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	9.0	8.8	8.3	8.4	7.9	7.7
Alcohol or substance-related disorders	†	†	†	†	†	--
Other mental disorders ²	7.0	6.5	6.5	10.3	10.0	9.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	46.4	46.4	46.1	45.3	46.1	46.0
Female	53.6	53.6	53.9	54.7	53.9	54.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	18.6	18.0	16.8	16.0	16.8	17.6
12 to 14 years	9.3	9.1	8.5	7.9	8.2	8.4
15 to 17 years	8.6	8.3	8.1	7.8	8.1	7.8
18 to 20 years	5.5	4.9	4.7	4.7	4.9	4.9
21 to 24 years	5.6	5.7	5.8	6.0	5.7	5.6
25 to 29 years	8.3	8.6	9.4	9.9	9.7	9.8
30 to 34 years	8.0	8.4	8.8	8.9	8.9	9.0
35 to 39 years	6.3	6.5	6.9	7.2	7.4	7.4
40 to 44 years	6.7	6.6	6.3	6.0	5.7	5.6
45 to 49 years	7.5	7.3	7.1	7.0	6.5	6.1
50 to 54 years	7.2	7.4	7.6	7.7	7.2	6.8
55 to 59 years	4.7	5.2	5.6	6.1	6.0	5.8
60 to 64 years	2.5	2.8	3.2	3.5	3.6	3.6
65 years and older	1.1	1.1	1.2	1.2	1.4	1.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.19b. Maryland individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	1.1	1.1	1.2	1.3	1.4
Black or African American	49.5	48.8	48.0	47.7	47.4	47.4
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	47.5	48.1	48.6	48.6	48.6	48.3
Other	1.5	1.5	1.7	1.8	2.1	2.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	91.2	90.9	90.6	90.1	89.5	88.8
Hispanic or Latino origin	8.8	9.1	9.4	9.9	10.5	11.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	3.4	3.1
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	96.6	96.9
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.20a. Massachusetts individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	29,226	32,046	33,556	33,921	31,659	26,383
Mental health diagnosis¹						
Trauma- and stressor-related disorders	2,523	2,659	2,833	1,425	1,776	712
Anxiety disorders	851	1,035	1,162	662	971	827
Attention deficit/hyperactivity disorder	768	850	909	245	400	318
Mood disorders	10,654	11,823	12,460	3,597	4,978	3,517
Bipolar disorders	6,482	6,894	7,383	1,543	2,062	1,656
Depressive disorders	4,172	4,929	5,077	2,054	2,916	1,861
Personality disorders	1,485	1,575	1,537	2	2	31
Schizophrenia and other psychotic disorders	10,360	10,842	10,934	2,877	3,721	3,322
Alcohol or substance-related disorders	--	--	--	--	--	--
Other mental disorders ²	1,208	1,241	1,187	1,243	1,320	521
Total	27,849	30,025	31,022	10,051	13,168	9,248
Gender						
Male	16,307	17,959	18,712	18,948	17,680	14,761
Female	12,918	14,086	14,843	14,973	13,979	11,622
Total	29,225	32,045	33,555	33,921	31,659	26,383
Age³						
0 to 11 years	1,392	1,722	1,965	1,925	1,681	1,266
12 to 14 years	1,187	1,283	1,469	1,614	1,394	984
15 to 17 years	1,510	1,619	1,866	1,821	1,619	1,010
18 to 20 years	1,324	1,392	1,431	1,514	1,386	1,022
21 to 24 years	2,203	2,271	2,245	2,141	1,957	1,549
25 to 29 years	2,662	2,915	2,994	3,153	2,849	2,180
30 to 34 years	2,433	2,619	2,799	2,876	2,674	2,149
35 to 39 years	2,017	2,230	2,417	2,495	2,437	2,008
40 to 44 years	2,351	2,460	2,385	2,280	2,065	1,758
45 to 49 years	2,956	3,043	2,940	2,861	2,606	2,119
50 to 54 years	3,368	3,651	3,710	3,583	3,317	2,776
55 to 59 years	2,822	3,291	3,405	3,484	3,323	3,094
60 to 64 years	1,708	2,043	2,240	2,353	2,419	2,414
65 years and older	1,293	1,507	1,689	1,821	1,932	2,052
Total	29,226	32,046	33,555	33,921	31,659	26,381

Continued. See notes at end of table.

**Table 9.20a. Massachusetts individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018 (continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	90	110	105	104	98	67
Asian ⁴	617	640	633	623	624	594
Black or African American	3,759	4,195	4,390	4,572	4,401	3,750
Native Hawaiian or Other Pacific Islander	13	16	20	19	12	10
White	20,485	22,142	22,555	22,366	20,703	16,409
Other	460	541	594	597	517	364
Total	25,424	27,644	28,297	28,281	26,355	21,194
Ethnicity						
Not of Hispanic or Latino origin	12,211	14,662	14,993	24,496	22,394	17,477
Hispanic or Latino origin	5,587	2,872	3,164	2,947	2,247	2,037
Total	17,798	17,534	18,157	27,443	24,641	19,514
Specific Hispanic or Latino origin						
Mexican	19	27	18	20	18	17
Puerto Rican	921	972	1,025	883	1,002	812
Cuban	--	--	--	--	--	--
Other specific	340	377	448	1,382	425	410
Total	1,280	1,376	1,491	2,285	1,445	1,239
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	674	569
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	21,387	19,259
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	22,061	19,828

n/c Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.20b. Massachusetts individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	9.1	8.9	9.1	14.2	13.5	7.7
Anxiety disorders	3.1	3.4	3.7	6.6	7.4	8.9
Attention deficit/hyperactivity disorder	2.8	2.8	2.9	2.4	3.0	3.4
Mood disorders	38.3	39.4	40.2	35.8	37.8	38.0
Bipolar disorders	23.3	23.0	23.8	15.4	15.7	17.9
Depressive disorders	15.0	16.4	16.4	20.4	22.1	20.1
Personality disorders	5.3	5.2	5.0	†	†	†
Schizophrenia and other psychotic disorders	37.2	36.1	35.2	28.6	28.3	35.9
Alcohol or substance-related disorders	--	--	--	--	--	--
Other mental disorders ²	4.3	4.1	3.8	12.4	10.0	5.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	55.8	56.0	55.8	55.9	55.8	55.9
Female	44.2	44.0	44.2	44.1	44.2	44.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	4.8	5.4	5.9	5.7	5.3	4.8
12 to 14 years	4.1	4.0	4.4	4.8	4.4	3.7
15 to 17 years	5.2	5.1	5.6	5.4	5.1	3.8
18 to 20 years	4.5	4.3	4.3	4.5	4.4	3.9
21 to 24 years	7.5	7.1	6.7	6.3	6.2	5.9
25 to 29 years	9.1	9.1	8.9	9.3	9.0	8.3
30 to 34 years	8.3	8.2	8.3	8.5	8.4	8.1
35 to 39 years	6.9	7.0	7.2	7.4	7.7	7.6
40 to 44 years	8.0	7.7	7.1	6.7	6.5	6.7
45 to 49 years	10.1	9.5	8.8	8.4	8.2	8.0
50 to 54 years	11.5	11.4	11.1	10.6	10.5	10.5
55 to 59 years	9.7	10.3	10.1	10.3	10.5	11.7
60 to 64 years	5.8	6.4	6.7	6.9	7.6	9.2
65 years and older	4.4	4.7	5.0	5.4	6.1	7.8
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.20b. Massachusetts individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018 (continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	2.4	2.3	2.2	2.2	2.4	2.8
Black or African American	14.8	15.2	15.5	16.2	16.7	17.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	80.6	80.1	79.7	79.1	78.6	77.4
Other	1.8	2.0	2.1	2.1	2.0	1.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	68.6	83.6	82.6	89.3	90.9	89.6
Hispanic or Latino origin	31.4	16.4	17.4	10.7	9.1	10.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	1.5	2.0	1.2	†	1.2	1.4
Puerto Rican	72.0	70.6	68.7	38.6	69.3	65.5
Cuban	--	--	--	--	--	--
Other specific	26.6	27.4	30.0	60.5	29.4	33.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	3.1	2.9
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	96.9	97.1
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.21a. Michigan individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	‡	‡	‡	‡	177,289	231,953
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	‡	‡	12,110	18,469
Anxiety disorders	‡	‡	‡	‡	9,657	13,515
Attention deficit/hyperactivity disorder	‡	‡	‡	‡	9,150	12,034
Mood disorders	‡	‡	‡	‡	60,753	81,840
Bipolar disorders	‡	‡	‡	‡	22,555	28,423
Depressive disorders	‡	‡	‡	‡	38,198	53,417
Personality disorders	‡	‡	‡	‡	961	1,302
Schizophrenia and other psychotic disorders	‡	‡	‡	‡	25,726	30,743
Alcohol or substance-related disorders	‡	‡	‡	‡	1,730	2,078
Other mental disorders ²	‡	‡	‡	‡	50,836	64,586
Total	‡	‡	‡	‡	170,923	224,567
Gender						
Male	‡	‡	‡	‡	92,096	119,663
Female	‡	‡	‡	‡	85,193	112,289
Total	‡	‡	‡	‡	177,289	231,952
Age³						
0 to 11 years	‡	‡	‡	‡	21,897	30,571
12 to 14 years	‡	‡	‡	‡	9,993	13,294
15 to 17 years	‡	‡	‡	‡	10,614	14,834
18 to 20 years	‡	‡	‡	‡	7,745	10,658
21 to 24 years	‡	‡	‡	‡	11,696	14,616
25 to 29 years	‡	‡	‡	‡	15,877	21,425
30 to 34 years	‡	‡	‡	‡	14,593	19,178
35 to 39 years	‡	‡	‡	‡	13,430	17,645
40 to 44 years	‡	‡	‡	‡	12,336	15,442
45 to 49 years	‡	‡	‡	‡	13,870	17,178
50 to 54 years	‡	‡	‡	‡	15,192	18,411
55 to 59 years	‡	‡	‡	‡	13,540	17,082
60 to 64 years	‡	‡	‡	‡	8,654	11,208
65 years and older	‡	‡	‡	‡	7,850	10,410
Total	‡	‡	‡	‡	177,287	231,952

Continued. See notes at end of table.

Table 9.21a. Michigan individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	‡	‡	‡	‡	1,129	1,411
Asian ⁴	‡	‡	‡	‡	755	974
Black or African American	‡	‡	‡	‡	34,396	46,251
Native Hawaiian or Other Pacific Islander	‡	‡	‡	‡	95	140
White	‡	‡	‡	‡	107,072	137,532
Other	‡	‡	‡	‡	32,534	43,811
Total	‡	‡	‡	‡	175,981	230,119
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	‡	151,247	198,513
Hispanic or Latino origin	‡	‡	‡	‡	9,322	13,498
Total	‡	‡	‡	‡	160,569	212,011
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	2,087	3,350
Puerto Rican	‡	‡	‡	‡	665	992
Cuban	‡	‡	‡	‡	98	150
Other specific	‡	‡	‡	‡	1,082	1,623
Total	‡	‡	‡	‡	3,932	6,115
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2,975	3,864
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	123,951	159,920
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	126,926	163,784

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.21b. Michigan individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	‡	‡	‡	‡	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	‡	‡	7.1	8.2
Anxiety disorders	‡	‡	‡	‡	5.6	6.0
Attention deficit/hyperactivity disorder	‡	‡	‡	‡	5.4	5.4
Mood disorders	‡	‡	‡	‡	35.5	36.4
Bipolar disorders	‡	‡	‡	‡	13.2	12.7
Depressive disorders	‡	‡	‡	‡	22.3	23.8
Personality disorders	‡	‡	‡	‡	†	†
Schizophrenia and other psychotic disorders	‡	‡	‡	‡	15.1	13.7
Alcohol or substance-related disorders	‡	‡	‡	‡	1.0	†
Other mental disorders ²	‡	‡	‡	‡	29.7	28.8
Total	‡	‡	‡	‡	100.0	100.0
Gender						
Male	‡	‡	‡	‡	51.9	51.6
Female	‡	‡	‡	‡	48.1	48.4
Total	‡	‡	‡	‡	100.0	100.0
Age³						
0 to 11 years	‡	‡	‡	‡	12.4	13.2
12 to 14 years	‡	‡	‡	‡	5.6	5.7
15 to 17 years	‡	‡	‡	‡	6.0	6.4
18 to 20 years	‡	‡	‡	‡	4.4	4.6
21 to 24 years	‡	‡	‡	‡	6.6	6.3
25 to 29 years	‡	‡	‡	‡	9.0	9.2
30 to 34 years	‡	‡	‡	‡	8.2	8.3
35 to 39 years	‡	‡	‡	‡	7.6	7.6
40 to 44 years	‡	‡	‡	‡	7.0	6.7
45 to 49 years	‡	‡	‡	‡	7.8	7.4
50 to 54 years	‡	‡	‡	‡	8.6	7.9
55 to 59 years	‡	‡	‡	‡	7.6	7.4
60 to 64 years	‡	‡	‡	‡	4.9	4.8
65 years and older	‡	‡	‡	‡	4.4	4.5
Total	‡	‡	‡	‡	100.0	100.0

Continued. See notes at end of table.

Table 9.21b. Michigan individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	‡	‡	‡	‡	†	†
Asian ⁴	‡	‡	‡	‡	†	†
Black or African American	‡	‡	‡	‡	19.5	20.1
Native Hawaiian or Other Pacific Islander	‡	‡	‡	‡	†	†
White	‡	‡	‡	‡	60.8	59.8
Other	‡	‡	‡	‡	18.5	19.0
Total	‡	‡	‡	‡	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	‡	94.2	93.6
Hispanic or Latino origin	‡	‡	‡	‡	5.8	6.4
Total	‡	‡	‡	‡	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	53.1	54.8
Puerto Rican	‡	‡	‡	‡	16.9	16.2
Cuban	‡	‡	‡	‡	2.5	2.5
Other specific	‡	‡	‡	‡	27.5	26.5
Total	‡	‡	‡	‡	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2.3	2.4
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	97.7	97.6
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.22a. Minnesota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	‡	205,841	225,786	259,024	270,651	287,962
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	39,250	44,738	51,213	52,900	59,657
Anxiety disorders	‡	28,571	34,963	43,596	58,572	61,843
Attention deficit/hyperactivity disorder	‡	20,054	21,940	24,463	25,165	23,686
Mood disorders	‡	66,940	70,511	70,565	60,262	73,746
Bipolar disorders	‡	16,082	15,949	14,357	11,982	15,009
Depressive disorders	‡	50,858	54,562	56,208	48,280	58,737
Personality disorders	‡	3,295	3,718	4,850	5,902	4,578
Schizophrenia and other psychotic disorders	‡	14,810	14,955	14,932	15,314	17,509
Alcohol or substance-related disorders	‡	323	335	318	36	114
Other mental disorders ²	‡	23,625	24,188	28,826	27,021	28,289
Total	‡	196,868	215,348	238,763	245,172	269,422
Gender						
Male	‡	93,739	102,433	117,157	121,791	129,777
Female	‡	112,101	123,353	141,867	148,860	158,185
Total	‡	205,840	225,786	259,024	270,651	287,962
Age³						
0 to 11 years	‡	29,523	31,867	40,303	41,822	45,166
12 to 14 years	‡	12,545	13,815	16,738	17,271	18,709
15 to 17 years	‡	12,452	13,619	16,552	17,471	18,926
18 to 20 years	‡	8,713	9,267	10,513	11,327	12,473
21 to 24 years	‡	12,479	13,247	14,912	15,082	15,957
25 to 29 years	‡	18,376	20,822	23,935	25,195	26,871
30 to 34 years	‡	18,702	20,749	23,376	24,306	25,603
35 to 39 years	‡	15,460	17,274	19,796	21,348	22,870
40 to 44 years	‡	14,851	15,905	17,030	17,399	17,991
45 to 49 years	‡	15,426	16,263	17,361	17,875	18,121
50 to 54 years	‡	16,514	17,998	19,220	19,100	19,098
55 to 59 years	‡	12,378	14,424	16,177	17,317	18,340
60 to 64 years	‡	7,937	9,502	10,930	11,817	12,911
65 years and older	‡	10,468	11,017	12,128	13,278	14,903
Total	‡	205,824	225,769	258,971	270,608	287,939

Continued. See notes at end of table.

Table 9.22a. Minnesota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	‡	8,368	8,988	9,702	9,861	10,566
Asian ⁴	‡	5,957	6,497	7,326	7,891	8,546
Black or African American	‡	28,380	29,328	32,494	33,622	35,148
Native Hawaiian or Other Pacific Islander	‡	184	216	244	278	301
White	‡	134,622	145,665	161,668	165,514	172,627
Other	‡	7,204	8,365	10,230	11,305	12,629
Total	‡	184,715	199,059	221,664	228,471	239,817
Ethnicity						
Not of Hispanic or Latino origin	‡	192,494	209,900	239,397	249,536	248,875
Hispanic or Latino origin	‡	11,191	12,954	15,378	16,244	16,618
Total	‡	203,685	222,854	254,775	265,780	265,493
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.22b. Minnesota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	‡	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	19.9	20.8	21.4	21.6	22.1
Anxiety disorders	‡	14.5	16.2	18.3	23.9	23.0
Attention deficit/hyperactivity disorder	‡	10.2	10.2	10.2	10.3	8.8
Mood disorders	‡	34.0	32.7	29.6	24.6	27.4
Bipolar disorders	‡	8.2	7.4	6.0	4.9	5.6
Depressive disorders	‡	25.8	25.3	23.5	19.7	21.8
Personality disorders	‡	1.7	1.7	2.0	2.4	1.7
Schizophrenia and other psychotic disorders	‡	7.5	6.9	6.3	6.2	6.5
Alcohol or substance-related disorders	‡	†	†	†	†	†
Other mental disorders ²	‡	12.0	11.2	12.1	11.0	10.5
Total	‡	100.0	100.0	100.0	100.0	100.0
Gender						
Male	‡	45.5	45.4	45.2	45.0	45.1
Female	‡	54.5	54.6	54.8	55.0	54.9
Total	‡	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	‡	14.3	14.1	15.6	15.5	15.7
12 to 14 years	‡	6.1	6.1	6.5	6.4	6.5
15 to 17 years	‡	6.0	6.0	6.4	6.5	6.6
18 to 20 years	‡	4.2	4.1	4.1	4.2	4.3
21 to 24 years	‡	6.1	5.9	5.8	5.6	5.5
25 to 29 years	‡	8.9	9.2	9.2	9.3	9.3
30 to 34 years	‡	9.1	9.2	9.0	9.0	8.9
35 to 39 years	‡	7.5	7.7	7.6	7.9	7.9
40 to 44 years	‡	7.2	7.0	6.6	6.4	6.2
45 to 49 years	‡	7.5	7.2	6.7	6.6	6.3
50 to 54 years	‡	8.0	8.0	7.4	7.1	6.6
55 to 59 years	‡	6.0	6.4	6.2	6.4	6.4
60 to 64 years	‡	3.9	4.2	4.2	4.4	4.5
65 years and older	‡	5.1	4.9	4.7	4.9	5.2
Total	‡	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.22b. Minnesota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	‡	4.5	4.5	4.4	4.3	4.4
Asian ⁴	‡	3.2	3.3	3.3	3.5	3.6
Black or African American	‡	15.4	14.7	14.7	14.7	14.7
Native Hawaiian or Other Pacific Islander	‡	†	†	†	†	†
White	‡	72.9	73.2	72.9	72.4	72.0
Other	‡	3.9	4.2	4.6	4.9	5.3
Total	‡	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	‡	94.5	94.2	94.0	93.9	93.7
Hispanic or Latino origin	‡	5.5	5.8	6.0	6.1	6.3
Total	‡	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.23a. Mississippi individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	87,379	85,962	90,279	83,432	73,867	85,449
Mental health diagnosis¹						
Trauma- and stressor-related disorders	7,584	5,024	6,776	6,158	5,742	7,577
Anxiety disorders	5,562	3,306	5,460	12,746	4,384	6,237
Attention deficit/hyperactivity disorder	13,292	7,061	13,535	13,398	10,599	13,776
Mood disorders	34,032	17,949	32,805	28,564	21,629	29,058
Bipolar disorders	11,876	6,142	11,155	9,840	6,692	8,855
Depressive disorders	22,156	11,807	21,650	18,724	14,937	20,203
Personality disorders	160	119	168	168	279	300
Schizophrenia and other psychotic disorders	10,682	4,175	11,920	10,439	7,565	10,365
Alcohol or substance-related disorders	4,183	2,881	2,644	2,327	2,396	6,773
Other mental disorders ²	11,275	6,704	9,811	7,327	9,284	10,832
Total	86,770	47,219	83,119	81,127	61,878	84,918
Gender						
Male	41,500	40,383	42,319	35,434	35,345	41,517
Female	45,860	45,534	47,922	40,366	38,479	43,874
Total	87,360	85,917	90,241	75,800	73,824	85,391
Age³						
0 to 11 years	12,745	15,166	14,147	13,410	13,900	15,515
12 to 14 years	7,192	7,560	7,562	6,969	6,624	7,569
15 to 17 years	7,381	7,088	7,829	7,248	6,454	7,053
18 to 20 years	4,954	3,652	4,655	4,486	3,362	3,990
21 to 24 years	4,319	4,148	4,312	3,905	3,225	4,013
25 to 29 years	5,553	5,495	5,863	5,259	4,759	6,248
30 to 34 years	6,123	5,936	6,346	5,659	4,857	6,227
35 to 39 years	5,654	5,412	6,008	5,615	4,942	6,230
40 to 44 years	5,739	5,523	5,638	5,126	4,176	5,100
45 to 49 years	6,098	5,935	6,014	5,395	4,595	5,137
50 to 54 years	6,833	6,593	6,969	6,247	4,969	5,394
55 to 59 years	5,925	5,734	6,252	5,784	4,872	5,312
60 to 64 years	3,706	3,531	4,006	3,861	3,382	3,741
65 years and older	5,157	4,188	4,678	4,468	3,749	3,920
Total	87,379	85,961	90,279	83,432	73,866	85,449

Continued. See notes at end of table.

Table 9.23a. Mississippi individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	197	178	176	131	142	212
Asian ⁴	143	125	136	125	108	145
Black or African American	41,259	43,645	46,397	37,101	38,963	42,637
Native Hawaiian or Other Pacific Islander	8	6	9	10	7	12
White	44,461	40,695	42,191	37,030	33,050	40,256
Other	934	933	928	840	913	1,218
Total	87,002	85,582	89,837	75,237	73,183	84,480
Ethnicity						
Not of Hispanic or Latino origin	86,081	84,675	88,809	73,447	69,903	79,507
Hispanic or Latino origin	806	779	838	868	1,102	1,341
Total	86,887	85,454	89,647	74,315	71,005	80,848
Specific Hispanic or Latino origin						
Mexican	408	371	398	391	412	539
Puerto Rican	84	79	86	78	72	81
Cuban	48	37	40	26	19	28
Other specific	--	--	--	--	--	--
Total	540	487	524	495	503	648
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	942	1,174
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	44,769	52,421
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	45,711	53,595

-- Quantity is zero; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.23b. Mississippi individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	8.7	10.6	8.2	7.6	9.3	8.9
Anxiety disorders	6.4	7.0	6.6	15.7	7.1	7.3
Attention deficit/hyperactivity disorder	15.3	15.0	16.3	16.5	17.1	16.2
Mood disorders	39.2	38.0	39.5	35.2	35.0	34.2
Bipolar disorders	13.7	13.0	13.4	12.1	10.8	10.4
Depressive disorders	25.5	25.0	26.0	23.1	24.1	23.8
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	12.3	8.8	14.3	12.9	12.2	12.2
Alcohol or substance-related disorders	4.8	6.1	3.2	2.9	3.9	8.0
Other mental disorders ²	13.0	14.2	11.8	9.0	15.0	12.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.5	47.0	46.9	46.7	47.9	48.6
Female	52.5	53.0	53.1	53.3	52.1	51.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	14.6	17.6	15.7	16.1	18.8	18.2
12 to 14 years	8.2	8.8	8.4	8.4	9.0	8.9
15 to 17 years	8.4	8.2	8.7	8.7	8.7	8.3
18 to 20 years	5.7	4.2	5.2	5.4	4.6	4.7
21 to 24 years	4.9	4.8	4.8	4.7	4.4	4.7
25 to 29 years	6.4	6.4	6.5	6.3	6.4	7.3
30 to 34 years	7.0	6.9	7.0	6.8	6.6	7.3
35 to 39 years	6.5	6.3	6.7	6.7	6.7	7.3
40 to 44 years	6.6	6.4	6.2	6.1	5.7	6.0
45 to 49 years	7.0	6.9	6.7	6.5	6.2	6.0
50 to 54 years	7.8	7.7	7.7	7.5	6.7	6.3
55 to 59 years	6.8	6.7	6.9	6.9	6.6	6.2
60 to 64 years	4.2	4.1	4.4	4.6	4.6	4.4
65 years and older	5.9	4.9	5.2	5.4	5.1	4.6
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.23b. Mississippi individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	47.4	51.0	51.6	49.3	53.2	50.5
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	51.1	47.6	47.0	49.2	45.2	47.7
Other	1.1	1.1	1.0	1.1	1.2	1.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	99.1	99.1	99.1	98.8	98.4	98.3
Hispanic or Latino origin	†	†	†	1.2	1.6	1.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	75.6	76.2	76.0	79.0	81.9	83.2
Puerto Rican	15.6	16.2	16.4	15.8	14.3	12.5
Cuban	8.9	7.6	7.6	5.3	3.8	4.3
Other specific	--	--	--	--	--	--
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2.1	2.2
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	97.9	97.8
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.24a. Missouri individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	75,707	74,670	75,472	76,269	77,497	76,409
Mental health diagnosis¹						
Trauma- and stressor-related disorders	6,209	6,154	6,491	1,846	2,510	3,028
Anxiety disorders	4,540	4,546	5,258	8,101	9,169	9,068
Attention deficit/hyperactivity disorder	5,657	5,503	5,511	5,898	6,835	6,764
Mood disorders	39,399	38,800	38,038	34,374	36,361	35,265
Bipolar disorders	18,144	17,578	17,094	14,021	14,105	12,909
Depressive disorders	21,255	21,222	20,944	20,353	22,256	22,356
Personality disorders	415	386	418	540	755	797
Schizophrenia and other psychotic disorders	13,745	13,471	13,419	13,138	13,563	13,187
Alcohol or substance-related disorders	2	2	4	--	--	--
Other mental disorders ²	2,954	3,006	3,180	4,910	4,828	5,019
Total	72,921	71,868	72,319	68,807	74,021	73,128
Gender						
Male	35,777	35,355	35,439	36,037	36,780	36,584
Female	39,930	39,315	40,033	40,232	40,717	39,825
Total	75,707	74,670	75,472	76,269	77,497	76,409
Age³						
0 to 11 years	7,083	7,009	7,145	7,600	8,049	8,138
12 to 14 years	4,293	4,344	4,444	4,594	4,660	4,873
15 to 17 years	4,063	3,966	4,264	4,497	4,600	4,290
18 to 20 years	2,554	2,427	2,387	2,495	2,650	2,661
21 to 24 years	4,074	3,758	3,515	3,562	3,526	3,382
25 to 29 years	6,247	5,946	5,788	5,677	5,693	5,587
30 to 34 years	6,971	6,902	6,637	6,450	6,535	6,136
35 to 39 years	6,348	6,392	6,489	6,628	6,821	6,534
40 to 44 years	6,818	6,607	6,549	6,163	6,045	5,868
45 to 49 years	7,712	7,234	7,044	6,937	6,869	6,644
50 to 54 years	8,253	8,255	8,496	8,280	8,004	7,448
55 to 59 years	5,944	6,187	6,566	6,796	6,956	7,243
60 to 64 years	3,182	3,335	3,693	3,955	4,269	4,620
65 years and older	2,164	2,308	2,455	2,635	2,820	2,985
Total	75,706	74,670	75,472	76,269	77,497	76,409

Continued. See notes at end of table.

Table 9.24a. Missouri individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	202	176	180	193	194	172
Asian ⁴	189	170	182	182	207	224
Black or African American	16,343	15,589	15,628	15,504	16,044	16,350
Native Hawaiian or Other Pacific Islander	36	27	30	37	39	32
White	55,185	55,106	55,429	56,105	56,370	54,880
Other	1,846	1,849	2,119	2,406	2,760	2,859
Total	73,801	72,917	73,568	74,427	75,614	74,517
Ethnicity						
Not of Hispanic or Latino origin	74,135	73,075	73,784	74,508	75,623	74,516
Hispanic or Latino origin	1,557	1,573	1,676	1,753	1,863	1,891
Total	75,692	74,648	75,460	76,261	77,486	76,407
Specific Hispanic or Latino origin						
Mexican	713	715	905	992	1,055	1,039
Puerto Rican	203	198	196	210	192	174
Cuban	48	46	56	55	52	55
Other specific	571	598	501	478	547	603
Total	1,535	1,557	1,658	1,735	1,846	1,871
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.24b. Missouri individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	8.5	8.6	9.0	2.7	3.4	4.1
Anxiety disorders	6.2	6.3	7.3	11.8	12.4	12.4
Attention deficit/hyperactivity disorder	7.8	7.7	7.6	8.6	9.2	9.2
Mood disorders	54.0	54.0	52.6	50.0	49.1	48.2
Bipolar disorders	24.9	24.5	23.6	20.4	19.1	17.7
Depressive disorders	29.1	29.5	29.0	29.6	30.1	30.6
Personality disorders	†	†	†	†	1.0	1.1
Schizophrenia and other psychotic disorders	18.8	18.7	18.6	19.1	18.3	18.0
Alcohol or substance-related disorders	†	†	†	--	--	--
Other mental disorders ²	4.1	4.2	4.4	7.1	6.5	6.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.3	47.3	47.0	47.2	47.5	47.9
Female	52.7	52.7	53.0	52.8	52.5	52.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	9.4	9.4	9.5	10.0	10.4	10.7
12 to 14 years	5.7	5.8	5.9	6.0	6.0	6.4
15 to 17 years	5.4	5.3	5.6	5.9	5.9	5.6
18 to 20 years	3.4	3.3	3.2	3.3	3.4	3.5
21 to 24 years	5.4	5.0	4.7	4.7	4.5	4.4
25 to 29 years	8.3	8.0	7.7	7.4	7.3	7.3
30 to 34 years	9.2	9.2	8.8	8.5	8.4	8.0
35 to 39 years	8.4	8.6	8.6	8.7	8.8	8.6
40 to 44 years	9.0	8.8	8.7	8.1	7.8	7.7
45 to 49 years	10.2	9.7	9.3	9.1	8.9	8.7
50 to 54 years	10.9	11.1	11.3	10.9	10.3	9.7
55 to 59 years	7.9	8.3	8.7	8.9	9.0	9.5
60 to 64 years	4.2	4.5	4.9	5.2	5.5	6.0
65 years and older	2.9	3.1	3.3	3.5	3.6	3.9
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.24b. Missouri individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	22.1	21.4	21.2	20.8	21.2	21.9
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	74.8	75.6	75.3	75.4	74.5	73.6
Other	2.5	2.5	2.9	3.2	3.7	3.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	97.9	97.9	97.8	97.7	97.6	97.5
Hispanic or Latino origin	2.1	2.1	2.2	2.3	2.4	2.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	46.4	45.9	54.6	57.2	57.2	55.5
Puerto Rican	13.2	12.7	11.8	12.1	10.4	9.3
Cuban	3.1	3.0	3.4	3.2	2.8	2.9
Other specific	37.2	38.4	30.2	27.6	29.6	32.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.25a. Montana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	41,812	43,792	46,288	38,476	50,515	63,052
Mental health diagnosis¹						
Trauma- and stressor-related disorders	6,128	6,625	7,543	5,906	8,789	9,788
Anxiety disorders	4,176	4,613	5,267	4,408	8,963	13,719
Attention deficit/hyperactivity disorder	2,593	2,689	2,659	2,093	3,364	4,220
Mood disorders	14,429	14,656	15,089	10,904	14,537	17,830
Bipolar disorders	4,836	4,750	4,722	3,529	3,692	3,759
Depressive disorders	9,593	9,906	10,367	7,375	10,845	14,071
Personality disorders	505	530	539	477	364	550
Schizophrenia and other psychotic disorders	3,116	3,128	3,091	2,407	2,804	3,370
Alcohol or substance-related disorders	697	721	742	511	191	14
Other mental disorders ²	5,643	5,808	5,829	4,399	10,226	11,698
Total	37,287	38,770	40,759	31,105	49,238	61,189
Gender						
Male	19,332	20,224	20,935	17,424	23,304	28,085
Female	22,480	23,568	25,353	21,051	27,211	34,941
Total	41,812	43,792	46,288	38,475	50,515	63,026
Age³						
0 to 11 years	9,752	10,552	11,093	9,113	13,589	13,315
12 to 14 years	3,657	3,928	4,413	3,814	5,220	5,660
15 to 17 years	3,285	3,779	4,090	3,639	4,876	5,479
18 to 20 years	1,784	1,765	1,912	1,636	2,094	3,079
21 to 24 years	1,973	2,007	1,960	1,518	2,163	3,365
25 to 29 years	2,887	2,937	3,095	2,508	3,469	5,372
30 to 34 years	2,826	2,902	3,121	2,627	3,489	5,250
35 to 39 years	2,278	2,371	2,737	2,349	3,150	4,615
40 to 44 years	2,254	2,256	2,308	1,812	2,387	3,462
45 to 49 years	2,319	2,154	2,182	1,892	2,330	3,148
50 to 54 years	2,533	2,554	2,528	2,006	2,376	3,024
55 to 59 years	2,126	2,275	2,351	2,014	2,319	2,923
60 to 64 years	1,359	1,477	1,580	1,364	1,542	2,131
65 years and older	2,778	2,834	2,916	2,183	1,511	2,028
Total	41,811	43,791	46,286	38,475	50,515	62,851

Continued. See notes at end of table.

Table 9.25a. Montana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	4,720	4,972	5,349	4,444	5,745	7,219
Asian ⁴	96	109	129	91	150	171
Black or African American	403	408	430	396	558	681
Native Hawaiian or Other Pacific Islander	35	21	23	21	33	52
White	34,436	35,663	37,017	30,579	39,116	48,140
Other	752	823	875	783	1,030	1,371
Total	40,442	41,996	43,823	36,314	46,632	57,634
Ethnicity						
Not of Hispanic or Latino origin	40,348	42,270	44,686	37,085	48,399	58,124
Hispanic or Latino origin	1,432	1,503	1,574	1,360	1,832	2,326
Total	41,780	43,773	46,260	38,445	50,231	60,450
Specific Hispanic or Latino origin						
Mexican	‡	‡	2	4	‡	‡
Puerto Rican	‡	‡	--	--	‡	‡
Cuban	‡	‡	--	--	‡	‡
Other specific	‡	‡	--	3	‡	‡
Total	‡	‡	2	7	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	205	172
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	8,140	8,238
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	8,345	8,410

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.25b. Montana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	16.4	17.1	18.5	19.0	17.9	16.0
Anxiety disorders	11.2	11.9	12.9	14.2	18.2	22.4
Attention deficit/hyperactivity disorder	7.0	6.9	6.5	6.7	6.8	6.9
Mood disorders	38.7	37.8	37.0	35.1	29.5	29.1
Bipolar disorders	13.0	12.3	11.6	11.3	7.5	6.1
Depressive disorders	25.7	25.6	25.4	23.7	22.0	23.0
Personality disorders	1.4	1.4	1.3	1.5	†	†
Schizophrenia and other psychotic disorders	8.4	8.1	7.6	7.7	5.7	5.5
Alcohol or substance-related disorders	1.9	1.9	1.8	1.6	†	†
Other mental disorders ²	15.1	15.0	14.3	14.1	20.8	19.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	46.2	46.2	45.2	45.3	46.1	44.6
Female	53.8	53.8	54.8	54.7	53.9	55.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	23.3	24.1	24.0	23.7	26.9	21.2
12 to 14 years	8.7	9.0	9.5	9.9	10.3	9.0
15 to 17 years	7.9	8.6	8.8	9.5	9.7	8.7
18 to 20 years	4.3	4.0	4.1	4.3	4.1	4.9
21 to 24 years	4.7	4.6	4.2	3.9	4.3	5.4
25 to 29 years	6.9	6.7	6.7	6.5	6.9	8.5
30 to 34 years	6.8	6.6	6.7	6.8	6.9	8.4
35 to 39 years	5.4	5.4	5.9	6.1	6.2	7.3
40 to 44 years	5.4	5.2	5.0	4.7	4.7	5.5
45 to 49 years	5.5	4.9	4.7	4.9	4.6	5.0
50 to 54 years	6.1	5.8	5.5	5.2	4.7	4.8
55 to 59 years	5.1	5.2	5.1	5.2	4.6	4.7
60 to 64 years	3.3	3.4	3.4	3.5	3.1	3.4
65 years and older	6.6	6.5	6.3	5.7	3.0	3.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.25b. Montana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	11.7	11.8	12.2	12.2	12.3	12.5
Asian ⁴	†	†	†	†	†	†
Black or African American	†	†	†	1.1	1.2	1.2
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	85.1	84.9	84.5	84.2	83.9	83.5
Other	1.9	2.0	2.0	2.2	2.2	2.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	96.6	96.6	96.6	96.5	96.4	96.2
Hispanic or Latino origin	3.4	3.4	3.4	3.5	3.6	3.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	100.0	57.1	‡	‡
Puerto Rican	‡	‡	--	--	‡	‡
Cuban	‡	‡	--	--	‡	‡
Other specific	‡	‡	--	42.9	‡	‡
Total	‡	‡	100.0	100.0	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2.5	2.0
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	97.5	98.0
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.26a. Nebraska individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	22,040	22,578	19,441	19,739	21,927	24,983
Mental health diagnosis¹						
Trauma- and stressor-related disorders	2,371	2,603	2,487	2,260	2,358	2,462
Anxiety disorders	1,062	1,150	1,203	1,533	1,741	1,760
Attention deficit/hyperactivity disorder	641	728	590	614	621	721
Mood disorders	11,740	11,733	10,091	8,837	9,318	9,179
Bipolar disorders	5,545	5,399	4,641	3,541	3,659	3,479
Depressive disorders	6,195	6,334	5,450	5,296	5,659	5,700
Personality disorders	149	129	137	65	73	116
Schizophrenia and other psychotic disorders	3,275	3,145	2,199	2,002	2,230	2,264
Alcohol or substance-related disorders	--	--	--	--	795	824
Other mental disorders ²	1,322	1,409	1,304	1,607	2,082	2,685
Total	20,560	20,897	18,011	16,918	19,218	20,011
Gender						
Male	10,978	11,248	9,636	9,866	10,993	12,339
Female	11,052	11,304	9,802	9,840	10,753	11,766
Total	22,030	22,552	19,438	19,706	21,746	24,105
Age³						
0 to 11 years	668	745	693	715	755	905
12 to 14 years	540	615	484	500	691	931
15 to 17 years	757	851	643	646	957	1,549
18 to 20 years	972	1,002	924	989	1,112	1,409
21 to 24 years	1,836	1,866	1,799	1,856	1,906	2,153
25 to 29 years	2,598	2,587	2,363	2,431	2,677	3,053
30 to 34 years	2,659	2,723	2,494	2,506	2,795	2,909
35 to 39 years	2,159	2,244	2,016	2,136	2,324	2,579
40 to 44 years	2,242	2,209	1,769	1,754	1,905	2,006
45 to 49 years	2,326	2,295	1,882	1,816	1,942	2,031
50 to 54 years	2,313	2,366	1,963	1,842	1,911	1,965
55 to 59 years	1,587	1,629	1,329	1,376	1,452	1,612
60 to 64 years	833	906	702	733	830	968
65 years and older	544	535	380	414	669	910
Total	22,034	22,573	19,441	19,714	21,926	24,980

Continued. See notes at end of table.

Table 9.26a. Nebraska individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	474	544	478	507	659	611
Asian ⁴	135	162	142	131	141	160
Black or African American	1,862	1,937	1,668	1,735	1,981	2,177
Native Hawaiian or Other Pacific Islander	125	144	122	129	64	89
White	19,109	19,443	16,733	16,496	17,543	18,376
Other	245	273	251	576	772	1,385
Total	21,950	22,503	19,394	19,574	21,160	22,798
Ethnicity						
Not of Hispanic or Latino origin	19,760	20,081	17,167	16,023	17,535	18,383
Hispanic or Latino origin	1,606	1,763	1,631	1,607	1,752	2,129
Total	21,366	21,844	18,798	17,630	19,287	20,512
Specific Hispanic or Latino origin						
Mexican	491	588	533	‡	‡	‡
Puerto Rican	51	56	45	‡	‡	‡
Cuban	22	30	25	‡	‡	‡
Other specific	248	237	200	‡	‡	‡
Total	812	911	803	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	497	401
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	17,115	17,653
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	17,612	18,054

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.26b. Nebraska individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11.5	12.5	13.8	13.4	12.3	12.3
Anxiety disorders	5.2	5.5	6.7	9.1	9.1	8.8
Attention deficit/hyperactivity disorder	3.1	3.5	3.3	3.6	3.2	3.6
Mood disorders	57.1	56.1	56.0	52.2	48.5	45.9
Bipolar disorders	27.0	25.8	25.8	20.9	19.0	17.4
Depressive disorders	30.1	30.3	30.3	31.3	29.4	28.5
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	15.9	15.1	12.2	11.8	11.6	11.3
Alcohol or substance-related disorders	--	--	--	--	4.1	4.1
Other mental disorders ²	6.4	6.7	7.2	9.5	10.8	13.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	49.8	49.9	49.6	50.1	50.6	51.2
Female	50.2	50.1	50.4	49.9	49.4	48.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	3.0	3.3	3.6	3.6	3.4	3.6
12 to 14 years	2.5	2.7	2.5	2.5	3.2	3.7
15 to 17 years	3.4	3.8	3.3	3.3	4.4	6.2
18 to 20 years	4.4	4.4	4.8	5.0	5.1	5.6
21 to 24 years	8.3	8.3	9.3	9.4	8.7	8.6
25 to 29 years	11.8	11.5	12.2	12.3	12.2	12.2
30 to 34 years	12.1	12.1	12.8	12.7	12.7	11.6
35 to 39 years	9.8	9.9	10.4	10.8	10.6	10.3
40 to 44 years	10.2	9.8	9.1	8.9	8.7	8.0
45 to 49 years	10.6	10.2	9.7	9.2	8.9	8.1
50 to 54 years	10.5	10.5	10.1	9.3	8.7	7.9
55 to 59 years	7.2	7.2	6.8	7.0	6.6	6.5
60 to 64 years	3.8	4.0	3.6	3.7	3.8	3.9
65 years and older	2.5	2.4	2.0	2.1	3.1	3.6
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.26b. Nebraska individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	2.2	2.4	2.5	2.6	3.1	2.7
Asian ⁴	†	†	†	†	†	†
Black or African American	8.5	8.6	8.6	8.9	9.4	9.5
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	87.1	86.4	86.3	84.3	82.9	80.6
Other	1.1	1.2	1.3	2.9	3.6	6.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	92.5	91.9	91.3	90.9	90.9	89.6
Hispanic or Latino origin	7.5	8.1	8.7	9.1	9.1	10.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	60.5	64.5	66.4	‡	‡	‡
Puerto Rican	6.3	6.1	5.6	‡	‡	‡
Cuban	2.7	3.3	3.1	‡	‡	‡
Other specific	30.5	26.0	24.9	‡	‡	‡
Total	100.0	100.0	100.0	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2.8	2.2
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	97.2	97.8
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.27a. Nevada individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	29,196	30,358	28,589	23,945	20,535	19,632
Mental health diagnosis¹						
Trauma- and stressor-related disorders	2,197	2,060	2,245	1,505	1,725	1,681
Anxiety disorders	1,670	1,668	1,604	1,571	1,619	1,589
Attention deficit/hyperactivity disorder	466	442	478	427	425	367
Mood disorders	14,533	14,204	12,935	3,188	3,311	3,030
Bipolar disorders	7,091	6,758	6,226	1,128	1,126	1,008
Depressive disorders	7,442	7,446	6,709	2,060	2,185	2,022
Personality disorders	42	26	26	559	519	437
Schizophrenia and other psychotic disorders	5,232	5,253	5,510	1,967	2,042	2,003
Alcohol or substance-related disorders	1,795	1,778	1,150	2,876	2,817	2,430
Other mental disorders ²	1,185	1,147	1,057	1,758	1,777	1,713
Total	27,120	26,578	25,005	13,851	14,235	13,250
Gender						
Male	13,881	14,277	13,760	11,739	10,135	9,824
Female	15,242	15,849	14,676	12,151	10,383	9,792
Total	29,123	30,126	28,436	23,890	20,518	19,616
Age³						
0 to 11 years	1,827	1,712	1,832	1,727	1,852	1,924
12 to 14 years	746	800	941	1,016	1,200	1,239
15 to 17 years	983	1,002	1,151	1,345	1,478	1,520
18 to 20 years	1,069	1,100	925	845	856	793
21 to 24 years	1,958	1,977	1,801	1,452	1,264	1,279
25 to 29 years	2,899	3,118	2,821	2,304	1,884	1,769
30 to 34 years	2,940	3,118	2,957	2,365	1,810	1,739
35 to 39 years	2,540	2,694	2,612	2,159	1,687	1,576
40 to 44 years	2,971	3,010	2,575	1,922	1,518	1,440
45 to 49 years	3,234	3,338	2,948	2,256	1,681	1,530
50 to 54 years	3,374	3,519	3,169	2,368	1,896	1,699
55 to 59 years	2,571	2,657	2,505	2,011	1,586	1,440
60 to 64 years	1,380	1,514	1,469	1,246	1,001	920
65 years and older	670	769	856	917	811	724
Total	29,162	30,328	28,562	23,933	20,524	19,592

Continued. See notes at end of table.

Table 9.27a. Nevada individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	327	370	340	295	348	362
Asian ⁴	573	563	519	469	436	470
Black or African American	3,842	3,894	3,680	3,004	2,585	2,540
Native Hawaiian or Other Pacific Islander	160	139	147	125	117	116
White	18,141	18,039	16,594	13,255	10,948	10,405
Other	3,193	3,190	3,101	2,470	1,966	1,901
Total	26,236	26,195	24,381	19,618	16,400	15,794
Ethnicity						
Not of Hispanic or Latino origin	21,544	20,345	18,468	15,043	12,171	11,953
Hispanic or Latino origin	3,847	3,722	3,771	3,373	3,038	2,990
Total	25,391	24,067	22,239	18,416	15,209	14,943
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	16	19
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	8,885	4,844
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	8,901	4,863

‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.27b. Nevada individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	8.1	7.8	9.0	10.9	12.1	12.7
Anxiety disorders	6.2	6.3	6.4	11.3	11.4	12.0
Attention deficit/hyperactivity disorder	1.7	1.7	1.9	3.1	3.0	2.8
Mood disorders	53.6	53.4	51.7	23.0	23.3	22.9
Bipolar disorders	26.1	25.4	24.9	8.1	7.9	7.6
Depressive disorders	27.4	28.0	26.8	14.9	15.3	15.3
Personality disorders	†	†	†	4.0	3.6	3.3
Schizophrenia and other psychotic disorders	19.3	19.8	22.0	14.2	14.3	15.1
Alcohol or substance-related disorders	6.6	6.7	4.6	20.8	19.8	18.3
Other mental disorders ²	4.4	4.3	4.2	12.7	12.5	12.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.7	47.4	48.4	49.1	49.4	50.1
Female	52.3	52.6	51.6	50.9	50.6	49.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	6.3	5.6	6.4	7.2	9.0	9.8
12 to 14 years	2.6	2.6	3.3	4.2	5.8	6.3
15 to 17 years	3.4	3.3	4.0	5.6	7.2	7.8
18 to 20 years	3.7	3.6	3.2	3.5	4.2	4.0
21 to 24 years	6.7	6.5	6.3	6.1	6.2	6.5
25 to 29 years	9.9	10.3	9.9	9.6	9.2	9.0
30 to 34 years	10.1	10.3	10.4	9.9	8.8	8.9
35 to 39 years	8.7	8.9	9.1	9.0	8.2	8.0
40 to 44 years	10.2	9.9	9.0	8.0	7.4	7.3
45 to 49 years	11.1	11.0	10.3	9.4	8.2	7.8
50 to 54 years	11.6	11.6	11.1	9.9	9.2	8.7
55 to 59 years	8.8	8.8	8.8	8.4	7.7	7.3
60 to 64 years	4.7	5.0	5.1	5.2	4.9	4.7
65 years and older	2.3	2.5	3.0	3.8	4.0	3.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.27b. Nevada individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	1.2	1.4	1.4	1.5	2.1	2.3
Asian ⁴	2.2	2.1	2.1	2.4	2.7	3.0
Black or African American	14.6	14.9	15.1	15.3	15.8	16.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	69.1	68.9	68.1	67.6	66.8	65.9
Other	12.2	12.2	12.7	12.6	12.0	12.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	84.8	84.5	83.0	81.7	80.0	80.0
Hispanic or Latino origin	15.2	15.5	17.0	18.3	20.0	20.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	†	†
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	99.8	99.6
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.28a. New Jersey individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	324,221	324,940	325,102	323,392	‡	‡
Mental health diagnosis¹						
Trauma- and stressor-related disorders	41,080	40,255	38,639	3,442	‡	‡
Anxiety disorders	19,535	19,695	19,274	1,179	‡	‡
Attention deficit/hyperactivity disorder	11,668	11,559	11,016	3,287	‡	‡
Mood disorders	89,971	87,844	86,614	2,139	‡	‡
Bipolar disorders	27,199	26,632	25,784	487	‡	‡
Depressive disorders	62,772	61,212	60,830	1,652	‡	‡
Personality disorders	2,090	2,061	2,043	25	‡	‡
Schizophrenia and other psychotic disorders	30,622	30,106	29,744	112	‡	‡
Alcohol or substance-related disorders	20,240	19,318	18,773	592	‡	‡
Other mental disorders ²	40,296	39,403	37,319	5,995	‡	‡
Total	255,502	250,241	243,422	16,771	‡	‡
Gender						
Male	153,450	154,435	154,484	153,430	‡	‡
Female	170,162	169,885	170,002	169,259	‡	‡
Total	323,612	324,320	324,486	322,689	‡	‡
Age³						
0 to 11 years	23,176	26,357	22,950	22,982	‡	‡
12 to 14 years	17,589	18,748	16,859	15,885	‡	‡
15 to 17 years	20,696	19,023	20,765	20,078	‡	‡
18 to 20 years	12,889	11,930	12,746	11,766	‡	‡
21 to 24 years	16,176	15,243	15,043	14,648	‡	‡
25 to 29 years	21,396	20,878	20,545	20,159	‡	‡
30 to 34 years	22,513	22,313	22,032	21,436	‡	‡
35 to 39 years	20,511	20,347	20,861	21,358	‡	‡
40 to 44 years	24,280	23,049	22,111	20,705	‡	‡
45 to 49 years	28,212	26,642	25,853	25,071	‡	‡
50 to 54 years	30,317	30,033	29,828	28,839	‡	‡
55 to 59 years	25,826	26,749	27,737	28,251	‡	‡
60 to 64 years	18,699	19,539	20,731	21,891	‡	‡
65 years and older	41,607	43,753	46,684	50,251	‡	‡
Total	323,887	324,604	324,745	323,320	‡	‡

Continued. See notes at end of table.

Table 9.28a. New Jersey individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	2,657	2,606	2,564	2,520	‡	‡
Asian ⁴	4,268	4,574	4,755	4,818	‡	‡
Black or African American	63,201	64,350	63,704	63,087	‡	‡
Native Hawaiian or Other Pacific Islander	42	46	44	24	‡	‡
White	171,457	170,702	170,128	168,286	‡	‡
Other	24,135	24,223	23,132	21,916	‡	‡
Total	265,760	266,501	264,327	260,651	‡	‡
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	‡	‡	‡
Hispanic or Latino origin	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.28b. New Jersey individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	‡	‡
Mental health diagnosis¹						
Trauma- and stressor-related disorders	16.1	16.1	15.9	20.5	‡	‡
Anxiety disorders	7.6	7.9	7.9	7.0	‡	‡
Attention deficit/hyperactivity disorder	4.6	4.6	4.5	19.6	‡	‡
Mood disorders	35.2	35.1	35.6	12.8	‡	‡
Bipolar disorders	10.6	10.6	10.6	2.9	‡	‡
Depressive disorders	24.6	24.5	25.0	9.9	‡	‡
Personality disorders	†	†	†	†	‡	‡
Schizophrenia and other psychotic disorders	12.0	12.0	12.2	†	‡	‡
Alcohol or substance-related disorders	7.9	7.7	7.7	3.5	‡	‡
Other mental disorders ²	15.8	15.7	15.3	35.7	‡	‡
Total	100.0	100.0	100.0	100.0	‡	‡
Gender						
Male	47.4	47.6	47.6	47.5	‡	‡
Female	52.6	52.4	52.4	52.5	‡	‡
Total	100.0	100.0	100.0	100.0	‡	‡
Age³						
0 to 11 years	7.2	8.1	7.1	7.1	‡	‡
12 to 14 years	5.4	5.8	5.2	4.9	‡	‡
15 to 17 years	6.4	5.9	6.4	6.2	‡	‡
18 to 20 years	4.0	3.7	3.9	3.6	‡	‡
21 to 24 years	5.0	4.7	4.6	4.5	‡	‡
25 to 29 years	6.6	6.4	6.3	6.2	‡	‡
30 to 34 years	7.0	6.9	6.8	6.6	‡	‡
35 to 39 years	6.3	6.3	6.4	6.6	‡	‡
40 to 44 years	7.5	7.1	6.8	6.4	‡	‡
45 to 49 years	8.7	8.2	8.0	7.8	‡	‡
50 to 54 years	9.4	9.3	9.2	8.9	‡	‡
55 to 59 years	8.0	8.2	8.5	8.7	‡	‡
60 to 64 years	5.8	6.0	6.4	6.8	‡	‡
65 years and older	12.8	13.5	14.4	15.5	‡	‡
Total	100.0	100.0	100.0	100.0	‡	‡

Continued. See notes at end of table.

Table 9.28b. New Jersey individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	†	†	†	†	‡	‡
Asian ⁴	1.6	1.7	1.8	1.8	‡	‡
Black or African American	23.8	24.1	24.1	24.2	‡	‡
Native Hawaiian or Other Pacific Islander	†	†	†	†	‡	‡
White	64.5	64.1	64.4	64.6	‡	‡
Other	9.1	9.1	8.8	8.4	‡	‡
Total	100.0	100.0	100.0	100.0	‡	‡
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	‡	‡	‡
Hispanic or Latino origin	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.29a. New Mexico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	80,772	93,300	121,603	131,220	147,669	149,503
Mental health diagnosis¹						
Trauma- and stressor-related disorders	21,370	21,759	23,462	29,202	33,836	25,568
Anxiety disorders	6,725	7,480	10,884	15,280	20,191	20,189
Attention deficit/hyperactivity disorder	7,062	6,632	6,819	7,191	7,835	7,559
Mood disorders	25,464	25,262	27,806	29,350	33,723	27,444
Bipolar disorders	9,607	9,782	9,170	8,172	8,656	6,705
Depressive disorders	15,857	15,480	18,636	21,178	25,067	20,739
Personality disorders	174	181	186	247	283	230
Schizophrenia and other psychotic disorders	4,787	5,669	4,566	5,139	6,610	5,557
Alcohol or substance-related disorders	4,445	6,740	13,576	18,216	25,538	26,528
Other mental disorders ²	8,343	13,259	17,848	21,352	16,020	36,423
Total	78,370	86,982	105,147	125,977	144,036	149,498
Gender						
Male	39,831	45,949	59,253	63,080	70,053	73,683
Female	40,917	47,331	62,179	68,062	77,554	75,602
Total	80,748	93,280	121,432	131,142	147,607	149,285
Age³						
0 to 11 years	20,132	25,064	29,082	30,204	25,265	33,849
12 to 14 years	8,808	8,879	9,161	9,891	10,113	10,108
15 to 17 years	8,789	8,790	9,016	10,135	10,471	9,859
18 to 20 years	3,726	3,547	4,781	5,332	5,992	5,739
21 to 24 years	3,997	4,216	6,513	7,253	8,798	7,395
25 to 29 years	5,675	6,293	10,124	11,298	14,291	12,839
30 to 34 years	5,803	6,829	10,057	11,233	13,843	13,043
35 to 39 years	4,513	5,487	8,378	9,730	12,169	11,868
40 to 44 years	4,235	4,927	7,144	7,603	9,378	9,116
45 to 49 years	4,020	4,699	6,858	7,179	9,017	8,445
50 to 54 years	4,130	4,983	7,306	7,430	9,212	8,342
55 to 59 years	3,216	3,965	5,782	6,094	8,032	7,678
60 to 64 years	2,019	2,678	3,896	3,976	5,334	5,449
65 years and older	1,707	2,942	3,502	3,861	5,753	5,772
Total	80,770	93,299	121,600	131,219	147,668	149,502

Continued. See notes at end of table.

Table 9.29a. New Mexico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	4,286	3,985	4,537	4,679	7,758	11,844
Asian ⁴	167	285	710	696	3,813	1,598
Black or African American	1,746	1,855	2,495	2,651	3,307	3,015
Native Hawaiian or Other Pacific Islander	121	113	136	85	115	141
White	43,760	48,914	75,059	81,789	101,867	104,095
Other	5,429	4,455	3,601	3,245	3,035	3,360
Total	55,509	59,607	86,538	93,145	119,895	124,053
Ethnicity						
Not of Hispanic or Latino origin	25,127	31,843	39,667	41,079	62,181	57,561
Hispanic or Latino origin	35,124	32,828	49,264	54,688	54,453	66,603
Total	60,251	64,671	88,931	95,767	116,634	124,164
Specific Hispanic or Latino origin						
Mexican	3,000	2,128	1,369	1,249	1,049	834
Puerto Rican	215	165	121	104	87	64
Cuban	128	95	63	54	55	44
Other specific	1,196	874	669	596	546	496
Total	4,539	3,262	2,222	2,003	1,737	1,438
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	206	699
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	5,011	92,375
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	5,217	93,074

n/c Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.29b. New Mexico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	27.3	25.0	22.3	23.2	23.5	17.1
Anxiety disorders	8.6	8.6	10.4	12.1	14.0	13.5
Attention deficit/hyperactivity disorder	9.0	7.6	6.5	5.7	5.4	5.1
Mood disorders	32.5	29.0	26.4	23.3	23.4	18.4
Bipolar disorders	12.3	11.2	8.7	6.5	6.0	4.5
Depressive disorders	20.2	17.8	17.7	16.8	17.4	13.9
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	6.1	6.5	4.3	4.1	4.6	3.7
Alcohol or substance-related disorders	5.7	7.7	12.9	14.5	17.7	17.7
Other mental disorders ²	10.6	15.2	17.0	16.9	11.1	24.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	49.3	49.3	48.8	48.1	47.5	49.4
Female	50.7	50.7	51.2	51.9	52.5	50.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	24.9	26.9	23.9	23.0	17.1	22.6
12 to 14 years	10.9	9.5	7.5	7.5	6.8	6.8
15 to 17 years	10.9	9.4	7.4	7.7	7.1	6.6
18 to 20 years	4.6	3.8	3.9	4.1	4.1	3.8
21 to 24 years	4.9	4.5	5.4	5.5	6.0	4.9
25 to 29 years	7.0	6.7	8.3	8.6	9.7	8.6
30 to 34 years	7.2	7.3	8.3	8.6	9.4	8.7
35 to 39 years	5.6	5.9	6.9	7.4	8.2	7.9
40 to 44 years	5.2	5.3	5.9	5.8	6.4	6.1
45 to 49 years	5.0	5.0	5.6	5.5	6.1	5.6
50 to 54 years	5.1	5.3	6.0	5.7	6.2	5.6
55 to 59 years	4.0	4.2	4.8	4.6	5.4	5.1
60 to 64 years	2.5	2.9	3.2	3.0	3.6	3.6
65 years and older	2.1	3.2	2.9	2.9	3.9	3.9
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.29b. New Mexico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	7.7	6.7	5.2	5.0	6.5	9.5
Asian ⁴	†	†	†	†	3.2	1.3
Black or African American	3.1	3.1	2.9	2.8	2.8	2.4
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	78.8	82.1	86.7	87.8	85.0	83.9
Other	9.8	7.5	4.2	3.5	2.5	2.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	41.7	49.2	44.6	42.9	53.3	46.4
Hispanic or Latino origin	58.3	50.8	55.4	57.1	46.7	53.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	66.1	65.2	61.6	62.4	60.4	58.0
Puerto Rican	4.7	5.1	5.4	5.2	5.0	4.5
Cuban	2.8	2.9	2.8	2.7	3.2	3.1
Other specific	26.3	26.8	30.1	29.8	31.4	34.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	3.9	†
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	96.1	99.2
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.30a. New York individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals¹	53,000	53,505	54,524	56,968	58,040	57,904
Mental health diagnosis²						
Trauma- and stressor-related disorders	5,450	5,880	6,179	6,945	7,316	7,579
Anxiety disorders	2,234	2,413	3,677	4,536	3,409	3,271
Attention deficit/hyperactivity disorder	1,486	1,557	1,724	2,074	2,004	1,870
Mood disorders	18,206	18,318	17,195	16,008	12,436	12,769
Bipolar disorders	9,519	9,523	7,667	6,327	5,005	5,166
Depressive disorders	8,687	8,795	9,528	9,681	7,431	7,603
Personality disorders	2,477	2,489	2,880	4,724	4,775	4,586
Schizophrenia and other psychotic disorders	20,381	19,950	19,763	19,372	19,398	19,316
Alcohol or substance-related disorders	--	--	--	181	--	--
Other mental disorders ³	1,539	1,574	1,455	1,997	7,243	7,118
Total	51,773	52,181	52,873	55,837	56,581	56,509
Gender						
Male	34,714	34,890	35,414	36,941	37,976	37,740
Female	18,277	18,613	19,109	20,024	20,062	20,163
Total	52,991	53,503	54,523	56,965	58,038	57,903
Age⁴						
0 to 11 years	1,758	1,905	1,982	2,195	2,266	2,280
12 to 14 years	1,801	1,952	2,087	2,219	2,341	2,412
15 to 17 years	2,285	2,457	2,645	2,897	3,010	3,030
18 to 20 years	1,467	1,412	1,365	1,540	1,483	1,453
21 to 24 years	2,858	2,865	3,027	3,112	3,119	2,939
25 to 29 years	4,383	4,405	4,604	5,053	5,367	5,304
30 to 34 years	4,790	4,811	4,839	5,132	5,243	5,381
35 to 39 years	4,232	4,359	4,594	4,893	5,247	5,332
40 to 44 years	4,824	4,556	4,347	4,397	4,343	4,420
45 to 49 years	5,946	5,629	5,476	5,347	5,243	4,924
50 to 54 years	6,215	6,115	6,106	6,124	5,898	5,735
55 to 59 years	4,859	5,190	5,291	5,412	5,573	5,562
60 to 64 years	3,413	3,553	3,656	3,845	4,055	4,091
65 years and older	4,168	4,295	4,505	4,801	4,852	5,036
Total	52,999	53,504	54,524	56,967	58,040	57,899

Continued. See notes at end of table.

Table 9.30a. New York individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	278	272	277	340	365	381
Asian ⁵	682	700	776	863	940	973
Black or African American	15,714	15,602	15,596	16,050	16,503	16,425
Native Hawaiian or Other Pacific Islander	56	57	69	69	54	51
White	26,839	27,209	27,611	28,853	29,408	29,277
Other	3,478	4,019	4,611	5,506	6,434	6,823
Total	47,047	47,859	48,940	51,681	53,704	53,930
Ethnicity						
Not of Hispanic or Latino origin	33,189	35,808	38,187	41,130	42,800	42,981
Hispanic or Latino origin	8,680	8,612	8,534	8,603	8,798	8,635
Total	41,869	44,420	46,721	49,733	51,598	51,616
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁶						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	362	336
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	13,595	12,675
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	13,957	13,011

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ New York reports only a subset of the total individuals served through their SMHAs.

² “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

³ “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

⁴ Age is based on the individual’s age at the midpoint of the reporting period.

⁵ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁶ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.30b. New York individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals¹	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis²						
Trauma- and stressor-related disorders	10.5	11.3	11.7	12.4	12.9	13.4
Anxiety disorders	4.3	4.6	7.0	8.1	6.0	5.8
Attention deficit/hyperactivity disorder	2.9	3.0	3.3	3.7	3.5	3.3
Mood disorders	35.2	35.1	32.5	28.7	22.0	22.6
Bipolar disorders	18.4	18.2	14.5	11.3	8.8	9.1
Depressive disorders	16.8	16.9	18.0	17.3	13.1	13.5
Personality disorders	4.8	4.8	5.4	8.5	8.4	8.1
Schizophrenia and other psychotic disorders	39.4	38.2	37.4	34.7	34.3	34.2
Alcohol or substance-related disorders	--	--	--	†	--	--
Other mental disorders ³	3.0	3.0	2.8	3.6	12.8	12.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	65.5	65.2	65.0	64.8	65.4	65.2
Female	34.5	34.8	35.0	35.2	34.6	34.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age⁴						
0 to 11 years	3.3	3.6	3.6	3.9	3.9	3.9
12 to 14 years	3.4	3.6	3.8	3.9	4.0	4.2
15 to 17 years	4.3	4.6	4.9	5.1	5.2	5.2
18 to 20 years	2.8	2.6	2.5	2.7	2.6	2.5
21 to 24 years	5.4	5.4	5.6	5.5	5.4	5.1
25 to 29 years	8.3	8.2	8.4	8.9	9.2	9.2
30 to 34 years	9.0	9.0	8.9	9.0	9.0	9.3
35 to 39 years	8.0	8.1	8.4	8.6	9.0	9.2
40 to 44 years	9.1	8.5	8.0	7.7	7.5	7.6
45 to 49 years	11.2	10.5	10.0	9.4	9.0	8.5
50 to 54 years	11.7	11.4	11.2	10.8	10.2	9.9
55 to 59 years	9.2	9.7	9.7	9.5	9.6	9.6
60 to 64 years	6.4	6.6	6.7	6.7	7.0	7.1
65 years and older	7.9	8.0	8.3	8.4	8.4	8.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.30b. New York individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁵	1.4	1.5	1.6	1.7	1.8	1.8
Black or African American	33.4	32.6	31.9	31.1	30.7	30.5
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	57.0	56.9	56.4	55.8	54.8	54.3
Other	7.4	8.4	9.4	10.7	12.0	12.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	79.3	80.6	81.7	82.7	82.9	83.3
Hispanic or Latino origin	20.7	19.4	18.3	17.3	17.1	16.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁶						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2.6	2.6
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	97.4	97.4
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ New York reports only a subset of the total individuals served through their SMHAs.

² “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

³ “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

⁴ Age is based on the individual’s age at the midpoint of the reporting period.

⁵ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁶ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.31a. North Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	223,440	219,448	225,452	138,079	142,665	101,046
Mental health diagnosis¹						
Trauma- and stressor-related disorders	25,936	24,040	26,502	10,791	15,541	11,858
Anxiety disorders	20,420	18,684	22,955	8,663	11,305	8,114
Attention deficit/hyperactivity disorder	20,459	15,393	14,810	2,601	4,227	2,439
Mood disorders	102,417	107,390	109,349	48,733	68,378	43,543
Bipolar disorders	42,350	46,274	41,700	16,653	23,957	14,038
Depressive disorders	60,067	61,116	67,649	32,080	44,421	29,505
Personality disorders	2,471	1,495	1,847	995	1,490	1,110
Schizophrenia and other psychotic disorders	21,329	22,349	22,853	11,250	17,546	13,345
Alcohol or substance-related disorders	--	--	--	--	--	--
Other mental disorders ²	24,098	22,498	20,045	7,785	13,737	9,766
Total	217,130	211,849	218,361	90,818	132,224	90,175
Gender						
Male	105,506	104,089	106,397	66,277	70,348	52,269
Female	117,934	115,359	119,054	71,767	72,279	48,760
Total	223,440	219,448	225,451	138,044	142,627	101,029
Age³						
0 to 11 years	29,351	27,476	25,245	4,806	4,075	3,577
12 to 14 years	16,096	15,562	14,410	5,106	4,519	3,471
15 to 17 years	16,758	16,307	15,833	7,125	6,551	4,208
18 to 20 years	11,033	10,871	11,130	6,028	6,948	4,567
21 to 24 years	15,163	14,915	16,113	10,867	11,794	8,938
25 to 29 years	19,328	19,823	22,167	15,602	17,277	13,374
30 to 34 years	19,163	19,062	20,774	14,768	15,872	11,875
35 to 39 years	16,933	17,024	18,481	13,673	14,780	10,600
40 to 44 years	17,807	17,220	17,601	12,816	12,889	9,232
45 to 49 years	17,974	17,112	17,547	13,655	13,739	9,546
50 to 54 years	17,640	17,578	18,108	13,997	13,531	8,904
55 to 59 years	12,334	12,560	13,325	10,326	10,714	6,765
60 to 64 years	6,754	6,949	7,482	5,618	5,808	3,727
65 years and older	7,104	6,989	7,236	3,689	4,167	2,261
Total	223,438	219,448	225,452	138,076	142,664	101,045

Continued. See notes at end of table.

**Table 9.31a. North Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018 (continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	3,977	4,701	5,477	2,258	2,455	1,768
Asian ⁴	791	788	842	627	708	561
Black or African American	74,113	71,807	71,737	42,131	44,334	32,233
Native Hawaiian or Other Pacific Islander	128	157	196	80	84	53
White	132,997	130,138	133,660	86,740	88,433	61,893
Other	10,657	10,949	11,924	5,625	6,022	4,227
Total	222,663	218,540	223,836	137,461	142,036	100,735
Ethnicity						
Not of Hispanic or Latino origin	212,036	207,861	213,138	112,014	115,515	96,321
Hispanic or Latino origin	9,523	9,773	10,606	4,814	5,428	4,407
Total	221,559	217,634	223,744	116,828	120,943	100,728
Specific Hispanic or Latino origin						
Mexican	4,218	4,256	4,505	1,846	1,915	1,356
Puerto Rican	888	879	906	462	510	382
Cuban	270	259	279	131	143	101
Other specific	4,124	4,360	4,887	2,310	2,810	2,522
Total	9,500	9,754	10,577	4,749	5,378	4,361
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2,069	1,682
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	124,358	87,432
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	126,427	89,114

-- Quantity is zero; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.31b. North Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11.9	11.3	12.1	11.9	11.8	13.1
Anxiety disorders	9.4	8.8	10.5	9.5	8.5	9.0
Attention deficit/hyperactivity disorder	9.4	7.3	6.8	2.9	3.2	2.7
Mood disorders	47.2	50.7	50.1	53.7	51.7	48.3
Bipolar disorders	19.5	21.8	19.1	18.3	18.1	15.6
Depressive disorders	27.7	28.8	31.0	35.3	33.6	32.7
Personality disorders	1.1	†	†	1.1	1.1	1.2
Schizophrenia and other psychotic disorders	9.8	10.5	10.5	12.4	13.3	14.8
Alcohol or substance-related disorders	--	--	--	--	--	--
Other mental disorders ²	11.1	10.6	9.2	8.6	10.4	10.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	47.2	47.4	47.2	48.0	49.3	51.7
Female	52.8	52.6	52.8	52.0	50.7	48.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	13.1	12.5	11.2	3.5	2.9	3.5
12 to 14 years	7.2	7.1	6.4	3.7	3.2	3.4
15 to 17 years	7.5	7.4	7.0	5.2	4.6	4.2
18 to 20 years	4.9	5.0	4.9	4.4	4.9	4.5
21 to 24 years	6.8	6.8	7.1	7.9	8.3	8.8
25 to 29 years	8.7	9.0	9.8	11.3	12.1	13.2
30 to 34 years	8.6	8.7	9.2	10.7	11.1	11.8
35 to 39 years	7.6	7.8	8.2	9.9	10.4	10.5
40 to 44 years	8.0	7.8	7.8	9.3	9.0	9.1
45 to 49 years	8.0	7.8	7.8	9.9	9.6	9.4
50 to 54 years	7.9	8.0	8.0	10.1	9.5	8.8
55 to 59 years	5.5	5.7	5.9	7.5	7.5	6.7
60 to 64 years	3.0	3.2	3.3	4.1	4.1	3.7
65 years and older	3.2	3.2	3.2	2.7	2.9	2.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.31b. North Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018 (continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	1.8	2.2	2.4	1.6	1.7	1.8
Asian ⁴	†	†	†	†	†	†
Black or African American	33.3	32.9	32.0	30.6	31.2	32.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	59.7	59.5	59.7	63.1	62.3	61.4
Other	4.8	5.0	5.3	4.1	4.2	4.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	95.7	95.5	95.3	95.9	95.5	95.6
Hispanic or Latino origin	4.3	4.5	4.7	4.1	4.5	4.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	44.4	43.6	42.6	38.9	35.6	31.1
Puerto Rican	9.3	9.0	8.6	9.7	9.5	8.8
Cuban	2.8	2.7	2.6	2.8	2.7	2.3
Other specific	43.4	44.7	46.2	48.6	52.2	57.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	1.6	1.9
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	98.4	98.1
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.32a. North Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	15,012	14,533	14,943	14,842	13,978	13,866
Mental health diagnosis¹						
Trauma- and stressor-related disorders	1,479	1,380	1,361	868	595	505
Anxiety disorders	1,572	1,678	1,705	1,189	922	776
Attention deficit/hyperactivity disorder	893	841	832	640	497	443
Mood disorders	5,328	5,288	5,263	4,062	3,201	2,778
Bipolar disorders	1,654	1,660	1,534	1,241	1,057	925
Depressive disorders	3,674	3,628	3,729	2,821	2,144	1,853
Personality disorders	163	95	117	114	108	87
Schizophrenia and other psychotic disorders	1,523	1,576	1,578	1,447	1,349	1,259
Alcohol or substance-related disorders	9	2	9	21	15	--
Other mental disorders ²	2,065	1,971	2,003	1,513	1,329	1,334
Total	13,032	12,831	12,868	9,854	8,016	7,182
Gender						
Male	7,316	7,021	7,291	7,259	6,885	7,039
Female	7,671	7,481	7,612	7,536	7,053	6,752
Total	14,987	14,502	14,903	14,795	13,938	13,791
Age³						
0 to 11 years	1,093	1,189	1,389	1,474	1,391	1,217
12 to 14 years	705	632	731	754	701	636
15 to 17 years	852	823	836	918	864	780
18 to 20 years	771	659	671	678	645	649
21 to 24 years	1,196	1,086	1,091	1,024	914	929
25 to 29 years	1,625	1,518	1,553	1,486	1,393	1,533
30 to 34 years	1,466	1,487	1,512	1,449	1,411	1,540
35 to 39 years	1,243	1,251	1,290	1,256	1,221	1,252
40 to 44 years	1,203	1,087	1,071	1,070	992	963
45 to 49 years	1,185	1,107	1,071	1,041	950	983
50 to 54 years	1,259	1,253	1,181	1,157	1,034	944
55 to 59 years	961	979	1,012	979	928	933
60 to 64 years	617	640	671	717	682	683
65 years and older	830	819	858	836	848	818
Total	15,006	14,530	14,937	14,839	13,974	13,860

Continued. See notes at end of table.

**Table 9.32a. North Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018 (continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	1,367	1,433	1,541	1,603	1,523	1,501
Asian ⁴	69	57	52	74	91	60
Black or African American	268	257	311	352	357	388
Native Hawaiian or Other Pacific Islander	18	11	16	22	17	14
White	12,892	12,384	12,578	12,239	11,447	11,315
Other	348	352	419	496	501	513
Total	14,962	14,494	14,917	14,786	13,936	13,791
Ethnicity						
Not of Hispanic or Latino origin	14,658	14,187	14,557	14,414	13,540	13,425
Hispanic or Latino origin	309	303	350	374	385	367
Total	14,967	14,490	14,907	14,788	13,925	13,792
Specific Hispanic or Latino origin						
Mexican	83	76	85	94	78	61
Puerto Rican	8	13	15	14	19	26
Cuban	5	5	3	4	2	2
Other specific	74	64	72	84	71	68
Total	170	158	175	196	170	157
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.32b. North Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11.3	10.8	10.6	8.8	7.4	7.0
Anxiety disorders	12.1	13.1	13.2	12.1	11.5	10.8
Attention deficit/hyperactivity disorder	6.9	6.6	6.5	6.5	6.2	6.2
Mood disorders	40.9	41.2	40.9	41.2	39.9	38.7
Bipolar disorders	12.7	12.9	11.9	12.6	13.2	12.9
Depressive disorders	28.2	28.3	29.0	28.6	26.7	25.8
Personality disorders	1.3	†	†	1.2	1.3	1.2
Schizophrenia and other psychotic disorders	11.7	12.3	12.3	14.7	16.8	17.5
Alcohol or substance-related disorders	†	†	†	†	†	--
Other mental disorders ²	15.8	15.4	15.6	15.4	16.6	18.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	48.8	48.4	48.9	49.1	49.4	51.0
Female	51.2	51.6	51.1	50.9	50.6	49.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	7.3	8.2	9.3	9.9	10.0	8.8
12 to 14 years	4.7	4.3	4.9	5.1	5.0	4.6
15 to 17 years	5.7	5.7	5.6	6.2	6.2	5.6
18 to 20 years	5.1	4.5	4.5	4.6	4.6	4.7
21 to 24 years	8.0	7.5	7.3	6.9	6.5	6.7
25 to 29 years	10.8	10.4	10.4	10.0	10.0	11.1
30 to 34 years	9.8	10.2	10.1	9.8	10.1	11.1
35 to 39 years	8.3	8.6	8.6	8.5	8.7	9.0
40 to 44 years	8.0	7.5	7.2	7.2	7.1	6.9
45 to 49 years	7.9	7.6	7.2	7.0	6.8	7.1
50 to 54 years	8.4	8.6	7.9	7.8	7.4	6.8
55 to 59 years	6.4	6.7	6.8	6.6	6.6	6.7
60 to 64 years	4.1	4.4	4.5	4.8	4.9	4.9
65 years and older	5.5	5.6	5.7	5.6	6.1	5.9
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.32b. North Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018 (continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	9.1	9.9	10.3	10.8	10.9	10.9
Asian ⁴	†	†	†	†	†	†
Black or African American	1.8	1.8	2.1	2.4	2.6	2.8
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	86.2	85.4	84.3	82.8	82.1	82.0
Other	2.3	2.4	2.8	3.4	3.6	3.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	97.9	97.9	97.7	97.5	97.2	97.3
Hispanic or Latino origin	2.1	2.1	2.3	2.5	2.8	2.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	48.8	48.1	48.6	48.0	45.9	38.9
Puerto Rican	4.7	8.2	8.6	7.1	11.2	16.6
Cuban	2.9	3.2	1.7	2.0	1.2	1.3
Other specific	43.5	40.5	41.1	42.9	41.8	43.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.33a. Ohio individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	323,540	372,965	415,094	430,145	531,703	538,536
Mental health diagnosis¹						
Trauma- and stressor-related disorders	51,257	56,696	63,277	69,878	81,945	81,609
Anxiety disorders	23,972	29,221	36,066	40,529	57,230	56,369
Attention deficit/hyperactivity disorder	36,461	38,717	41,678	42,247	48,854	46,037
Mood disorders	119,160	137,221	152,946	150,711	145,370	160,706
Bipolar disorders	47,037	52,876	57,789	52,505	48,193	50,839
Depressive disorders	72,123	84,345	95,157	98,206	97,177	109,867
Personality disorders	1,276	1,395	1,601	1,685	3,059	2,835
Schizophrenia and other psychotic disorders	26,975	29,170	28,773	29,290	37,068	39,583
Alcohol or substance-related disorders	19,292	29,718	20,278	26,134	203	111
Other mental disorders ²	36,858	39,064	44,780	51,587	59,177	61,890
Total	315,251	361,202	389,399	412,061	432,906	449,140
Gender						
Male	150,624	174,875	195,810	200,320	259,217	260,267
Female	171,333	195,718	214,836	222,719	261,984	266,619
Total	321,957	370,593	410,646	423,039	521,201	526,886
Age³						
0 to 11 years	66,050	68,703	75,139	77,976	82,573	80,378
12 to 14 years	30,730	32,703	35,906	36,243	38,023	38,352
15 to 17 years	28,276	30,719	35,223	37,338	41,498	40,300
18 to 20 years	14,688	15,560	18,126	19,401	24,395	23,877
21 to 24 years	15,739	19,539	23,555	24,457	33,843	31,748
25 to 29 years	22,880	27,600	32,489	34,530	52,418	52,461
30 to 34 years	23,938	29,060	33,035	34,345	50,120	50,658
35 to 39 years	20,630	25,525	29,534	31,507	44,649	46,371
40 to 44 years	20,399	24,876	27,473	27,415	34,849	35,232
45 to 49 years	20,722	25,475	27,137	27,497	33,946	34,570
50 to 54 years	21,974	27,343	29,383	28,938	33,714	33,615
55 to 59 years	17,288	21,676	23,947	24,702	29,877	31,738
60 to 64 years	9,980	12,240	13,530	14,592	18,173	21,130
65 years and older	10,243	11,939	10,614	11,199	13,619	18,104
Total	323,537	372,958	415,091	430,140	531,697	538,534

Continued. See notes at end of table.

Table 9.33a. Ohio individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	244	360	470	830	1,477	1,673
Asian ⁴	467	649	1,163	1,431	2,051	2,326
Black or African American	82,206	91,737	96,664	86,900	115,983	118,653
Native Hawaiian or Other Pacific Islander	86	101	143	154	379	459
White	225,692	257,077	279,089	286,219	347,744	350,530
Other	3,354	5,404	3,368	2,077	3,257	2,722
Total	312,049	355,328	380,897	377,611	470,891	476,363
Ethnicity						
Not of Hispanic or Latino origin	209,539	181,548	291,822	180,799	478,582	485,761
Hispanic or Latino origin	6,230	5,212	7,433	6,458	14,112	14,603
Total	215,769	186,760	299,255	187,257	492,694	500,364
Specific Hispanic or Latino origin						
Mexican	821	617	341	536	560	511
Puerto Rican	1,530	1,120	2,813	1,352	576	513
Cuban	45	42	24	30	22	26
Other specific	1,994	1,541	1,348	1,189	3,169	2,317
Total	4,390	3,320	4,526	3,107	4,327	3,367
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.33b. Ohio individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	16.3	15.7	16.2	17.0	18.9	18.2
Anxiety disorders	7.6	8.1	9.3	9.8	13.2	12.6
Attention deficit/hyperactivity disorder	11.6	10.7	10.7	10.3	11.3	10.3
Mood disorders	37.8	38.0	39.3	36.6	33.6	35.8
Bipolar disorders	14.9	14.6	14.8	12.7	11.1	11.3
Depressive disorders	22.9	23.4	24.4	23.8	22.4	24.5
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	8.6	8.1	7.4	7.1	8.6	8.8
Alcohol or substance-related disorders	6.1	8.2	5.2	6.3	†	†
Other mental disorders ²	11.7	10.8	11.5	12.5	13.7	13.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	46.8	47.2	47.7	47.4	49.7	49.4
Female	53.2	52.8	52.3	52.6	50.3	50.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	20.4	18.4	18.1	18.1	15.5	14.9
12 to 14 years	9.5	8.8	8.7	8.4	7.2	7.1
15 to 17 years	8.7	8.2	8.5	8.7	7.8	7.5
18 to 20 years	4.5	4.2	4.4	4.5	4.6	4.4
21 to 24 years	4.9	5.2	5.7	5.7	6.4	5.9
25 to 29 years	7.1	7.4	7.8	8.0	9.9	9.7
30 to 34 years	7.4	7.8	8.0	8.0	9.4	9.4
35 to 39 years	6.4	6.8	7.1	7.3	8.4	8.6
40 to 44 years	6.3	6.7	6.6	6.4	6.6	6.5
45 to 49 years	6.4	6.8	6.5	6.4	6.4	6.4
50 to 54 years	6.8	7.3	7.1	6.7	6.3	6.2
55 to 59 years	5.3	5.8	5.8	5.7	5.6	5.9
60 to 64 years	3.1	3.3	3.3	3.4	3.4	3.9
65 years and older	3.2	3.2	2.6	2.6	2.6	3.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.33b. Ohio individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	26.3	25.8	25.4	23.0	24.6	24.9
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	72.3	72.3	73.3	75.8	73.8	73.6
Other	1.1	1.5	†	†	†	†
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	97.1	97.2	97.5	96.6	97.1	97.1
Hispanic or Latino origin	2.9	2.8	2.5	3.4	2.9	2.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	18.7	18.6	7.5	17.3	12.9	15.2
Puerto Rican	34.9	33.7	62.2	43.5	13.3	15.2
Cuban	1.0	1.3	†	†	†	†
Other specific	45.4	46.4	29.8	38.3	73.2	68.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual’s age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.34a. Oklahoma individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	77,409	79,751	74,836	92,310	96,368	101,831
Mental health diagnosis¹						
Trauma- and stressor-related disorders	7,770	8,438	8,231	11,446	12,600	14,598
Anxiety disorders	8,088	5,104	4,913	8,082	9,392	9,960
Attention deficit/hyperactivity disorder	2,512	2,655	2,302	3,506	3,604	3,580
Mood disorders	33,353	40,345	34,067	41,342	42,591	42,477
Bipolar disorders	12,688	15,956	13,087	12,296	11,417	10,429
Depressive disorders	20,665	24,389	20,980	29,046	31,174	32,048
Personality disorders	1,545	66	64	108	103	141
Schizophrenia and other psychotic disorders	7,789	8,545	7,179	9,486	9,803	9,690
Alcohol or substance-related disorders	--	--	--	--	--	--
Other mental disorders ²	7,445	6,983	9,038	8,925	7,033	7,066
Total	68,502	72,136	65,794	82,895	85,126	87,512
Gender						
Male	35,062	36,363	34,903	42,394	44,570	47,871
Female	42,347	43,388	39,933	49,916	51,798	53,960
Total	77,409	79,751	74,836	92,310	96,368	101,831
Age³						
0 to 11 years	4,587	8,911	9,010	11,173	11,704	12,845
12 to 14 years	3,099	4,734	4,654	5,350	5,544	6,246
15 to 17 years	3,664	4,804	5,001	5,965	6,084	6,598
18 to 20 years	3,294	3,396	3,467	4,179	4,553	5,006
21 to 24 years	5,487	5,572	5,334	6,246	6,295	6,625
25 to 29 years	8,040	7,557	7,249	8,889	9,562	10,260
30 to 34 years	8,887	8,118	7,688	9,050	9,597	9,973
35 to 39 years	7,540	6,777	6,526	8,240	8,934	9,464
40 to 44 years	7,335	6,766	5,999	7,224	7,433	7,576
45 to 49 years	7,314	6,907	5,815	7,214	7,426	7,527
50 to 54 years	7,865	7,131	6,093	7,644	7,358	7,182
55 to 59 years	5,610	5,024	4,483	6,077	6,265	6,491
60 to 64 years	2,917	2,558	2,232	3,174	3,450	3,710
65 years and older	1,769	1,496	1,236	1,884	2,161	2,327
Total	77,408	79,751	74,787	92,309	96,366	101,830

Continued. See notes at end of table.

Table 9.34a. Oklahoma individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	4,786	6,081	6,201	7,728	8,272	8,954
Asian ⁴	303	430	433	492	549	553
Black or African American	8,597	10,212	9,598	12,231	12,715	13,138
Native Hawaiian or Other Pacific Islander	83	228	233	358	362	404
White	60,952	58,791	54,655	66,971	70,015	72,979
Other	2,666	3,997	3,696	4,515	4,448	5,799
Total	77,387	79,739	74,816	92,295	96,361	101,827
Ethnicity						
Not of Hispanic or Latino origin	72,835	75,418	69,645	85,379	89,080	93,898
Hispanic or Latino origin	4,542	4,326	5,184	6,921	7,278	7,919
Total	77,377	79,744	74,829	92,300	96,358	101,817
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	1,512	1,645
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	57,655	60,445
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	59,167	62,090

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.34b. Oklahoma individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11.3	11.7	12.5	13.8	14.8	16.7
Anxiety disorders	11.8	7.1	7.5	9.7	11.0	11.4
Attention deficit/hyperactivity disorder	3.7	3.7	3.5	4.2	4.2	4.1
Mood disorders	48.7	55.9	51.8	49.9	50.0	48.5
Bipolar disorders	18.5	22.1	19.9	14.8	13.4	11.9
Depressive disorders	30.2	33.8	31.9	35.0	36.6	36.6
Personality disorders	2.3	†	†	†	†	†
Schizophrenia and other psychotic disorders	11.4	11.8	10.9	11.4	11.5	11.1
Alcohol or substance-related disorders	--	--	--	--	--	--
Other mental disorders ²	10.9	9.7	13.7	10.8	8.3	8.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	45.3	45.6	46.6	45.9	46.2	47.0
Female	54.7	54.4	53.4	54.1	53.8	53.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	5.9	11.2	12.0	12.1	12.1	12.6
12 to 14 years	4.0	5.9	6.2	5.8	5.8	6.1
15 to 17 years	4.7	6.0	6.7	6.5	6.3	6.5
18 to 20 years	4.3	4.3	4.6	4.5	4.7	4.9
21 to 24 years	7.1	7.0	7.1	6.8	6.5	6.5
25 to 29 years	10.4	9.5	9.7	9.6	9.9	10.1
30 to 34 years	11.5	10.2	10.3	9.8	10.0	9.8
35 to 39 years	9.7	8.5	8.7	8.9	9.3	9.3
40 to 44 years	9.5	8.5	8.0	7.8	7.7	7.4
45 to 49 years	9.4	8.7	7.8	7.8	7.7	7.4
50 to 54 years	10.2	8.9	8.1	8.3	7.6	7.1
55 to 59 years	7.2	6.3	6.0	6.6	6.5	6.4
60 to 64 years	3.8	3.2	3.0	3.4	3.6	3.6
65 years and older	2.3	1.9	1.7	2.0	2.2	2.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.34b. Oklahoma individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	6.2	7.6	8.3	8.4	8.6	8.8
Asian ⁴	†	†	†	†	†	†
Black or African American	11.1	12.8	12.8	13.3	13.2	12.9
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	78.8	73.7	73.1	72.6	72.7	71.7
Other	3.4	5.0	4.9	4.9	4.6	5.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	94.1	94.6	93.1	92.5	92.4	92.2
Hispanic or Latino origin	5.9	5.4	6.9	7.5	7.6	7.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2.6	2.6
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	97.4	97.4
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.35a. Oregon individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	129,245	120,564	131,581	126,319	126,493	134,743
Mental health diagnosis¹						
Trauma- and stressor-related disorders	25,323	24,297	30,886	42,457	41,374	42,073
Anxiety disorders	9,678	8,060	13,826	18,366	27,102	25,243
Attention deficit/hyperactivity disorder	7,524	10,203	258	8,900	--	7,672
Mood disorders	25,251	21,511	35,088	35,398	35,927	34,660
Bipolar disorders	8,004	5,299	9,757	9,926	9,182	8,850
Depressive disorders	17,247	16,212	25,331	25,472	26,745	25,810
Personality disorders	231	291	338	3,143	794	691
Schizophrenia and other psychotic disorders	9,076	6,921	10,314	11,282	11,794	11,182
Alcohol or substance-related disorders	2	1	--	25	8	1
Other mental disorders ²	5,919	7,619	4,754	6,449	9,193	13,115
Total	83,004	78,903	95,464	126,020	126,192	134,637
Gender						
Male	57,523	53,822	56,142	53,793	51,201	57,158
Female	71,719	66,742	75,439	72,482	75,271	77,583
Total	129,242	120,564	131,581	126,275	126,472	134,741
Age³						
0 to 11 years	19,338	20,793	19,432	20,114	16,290	20,990
12 to 14 years	10,038	9,984	10,062	9,432	7,664	9,180
15 to 17 years	10,888	10,060	10,961	10,347	9,886	11,082
18 to 20 years	6,575	5,421	6,040	5,041	5,426	6,555
21 to 24 years	7,764	7,066	7,710	6,916	7,252	7,120
25 to 29 years	9,830	9,165	11,088	11,467	12,942	12,835
30 to 34 years	10,142	9,554	11,218	11,395	12,426	12,642
35 to 39 years	8,656	8,103	9,767	9,806	10,685	11,194
40 to 44 years	8,731	7,973	9,275	8,491	8,853	8,747
45 to 49 years	8,690	7,844	8,900	8,441	8,887	8,574
50 to 54 years	8,899	8,054	9,331	8,798	8,935	8,192
55 to 59 years	6,825	6,197	7,399	7,303	7,657	7,731
60 to 64 years	4,511	3,966	4,921	4,741	5,238	5,477
65 years and older	8,279	6,383	5,476	4,026	4,351	4,424
Total	129,166	120,563	131,580	126,318	126,492	134,743

Continued. See notes at end of table.

Table 9.35a. Oregon individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	4,178	3,769	3,873	4,929	4,103	4,396
Asian ⁴	2,214	2,041	2,102	6,639	3,668	2,075
Black or African American	5,818	5,326	5,460	4,565	4,383	4,929
Native Hawaiian or Other Pacific Islander	347	318	455	322	339	463
White	100,789	93,917	100,385	94,307	90,835	101,994
Other	20	35	58	105	6,346	5,063
Total	113,366	105,406	112,333	110,867	109,674	118,920
Ethnicity						
Not of Hispanic or Latino origin	75,215	73,313	90,760	107,047	99,599	97,478
Hispanic or Latino origin	9,606	10,076	13,638	17,961	17,865	14,914
Total	84,821	83,389	104,398	125,008	117,464	112,392
Specific Hispanic or Latino origin						
Mexican	259	280	406	1,426	5,402	2,522
Puerto Rican	47	56	73	136	346	194
Cuban	44	50	57	107	373	264
Other specific	266	314	498	996	2,734	1,356
Total	616	700	1,034	2,665	8,855	4,336
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2,026	2,303
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	48,344	61,869
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	50,370	64,172

-- Quantity is zero; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.35b. Oregon individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	30.5	30.8	32.4	33.7	32.8	31.2
Anxiety disorders	11.7	10.2	14.5	14.6	21.5	18.7
Attention deficit/hyperactivity disorder	9.1	12.9	†	7.1	--	5.7
Mood disorders	30.4	27.3	36.8	28.1	28.5	25.7
Bipolar disorders	9.6	6.7	10.2	7.9	7.3	6.6
Depressive disorders	20.8	20.5	26.5	20.2	21.2	19.2
Personality disorders	†	†	†	2.5	†	†
Schizophrenia and other psychotic disorders	10.9	8.8	10.8	9.0	9.3	8.3
Alcohol or substance-related disorders	†	†	--	†	†	†
Other mental disorders ²	7.1	9.7	5.0	5.1	7.3	9.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	44.5	44.6	42.7	42.6	40.5	42.4
Female	55.5	55.4	57.3	57.4	59.5	57.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	15.0	17.2	14.8	15.9	12.9	15.6
12 to 14 years	7.8	8.3	7.6	7.5	6.1	6.8
15 to 17 years	8.4	8.3	8.3	8.2	7.8	8.2
18 to 20 years	5.1	4.5	4.6	4.0	4.3	4.9
21 to 24 years	6.0	5.9	5.9	5.5	5.7	5.3
25 to 29 years	7.6	7.6	8.4	9.1	10.2	9.5
30 to 34 years	7.9	7.9	8.5	9.0	9.8	9.4
35 to 39 years	6.7	6.7	7.4	7.8	8.4	8.3
40 to 44 years	6.8	6.6	7.0	6.7	7.0	6.5
45 to 49 years	6.7	6.5	6.8	6.7	7.0	6.4
50 to 54 years	6.9	6.7	7.1	7.0	7.1	6.1
55 to 59 years	5.3	5.1	5.6	5.8	6.1	5.7
60 to 64 years	3.5	3.3	3.7	3.8	4.1	4.1
65 years and older	6.4	5.3	4.2	3.2	3.4	3.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.35b. Oregon individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	3.7	3.6	3.4	4.4	3.7	3.7
Asian ⁴	2.0	1.9	1.9	6.0	3.3	1.7
Black or African American	5.1	5.1	4.9	4.1	4.0	4.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	88.9	89.1	89.4	85.1	82.8	85.8
Other	†	†	†	†	5.8	4.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	88.7	87.9	86.9	85.6	84.8	86.7
Hispanic or Latino origin	11.3	12.1	13.1	14.4	15.2	13.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	42.0	40.0	39.3	53.5	61.0	58.2
Puerto Rican	7.6	8.0	7.1	5.1	3.9	4.5
Cuban	7.1	7.1	5.5	4.0	4.2	6.1
Other specific	43.2	44.9	48.2	37.4	30.9	31.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	4.0	3.6
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	96.0	96.4
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.36a. Pennsylvania individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	646,605	615,570	‡	543,618	573,439	565,437
Mental health diagnosis¹						
Trauma- and stressor-related disorders	47,739	48,458	‡	58,157	68,347	73,952
Anxiety disorders	27,382	29,742	‡	37,800	47,883	53,963
Attention deficit/hyperactivity disorder	50,211	55,251	‡	60,275	62,899	60,220
Mood disorders	139,649	175,461	‡	200,814	205,200	182,894
Bipolar disorders	48,340	68,006	‡	76,668	69,524	53,537
Depressive disorders	91,309	107,455	‡	124,146	135,676	129,357
Personality disorders	527	711	‡	1,037	1,365	1,714
Schizophrenia and other psychotic disorders	19,233	34,573	‡	37,886	39,264	38,221
Alcohol or substance-related disorders	68,855	72,543	‡	--	3	1
Other mental disorders ²	62,109	65,846	‡	64,534	73,925	76,434
Total	415,705	482,585	‡	460,503	498,886	487,399
Gender						
Male	323,328	307,288	‡	255,645	275,196	272,294
Female	323,277	307,756	‡	287,972	298,243	293,143
Total	646,605	615,044	‡	543,617	573,439	565,437
Age³						
0 to 11 years	104,284	107,728	‡	96,367	98,112	95,591
12 to 14 years	43,060	40,830	‡	39,503	41,076	39,897
15 to 17 years	46,276	39,115	‡	39,155	40,896	39,764
18 to 20 years	38,766	31,524	‡	25,524	25,703	25,073
21 to 24 years	44,368	41,611	‡	31,497	31,840	30,964
25 to 29 years	56,494	54,378	‡	45,579	49,540	49,947
30 to 34 years	51,832	49,843	‡	42,999	46,701	47,122
35 to 39 years	42,317	40,311	‡	36,355	40,703	41,805
40 to 44 years	46,182	43,830	‡	33,706	35,096	34,136
45 to 49 years	49,691	47,302	‡	36,755	38,401	36,777
50 to 54 years	46,868	44,970	‡	39,555	41,707	39,188
55 to 59 years	33,697	32,858	‡	32,253	34,897	34,826
60 to 64 years	19,664	18,444	‡	20,812	23,356	23,967
65 years and older	23,100	22,823	‡	23,556	25,408	26,378
Total	646,599	615,567	‡	543,616	573,436	565,435

Continued. See notes at end of table.

**Table 9.36a. Pennsylvania individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018 (continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	1,257	1,399	‡	960	1,026	1,114
Asian ⁴	4,481	4,327	‡	6,226	6,901	7,726
Black or African American	141,631	130,941	‡	120,988	125,986	123,194
Native Hawaiian or Other Pacific Islander	277	265	‡	269	263	289
White	427,298	413,247	‡	348,975	369,781	366,961
Other	70,316	63,073	‡	62,700	63,897	58,710
Total	645,260	613,252	‡	540,118	567,854	557,994
Ethnicity						
Not of Hispanic or Latino origin	568,731	543,982	‡	469,278	497,799	494,529
Hispanic or Latino origin	77,692	71,029	‡	72,766	74,316	68,607
Total	646,423	615,011	‡	542,044	572,115	563,136
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.36b. Pennsylvania individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	‡	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11.5	10.0	‡	12.6	13.7	15.2
Anxiety disorders	6.6	6.2	‡	8.2	9.6	11.1
Attention deficit/hyperactivity disorder	12.1	11.4	‡	13.1	12.6	12.4
Mood disorders	33.6	36.4	‡	43.6	41.1	37.5
Bipolar disorders	11.6	14.1	‡	16.6	13.9	11.0
Depressive disorders	22.0	22.3	‡	27.0	27.2	26.5
Personality disorders	†	†	‡	†	†	†
Schizophrenia and other psychotic disorders	4.6	7.2	‡	8.2	7.9	7.8
Alcohol or substance-related disorders	16.6	15.0	‡	--	†	†
Other mental disorders ²	14.9	13.6	‡	14.0	14.8	15.7
Total	100.0	100.0	‡	100.0	100.0	100.0
Gender						
Male	50.0	50.0	‡	47.0	48.0	48.2
Female	50.0	50.0	‡	53.0	52.0	51.8
Total	100.0	100.0	‡	100.0	100.0	100.0
Age³						
0 to 11 years	16.1	17.5	‡	17.7	17.1	16.9
12 to 14 years	6.7	6.6	‡	7.3	7.2	7.1
15 to 17 years	7.2	6.4	‡	7.2	7.1	7.0
18 to 20 years	6.0	5.1	‡	4.7	4.5	4.4
21 to 24 years	6.9	6.8	‡	5.8	5.6	5.5
25 to 29 years	8.7	8.8	‡	8.4	8.6	8.8
30 to 34 years	8.0	8.1	‡	7.9	8.1	8.3
35 to 39 years	6.5	6.5	‡	6.7	7.1	7.4
40 to 44 years	7.1	7.1	‡	6.2	6.1	6.0
45 to 49 years	7.7	7.7	‡	6.8	6.7	6.5
50 to 54 years	7.2	7.3	‡	7.3	7.3	6.9
55 to 59 years	5.2	5.3	‡	5.9	6.1	6.2
60 to 64 years	3.0	3.0	‡	3.8	4.1	4.2
65 years and older	3.6	3.7	‡	4.3	4.4	4.7
Total	100.0	100.0	‡	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.36b. Pennsylvania individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018 (continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	‡	†	†	†
Asian ⁴	†	†	‡	1.2	1.2	1.4
Black or African American	21.9	21.4	‡	22.4	22.2	22.1
Native Hawaiian or Other Pacific Islander	†	†	‡	†	†	†
White	66.2	67.4	‡	64.6	65.1	65.8
Other	10.9	10.3	‡	11.6	11.3	10.5
Total	100.0	100.0	‡	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	88.0	88.5	‡	86.6	87.0	87.8
Hispanic or Latino origin	12.0	11.5	‡	13.4	13.0	12.2
Total	100.0	100.0	‡	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.37a. Puerto Rico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	‡	21,387	23,194	17,567	25,529	2,918
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	2	5	5	88	105
Anxiety disorders	‡	1	3	3	49	101
Attention deficit/hyperactivity disorder	‡	31	58	57	262	193
Mood disorders	‡	5	17	17	481	898
Bipolar disorders	‡	--	3	3	93	180
Depressive disorders	‡	5	14	14	388	718
Personality disorders	‡	--	--	--	2	19
Schizophrenia and other psychotic disorders	‡	2	8	7	289	670
Alcohol or substance-related disorders	‡	--	8	8	269	328
Other mental disorders ²	‡	13	42	42	227	355
Total	‡	54	141	139	1,667	2,669
Gender						
Male	‡	11,044	12,054	9,098	13,490	1,700
Female	‡	10,341	11,139	8,468	12,037	1,218
Total	‡	21,385	23,193	17,566	25,527	2,918
Age³						
0 to 11 years	‡	1,062	1,190	650	1,065	238
12 to 14 years	‡	658	808	439	806	198
15 to 17 years	‡	855	1,064	622	1,110	214
18 to 20 years	‡	606	731	729	1,110	79
21 to 24 years	‡	1,048	1,003	664	1,005	117
25 to 29 years	‡	1,295	1,398	1,129	1,560	191
30 to 34 years	‡	1,418	1,505	1,141	1,595	196
35 to 39 years	‡	1,491	1,595	1,188	1,746	193
40 to 44 years	‡	1,614	1,696	1,237	1,738	185
45 to 49 years	‡	2,148	2,125	1,502	2,050	216
50 to 54 years	‡	2,396	2,536	1,938	2,715	287
55 to 59 years	‡	2,391	2,528	1,951	2,714	268
60 to 64 years	‡	1,880	2,081	1,685	2,427	204
65 years and older	‡	2,525	2,933	2,692	3,888	332
Total	‡	21,387	23,193	17,567	25,529	2,918

Continued. See notes at end of table.

Table 9.37a. Puerto Rico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	‡	17	36	11	66	30
Asian ⁴	‡	--	4	--	13	7
Black or African American	‡	223	260	205	309	49
Native Hawaiian or Other Pacific Islander	‡	--	--	--	10	5
White	‡	982	1,168	955	1,685	441
Other	‡	14,233	15,322	11,933	16,948	2,201
Total	‡	15,455	16,790	13,104	19,031	2,733
Ethnicity						
Not of Hispanic or Latino origin	‡	96	108	77	109	19
Hispanic or Latino origin	‡	21,054	22,865	17,297	25,177	2,881
Total	‡	21,150	22,973	17,374	25,286	2,900
Specific Hispanic or Latino origin						
Mexican	‡	26	25	23	31	4
Puerto Rican	‡	20,458	22,217	16,726	24,477	2,787
Cuban	‡	43	45	36	48	8
Other specific	‡	482	513	464	531	42
Total	‡	21,009	22,800	17,249	25,087	2,841
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	354	84
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	17,700	1,914
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	18,054	1,998

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.37b. Puerto Rico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	‡	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	3.7	3.5	3.6	5.3	3.9
Anxiety disorders	‡	1.9	2.1	2.2	2.9	3.8
Attention deficit/hyperactivity disorder	‡	57.4	41.1	41.0	15.7	7.2
Mood disorders	‡	9.3	12.1	12.2	28.9	33.6
Bipolar disorders	‡	--	2.1	2.2	5.6	6.7
Depressive disorders	‡	9.3	9.9	10.1	23.3	26.9
Personality disorders	‡	--	--	--	†	†
Schizophrenia and other psychotic disorders	‡	3.7	5.7	5.0	17.3	25.1
Alcohol or substance-related disorders	‡	--	5.7	5.8	16.1	12.3
Other mental disorders ²	‡	24.1	29.8	30.2	13.6	13.3
Total	‡	100.0	100.0	100.0	100.0	100.0
Gender						
Male	‡	51.6	52.0	51.8	52.8	58.3
Female	‡	48.4	48.0	48.2	47.2	41.7
Total	‡	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	‡	5.0	5.1	3.7	4.2	8.2
12 to 14 years	‡	3.1	3.5	2.5	3.2	6.8
15 to 17 years	‡	4.0	4.6	3.5	4.3	7.3
18 to 20 years	‡	2.8	3.2	4.1	4.3	2.7
21 to 24 years	‡	4.9	4.3	3.8	3.9	4.0
25 to 29 years	‡	6.1	6.0	6.4	6.1	6.5
30 to 34 years	‡	6.6	6.5	6.5	6.2	6.7
35 to 39 years	‡	7.0	6.9	6.8	6.8	6.6
40 to 44 years	‡	7.5	7.3	7.0	6.8	6.3
45 to 49 years	‡	10.0	9.2	8.6	8.0	7.4
50 to 54 years	‡	11.2	10.9	11.0	10.6	9.8
55 to 59 years	‡	11.2	10.9	11.1	10.6	9.2
60 to 64 years	‡	8.8	9.0	9.6	9.5	7.0
65 years and older	‡	11.8	12.6	15.3	15.2	11.4
Total	‡	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.37b. Puerto Rico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	‡	†	†	†	†	1.1
Asian ⁴	‡	--	†	--	†	†
Black or African American	‡	1.4	1.5	1.6	1.6	1.8
Native Hawaiian or Other Pacific Islander	‡	--	--	--	†	†
White	‡	6.4	7.0	7.3	8.9	16.1
Other	‡	92.1	91.3	91.1	89.1	80.5
Total	‡	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	‡	†	†	†	†	†
Hispanic or Latino origin	‡	99.5	99.5	99.6	99.6	99.3
Total	‡	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	†	†	†	†	†
Puerto Rican	‡	97.4	97.4	97.0	97.6	98.1
Cuban	‡	†	†	†	†	†
Other specific	‡	2.3	2.3	2.7	2.1	1.5
Total	‡	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2.0	4.2
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	98.0	95.8
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.38a. Rhode Island individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	32,310	30,912	32,164	31,663	31,184	26,454
Mental health diagnosis¹						
Trauma- and stressor-related disorders	3,224	2,974	2,537	2,895	3,183	2,447
Anxiety disorders	2,031	1,907	1,437	2,039	2,182	1,671
Attention deficit/hyperactivity disorder	1,470	1,325	994	1,419	1,461	1,223
Mood disorders	12,364	10,349	8,795	11,722	12,417	10,305
Bipolar disorders	5,270	4,455	3,513	3,223	3,305	2,860
Depressive disorders	7,094	5,894	5,282	8,499	9,112	7,445
Personality disorders	35	29	42	272	268	233
Schizophrenia and other psychotic disorders	2,376	2,031	1,656	3,757	3,730	3,414
Alcohol or substance-related disorders	45	36	46	--	--	--
Other mental disorders ²	1,373	1,277	990	1,711	1,671	1,344
Total	22,918	19,928	16,497	23,815	24,912	20,637
Gender						
Male	15,969	15,449	16,078	15,850	15,496	12,992
Female	16,341	15,463	16,086	15,813	15,688	13,462
Total	32,310	30,912	32,164	31,663	31,184	26,454
Age³						
0 to 11 years	3,077	3,234	3,209	3,161	3,101	2,482
12 to 14 years	1,797	1,752	1,802	1,737	1,768	1,439
15 to 17 years	2,253	2,082	2,239	2,118	2,023	1,676
18 to 20 years	1,612	1,362	1,485	1,558	1,556	1,268
21 to 24 years	1,957	1,806	1,760	1,785	1,757	1,380
25 to 29 years	2,328	2,178	2,387	2,436	2,539	1,970
30 to 34 years	2,486	2,307	2,401	2,336	2,334	2,010
35 to 39 years	2,154	2,078	2,250	2,214	2,297	1,962
40 to 44 years	2,697	2,499	2,403	2,216	2,045	1,711
45 to 49 years	3,021	2,920	2,941	2,768	2,645	2,198
50 to 54 years	3,070	3,060	3,180	3,178	2,883	2,526
55 to 59 years	2,352	2,399	2,603	2,631	2,618	2,436
60 to 64 years	1,412	1,449	1,607	1,670	1,729	1,744
65 years and older	2,094	1,786	1,896	1,855	1,889	1,652
Total	32,310	30,912	32,163	31,663	31,184	26,454

Continued. See notes at end of table.

**Table 9.38a. Rhode Island individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018 (continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	251	243	241	245	231	215
Asian ⁴	347	339	375	352	339	286
Black or African American	3,044	3,082	3,350	3,503	3,551	3,079
Native Hawaiian or Other Pacific Islander	44	54	43	43	52	58
White	22,879	21,669	22,411	22,131	21,662	18,200
Other	354	321	276	255	259	256
Total	26,919	25,708	26,696	26,529	26,094	22,094
Ethnicity						
Not of Hispanic or Latino origin	24,719	22,827	23,115	22,782	22,535	18,962
Hispanic or Latino origin	5,661	6,036	6,715	6,792	6,709	5,736
Total	30,380	28,863	29,830	29,574	29,244	24,698
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	685	550
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	20,298	17,453
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	20,983	18,003

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.38b. Rhode Island individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	14.1	14.9	15.4	12.2	12.8	11.9
Anxiety disorders	8.9	9.6	8.7	8.6	8.8	8.1
Attention deficit/hyperactivity disorder	6.4	6.6	6.0	6.0	5.9	5.9
Mood disorders	53.9	51.9	53.3	49.2	49.8	49.9
Bipolar disorders	23.0	22.4	21.3	13.5	13.3	13.9
Depressive disorders	31.0	29.6	32.0	35.7	36.6	36.1
Personality disorders	†	†	†	1.1	1.1	1.1
Schizophrenia and other psychotic disorders	10.4	10.2	10.0	15.8	15.0	16.5
Alcohol or substance-related disorders	†	†	†	--	--	--
Other mental disorders ²	6.0	6.4	6.0	7.2	6.7	6.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	49.4	50.0	50.0	50.1	49.7	49.1
Female	50.6	50.0	50.0	49.9	50.3	50.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	9.5	10.5	10.0	10.0	9.9	9.4
12 to 14 years	5.6	5.7	5.6	5.5	5.7	5.4
15 to 17 years	7.0	6.7	7.0	6.7	6.5	6.3
18 to 20 years	5.0	4.4	4.6	4.9	5.0	4.8
21 to 24 years	6.1	5.8	5.5	5.6	5.6	5.2
25 to 29 years	7.2	7.0	7.4	7.7	8.1	7.4
30 to 34 years	7.7	7.5	7.5	7.4	7.5	7.6
35 to 39 years	6.7	6.7	7.0	7.0	7.4	7.4
40 to 44 years	8.3	8.1	7.5	7.0	6.6	6.5
45 to 49 years	9.4	9.4	9.1	8.7	8.5	8.3
50 to 54 years	9.5	9.9	9.9	10.0	9.2	9.5
55 to 59 years	7.3	7.8	8.1	8.3	8.4	9.2
60 to 64 years	4.4	4.7	5.0	5.3	5.5	6.6
65 years and older	6.5	5.8	5.9	5.9	6.1	6.2
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.38b. Rhode Island individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018 (continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	1.3	1.3	1.4	1.3	1.3	1.3
Black or African American	11.3	12.0	12.5	13.2	13.6	13.9
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	85.0	84.3	83.9	83.4	83.0	82.4
Other	1.3	1.2	1.0	†	†	1.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	81.4	79.1	77.5	77.0	77.1	76.8
Hispanic or Latino origin	18.6	20.9	22.5	23.0	22.9	23.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	3.3	3.1
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	96.7	96.9
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.39a. South Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	79,301	80,982	83,286	85,254	85,754	87,192
Mental health diagnosis¹						
Trauma- and stressor-related disorders	6,864	7,381	7,964	8,336	9,105	10,564
Anxiety disorders	4,160	4,411	4,955	5,336	5,650	6,100
Attention deficit/hyperactivity disorder	9,814	9,830	10,122	10,242	10,216	10,116
Mood disorders	30,612	31,354	32,604	30,573	29,667	30,460
Bipolar disorders	12,133	12,239	12,204	10,534	10,037	10,409
Depressive disorders	18,479	19,115	20,400	20,039	19,630	20,051
Personality disorders	917	910	850	768	722	872
Schizophrenia and other psychotic disorders	15,026	15,231	15,051	14,787	14,778	15,733
Alcohol or substance-related disorders	2,098	2,091	1,841	1,635	1,666	1,912
Other mental disorders ²	7,187	7,066	6,726	8,214	8,666	8,565
Total	76,678	78,274	80,113	79,891	80,470	84,322
Gender						
Male	37,147	37,880	38,692	39,471	39,724	40,711
Female	41,992	43,088	44,459	45,770	46,022	46,472
Total	79,139	80,968	83,151	85,241	85,746	87,183
Age³						
0 to 11 years	11,289	11,478	11,657	12,159	12,622	13,252
12 to 14 years	6,912	7,048	7,234	7,336	7,338	7,765
15 to 17 years	7,117	7,411	7,729	7,912	8,003	8,197
18 to 20 years	2,839	2,799	3,089	3,252	3,301	3,451
21 to 24 years	3,774	3,874	3,972	4,059	3,921	3,964
25 to 29 years	5,179	5,289	5,624	5,768	6,068	6,042
30 to 34 years	5,520	5,673	5,749	5,937	5,758	5,810
35 to 39 years	5,184	5,335	5,529	5,797	5,776	5,782
40 to 44 years	5,791	5,841	5,747	5,607	5,304	5,365
45 to 49 years	6,522	6,370	6,383	6,249	6,157	5,815
50 to 54 years	6,966	7,140	7,165	7,129	6,746	6,432
55 to 59 years	5,527	5,750	5,961	6,132	6,240	6,226
60 to 64 years	3,503	3,634	3,817	3,994	4,227	4,463
65 years and older	3,149	3,330	3,620	3,916	4,290	4,624
Total	79,272	80,972	83,276	85,247	85,751	87,188

Continued. See notes at end of table.

**Table 9.39a. South Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018 (continued)**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	229	230	258	274	272	280
Asian ⁴	245	268	256	286	273	61
Black or African American	31,551	32,361	33,092	33,759	34,076	34,217
Native Hawaiian or Other Pacific Islander	14	12	12	15	13	16
White	43,804	44,103	45,170	46,045	45,729	45,680
Other	2,482	2,804	3,089	3,248	4,047	4,376
Total	78,325	79,778	81,877	83,627	84,410	84,630
Ethnicity						
Not of Hispanic or Latino origin	58,450	60,309	61,718	63,530	64,560	65,519
Hispanic or Latino origin	2,112	2,412	2,490	2,776	2,951	3,142
Total	60,562	62,721	64,208	66,306	67,511	68,661
Specific Hispanic or Latino origin						
Mexican	631	660	631	847	958	940
Puerto Rican	171	156	156	210	239	281
Cuban	431	566	579	506	417	368
Other specific	854	1,014	1,106	1,183	1,315	1,532
Total	2,087	2,396	2,472	2,746	2,929	3,121
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	906	836
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	36,914	57,085
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	37,820	57,921

n/c Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.39b. South Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	9.0	9.4	9.9	10.4	11.3	12.5
Anxiety disorders	5.4	5.6	6.2	6.7	7.0	7.2
Attention deficit/hyperactivity disorder	12.8	12.6	12.6	12.8	12.7	12.0
Mood disorders	39.9	40.1	40.7	38.3	36.9	36.1
Bipolar disorders	15.8	15.6	15.2	13.2	12.5	12.3
Depressive disorders	24.1	24.4	25.5	25.1	24.4	23.8
Personality disorders	1.2	1.2	1.1	†	†	1.0
Schizophrenia and other psychotic disorders	19.6	19.5	18.8	18.5	18.4	18.7
Alcohol or substance-related disorders	2.7	2.7	2.3	2.0	2.1	2.3
Other mental disorders ²	9.4	9.0	8.4	10.3	10.8	10.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	46.9	46.8	46.5	46.3	46.3	46.7
Female	53.1	53.2	53.5	53.7	53.7	53.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	14.2	14.2	14.0	14.3	14.7	15.2
12 to 14 years	8.7	8.7	8.7	8.6	8.6	8.9
15 to 17 years	9.0	9.2	9.3	9.3	9.3	9.4
18 to 20 years	3.6	3.5	3.7	3.8	3.8	4.0
21 to 24 years	4.8	4.8	4.8	4.8	4.6	4.5
25 to 29 years	6.5	6.5	6.8	6.8	7.1	6.9
30 to 34 years	7.0	7.0	6.9	7.0	6.7	6.7
35 to 39 years	6.5	6.6	6.6	6.8	6.7	6.6
40 to 44 years	7.3	7.2	6.9	6.6	6.2	6.2
45 to 49 years	8.2	7.9	7.7	7.3	7.2	6.7
50 to 54 years	8.8	8.8	8.6	8.4	7.9	7.4
55 to 59 years	7.0	7.1	7.2	7.2	7.3	7.1
60 to 64 years	4.4	4.5	4.6	4.7	4.9	5.1
65 years and older	4.0	4.1	4.3	4.6	5.0	5.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.39b. South Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018 (continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	40.3	40.6	40.4	40.4	40.4	40.4
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	55.9	55.3	55.2	55.1	54.2	54.0
Other	3.2	3.5	3.8	3.9	4.8	5.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	96.5	96.2	96.1	95.8	95.6	95.4
Hispanic or Latino origin	3.5	3.8	3.9	4.2	4.4	4.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	30.2	27.5	25.5	30.8	32.7	30.1
Puerto Rican	8.2	6.5	6.3	7.6	8.2	9.0
Cuban	20.7	23.6	23.4	18.4	14.2	11.8
Other specific	40.9	42.3	44.7	43.1	44.9	49.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2.4	1.4
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	97.6	98.6
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

† Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.40a. South Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	14,438	14,817	15,928	16,055	15,726	16,278
Mental health diagnosis¹						
Trauma- and stressor-related disorders	3,328	3,466	3,782	3,815	3,542	3,932
Anxiety disorders	947	1,054	1,174	1,216	1,365	1,561
Attention deficit/hyperactivity disorder	1,233	1,257	1,286	1,251	1,210	1,225
Mood disorders	5,179	5,340	5,739	5,039	4,952	5,290
Bipolar disorders	1,785	1,820	1,882	1,572	1,476	1,490
Depressive disorders	3,394	3,520	3,857	3,467	3,476	3,800
Personality disorders	7	8	7	8	5	5
Schizophrenia and other psychotic disorders	1,679	1,639	1,788	1,465	1,408	1,442
Alcohol or substance-related disorders	79	78	76	34	21	19
Other mental disorders ²	1,931	1,901	1,985	2,339	2,214	1,836
Total	14,383	14,743	15,837	15,167	14,717	15,310
Gender						
Male	7,051	7,205	7,709	7,569	7,339	7,509
Female	7,387	7,612	8,219	8,486	8,387	8,769
Total	14,438	14,817	15,928	16,055	15,726	16,278
Age³						
0 to 11 years	2,298	2,439	2,544	2,584	2,389	2,508
12 to 14 years	1,195	1,205	1,259	1,291	1,214	1,485
15 to 17 years	1,283	1,249	1,415	1,412	1,429	1,445
18 to 20 years	695	721	690	793	732	816
21 to 24 years	872	853	912	946	929	935
25 to 29 years	1,077	1,137	1,178	1,210	1,217	1,156
30 to 34 years	1,083	1,160	1,242	1,249	1,194	1,179
35 to 39 years	882	926	1,089	1,130	1,142	1,217
40 to 44 years	898	910	996	902	886	896
45 to 49 years	905	888	913	871	903	933
50 to 54 years	1,089	1,124	1,179	1,080	957	930
55 to 59 years	794	837	922	972	1,011	1,022
60 to 64 years	528	555	654	644	673	698
65 years and older	839	813	935	971	1,050	1,058
Total	14,438	14,817	15,928	16,055	15,726	16,278

Continued. See notes at end of table.

**Table 9.40a. South Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018 (continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	259	234	295	292	225	221
Asian ⁴	55	81	109	96	113	120
Black or African American	294	338	430	446	479	479
Native Hawaiian or Other Pacific Islander	26	23	31	17	28	18
White	10,882	11,125	11,689	11,630	11,310	11,547
Other	2,902	3,000	3,325	3,547	3,547	3,844
Total	14,418	14,801	15,879	16,028	15,702	16,229
Ethnicity						
Not of Hispanic or Latino origin	13,927	14,278	15,099	15,351	15,012	15,444
Hispanic or Latino origin	486	522	645	641	669	769
Total	14,413	14,800	15,744	15,992	15,681	16,213
Specific Hispanic or Latino origin						
Mexican	162	164	210	208	205	250
Puerto Rican	34	38	50	39	64	65
Cuban	5	9	14	7	14	16
Other specific	78	77	107	92	100	119
Total	279	288	381	346	383	450
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.40b. South Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	23.1	23.5	23.9	25.2	24.1	25.7
Anxiety disorders	6.6	7.1	7.4	8.0	9.3	10.2
Attention deficit/hyperactivity disorder	8.6	8.5	8.1	8.2	8.2	8.0
Mood disorders	36.0	36.2	36.2	33.2	33.6	34.6
Bipolar disorders	12.4	12.3	11.9	10.4	10.0	9.7
Depressive disorders	23.6	23.9	24.4	22.9	23.6	24.8
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	11.7	11.1	11.3	9.7	9.6	9.4
Alcohol or substance-related disorders	†	†	†	†	†	†
Other mental disorders ²	13.4	12.9	12.5	15.4	15.0	12.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	48.8	48.6	48.4	47.1	46.7	46.1
Female	51.2	51.4	51.6	52.9	53.3	53.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	15.9	16.5	16.0	16.1	15.2	15.4
12 to 14 years	8.3	8.1	7.9	8.0	7.7	9.1
15 to 17 years	8.9	8.4	8.9	8.8	9.1	8.9
18 to 20 years	4.8	4.9	4.3	4.9	4.7	5.0
21 to 24 years	6.0	5.8	5.7	5.9	5.9	5.7
25 to 29 years	7.5	7.7	7.4	7.5	7.7	7.1
30 to 34 years	7.5	7.8	7.8	7.8	7.6	7.2
35 to 39 years	6.1	6.2	6.8	7.0	7.3	7.5
40 to 44 years	6.2	6.1	6.3	5.6	5.6	5.5
45 to 49 years	6.3	6.0	5.7	5.4	5.7	5.7
50 to 54 years	7.5	7.6	7.4	6.7	6.1	5.7
55 to 59 years	5.5	5.6	5.8	6.1	6.4	6.3
60 to 64 years	3.7	3.7	4.1	4.0	4.3	4.3
65 years and older	5.8	5.5	5.9	6.0	6.7	6.5
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.40b. South Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018 (continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	1.8	1.6	1.9	1.8	1.4	1.4
Asian ⁴	†	†	†	†	†	†
Black or African American	2.0	2.3	2.7	2.8	3.1	3.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	75.5	75.2	73.6	72.6	72.0	71.2
Other	20.1	20.3	20.9	22.1	22.6	23.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	96.6	96.5	95.9	96.0	95.7	95.3
Hispanic or Latino origin	3.4	3.5	4.1	4.0	4.3	4.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	58.1	56.9	55.1	60.1	53.5	55.6
Puerto Rican	12.2	13.2	13.1	11.3	16.7	14.4
Cuban	1.8	3.1	3.7	2.0	3.7	3.6
Other specific	28.0	26.7	28.1	26.6	26.1	26.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.41a. Tennessee individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals¹	95,636	111,265	115,168	118,609	108,269	117,555
Mental health diagnosis²						
Trauma- and stressor-related disorders	7,479	8,268	7,560	7,798	5,860	6,604
Anxiety disorders	10,056	9,896	8,091	7,661	6,970	7,804
Attention deficit/hyperactivity disorder	5,805	5,117	4,750	5,204	2,489	2,728
Mood disorders	51,175	60,805	52,854	38,921	35,557	34,382
Bipolar disorders	23,132	26,789	22,418	13,472	12,652	12,094
Depressive disorders	28,043	34,016	30,436	25,449	22,905	22,288
Personality disorders	415	365	407	381	287	479
Schizophrenia and other psychotic disorders	7,644	11,567	9,180	7,626	7,425	7,150
Alcohol or substance-related disorders	46	6	1	--	--	--
Other mental disorders ³	4,440	9,314	4,980	8,240	4,752	4,890
Total	87,060	105,338	87,823	75,831	63,340	64,037
Gender						
Male	42,384	50,014	53,099	54,977	50,047	56,091
Female	53,196	58,957	61,449	63,050	57,823	61,100
Total	95,580	108,971	114,548	118,027	107,870	117,191
Age⁴						
0 to 11 years	7,788	8,088	8,506	8,596	3,399	7,080
12 to 14 years	4,047	5,480	5,920	6,192	5,144	6,106
15 to 17 years	4,068	6,204	6,995	7,606	7,101	6,977
18 to 20 years	3,515	4,574	4,896	5,548	5,485	5,680
21 to 24 years	7,101	8,209	8,388	8,511	8,168	8,448
25 to 29 years	9,973	11,297	11,638	12,341	12,375	13,231
30 to 34 years	10,339	11,404	11,632	12,096	11,520	12,214
35 to 39 years	9,083	10,153	10,674	11,029	10,917	11,671
40 to 44 years	9,850	10,602	10,482	10,098	9,336	9,842
45 to 49 years	10,294	10,883	10,506	10,442	9,936	10,425
50 to 54 years	9,167	10,555	10,797	10,625	9,738	9,960
55 to 59 years	6,013	7,279	7,561	7,813	7,557	8,004
60 to 64 years	2,714	3,599	3,975	4,225	4,225	4,640
65 years and older	1,667	2,934	3,197	3,486	3,368	3,275
Total	95,619	111,261	115,167	118,608	108,269	117,553

Continued. See notes at end of table.

Table 9.41a. Tennessee individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	716	515	466	406	311	280
Asian ⁵	173	258	264	229	4,845	11,378
Black or African American	16,316	17,407	21,077	22,264	21,729	31,714
Native Hawaiian or Other Pacific Islander	63	46	71	71	53	63
White	69,961	76,278	78,123	78,982	67,282	65,379
Other	764	728	878	970	1,054	1,011
Total	87,993	95,232	100,879	102,922	95,274	109,825
Ethnicity						
Not of Hispanic or Latino origin	57,147	60,855	68,245	72,370	64,690	74,513
Hispanic or Latino origin	1,986	1,959	2,261	2,559	1,726	1,743
Total	59,133	62,814	70,506	74,929	66,416	76,256
Specific Hispanic or Latino origin						
Mexican	39	34	52	42	36	23
Puerto Rican	6	10	10	9	15	10
Cuban	9	6	8	2	3	1
Other specific	43	49	43	69	49	41
Total	97	99	113	122	103	75
Veteran status⁶						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	1,269	1,322
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	31,701	36,033
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	32,970	37,355

-- Quantity is zero; *n/c* Not collected.

¹ Tennessee reports only a subset of the total individuals served through their SMHAs.

² “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

³ “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

⁴ Age is based on the individual’s age at the midpoint of the reporting period.

⁵ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁶ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.41b. Tennessee individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals¹	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis²						
Trauma- and stressor-related disorders	8.6	7.8	8.6	10.3	9.3	10.3
Anxiety disorders	11.6	9.4	9.2	10.1	11.0	12.2
Attention deficit/hyperactivity disorder	6.7	4.9	5.4	6.9	3.9	4.3
Mood disorders	58.8	57.7	60.2	51.3	56.1	53.7
Bipolar disorders	26.6	25.4	25.5	17.8	20.0	18.9
Depressive disorders	32.2	32.3	34.7	33.6	36.2	34.8
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	8.8	11.0	10.5	10.1	11.7	11.2
Alcohol or substance-related disorders	†	†	†	--	--	--
Other mental disorders ³	5.1	8.8	5.7	10.9	7.5	7.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	44.3	45.9	46.4	46.6	46.4	47.9
Female	55.7	54.1	53.6	53.4	53.6	52.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age⁴						
0 to 11 years	8.1	7.3	7.4	7.2	3.1	6.0
12 to 14 years	4.2	4.9	5.1	5.2	4.8	5.2
15 to 17 years	4.3	5.6	6.1	6.4	6.6	5.9
18 to 20 years	3.7	4.1	4.3	4.7	5.1	4.8
21 to 24 years	7.4	7.4	7.3	7.2	7.5	7.2
25 to 29 years	10.4	10.2	10.1	10.4	11.4	11.3
30 to 34 years	10.8	10.2	10.1	10.2	10.6	10.4
35 to 39 years	9.5	9.1	9.3	9.3	10.1	9.9
40 to 44 years	10.3	9.5	9.1	8.5	8.6	8.4
45 to 49 years	10.8	9.8	9.1	8.8	9.2	8.9
50 to 54 years	9.6	9.5	9.4	9.0	9.0	8.5
55 to 59 years	6.3	6.5	6.6	6.6	7.0	6.8
60 to 64 years	2.8	3.2	3.5	3.6	3.9	3.9
65 years and older	1.7	2.6	2.8	2.9	3.1	2.8
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.41b. Tennessee individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁵	†	†	†	†	5.1	10.4
Black or African American	18.5	18.3	20.9	21.6	22.8	28.9
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	79.5	80.1	77.4	76.7	70.6	59.5
Other	†	†	†	†	1.1	†
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	96.6	96.9	96.8	96.6	97.4	97.7
Hispanic or Latino origin	3.4	3.1	3.2	3.4	2.6	2.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	40.2	34.3	46.0	34.4	35.0	30.7
Puerto Rican	6.2	10.1	8.8	7.4	14.6	13.3
Cuban	9.3	6.1	7.1	1.6	2.9	1.3
Other specific	44.3	49.5	38.1	56.6	47.6	54.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁶						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	3.8	3.5
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	96.2	96.5
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals; *n/c* Not collected.

¹ Tennessee reports only a subset of the total individuals served through their SMHAs.

² “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

³ “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

⁴ Age is based on the individual’s age at the midpoint of the reporting period.

⁵ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁶ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.42a. Texas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	311,228	324,439	348,147	368,722	379,956	400,154
Mental health diagnosis¹						
Trauma- and stressor-related disorders	6,756	5,634	5,751	8,606	9,828	10,894
Anxiety disorders	2,799	2,781	2,997	4,281	4,511	5,238
Attention deficit/hyperactivity disorder	21,562	21,379	15,564	25,801	27,435	28,366
Mood disorders	144,535	144,611	111,636	151,769	156,704	158,547
Bipolar disorders	73,366	72,400	52,519	67,204	67,143	66,516
Depressive disorders	71,169	72,211	59,117	84,565	89,561	92,031
Personality disorders	501	487	474	364	312	297
Schizophrenia and other psychotic disorders	47,117	47,317	30,722	49,248	50,334	49,651
Alcohol or substance-related disorders	--	--	--	--	--	--
Other mental disorders ²	11,904	11,567	8,489	15,613	15,098	15,748
Total	235,174	233,776	175,633	255,682	264,222	268,741
Gender						
Male	155,380	161,633	173,549	185,317	190,881	200,986
Female	155,845	162,804	174,597	183,404	189,074	199,167
Total	311,225	324,437	348,146	368,721	379,955	400,153
Age³						
0 to 11 years	30,087	30,135	32,239	34,938	36,013	38,926
12 to 14 years	16,819	18,254	20,422	21,903	22,382	24,985
15 to 17 years	18,386	20,110	22,592	24,359	25,250	27,088
18 to 20 years	14,387	14,461	15,671	17,040	18,213	20,027
21 to 24 years	21,919	22,189	23,763	24,919	25,196	26,642
25 to 29 years	30,044	30,741	33,045	35,655	37,162	39,416
30 to 34 years	30,429	31,540	33,406	35,509	36,276	38,271
35 to 39 years	25,527	26,809	29,335	32,320	33,719	36,120
40 to 44 years	26,813	26,953	27,607	27,964	28,193	28,941
45 to 49 years	28,812	28,367	28,739	29,595	29,643	29,686
50 to 54 years	28,801	30,088	31,165	31,699	31,110	30,001
55 to 59 years	20,625	22,403	24,695	26,011	27,092	27,543
60 to 64 years	10,801	12,394	13,943	15,105	16,386	17,548
65 years and older	7,774	9,993	11,521	11,703	13,319	14,957
Total	311,224	324,437	348,143	368,720	379,954	400,151

Continued. See notes at end of table.

Table 9.42a. Texas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	831	869	966	1,092	1,158	1,277
Asian ⁴	2,548	2,495	2,704	2,818	2,947	3,297
Black or African American	75,859	77,606	81,617	84,826	84,752	86,400
Native Hawaiian or Other Pacific Islander	80	129	155	185	250	309
White	221,762	232,532	250,913	266,234	274,229	291,333
Other	10,134	10,798	11,777	13,557	16,607	17,525
Total	311,214	324,429	348,132	368,712	379,943	400,141
Ethnicity						
Not of Hispanic or Latino origin	229,452	240,103	256,543	272,116	281,244	295,056
Hispanic or Latino origin	81,754	84,311	91,576	96,582	98,684	105,075
Total	311,206	324,414	348,119	368,698	379,928	400,131
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.42b. Texas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	2.9	2.4	3.3	3.4	3.7	4.1
Anxiety disorders	1.2	1.2	1.7	1.7	1.7	1.9
Attention deficit/hyperactivity disorder	9.2	9.1	8.9	10.1	10.4	10.6
Mood disorders	61.5	61.9	63.6	59.4	59.3	59.0
Bipolar disorders	31.2	31.0	29.9	26.3	25.4	24.8
Depressive disorders	30.3	30.9	33.7	33.1	33.9	34.2
Personality disorders	†	†	†	†	†	†
Schizophrenia and other psychotic disorders	20.0	20.2	17.5	19.3	19.0	18.5
Alcohol or substance-related disorders	--	--	--	--	--	--
Other mental disorders ²	5.1	4.9	4.8	6.1	5.7	5.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	49.9	49.8	49.8	50.3	50.2	50.2
Female	50.1	50.2	50.2	49.7	49.8	49.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	9.7	9.3	9.3	9.5	9.5	9.7
12 to 14 years	5.4	5.6	5.9	5.9	5.9	6.2
15 to 17 years	5.9	6.2	6.5	6.6	6.6	6.8
18 to 20 years	4.6	4.5	4.5	4.6	4.8	5.0
21 to 24 years	7.0	6.8	6.8	6.8	6.6	6.7
25 to 29 years	9.7	9.5	9.5	9.7	9.8	9.9
30 to 34 years	9.8	9.7	9.6	9.6	9.5	9.6
35 to 39 years	8.2	8.3	8.4	8.8	8.9	9.0
40 to 44 years	8.6	8.3	7.9	7.6	7.4	7.2
45 to 49 years	9.3	8.7	8.3	8.0	7.8	7.4
50 to 54 years	9.3	9.3	9.0	8.6	8.2	7.5
55 to 59 years	6.6	6.9	7.1	7.1	7.1	6.9
60 to 64 years	3.5	3.8	4.0	4.1	4.3	4.4
65 years and older	2.5	3.1	3.3	3.2	3.5	3.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.42b. Texas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	†	†	†	†	†	†
Black or African American	24.4	23.9	23.4	23.0	22.3	21.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	71.3	71.7	72.1	72.2	72.2	72.8
Other	3.3	3.3	3.4	3.7	4.4	4.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	73.7	74.0	73.7	73.8	74.0	73.7
Hispanic or Latino origin	26.3	26.0	26.3	26.2	26.0	26.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.43b. Utah individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11.4	12.7	13.9	13.5	14.8	14.5
Anxiety disorders	14.6	15.8	15.1	15.5	17.6	17.5
Attention deficit/hyperactivity disorder	3.1	3.1	3.3	4.1	4.2	3.8
Mood disorders	45.2	43.7	43.1	32.5	37.0	38.2
Bipolar disorders	17.8	17.0	16.1	7.6	8.3	8.6
Depressive disorders	27.4	26.8	27.0	24.9	28.6	29.5
Personality disorders	1.2	1.1	1.5	1.9	1.4	1.2
Schizophrenia and other psychotic disorders	12.0	11.7	11.0	9.6	11.0	11.5
Alcohol or substance-related disorders	--	--	†	9.0	†	†
Other mental disorders ²	12.6	11.9	11.4	14.0	14.0	13.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	46.9	47.4	47.7	47.7	47.7	48.2
Female	53.1	52.6	52.3	52.3	52.3	51.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	22.9	22.6	22.9	22.6	22.4	21.5
12 to 14 years	8.7	9.2	9.3	9.6	9.6	10.0
15 to 17 years	6.2	6.5	7.2	7.6	7.8	8.0
18 to 20 years	3.3	3.2	3.3	3.2	3.4	3.7
21 to 24 years	5.4	5.6	5.4	5.4	5.4	5.2
25 to 29 years	8.3	8.3	8.0	8.0	8.1	8.0
30 to 34 years	8.8	8.6	8.5	8.3	8.4	8.1
35 to 39 years	7.1	7.2	7.4	7.5	7.9	7.9
40 to 44 years	6.2	6.1	6.0	6.0	5.8	5.9
45 to 49 years	6.0	5.5	5.2	5.1	5.1	5.3
50 to 54 years	5.9	5.8	5.6	5.3	4.8	4.7
55 to 59 years	4.4	4.4	4.3	4.3	4.4	4.4
60 to 64 years	2.8	2.8	2.7	2.8	2.9	3.0
65 years and older	4.1	4.1	4.1	4.2	4.2	4.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.43b. Utah individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	2.1	2.1	2.1	2.1	1.9	2.6
Asian ⁴	†	1.0	†	†	†	†
Black or African American	3.1	3.0	3.1	3.0	2.5	2.4
Native Hawaiian or Other Pacific Islander	8.0	8.8	10.0	12.2	†	2.0
White	75.6	74.8	74.0	71.7	82.6	81.3
Other	10.3	10.3	9.9	10.0	11.5	11.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	85.6	85.4	85.1	84.3	84.3	84.2
Hispanic or Latino origin	14.4	14.6	14.9	15.7	15.7	15.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	2.6	2.7
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	97.4	97.3
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.43a. Utah individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	45,809	48,567	51,080	53,708	55,525	56,095
Mental health diagnosis¹						
Trauma- and stressor-related disorders	4,699	5,383	6,171	6,428	7,163	7,028
Anxiety disorders	6,030	6,662	6,693	7,406	8,549	8,436
Attention deficit/hyperactivity disorder	1,262	1,320	1,455	1,952	2,016	1,859
Mood disorders	18,657	18,473	19,156	15,499	17,939	18,444
Bipolar disorders	7,340	7,170	7,139	3,631	4,048	4,168
Depressive disorders	11,317	11,303	12,017	11,868	13,891	14,276
Personality disorders	513	457	654	900	694	561
Schizophrenia and other psychotic disorders	4,950	4,930	4,883	4,574	5,336	5,575
Alcohol or substance-related disorders	--	--	345	4,287	17	22
Other mental disorders ²	5,204	5,027	5,089	6,662	6,785	6,394
Total	41,315	42,252	44,446	47,708	48,499	48,319
Gender						
Male	21,493	23,011	24,379	25,645	26,496	27,059
Female	24,316	25,556	26,701	28,063	29,029	29,036
Total	45,809	48,567	51,080	53,708	55,525	56,095
Age³						
0 to 11 years	10,487	10,984	11,695	12,158	12,433	12,034
12 to 14 years	3,966	4,485	4,766	5,171	5,357	5,586
15 to 17 years	2,844	3,165	3,674	4,081	4,327	4,513
18 to 20 years	1,525	1,578	1,693	1,744	1,868	2,055
21 to 24 years	2,489	2,697	2,742	2,924	2,972	2,912
25 to 29 years	3,807	4,045	4,078	4,299	4,476	4,501
30 to 34 years	4,013	4,172	4,364	4,438	4,652	4,556
35 to 39 years	3,241	3,515	3,780	4,013	4,367	4,405
40 to 44 years	2,834	2,952	3,067	3,249	3,202	3,313
45 to 49 years	2,744	2,671	2,674	2,729	2,828	2,946
50 to 54 years	2,699	2,805	2,843	2,839	2,676	2,662
55 to 59 years	2,024	2,127	2,213	2,327	2,439	2,472
60 to 64 years	1,260	1,364	1,387	1,500	1,588	1,704
65 years and older	1,875	2,007	2,103	2,236	2,337	2,435
Total	45,808	48,567	51,079	53,708	55,522	56,094

Continued. See notes at end of table.

Table 9.43a. Utah individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	908	926	991	1,048	977	1,346
Asian ⁴	422	457	466	477	387	395
Black or African American	1,339	1,336	1,439	1,508	1,278	1,228
Native Hawaiian or Other Pacific Islander	3,480	3,908	4,656	6,068	347	1,023
White	33,038	33,130	34,514	35,550	42,165	41,902
Other	4,493	4,543	4,606	4,951	5,868	5,666
Total	43,680	44,300	46,672	49,602	51,022	51,560
Ethnicity						
Not of Hispanic or Latino origin	37,830	38,310	40,195	42,188	43,090	42,977
Hispanic or Latino origin	6,369	6,542	7,034	7,837	8,044	8,065
Total	44,199	44,852	47,229	50,025	51,134	51,042
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	757	780
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	28,498	28,460
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	29,255	29,240

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.44a. Vermont individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	24,435	24,613	24,580	31,244	30,924	31,263
Mental health diagnosis¹						
Trauma- and stressor-related disorders	6,255	5,821	5,519	6,272	6,749	7,648
Anxiety disorders	2,147	2,161	1,926	2,331	2,507	2,699
Attention deficit/hyperactivity disorder	1,476	1,458	1,263	1,554	1,540	1,515
Mood disorders	6,225	5,558	4,891	4,350	4,482	4,769
Bipolar disorders	1,717	1,554	1,406	1,048	1,080	1,085
Depressive disorders	4,508	4,004	3,485	3,302	3,402	3,684
Personality disorders	113	91	98	350	344	355
Schizophrenia and other psychotic disorders	1,645	1,250	1,139	1,584	1,603	1,640
Alcohol or substance-related disorders	188	12	22	1,030	1,191	4,871
Other mental disorders ²	2,343	2,184	1,833	2,638	2,594	5,531
Total	20,392	18,535	16,691	20,109	21,010	29,028
Gender						
Male	12,263	12,290	12,234	16,182	16,112	16,332
Female	12,152	12,306	12,324	14,989	14,772	14,865
Total	24,415	24,596	24,558	31,171	30,884	31,197
Age³						
0 to 11 years	5,196	5,386	5,220	5,324	5,279	5,609
12 to 14 years	2,180	2,161	2,232	2,207	2,232	2,283
15 to 17 years	2,194	2,216	2,279	2,507	2,433	2,657
18 to 20 years	1,470	1,445	1,408	1,783	1,764	1,765
21 to 24 years	1,438	1,410	1,433	2,114	2,008	1,996
25 to 29 years	1,641	1,545	1,587	2,825	2,712	2,620
30 to 34 years	1,500	1,557	1,524	2,655	2,625	2,463
35 to 39 years	1,231	1,243	1,299	1,986	2,093	2,091
40 to 44 years	1,315	1,279	1,197	1,665	1,606	1,575
45 to 49 years	1,402	1,334	1,320	1,714	1,690	1,687
50 to 54 years	1,466	1,494	1,384	1,820	1,764	1,690
55 to 59 years	1,179	1,245	1,235	1,616	1,657	1,623
60 to 64 years	885	853	941	1,100	1,152	1,212
65 years and older	1,312	1,436	1,495	1,928	1,908	1,992
Total	24,409	24,604	24,554	31,244	30,923	31,263

Continued. See notes at end of table.

Table 9.44a. Vermont individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	268	271	278	253	244	250
Asian ⁴	137	170	159	162	160	189
Black or African American	550	555	612	585	593	564
Native Hawaiian or Other Pacific Islander	--	--	--	--	--	--
White	21,686	21,882	21,674	20,520	20,097	20,107
Other	499	538	516	502	542	608
Total	23,140	23,416	23,239	22,022	21,636	21,718
Ethnicity						
Not of Hispanic or Latino origin	21,324	21,470	21,289	19,792	19,093	19,086
Hispanic or Latino origin	782	696	706	605	588	642
Total	22,106	22,166	21,995	20,397	19,681	19,728
Specific Hispanic or Latino origin						
Mexican	102	107	113	94	106	123
Puerto Rican	117	124	120	123	123	130
Cuban	15	22	22	25	22	23
Other specific	150	156	181	173	186	178
Total	384	409	436	415	437	454
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	555	518
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	10,800	10,717
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	11,355	11,235

-- Quantity is zero; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.44b. Vermont individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	30.7	31.4	33.1	31.2	32.1	26.3
Anxiety disorders	10.5	11.7	11.5	11.6	11.9	9.3
Attention deficit/hyperactivity disorder	7.2	7.9	7.6	7.7	7.3	5.2
Mood disorders	30.5	30.0	29.3	21.6	21.3	16.4
Bipolar disorders	8.4	8.4	8.4	5.2	5.1	3.7
Depressive disorders	22.1	21.6	20.9	16.4	16.2	12.7
Personality disorders	†	†	†	1.7	1.6	1.2
Schizophrenia and other psychotic disorders	8.1	6.7	6.8	7.9	7.6	5.6
Alcohol or substance-related disorders	†	†	†	5.1	5.7	16.8
Other mental disorders ²	11.5	11.8	11.0	13.1	12.3	19.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	50.2	50.0	49.8	51.9	52.2	52.4
Female	49.8	50.0	50.2	48.1	47.8	47.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	21.3	21.9	21.3	17.0	17.1	17.9
12 to 14 years	8.9	8.8	9.1	7.1	7.2	7.3
15 to 17 years	9.0	9.0	9.3	8.0	7.9	8.5
18 to 20 years	6.0	5.9	5.7	5.7	5.7	5.6
21 to 24 years	5.9	5.7	5.8	6.8	6.5	6.4
25 to 29 years	6.7	6.3	6.5	9.0	8.8	8.4
30 to 34 years	6.1	6.3	6.2	8.5	8.5	7.9
35 to 39 years	5.0	5.1	5.3	6.4	6.8	6.7
40 to 44 years	5.4	5.2	4.9	5.3	5.2	5.0
45 to 49 years	5.7	5.4	5.4	5.5	5.5	5.4
50 to 54 years	6.0	6.1	5.6	5.8	5.7	5.4
55 to 59 years	4.8	5.1	5.0	5.2	5.4	5.2
60 to 64 years	3.6	3.5	3.8	3.5	3.7	3.9
65 years and older	5.4	5.8	6.1	6.2	6.2	6.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.44b. Vermont individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	1.2	1.2	1.2	1.1	1.1	1.2
Asian ⁴	†	†	†	†	†	†
Black or African American	2.4	2.4	2.6	2.7	2.7	2.6
Native Hawaiian or Other Pacific Islander	--	--	--	--	--	--
White	93.7	93.4	93.3	93.2	92.9	92.6
Other	2.2	2.3	2.2	2.3	2.5	2.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	96.5	96.9	96.8	97.0	97.0	96.7
Hispanic or Latino origin	3.5	3.1	3.2	3.0	3.0	3.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	26.6	26.2	25.9	22.7	24.3	27.1
Puerto Rican	30.5	30.3	27.5	29.6	28.1	28.6
Cuban	3.9	5.4	5.0	6.0	5.0	5.1
Other specific	39.1	38.1	41.5	41.7	42.6	39.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	4.9	4.6
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	95.1	95.4
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.45a. Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	109,699	112,577	115,824	117,415	122,624	123,101
Mental health diagnosis¹						
Trauma- and stressor-related disorders	10,441	11,074	11,820	11,798	13,114	15,114
Anxiety disorders	10,897	12,057	12,738	12,910	13,543	14,530
Attention deficit/hyperactivity disorder	10,010	10,438	10,229	11,355	11,423	11,621
Mood disorders	40,390	41,232	42,399	36,521	37,556	38,957
Bipolar disorders	17,646	18,370	18,507	13,077	13,088	13,368
Depressive disorders	22,744	22,862	23,892	23,444	24,468	25,589
Personality disorders	4,560	4,360	4,288	3,731	3,550	3,583
Schizophrenia and other psychotic disorders	15,652	15,829	16,037	16,628	16,945	17,143
Alcohol or substance-related disorders	--	--	--	--	2	2
Other mental disorders ²	11,934	11,820	12,353	18,281	17,425	16,311
Total	103,884	106,810	109,864	111,224	113,558	117,261
Gender						
Male	54,777	56,379	58,354	59,001	61,403	61,867
Female	54,901	56,183	57,452	58,400	61,187	61,189
Total	109,678	112,562	115,806	117,401	122,590	123,056
Age³						
0 to 11 years	14,609	15,054	15,219	15,290	15,993	15,730
12 to 14 years	7,961	8,334	8,594	8,459	8,847	9,075
15 to 17 years	8,626	9,279	10,153	10,690	10,783	10,745
18 to 20 years	4,582	4,618	4,710	4,814	5,317	5,366
21 to 24 years	5,859	6,113	6,121	6,109	6,416	6,447
25 to 29 years	8,460	8,577	8,987	9,159	9,850	9,915
30 to 34 years	8,428	8,471	8,807	9,160	9,711	9,824
35 to 39 years	7,253	7,590	7,866	8,229	8,939	9,271
40 to 44 years	8,123	7,998	7,834	7,604	7,684	7,832
45 to 49 years	9,132	9,007	8,891	8,619	8,734	8,504
50 to 54 years	9,438	9,555	9,737	9,732	9,717	9,268
55 to 59 years	7,342	7,738	8,038	8,133	8,509	8,623
60 to 64 years	4,686	4,932	5,218	5,512	5,844	5,941
65 years and older	5,197	5,305	5,640	5,891	6,276	6,553
Total	109,696	112,571	115,815	117,401	122,620	123,094

Continued. See notes at end of table.

Table 9.45a. Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	254	264	273	279	227	389
Asian ⁴	1,159	1,185	1,256	1,270	1,352	1,421
Black or African American	30,957	31,921	33,011	33,584	34,890	33,967
Native Hawaiian or Other Pacific Islander	84	78	89	85	82	93
White	68,268	69,389	70,767	71,323	73,594	74,726
Other	8,050	8,785	9,437	9,916	10,665	10,945
Total	108,772	111,622	114,833	116,457	120,810	121,541
Ethnicity						
Not of Hispanic or Latino origin	101,103	103,281	106,063	107,310	110,782	110,838
Hispanic or Latino origin	6,480	6,847	7,325	7,522	8,232	8,815
Total	107,583	110,128	113,388	114,832	119,014	119,653
Specific Hispanic or Latino origin						
Mexican	847	912	887	916	973	1,015
Puerto Rican	861	916	922	895	975	1,016
Cuban	127	123	124	127	125	115
Other specific	3,514	3,626	3,953	3,962	4,297	4,529
Total	5,349	5,577	5,886	5,900	6,370	6,675
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.45b. Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	10.1	10.4	10.8	10.6	11.5	12.9
Anxiety disorders	10.5	11.3	11.6	11.6	11.9	12.4
Attention deficit/hyperactivity disorder	9.6	9.8	9.3	10.2	10.1	9.9
Mood disorders	38.9	38.6	38.6	32.8	33.1	33.2
Bipolar disorders	17.0	17.2	16.8	11.8	11.5	11.4
Depressive disorders	21.9	21.4	21.7	21.1	21.5	21.8
Personality disorders	4.4	4.1	3.9	3.4	3.1	3.1
Schizophrenia and other psychotic disorders	15.1	14.8	14.6	15.0	14.9	14.6
Alcohol or substance-related disorders	--	--	--	--	†	†
Other mental disorders ²	11.5	11.1	11.2	16.4	15.3	13.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	49.9	50.1	50.4	50.3	50.1	50.3
Female	50.1	49.9	49.6	49.7	49.9	49.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	13.3	13.4	13.1	13.0	13.0	12.8
12 to 14 years	7.3	7.4	7.4	7.2	7.2	7.4
15 to 17 years	7.9	8.2	8.8	9.1	8.8	8.7
18 to 20 years	4.2	4.1	4.1	4.1	4.3	4.4
21 to 24 years	5.3	5.4	5.3	5.2	5.2	5.2
25 to 29 years	7.7	7.6	7.8	7.8	8.0	8.1
30 to 34 years	7.7	7.5	7.6	7.8	7.9	8.0
35 to 39 years	6.6	6.7	6.8	7.0	7.3	7.5
40 to 44 years	7.4	7.1	6.8	6.5	6.3	6.4
45 to 49 years	8.3	8.0	7.7	7.3	7.1	6.9
50 to 54 years	8.6	8.5	8.4	8.3	7.9	7.5
55 to 59 years	6.7	6.9	6.9	6.9	6.9	7.0
60 to 64 years	4.3	4.4	4.5	4.7	4.8	4.8
65 years and older	4.7	4.7	4.9	5.0	5.1	5.3
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.45b. Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	†	†	†	†	†	†
Asian ⁴	1.1	1.1	1.1	1.1	1.1	1.2
Black or African American	28.5	28.6	28.7	28.8	28.9	27.9
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	62.8	62.2	61.6	61.2	60.9	61.5
Other	7.4	7.9	8.2	8.5	8.8	9.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	94.0	93.8	93.5	93.4	93.1	92.6
Hispanic or Latino origin	6.0	6.2	6.5	6.6	6.9	7.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	15.8	16.4	15.1	15.5	15.3	15.2
Puerto Rican	16.1	16.4	15.7	15.2	15.3	15.2
Cuban	2.4	2.2	2.1	2.2	2.0	1.7
Other specific	65.7	65.0	67.2	67.2	67.5	67.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

- - Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.46a. Washington individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	177,858	194,591	224,463	249,446	278,556	290,864
Mental health diagnosis¹						
Trauma- and stressor-related disorders	28,154	31,135	36,355	42,452	51,660	56,177
Anxiety disorders	19,189	23,567	29,522	41,653	51,532	56,200
Attention deficit/hyperactivity disorder	14,414	15,246	16,167	17,921	20,092	20,445
Mood disorders	69,858	78,991	94,341	93,117	90,572	88,085
Bipolar disorders	19,933	21,275	22,750	19,722	18,826	18,161
Depressive disorders	49,925	57,716	71,591	73,395	71,746	69,924
Personality disorders	259	230	238	1,080	3,422	4,059
Schizophrenia and other psychotic disorders	18,186	18,566	19,671	19,791	20,283	20,155
Alcohol or substance-related disorders	2,154	2,258	2,378	2,638	3,716	3,850
Other mental disorders ²	9,091	8,926	9,111	16,402	19,187	20,505
Total	161,305	178,919	207,783	235,054	260,464	269,476
Gender						
Male	83,180	89,630	101,915	110,832	123,232	129,591
Female	94,430	104,829	122,344	137,639	153,966	159,786
Total	177,610	194,459	224,259	248,471	277,198	289,377
Age³						
0 to 11 years	26,610	26,940	28,448	30,750	33,504	34,133
12 to 14 years	14,188	14,622	15,479	16,513	18,313	20,187
15 to 17 years	15,212	16,382	18,150	19,904	21,652	23,052
18 to 20 years	9,505	10,278	11,904	13,722	15,880	17,341
21 to 24 years	9,398	10,270	12,631	14,738	17,126	18,427
25 to 29 years	13,808	16,033	20,357	23,581	27,500	28,729
30 to 34 years	14,247	16,513	20,616	23,611	27,007	28,151
35 to 39 years	11,848	13,905	17,674	20,579	23,469	24,850
40 to 44 years	11,802	12,870	14,545	15,757	17,749	18,677
45 to 49 years	11,837	13,078	15,380	16,636	17,715	17,391
50 to 54 years	12,815	14,372	16,178	16,595	17,342	17,010
55 to 59 years	9,950	11,311	13,257	14,759	16,390	16,845
60 to 64 years	6,366	7,501	8,922	10,260	11,476	12,169
65 years and older	10,063	10,427	10,766	11,317	12,342	12,623
Total	177,649	194,502	224,307	248,722	277,465	289,585

Continued. See notes at end of table.

Table 9.46a. Washington individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	6,293	6,920	7,879	8,452	9,221	8,704
Asian ⁴	3,706	4,238	5,009	5,466	6,443	6,691
Black or African American	15,371	15,994	17,890	18,861	20,213	20,810
Native Hawaiian or Other Pacific Islander	2,135	2,419	2,945	3,357	3,814	4,141
White	128,800	141,743	163,919	182,033	201,573	209,195
Other	19,024	20,819	24,027	27,183	31,189	33,937
Total	175,329	192,133	221,669	245,352	272,453	283,478
Ethnicity						
Not of Hispanic or Latino origin	144,866	158,214	181,745	199,850	219,238	225,363
Hispanic or Latino origin	28,691	31,632	37,072	41,481	47,311	51,700
Total	173,557	189,846	218,817	241,331	266,549	277,063
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	4,349	4,386
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	99,408	105,086
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	103,757	109,472

‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.46b. Washington individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	17.5	17.4	17.5	18.1	19.8	20.8
Anxiety disorders	11.9	13.2	14.2	17.7	19.8	20.9
Attention deficit/hyperactivity disorder	8.9	8.5	7.8	7.6	7.7	7.6
Mood disorders	43.3	44.1	45.4	39.6	34.8	32.7
Bipolar disorders	12.4	11.9	10.9	8.4	7.2	6.7
Depressive disorders	31.0	32.3	34.5	31.2	27.5	25.9
Personality disorders	†	†	†	†	1.3	1.5
Schizophrenia and other psychotic disorders	11.3	10.4	9.5	8.4	7.8	7.5
Alcohol or substance-related disorders	1.3	1.3	1.1	1.1	1.4	1.4
Other mental disorders ²	5.6	5.0	4.4	7.0	7.4	7.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	46.8	46.1	45.4	44.6	44.5	44.8
Female	53.2	53.9	54.6	55.4	55.5	55.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	15.0	13.9	12.7	12.4	12.1	11.8
12 to 14 years	8.0	7.5	6.9	6.6	6.6	7.0
15 to 17 years	8.6	8.4	8.1	8.0	7.8	8.0
18 to 20 years	5.4	5.3	5.3	5.5	5.7	6.0
21 to 24 years	5.3	5.3	5.6	5.9	6.2	6.4
25 to 29 years	7.8	8.2	9.1	9.5	9.9	9.9
30 to 34 years	8.0	8.5	9.2	9.5	9.7	9.7
35 to 39 years	6.7	7.1	7.9	8.3	8.5	8.6
40 to 44 years	6.6	6.6	6.5	6.3	6.4	6.4
45 to 49 years	6.7	6.7	6.9	6.7	6.4	6.0
50 to 54 years	7.2	7.4	7.2	6.7	6.3	5.9
55 to 59 years	5.6	5.8	5.9	5.9	5.9	5.8
60 to 64 years	3.6	3.9	4.0	4.1	4.1	4.2
65 years and older	5.7	5.4	4.8	4.6	4.4	4.4
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.46b. Washington individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	3.6	3.6	3.6	3.4	3.4	3.1
Asian ⁴	2.1	2.2	2.3	2.2	2.4	2.4
Black or African American	8.8	8.3	8.1	7.7	7.4	7.3
Native Hawaiian or Other Pacific Islander	1.2	1.3	1.3	1.4	1.4	1.5
White	73.5	73.8	73.9	74.2	74.0	73.8
Other	10.9	10.8	10.8	11.1	11.4	12.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	83.5	83.3	83.1	82.8	82.3	81.3
Hispanic or Latino origin	16.5	16.7	16.9	17.2	17.7	18.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	4.2	4.0
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	95.8	96.0
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual’s age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.47a. West Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	‡	‡	‡	‡	‡	50,605
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	‡	‡	‡	3,657
Anxiety disorders	‡	‡	‡	‡	‡	7,412
Attention deficit/hyperactivity disorder	‡	‡	‡	‡	‡	3,302
Mood disorders	‡	‡	‡	‡	‡	18,224
Bipolar disorders	‡	‡	‡	‡	‡	5,767
Depressive disorders	‡	‡	‡	‡	‡	12,457
Personality disorders	‡	‡	‡	‡	‡	254
Schizophrenia and other psychotic disorders	‡	‡	‡	‡	‡	2,262
Alcohol or substance-related disorders	‡	‡	‡	‡	‡	4,205
Other mental disorders ²	‡	‡	‡	‡	‡	3,666
Total	‡	‡	‡	‡	‡	42,982
Gender						
Male	‡	‡	‡	‡	‡	22,982
Female	‡	‡	‡	‡	‡	27,606
Total	‡	‡	‡	‡	‡	50,588
Age³						
0 to 11 years	‡	‡	‡	‡	‡	3,478
12 to 14 years	‡	‡	‡	‡	‡	2,322
15 to 17 years	‡	‡	‡	‡	‡	2,947
18 to 20 years	‡	‡	‡	‡	‡	2,726
21 to 24 years	‡	‡	‡	‡	‡	3,265
25 to 29 years	‡	‡	‡	‡	‡	5,186
30 to 34 years	‡	‡	‡	‡	‡	4,798
35 to 39 years	‡	‡	‡	‡	‡	4,896
40 to 44 years	‡	‡	‡	‡	‡	4,292
45 to 49 years	‡	‡	‡	‡	‡	4,036
50 to 54 years	‡	‡	‡	‡	‡	3,820
55 to 59 years	‡	‡	‡	‡	‡	3,525
60 to 64 years	‡	‡	‡	‡	‡	2,515
65 years and older	‡	‡	‡	‡	‡	2,799
Total	‡	‡	‡	‡	‡	50,605

Continued. See notes at end of table.

**Table 9.47a. West Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018 (continued)**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	‡	‡	‡	‡	‡	68
Asian ⁴	‡	‡	‡	‡	‡	55
Black or African American	‡	‡	‡	‡	‡	1,967
Native Hawaiian or Other Pacific Islander	‡	‡	‡	‡	‡	26
White	‡	‡	‡	‡	‡	42,516
Other	‡	‡	‡	‡	‡	5,225
Total	‡	‡	‡	‡	‡	49,857
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	‡	‡	‡
Hispanic or Latino origin	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	972
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	34,658
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	35,630

‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.47b. West Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	‡	‡	‡	‡	‡	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	‡	‡	‡	‡	‡	8.5
Anxiety disorders	‡	‡	‡	‡	‡	17.2
Attention deficit/hyperactivity disorder	‡	‡	‡	‡	‡	7.7
Mood disorders	‡	‡	‡	‡	‡	42.4
Bipolar disorders	‡	‡	‡	‡	‡	13.4
Depressive disorders	‡	‡	‡	‡	‡	29.0
Personality disorders	‡	‡	‡	‡	‡	†
Schizophrenia and other psychotic disorders	‡	‡	‡	‡	‡	5.3
Alcohol or substance-related disorders	‡	‡	‡	‡	‡	9.8
Other mental disorders ²	‡	‡	‡	‡	‡	8.5
Total	‡	‡	‡	‡	‡	100.0
Gender						
Male	‡	‡	‡	‡	‡	45.4
Female	‡	‡	‡	‡	‡	54.6
Total	‡	‡	‡	‡	‡	100.0
Age³						
0 to 11 years	‡	‡	‡	‡	‡	6.9
12 to 14 years	‡	‡	‡	‡	‡	4.6
15 to 17 years	‡	‡	‡	‡	‡	5.8
18 to 20 years	‡	‡	‡	‡	‡	5.4
21 to 24 years	‡	‡	‡	‡	‡	6.5
25 to 29 years	‡	‡	‡	‡	‡	10.2
30 to 34 years	‡	‡	‡	‡	‡	9.5
35 to 39 years	‡	‡	‡	‡	‡	9.7
40 to 44 years	‡	‡	‡	‡	‡	8.5
45 to 49 years	‡	‡	‡	‡	‡	8.0
50 to 54 years	‡	‡	‡	‡	‡	7.5
55 to 59 years	‡	‡	‡	‡	‡	7.0
60 to 64 years	‡	‡	‡	‡	‡	5.0
65 years and older	‡	‡	‡	‡	‡	5.5
Total	‡	‡	‡	‡	‡	100.0

Continued. See notes at end of table.

**Table 9.47b. West Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018 (continued)**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	‡	‡	‡	‡	‡	†
Asian ⁴	‡	‡	‡	‡	‡	†
Black or African American	‡	‡	‡	‡	‡	3.9
Native Hawaiian or Other Pacific Islander	‡	‡	‡	‡	‡	†
White	‡	‡	‡	‡	‡	85.3
Other	‡	‡	‡	‡	‡	10.5
Total	‡	‡	‡	‡	‡	100.0
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	‡	‡	‡
Hispanic or Latino origin	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	2.7
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	97.3
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	100.0

‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.48a. Wisconsin individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	97,812	100,684	72,929	77,345	65,079	71,193
Mental health diagnosis¹						
Trauma- and stressor-related disorders	11,342	11,593	8,208	8,149	6,531	6,785
Anxiety disorders	7,139	7,841	5,760	7,506	9,359	10,509
Attention deficit/hyperactivity disorder	3,514	3,645	2,603	2,717	2,513	2,811
Mood disorders	39,519	39,469	25,078	23,535	16,770	16,940
Bipolar disorders	15,380	15,246	10,433	9,419	5,785	5,714
Depressive disorders	24,139	24,223	14,645	14,116	10,985	11,226
Personality disorders	971	981	388	2,022	3,198	6,150
Schizophrenia and other psychotic disorders	15,106	14,996	10,271	10,049	8,561	9,134
Alcohol or substance-related disorders	1,714	1,396	828	628	241	127
Other mental disorders ²	9,731	11,739	12,392	14,789	10,177	11,125
Total	89,036	91,660	65,528	69,395	57,350	63,581
Gender						
Male	48,806	49,728	36,195	38,405	32,093	35,462
Female	49,006	50,951	36,727	38,927	32,974	35,723
Total	97,812	100,679	72,922	77,332	65,067	71,185
Age³						
0 to 11 years	3,264	3,302	3,088	3,425	3,075	3,432
12 to 14 years	3,171	3,418	3,402	3,474	3,156	3,635
15 to 17 years	4,645	4,851	4,766	5,261	4,806	5,405
18 to 20 years	4,566	4,513	3,494	3,865	3,421	3,966
21 to 24 years	6,890	6,790	5,239	5,477	4,567	4,910
25 to 29 years	9,501	9,519	7,033	7,419	6,116	6,745
30 to 34 years	9,320	9,710	7,028	7,309	6,051	6,559
35 to 39 years	8,083	8,261	6,036	6,518	5,348	6,001
40 to 44 years	8,795	8,882	5,893	6,021	4,730	4,961
45 to 49 years	9,780	9,503	6,289	6,362	5,104	5,387
50 to 54 years	10,101	10,539	7,146	7,198	5,661	5,671
55 to 59 years	7,778	8,275	5,549	6,007	5,105	5,546
60 to 64 years	4,890	5,410	3,635	3,987	3,450	3,907
65 years and older	7,028	7,711	4,331	5,022	4,488	5,067
Total	97,812	100,684	72,929	77,345	65,078	71,192

Continued. See notes at end of table.

Table 9.48a. Wisconsin individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	1,549	1,724	1,525	1,525	1,292	1,383
Asian ⁴	1,275	1,278	909	947	786	929
Black or African American	11,849	10,974	6,983	7,527	5,982	7,154
Native Hawaiian or Other Pacific Islander	59	66	85	65	65	84
White	82,245	84,336	60,582	63,259	51,983	55,001
Other	502	887	1,036	1,382	1,497	2,394
Total	97,479	99,265	71,120	74,705	61,605	66,945
Ethnicity						
Not of Hispanic or Latino origin	92,595	91,345	63,803	65,824	53,268	59,458
Hispanic or Latino origin	3,991	3,981	2,704	3,048	2,499	3,061
Total	96,586	95,326	66,507	68,872	55,767	62,519
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.48b. Wisconsin individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	12.7	12.6	12.5	11.7	11.4	10.7
Anxiety disorders	8.0	8.6	8.8	10.8	16.3	16.5
Attention deficit/hyperactivity disorder	3.9	4.0	4.0	3.9	4.4	4.4
Mood disorders	44.4	43.1	38.3	33.9	29.2	26.6
Bipolar disorders	17.3	16.6	15.9	13.6	10.1	9.0
Depressive disorders	27.1	26.4	22.3	20.3	19.2	17.7
Personality disorders	1.1	1.1	†	2.9	5.6	9.7
Schizophrenia and other psychotic disorders	17.0	16.4	15.7	14.5	14.9	14.4
Alcohol or substance-related disorders	1.9	1.5	1.3	†	†	†
Other mental disorders ²	10.9	12.8	18.9	21.3	17.7	17.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	49.9	49.4	49.6	49.7	49.3	49.8
Female	50.1	50.6	50.4	50.3	50.7	50.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	3.3	3.3	4.2	4.4	4.7	4.8
12 to 14 years	3.2	3.4	4.7	4.5	4.8	5.1
15 to 17 years	4.7	4.8	6.5	6.8	7.4	7.6
18 to 20 years	4.7	4.5	4.8	5.0	5.3	5.6
21 to 24 years	7.0	6.7	7.2	7.1	7.0	6.9
25 to 29 years	9.7	9.5	9.6	9.6	9.4	9.5
30 to 34 years	9.5	9.6	9.6	9.4	9.3	9.2
35 to 39 years	8.3	8.2	8.3	8.4	8.2	8.4
40 to 44 years	9.0	8.8	8.1	7.8	7.3	7.0
45 to 49 years	10.0	9.4	8.6	8.2	7.8	7.6
50 to 54 years	10.3	10.5	9.8	9.3	8.7	8.0
55 to 59 years	8.0	8.2	7.6	7.8	7.8	7.8
60 to 64 years	5.0	5.4	5.0	5.2	5.3	5.5
65 years and older	7.2	7.7	5.9	6.5	6.9	7.1
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.48b. Wisconsin individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	1.6	1.7	2.1	2.0	2.1	2.1
Asian ⁴	1.3	1.3	1.3	1.3	1.3	1.4
Black or African American	12.2	11.1	9.8	10.1	9.7	10.7
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	84.4	85.0	85.2	84.7	84.4	82.2
Other	†	†	1.5	1.8	2.4	3.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	95.9	95.8	95.9	95.6	95.5	95.1
Hispanic or Latino origin	4.1	4.2	4.1	4.4	4.5	4.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	‡

‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.49a. Wyoming individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	16,582	17,142	17,798	17,921	17,365	16,256
Mental health diagnosis¹						
Trauma- and stressor-related disorders	3,117	3,160	3,361	3,527	3,299	2,971
Anxiety disorders	1,693	1,889	2,116	2,277	2,462	2,298
Attention deficit/hyperactivity disorder	986	1,080	1,084	992	956	862
Mood disorders	6,888	7,142	7,276	7,032	6,300	6,167
Bipolar disorders	2,104	2,163	2,177	1,909	1,544	1,487
Depressive disorders	4,784	4,979	5,099	5,123	4,756	4,680
Personality disorders	137	203	249	255	308	316
Schizophrenia and other psychotic disorders	796	941	961	963	937	939
Alcohol or substance-related disorders	--	--	--	--	--	--
Other mental disorders ²	2,597	2,289	2,233	2,191	2,394	2,209
Total	16,214	16,704	17,280	17,237	16,656	15,762
Gender						
Male	7,424	7,695	7,938	7,940	7,738	7,316
Female	9,158	9,447	9,860	9,981	9,627	8,940
Total	16,582	17,142	17,798	17,921	17,365	16,256
Age³						
0 to 11 years	2,304	2,274	2,365	2,333	2,143	1,913
12 to 14 years	1,145	1,174	1,218	1,200	1,227	1,214
15 to 17 years	1,366	1,347	1,486	1,394	1,431	1,392
18 to 20 years	746	739	747	827	852	832
21 to 24 years	1,180	1,155	1,240	1,245	1,164	1,029
25 to 29 years	1,600	1,663	1,672	1,794	1,735	1,535
30 to 34 years	1,617	1,644	1,635	1,746	1,610	1,443
35 to 39 years	1,308	1,347	1,480	1,516	1,495	1,421
40 to 44 years	1,180	1,247	1,277	1,235	1,159	1,085
45 to 49 years	1,073	1,091	1,098	1,089	1,067	1,041
50 to 54 years	1,090	1,199	1,172	1,113	1,012	873
55 to 59 years	842	945	972	967	927	880
60 to 64 years	534	601	689	668	663	675
65 years and older	597	715	747	794	880	922
Total	16,582	17,141	17,798	17,921	17,365	16,255

Continued. See notes at end of table.

Table 9.49a. Wyoming individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2013–2018 (continued)

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	371	354	383	372	344	325
Asian ⁴	81	67	75	72	77	70
Black or African American	210	256	258	276	274	251
Native Hawaiian or Other Pacific Islander	36	26	33	41	41	28
White	14,470	15,025	15,541	15,563	15,040	14,104
Other	544	535	564	652	662	619
Total	15,712	16,263	16,854	16,976	16,438	15,397
Ethnicity						
Not of Hispanic or Latino origin	14,297	14,883	15,543	15,557	14,965	14,069
Hispanic or Latino origin	1,471	1,576	1,712	1,833	1,805	1,600
Total	15,768	16,459	17,255	17,390	16,770	15,669
Specific Hispanic or Latino origin						
Mexican	788	830	924	952	928	781
Puerto Rican	46	50	55	55	77	64
Cuban	13	17	24	24	25	11
Other specific	592	659	689	767	739	710
Total	1,439	1,556	1,692	1,798	1,769	1,566
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	249	218
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	4,299	3,814
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	4,548	4,032

-- Quantity is zero; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 9.49b. Wyoming individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	19.2	18.9	19.5	20.5	19.8	18.8
Anxiety disorders	10.4	11.3	12.2	13.2	14.8	14.6
Attention deficit/hyperactivity disorder	6.1	6.5	6.3	5.8	5.7	5.5
Mood disorders	42.5	42.8	42.1	40.8	37.8	39.1
Bipolar disorders	13.0	12.9	12.6	11.1	9.3	9.4
Depressive disorders	29.5	29.8	29.5	29.7	28.6	29.7
Personality disorders	†	1.2	1.4	1.5	1.8	2.0
Schizophrenia and other psychotic disorders	4.9	5.6	5.6	5.6	5.6	6.0
Alcohol or substance-related disorders	--	--	--	--	--	--
Other mental disorders ²	16.0	13.7	12.9	12.7	14.4	14.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	44.8	44.9	44.6	44.3	44.6	45.0
Female	55.2	55.1	55.4	55.7	55.4	55.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	13.9	13.3	13.3	13.0	12.3	11.8
12 to 14 years	6.9	6.8	6.8	6.7	7.1	7.5
15 to 17 years	8.2	7.9	8.3	7.8	8.2	8.6
18 to 20 years	4.5	4.3	4.2	4.6	4.9	5.1
21 to 24 years	7.1	6.7	7.0	6.9	6.7	6.3
25 to 29 years	9.6	9.7	9.4	10.0	10.0	9.4
30 to 34 years	9.8	9.6	9.2	9.7	9.3	8.9
35 to 39 years	7.9	7.9	8.3	8.5	8.6	8.7
40 to 44 years	7.1	7.3	7.2	6.9	6.7	6.7
45 to 49 years	6.5	6.4	6.2	6.1	6.1	6.4
50 to 54 years	6.6	7.0	6.6	6.2	5.8	5.4
55 to 59 years	5.1	5.5	5.5	5.4	5.3	5.4
60 to 64 years	3.2	3.5	3.9	3.7	3.8	4.2
65 years and older	3.6	4.2	4.2	4.4	5.1	5.7
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 9.49b. Wyoming individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2013–2018 (continued)

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	2.4	2.2	2.3	2.2	2.1	2.1
Asian ⁴	†	†	†	†	†	†
Black or African American	1.3	1.6	1.5	1.6	1.7	1.6
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†
White	92.1	92.4	92.2	91.7	91.5	91.6
Other	3.5	3.3	3.3	3.8	4.0	4.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	90.7	90.4	90.1	89.5	89.2	89.8
Hispanic or Latino origin	9.3	9.6	9.9	10.5	10.8	10.2
Total	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	54.8	53.3	54.6	52.9	52.5	49.9
Puerto Rican	3.2	3.2	3.3	3.1	4.4	4.1
Cuban	†	1.1	1.4	1.3	1.4	†
Other specific	41.1	42.4	40.7	42.7	41.8	45.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	5.5	5.4
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	94.5	94.6
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.50a. Other jurisdictions individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	206	182	175	376	389	392
Mental health diagnosis¹						
Trauma- and stressor-related disorders	7	5	5	10	21	8
Anxiety disorders	5	1	2	12	9	9
Attention deficit/hyperactivity disorder	9	4	2	14	7	10
Mood disorders	30	21	16	39	35	20
Bipolar disorders	8	6	6	18	15	4
Depressive disorders	22	15	10	21	20	16
Personality disorders	2	--	1	--	--	--
Schizophrenia and other psychotic disorders	24	24	14	99	87	112
Alcohol or substance-related disorders	15	14	22	46	54	95
Other mental disorders ²	9	6	5	38	21	24
Total	101	75	67	258	234	278
Gender						
Male	136	123	118	243	247	265
Female	70	59	57	133	142	127
Total	206	182	175	376	389	392
Age³						
0 to 11 years	10	--	--	4	13	--
12 to 14 years	5	5	3	8	9	3
15 to 17 years	11	8	5	15	5	7
18 to 20 years	6	4	12	15	18	14
21 to 24 years	18	19	18	30	33	31
25 to 29 years	23	26	26	45	56	53
30 to 34 years	26	27	27	40	48	56
35 to 39 years	23	20	21	30	46	58
40 to 44 years	34	21	13	42	35	38
45 to 49 years	26	15	16	36	23	39
50 to 54 years	11	16	11	43	39	40
55 to 59 years	9	11	13	35	27	22
60 to 64 years	3	6	4	19	22	19
65 years and older	--	--	--	10	13	6
Total	205	178	169	372	387	386

Continued. See notes at end of table.

**Table 9.50a. Other jurisdictions individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Number, 2013–2018 (continued)**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Race						
American Indian or Alaska Native	--	--	--	--	--	--
Asian ⁴	30	15	16	241	209	200
Black or African American	--	--	--	--	--	1
Native Hawaiian or Other Pacific Islander	--	--	--	1	--	--
White	3	2	8	13	9	6
Other	46	37	46	38	42	51
Total	79	54	70	293	260	258
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	234	178	182
Hispanic or Latino origin	‡	‡	‡	--	--	--
Total	‡	‡	‡	234	178	182
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	4
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	46
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	50

-- Quantity is zero; ‡ No data submitted; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

**Table 9.50b. Other jurisdictions individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018**

<i>Selected characteristics</i>	2013	2014	2015	2016	2017	2018
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0
Mental health diagnosis¹						
Trauma- and stressor-related disorders	6.9	6.7	7.5	3.9	9.0	2.9
Anxiety disorders	5.0	1.3	3.0	4.7	3.8	3.2
Attention deficit/hyperactivity disorder	8.9	5.3	3.0	5.4	3.0	3.6
Mood disorders	29.7	28.0	23.9	15.1	15.0	7.2
Bipolar disorders	7.9	8.0	9.0	7.0	6.4	1.4
Depressive disorders	21.8	20.0	14.9	8.1	8.5	5.8
Personality disorders	2.0	--	1.5	--	--	--
Schizophrenia and other psychotic disorders	23.8	32.0	20.9	38.4	37.2	40.3
Alcohol or substance-related disorders	14.9	18.7	32.8	17.8	23.1	34.2
Other mental disorders ²	8.9	8.0	7.5	14.7	9.0	8.6
Total	100.0	100.0	100.0	100.0	100.0	100.0
Gender						
Male	66.0	67.6	67.4	64.6	63.5	67.6
Female	34.0	32.4	32.6	35.4	36.5	32.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Age³						
0 to 11 years	4.9	--	--	1.1	3.4	--
12 to 14 years	2.4	2.8	1.8	2.2	2.3	†
15 to 17 years	5.4	4.5	3.0	4.0	1.3	1.8
18 to 20 years	2.9	2.2	7.1	4.0	4.7	3.6
21 to 24 years	8.8	10.7	10.7	8.1	8.5	8.0
25 to 29 years	11.2	14.6	15.4	12.1	14.5	13.7
30 to 34 years	12.7	15.2	16.0	10.8	12.4	14.5
35 to 39 years	11.2	11.2	12.4	8.1	11.9	15.0
40 to 44 years	16.6	11.8	7.7	11.3	9.0	9.8
45 to 49 years	12.7	8.4	9.5	9.7	5.9	10.1
50 to 54 years	5.4	9.0	6.5	11.6	10.1	10.4
55 to 59 years	4.4	6.2	7.7	9.4	7.0	5.7
60 to 64 years	1.5	3.4	2.4	5.1	5.7	4.9
65 years and older	--	--	--	2.7	3.4	1.6
Total	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

**Table 9.50b. Other jurisdictions individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status:
Column percent distribution, 2013–2018 (continued)**

<i>Selected characteristics</i>	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Race						
American Indian or Alaska Native	--	--	--	--	--	--
Asian ⁴	38.0	27.8	22.9	82.3	80.4	77.5
Black or African American	--	--	--	--	--	†
Native Hawaiian or Other Pacific Islander	--	--	--	†	--	--
White	3.8	3.7	11.4	4.4	3.5	2.3
Other	58.2	68.5	65.7	13.0	16.2	19.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity						
Not of Hispanic or Latino origin	‡	‡	‡	100.0	100.0	100.0
Hispanic or Latino origin	‡	‡	‡	--	--	--
Total	‡	‡	‡	100.0	100.0	100.0
Specific Hispanic or Latino origin						
Mexican	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡
Veteran status⁵						
Veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	8.0
Not a veteran	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	92.0
Total	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	‡	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals; *n/c* Not collected.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

² “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes alcohol- and substance-related diagnoses. See Appendix E.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTES: Percentages may not sum to 100 percent because of rounding. Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Table 10.1a. Alabama individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	101,599	8,158	5,900	13,669	11,831	27,591	437	17,752	16,261	
Gender										
Male	48,090	3,711	2,218	9,735	3,919	8,798	184	10,516	9,009	
Female	53,505	4,447	3,682	3,932	7,912	18,791	253	7,236	7,252	
Total	101,595	8,158	5,900	13,667	11,831	27,589	437	17,752	16,261	
Age²										
0 to 11 years	15,969	3,003	751	7,749	21	464	1	23	3,957	
12 to 14 years	8,388	1,342	575	2,921	72	1,252	3	42	2,181	
15 to 17 years	8,689	1,088	745	1,970	217	2,207	5	100	2,357	
18 to 20 years	4,420	333	391	578	409	1,456	16	302	935	
21 to 24 years	4,496	235	329	137	684	1,507	42	797	765	
25 to 29 years	7,061	319	495	108	1,221	2,269	67	1,554	1,028	
30 to 34 years	7,063	307	467	55	1,309	2,332	56	1,739	798	
35 to 39 years	7,442	337	400	44	1,430	2,344	59	2,024	804	
40 to 44 years	6,525	237	341	36	1,307	2,232	57	1,656	659	
45 to 49 years	6,814	240	314	25	1,317	2,474	32	1,770	642	
50 to 54 years	7,065	203	309	12	1,256	2,736	34	1,947	568	
55 to 59 years	6,914	167	282	15	1,118	2,601	29	2,154	548	
60 to 64 years	5,081	126	207	14	736	1,803	22	1,807	366	
65 years and older	5,666	220	293	4	733	1,913	14	1,836	653	
Total	101,593	8,157	5,899	13,668	11,830	27,590	437	17,751	16,261	
Race										
American Indian or Alaska Native	290	31	21	17	45	69	7	56	44	
Asian ³	209	24	13	18	10	50	--	55	39	
Black or African American	37,814	2,613	1,247	5,689	2,380	8,816	153	10,445	6,471	
Native Hawaiian or Other Pacific Islander	57	4	1	8	7	21	--	4	12	
White	55,715	4,608	4,069	6,605	8,769	16,949	258	6,391	8,066	
Other	6,232	741	440	1,130	460	1,313	18	577	1,553	
Total	100,317	8,021	5,791	13,467	11,671	27,218	436	17,528	16,185	

Continued. See notes at end of table.

Table 10.1a. Alabama individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	98,877	7,787	5,679	13,165	11,613	26,895	428	17,530	15,780
Hispanic or Latino origin	2,717	371	221	503	216	695	9	222	480
Total	101,594	8,158	5,900	13,668	11,829	27,590	437	17,752	16,260
Specific Hispanic or Latino origin									
Mexican	1,095	181	91	223	50	274	1	60	215
Puerto Rican	362	28	38	46	41	105	2	42	60
Cuban	45	1	1	4	9	14	--	11	5
Other specific	1,198	161	91	230	111	300	6	104	195
Total	2,700	371	221	503	211	693	9	217	475
Veteran status⁴									
Veteran	981	85	54	7	175	301	3	239	117
Not a veteran	53,626	2,440	3,386	823	9,247	19,931	362	10,729	6,708
Total	54,607	2,525	3,440	830	9,422	20,232	365	10,968	6,825

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.1b. Alabama individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	47.3	45.5	37.6	71.2	33.1	31.9	42.1	59.2	55.4	
Female	52.7	54.5	62.4	28.8	66.9	68.1	57.9	40.8	44.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	15.7	36.8	12.7	56.7	†	1.7	†	†	24.3	
12 to 14 years	8.3	16.5	9.7	21.4	†	4.5	†	†	13.4	
15 to 17 years	8.6	13.3	12.6	14.4	1.8	8.0	1.1	†	14.5	
18 to 20 years	4.4	4.1	6.6	4.2	3.5	5.3	3.7	1.7	5.7	
21 to 24 years	4.4	2.9	5.6	1.0	5.8	5.5	9.6	4.5	4.7	
25 to 29 years	7.0	3.9	8.4	†	10.3	8.2	15.3	8.8	6.3	
30 to 34 years	7.0	3.8	7.9	†	11.1	8.5	12.8	9.8	4.9	
35 to 39 years	7.3	4.1	6.8	†	12.1	8.5	13.5	11.4	4.9	
40 to 44 years	6.4	2.9	5.8	†	11.0	8.1	13.0	9.3	4.1	
45 to 49 years	6.7	2.9	5.3	†	11.1	9.0	7.3	10.0	3.9	
50 to 54 years	7.0	2.5	5.2	†	10.6	9.9	7.8	11.0	3.5	
55 to 59 years	6.8	2.0	4.8	†	9.5	9.4	6.6	12.1	3.4	
60 to 64 years	5.0	1.5	3.5	†	6.2	6.5	5.0	10.2	2.3	
65 years and older	5.6	2.7	5.0	†	6.2	6.9	3.2	10.3	4.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	1.6	†	†	
Asian ³	†	†	†	†	†	†	--	†	†	
Black or African American	37.7	32.6	21.5	42.2	20.4	32.4	35.1	59.6	40.0	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†	
White	55.5	57.4	70.3	49.0	75.1	62.3	59.2	36.5	49.8	
Other	6.2	9.2	7.6	8.4	3.9	4.8	4.1	3.3	9.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.1b. Alabama individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	97.3	95.5	96.3	96.3	98.2	97.5	97.9	98.7	97.0
Hispanic or Latino origin	2.7	4.5	3.7	3.7	1.8	2.5	2.1	1.3	3.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	40.6	48.8	41.2	44.3	23.7	39.5	11.1	27.6	45.3
Puerto Rican	13.4	7.5	17.2	9.1	19.4	15.2	22.2	19.4	12.6
Cuban	1.7	†	†	†	4.3	2.0	- -	5.1	1.1
Other specific	44.4	43.4	41.2	45.7	52.6	43.3	66.7	47.9	41.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	1.8	3.4	1.6	†	1.9	1.5	†	2.2	1.7
Not a veteran	98.2	96.6	98.4	99.2	98.1	98.5	99.2	97.8	98.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

- - Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.2a. Arizona individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	169,861	12,659	29,586	20,587	28,143	42,320	1,319	21,951	13,296
Gender									
Male	80,231	4,415	11,902	14,987	11,084	14,820	388	13,607	9,028
Female	89,630	8,244	17,684	5,600	17,059	27,500	931	8,344	4,268
Total	169,861	12,659	29,586	20,587	28,143	42,320	1,319	21,951	13,296
Age²									
0 to 11 years	21,317	1,636	2,584	10,771	483	741	12	82	5,008
12 to 14 years	12,182	887	1,947	4,346	709	2,084	15	113	2,081
15 to 17 years	12,910	985	2,408	2,645	1,298	3,593	37	212	1,732
18 to 20 years	7,882	638	1,639	692	1,264	2,381	90	520	658
21 to 24 years	10,047	791	2,262	363	1,761	2,799	134	1,285	652
25 to 29 years	15,620	1,196	3,684	572	2,879	4,207	184	2,157	741
30 to 34 years	14,985	1,168	3,378	411	2,934	3,929	186	2,421	558
35 to 39 years	13,859	1,203	2,749	264	2,982	3,578	163	2,452	468
40 to 44 years	11,722	977	2,047	179	2,658	3,098	123	2,340	300
45 to 49 years	12,107	932	1,946	140	2,773	3,467	109	2,450	290
50 to 54 years	12,385	879	1,782	105	2,794	3,870	102	2,562	291
55 to 59 years	11,889	769	1,626	66	2,643	3,978	93	2,469	245
60 to 64 years	8,254	443	1,140	32	1,716	3,062	48	1,615	198
65 years and older	4,701	155	394	1	1,249	1,533	23	1,272	74
Total	169,860	12,659	29,586	20,587	28,143	42,320	1,319	21,950	13,296
Race									
American Indian or Alaska Native	4,220	484	612	399	423	1,246	31	644	381
Asian ³	1,382	117	213	95	180	318	8	336	115
Black or African American	13,071	1,013	1,520	1,873	2,055	2,670	67	2,798	1,075
Native Hawaiian or Other Pacific Islander	451	35	78	47	59	113	3	75	41
White	147,209	10,687	26,595	17,579	24,959	37,115	1,190	17,722	11,362
Other	3,116	302	497	585	396	729	18	283	306
Total	169,449	12,638	29,515	20,578	28,072	42,191	1,317	21,858	13,280

Continued. See notes at end of table.

Table 10.2a. Arizona individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	122,163	9,461	20,584	13,064	22,745	30,013	1,108	16,707	8,481
Hispanic or Latino origin	47,210	3,176	8,931	7,513	5,319	12,176	208	5,090	4,797
Total	169,373	12,637	29,515	20,577	28,064	42,189	1,316	21,797	13,278
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	144	25	25	1	22	44	--	26	1
Not a veteran	6,898	598	1,491	179	1,189	2,239	74	843	285
Total	7,042	623	1,516	180	1,211	2,283	74	869	286

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.2b. Arizona individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	47.2	34.9	40.2	72.8	39.4	35.0	29.4	62.0	67.9	
Female	52.8	65.1	59.8	27.2	60.6	65.0	70.6	38.0	32.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	12.5	12.9	8.7	52.3	1.7	1.8	†	†	37.7	
12 to 14 years	7.2	7.0	6.6	21.1	2.5	4.9	1.1	†	15.7	
15 to 17 years	7.6	7.8	8.1	12.8	4.6	8.5	2.8	†	13.0	
18 to 20 years	4.6	5.0	5.5	3.4	4.5	5.6	6.8	2.4	4.9	
21 to 24 years	5.9	6.2	7.6	1.8	6.3	6.6	10.2	5.9	4.9	
25 to 29 years	9.2	9.4	12.5	2.8	10.2	9.9	13.9	9.8	5.6	
30 to 34 years	8.8	9.2	11.4	2.0	10.4	9.3	14.1	11.0	4.2	
35 to 39 years	8.2	9.5	9.3	1.3	10.6	8.5	12.4	11.2	3.5	
40 to 44 years	6.9	7.7	6.9	†	9.4	7.3	9.3	10.7	2.3	
45 to 49 years	7.1	7.4	6.6	†	9.9	8.2	8.3	11.2	2.2	
50 to 54 years	7.3	6.9	6.0	†	9.9	9.1	7.7	11.7	2.2	
55 to 59 years	7.0	6.1	5.5	†	9.4	9.4	7.1	11.2	1.8	
60 to 64 years	4.9	3.5	3.9	†	6.1	7.2	3.6	7.4	1.5	
65 years and older	2.8	1.2	1.3	†	4.4	3.6	1.7	5.8	†	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	2.5	3.8	2.1	1.9	1.5	3.0	2.4	2.9	2.9	
Asian ³	†	†	†	†	†	†	†	1.5	†	
Black or African American	7.7	8.0	5.1	9.1	7.3	6.3	5.1	12.8	8.1	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	86.9	84.6	90.1	85.4	88.9	88.0	90.4	81.1	85.6	
Other	1.8	2.4	1.7	2.8	1.4	1.7	1.4	1.3	2.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.2b. Arizona individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	72.1	74.9	69.7	63.5	81.0	71.1	84.2	76.6	63.9
Hispanic or Latino origin	27.9	25.1	30.3	36.5	19.0	28.9	15.8	23.4	36.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	2.0	4.0	1.6	†	1.8	1.9	--	3.0	†
Not a veteran	98.0	96.0	98.4	99.4	98.2	98.1	100.0	97.0	99.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.3a. Arkansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	59,565	10,359	7,699	1,075	6,044	18,416	418	6,758	8,796
Gender									
Male	27,107	4,623	3,288	741	2,073	6,559	153	4,151	5,519
Female	32,408	5,731	4,406	333	3,969	11,848	265	2,601	3,255
Total	59,515	10,354	7,694	1,074	6,042	18,407	418	6,752	8,774
Age²									
0 to 11 years	8,583	3,249	1,282	489	49	560	8	52	2,894
12 to 14 years	4,282	1,184	743	155	17	798	1	12	1,372
15 to 17 years	4,865	1,047	748	136	68	1,548	5	35	1,278
18 to 20 years	3,406	576	561	75	159	1,242	10	94	689
21 to 24 years	3,066	430	440	41	345	1,162	44	310	294
25 to 29 years	4,600	645	596	50	569	1,655	72	632	381
30 to 34 years	4,768	670	637	41	658	1,655	61	704	342
35 to 39 years	4,940	673	666	22	771	1,626	69	824	289
40 to 44 years	4,230	458	498	17	719	1,501	45	742	250
45 to 49 years	4,087	421	402	13	742	1,592	40	661	216
50 to 54 years	3,969	386	345	12	625	1,586	33	746	236
55 to 59 years	3,777	307	353	11	583	1,502	15	799	207
60 to 64 years	2,520	147	202	6	418	998	7	598	144
65 years and older	2,472	166	226	7	321	991	8	549	204
Total	59,565	10,359	7,699	1,075	6,044	18,416	418	6,758	8,796
Race									
American Indian or Alaska Native	245	46	38	3	28	80	1	20	29
Asian ³	205	37	25	1	14	73	--	38	17
Black or African American	10,756	1,308	815	138	757	3,085	75	2,609	1,969
Native Hawaiian or Other Pacific Islander	52	7	7	2	6	14	--	6	10
White	37,707	6,817	5,326	688	4,643	11,845	280	3,394	4,714
Other	828	190	124	37	51	202	8	39	177
Total	49,793	8,405	6,335	869	5,499	15,299	364	6,106	6,916

Continued. See notes at end of table.

Table 10.3a. Arkansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	50,887	8,552	6,466	856	5,490	15,707	381	6,152	7,283
Hispanic or Latino origin	2,630	621	392	86	113	725	7	135	551
Total	53,517	9,173	6,858	942	5,603	16,432	388	6,287	7,834
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.3b. Arkansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	45.5	44.6	42.7	69.0	34.3	35.6	36.6	61.5	62.9	
Female	54.5	55.4	57.3	31.0	65.7	64.4	63.4	38.5	37.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Age²										
0 to 11 years	14.4	31.4	16.7	45.5	†	3.0	1.9	†	32.9	
12 to 14 years	7.2	11.4	9.7	14.4	†	4.3	†	†	15.6	
15 to 17 years	8.2	10.1	9.7	12.7	1.1	8.4	1.2	†	14.5	
18 to 20 years	5.7	5.6	7.3	7.0	2.6	6.7	2.4	1.4	7.8	
21 to 24 years	5.1	4.2	5.7	3.8	5.7	6.3	10.5	4.6	3.3	
25 to 29 years	7.7	6.2	7.7	4.7	9.4	9.0	17.2	9.4	4.3	
30 to 34 years	8.0	6.5	8.3	3.8	10.9	9.0	14.6	10.4	3.9	
35 to 39 years	8.3	6.5	8.7	2.0	12.8	8.8	16.5	12.2	3.3	
40 to 44 years	7.1	4.4	6.5	1.6	11.9	8.2	10.8	11.0	2.8	
45 to 49 years	6.9	4.1	5.2	1.2	12.3	8.6	9.6	9.8	2.5	
50 to 54 years	6.7	3.7	4.5	1.1	10.3	8.6	7.9	11.0	2.7	
55 to 59 years	6.3	3.0	4.6	1.0	9.6	8.2	3.6	11.8	2.4	
60 to 64 years	4.2	1.4	2.6	†	6.9	5.4	1.7	8.8	1.6	
65 years and older	4.2	1.6	2.9	†	5.3	5.4	1.9	8.1	2.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†	
Asian ³	†	†	†	†	†	†	--	†	†	
Black or African American	21.6	15.6	12.9	15.9	13.8	20.2	20.6	42.7	28.5	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†	
White	75.7	81.1	84.1	79.2	84.4	77.4	76.9	55.6	68.2	
Other	1.7	2.3	2.0	4.3	†	1.3	2.2	†	2.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	

Continued. See notes at end of table.

Table 10.3b. Arkansas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	95.1	93.2	94.3	90.9	98.0	95.6	98.2	97.9	93.0
Hispanic or Latino origin	4.9	6.8	5.7	9.1	2.0	4.4	1.8	2.1	7.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.4a. California individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	660,567	75,912	54,673	28,971	49,619	160,164	1,057	121,780	168,391
Gender									
Male	345,388	37,009	25,313	23,028	20,041	59,479	605	78,775	101,138
Female	314,414	38,841	29,309	5,934	29,534	100,493	451	42,920	66,932
Total	659,802	75,850	54,622	28,962	49,575	159,972	1,056	121,695	168,070
Age²									
0 to 11 years	110,640	31,976	17,201	18,408	567	9,034	16	424	33,014
12 to 14 years	53,814	9,305	6,630	5,648	906	15,032	7	619	15,667
15 to 17 years	63,045	8,284	6,566	3,031	1,859	21,127	22	1,335	20,821
18 to 20 years	30,424	3,082	2,914	758	2,070	9,003	61	3,337	9,199
21 to 24 years	35,138	2,758	2,446	266	3,211	8,360	99	8,277	9,721
25 to 29 years	51,479	3,640	3,440	295	5,184	11,257	162	13,409	14,092
30 to 34 years	48,191	3,361	3,121	158	5,142	10,191	162	13,692	12,364
35 to 39 years	44,567	2,891	2,533	116	5,008	9,858	103	13,067	10,991
40 to 44 years	37,176	2,258	2,034	84	4,341	9,052	95	10,931	8,381
45 to 49 years	41,379	2,120	1,962	66	5,068	10,849	79	12,181	9,054
50 to 54 years	45,409	2,022	1,810	47	5,353	13,263	79	13,674	9,161
55 to 59 years	42,998	1,726	1,608	39	4,891	13,783	65	13,159	7,727
60 to 64 years	29,515	1,117	1,172	29	3,317	10,180	55	9,217	4,428
65 years and older	24,759	1,051	1,079	11	2,607	8,635	42	7,940	3,394
Total	658,534	75,591	54,516	28,956	49,524	159,624	1,047	121,262	168,014
Race									
American Indian or Alaska Native	4,936	592	394	131	456	1,234	10	848	1,271
Asian ³	24,907	2,066	1,436	517	1,561	7,587	22	7,741	3,977
Black or African American	90,793	9,946	4,415	4,133	6,292	18,896	184	24,614	22,313
Native Hawaiian or Other Pacific Islander	1,979	239	147	66	137	449	5	431	505
White	200,629	17,820	14,550	6,219	23,497	46,917	426	38,856	52,344
Other	229,806	31,541	23,579	13,543	11,596	59,104	197	29,752	60,494
Total	553,050	62,204	44,521	24,609	43,539	134,187	844	102,242	140,904

Continued. See notes at end of table.

Table 10.4a. California individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	298,616	28,496	19,940	10,521	29,870	71,469	472	63,379	74,469
Hispanic or Latino origin	234,967	33,176	24,315	14,376	11,241	60,157	194	29,674	61,834
Total	533,583	61,672	44,255	24,897	41,111	131,626	666	93,053	136,303
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	423	5	--	--	36	15	6	192	169
Not a veteran	6,183	181	28	4	564	551	137	3,696	1,022
Total	6,606	186	28	4	600	566	143	3,888	1,191

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.4b. California individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	52.3	48.8	46.3	79.5	40.4	37.2	57.3	64.7	60.2	
Female	47.7	51.2	53.7	20.5	59.6	62.8	42.7	35.3	39.8	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	16.8	42.3	31.6	63.6	1.1	5.7	1.5	†	19.6	
12 to 14 years	8.2	12.3	12.2	19.5	1.8	9.4	†	†	9.3	
15 to 17 years	9.6	11.0	12.0	10.5	3.8	13.2	2.1	1.1	12.4	
18 to 20 years	4.6	4.1	5.3	2.6	4.2	5.6	5.8	2.8	5.5	
21 to 24 years	5.3	3.6	4.5	†	6.5	5.2	9.5	6.8	5.8	
25 to 29 years	7.8	4.8	6.3	1.0	10.5	7.1	15.5	11.1	8.4	
30 to 34 years	7.3	4.4	5.7	†	10.4	6.4	15.5	11.3	7.4	
35 to 39 years	6.8	3.8	4.6	†	10.1	6.2	9.8	10.8	6.5	
40 to 44 years	5.6	3.0	3.7	†	8.8	5.7	9.1	9.0	5.0	
45 to 49 years	6.3	2.8	3.6	†	10.2	6.8	7.5	10.0	5.4	
50 to 54 years	6.9	2.7	3.3	†	10.8	8.3	7.5	11.3	5.5	
55 to 59 years	6.5	2.3	2.9	†	9.9	8.6	6.2	10.9	4.6	
60 to 64 years	4.5	1.5	2.1	†	6.7	6.4	5.3	7.6	2.6	
65 years and older	3.8	1.4	2.0	†	5.3	5.4	4.0	6.5	2.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	1.0	†	1.2	†	†	
Asian ³	4.5	3.3	3.2	2.1	3.6	5.7	2.6	7.6	2.8	
Black or African American	16.4	16.0	9.9	16.8	14.5	14.1	21.8	24.1	15.8	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	36.3	28.6	32.7	25.3	54.0	35.0	50.5	38.0	37.1	
Other	41.6	50.7	53.0	55.0	26.6	44.0	23.3	29.1	42.9	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.4b. California individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	56.0	46.2	45.1	42.3	72.7	54.3	70.9	68.1	54.6
Hispanic or Latino origin	44.0	53.8	54.9	57.7	27.3	45.7	29.1	31.9	45.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	6.4	2.7	--	--	6.0	2.7	4.2	4.9	14.2
Not a veteran	93.6	97.3	100.0	100.0	94.0	97.3	95.8	95.1	85.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.5a. Colorado individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	148,846	36,755	18,175	7,276	15,560	39,107	974	12,703	18,296
Gender									
Male	68,163	14,967	7,192	5,298	6,149	14,405	355	8,405	11,392
Female	80,682	21,787	10,983	1,978	9,411	24,702	619	4,298	6,904
Total	148,845	36,754	18,175	7,276	15,560	39,107	974	12,703	18,296
Age²									
0 to 11 years	18,766	9,390	2,142	3,157	38	968	1	29	3,041
12 to 14 years	12,043	4,082	1,476	1,466	123	3,019	8	64	1,805
15 to 17 years	13,572	3,486	1,836	889	412	4,641	8	188	2,112
18 to 20 years	7,140	1,479	1,049	277	591	2,490	42	374	838
21 to 24 years	8,543	1,573	1,188	217	1,019	2,561	74	861	1,050
25 to 29 years	14,356	2,879	2,023	310	1,865	3,792	142	1,504	1,841
30 to 34 years	13,784	2,881	1,849	289	2,002	3,445	159	1,390	1,769
35 to 39 years	12,869	2,739	1,596	210	1,890	3,298	120	1,509	1,507
40 to 44 years	9,866	1,993	1,144	126	1,577	2,596	95	1,246	1,089
45 to 49 years	9,272	1,785	1,041	112	1,497	2,669	87	1,193	888
50 to 54 years	8,865	1,605	868	83	1,486	2,687	97	1,205	834
55 to 59 years	8,345	1,316	795	67	1,345	2,792	71	1,248	711
60 to 64 years	5,784	806	548	41	878	2,056	38	976	441
65 years and older	5,640	740	620	32	837	2,093	32	916	370
Total	148,845	36,754	18,175	7,276	15,560	39,107	974	12,703	18,296
Race									
American Indian or Alaska Native	3,353	930	394	132	293	876	23	296	409
Asian ³	1,519	352	188	40	109	415	5	268	142
Black or African American	8,695	2,075	627	407	859	1,961	78	1,585	1,103
Native Hawaiian or Other Pacific Islander	434	124	51	16	37	101	2	41	62
White	102,254	24,111	13,109	5,014	11,476	26,752	726	8,248	12,818
Other	15,314	4,008	1,652	949	1,595	4,246	75	1,126	1,663
Total	131,569	31,600	16,021	6,558	14,369	34,351	909	11,564	16,197

Continued. See notes at end of table.

Table 10.5a. Colorado individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	100,970	22,875	12,182	4,636	12,288	26,615	766	9,536	12,072
Hispanic or Latino origin	37,295	10,944	4,706	2,072	2,328	9,779	156	2,435	4,875
Total	138,265	33,819	16,888	6,708	14,616	36,394	922	11,971	16,947
Specific Hispanic or Latino origin									
Mexican	23,554	7,001	2,972	1,406	1,322	6,227	92	1,412	3,122
Puerto Rican	879	267	110	53	101	197	6	47	98
Cuban	399	95	53	16	34	113	3	32	53
Other specific	12,341	3,575	1,568	594	864	3,230	53	875	1,582
Total	37,173	10,938	4,703	2,069	2,321	9,767	154	2,366	4,855
Veteran status⁴									
Veteran	3,139	797	303	54	376	862	29	344	374
Not a veteran	100,914	18,978	12,405	1,709	14,566	29,581	907	11,846	10,922
Total	104,053	19,775	12,708	1,763	14,942	30,443	936	12,190	11,296

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting periods. Data as of 07.08.2019.

Table 10.5b. Colorado individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	45.8	40.7	39.6	72.8	39.5	36.8	36.4	66.2	62.3	
Female	54.2	59.3	60.4	27.2	60.5	63.2	63.6	33.8	37.7	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	12.6	25.5	11.8	43.4	†	2.5	†	†	16.6	
12 to 14 years	8.1	11.1	8.1	20.1	†	7.7	†	†	9.9	
15 to 17 years	9.1	9.5	10.1	12.2	2.6	11.9	†	1.5	11.5	
18 to 20 years	4.8	4.0	5.8	3.8	3.8	6.4	4.3	2.9	4.6	
21 to 24 years	5.7	4.3	6.5	3.0	6.5	6.5	7.6	6.8	5.7	
25 to 29 years	9.6	7.8	11.1	4.3	12.0	9.7	14.6	11.8	10.1	
30 to 34 years	9.3	7.8	10.2	4.0	12.9	8.8	16.3	10.9	9.7	
35 to 39 years	8.6	7.5	8.8	2.9	12.1	8.4	12.3	11.9	8.2	
40 to 44 years	6.6	5.4	6.3	1.7	10.1	6.6	9.8	9.8	6.0	
45 to 49 years	6.2	4.9	5.7	1.5	9.6	6.8	8.9	9.4	4.9	
50 to 54 years	6.0	4.4	4.8	1.1	9.6	6.9	10.0	9.5	4.6	
55 to 59 years	5.6	3.6	4.4	†	8.6	7.1	7.3	9.8	3.9	
60 to 64 years	3.9	2.2	3.0	†	5.6	5.3	3.9	7.7	2.4	
65 years and older	3.8	2.0	3.4	†	5.4	5.4	3.3	7.2	2.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	2.5	2.9	2.5	2.0	2.0	2.6	2.5	2.6	2.5	
Asian ³	1.2	1.1	1.2	†	†	1.2	†	2.3	†	
Black or African American	6.6	6.6	3.9	6.2	6.0	5.7	8.6	13.7	6.8	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	77.7	76.3	81.8	76.5	79.9	77.9	79.9	71.3	79.1	
Other	11.6	12.7	10.3	14.5	11.1	12.4	8.3	9.7	10.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.5b. Colorado individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

<i>Selected characteristics</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Ethnicity									
Not of Hispanic or Latino origin	73.0	67.6	72.1	69.1	84.1	73.1	83.1	79.7	71.2
Hispanic or Latino origin	27.0	32.4	27.9	30.9	15.9	26.9	16.9	20.3	28.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	63.4	64.0	63.2	68.0	57.0	63.8	59.7	59.7	64.3
Puerto Rican	2.4	2.4	2.3	2.6	4.4	2.0	3.9	2.0	2.0
Cuban	1.1	†	1.1	†	1.5	1.2	1.9	1.4	1.1
Other specific	33.2	32.7	33.3	28.7	37.2	33.1	34.4	37.0	32.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	3.0	4.0	2.4	3.1	2.5	2.8	3.1	2.8	3.3
Not a veteran	97.0	96.0	97.6	96.9	97.5	97.2	96.9	97.2	96.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.6a. Connecticut individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	77,860	13,690	8,023	6,462	7,914	18,344	506	10,517	12,404
Gender									
Male	39,447	6,316	3,468	4,826	3,228	7,193	221	6,746	7,449
Female	38,369	7,369	4,549	1,636	4,683	11,139	283	3,762	4,948
Total	77,816	13,685	8,017	6,462	7,911	18,332	504	10,508	12,397
Age²									
0 to 11 years	14,345	5,128	1,287	3,671	25	713	1	55	3,465
12 to 14 years	8,079	1,962	1,013	1,278	59	1,677	2	33	2,055
15 to 17 years	8,479	1,694	1,142	757	130	2,225	14	93	2,424
18 to 20 years	3,295	514	392	223	254	838	32	234	808
21 to 24 years	3,112	398	372	117	414	762	47	486	516
25 to 29 years	4,352	586	500	121	676	1,048	55	823	543
30 to 34 years	4,325	578	494	92	708	1,022	66	943	422
35 to 39 years	4,080	554	444	67	746	1,032	39	882	316
40 to 44 years	3,599	427	362	32	689	1,001	42	773	273
45 to 49 years	4,536	489	411	39	904	1,343	47	993	310
50 to 54 years	5,649	487	482	20	1,048	1,800	55	1,371	386
55 to 59 years	5,649	424	446	24	979	1,901	49	1,478	348
60 to 64 years	4,268	240	326	14	702	1,541	28	1,180	237
65 years and older	4,091	209	352	7	580	1,441	29	1,173	300
Total	77,859	13,690	8,023	6,462	7,914	18,344	506	10,517	12,403
Race									
American Indian or Alaska Native	377	67	31	24	51	102	4	48	50
Asian ³	783	106	79	32	57	217	3	180	109
Black or African American	13,233	2,464	737	1,156	1,059	2,429	94	2,902	2,392
Native Hawaiian or Other Pacific Islander	206	34	12	10	24	63	1	36	26
White	46,935	7,558	5,613	3,438	5,561	11,699	333	5,783	6,950
Other	12,186	2,325	1,064	1,238	1,013	3,035	59	1,396	2,056
Total	73,720	12,554	7,536	5,898	7,765	17,545	494	10,345	11,583

Continued. See notes at end of table.

Table 10.6a. Connecticut individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	53,402	8,262	5,721	3,539	6,199	12,696	386	8,450	8,149
Hispanic or Latino origin	21,078	4,575	1,946	2,595	1,494	4,879	90	1,812	3,687
Total	74,480	12,837	7,667	6,134	7,693	17,575	476	10,262	11,836
Specific Hispanic or Latino origin									
Mexican	309	59	38	25	11	92	4	28	52
Puerto Rican	6,929	887	604	502	888	2,169	54	1,051	774
Cuban	91	10	10	4	17	22	--	21	7
Other specific	3,834	529	404	115	516	1,194	23	651	402
Total	11,163	1,485	1,056	646	1,432	3,477	81	1,751	1,235
Veteran status⁴									
Veteran	1,336	174	82	6	186	321	23	451	93
Not a veteran	38,575	3,895	3,914	533	6,690	11,270	345	8,902	3,026
Total	39,911	4,069	3,996	539	6,876	11,591	368	9,353	3,119

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.6b. Connecticut individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	50.7	46.2	43.3	74.7	40.8	39.2	43.8	64.2	60.1	
Female	49.3	53.8	56.7	25.3	59.2	60.8	56.2	35.8	39.9	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	18.4	37.5	16.0	56.8	†	3.9	†	†	27.9	
12 to 14 years	10.4	14.3	12.6	19.8	†	9.1	†	†	16.6	
15 to 17 years	10.9	12.4	14.2	11.7	1.6	12.1	2.8	†	19.5	
18 to 20 years	4.2	3.8	4.9	3.5	3.2	4.6	6.3	2.2	6.5	
21 to 24 years	4.0	2.9	4.6	1.8	5.2	4.2	9.3	4.6	4.2	
25 to 29 years	5.6	4.3	6.2	1.9	8.5	5.7	10.9	7.8	4.4	
30 to 34 years	5.6	4.2	6.2	1.4	8.9	5.6	13.0	9.0	3.4	
35 to 39 years	5.2	4.0	5.5	1.0	9.4	5.6	7.7	8.4	2.5	
40 to 44 years	4.6	3.1	4.5	†	8.7	5.5	8.3	7.4	2.2	
45 to 49 years	5.8	3.6	5.1	†	11.4	7.3	9.3	9.4	2.5	
50 to 54 years	7.3	3.6	6.0	†	13.2	9.8	10.9	13.0	3.1	
55 to 59 years	7.3	3.1	5.6	†	12.4	10.4	9.7	14.1	2.8	
60 to 64 years	5.5	1.8	4.1	†	8.9	8.4	5.5	11.2	1.9	
65 years and older	5.3	1.5	4.4	†	7.3	7.9	5.7	11.2	2.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†	
Asian ³	1.1	†	1.0	†	†	1.2	†	1.7	†	
Black or African American	18.0	19.6	9.8	19.6	13.6	13.8	19.0	28.1	20.7	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	63.7	60.2	74.5	58.3	71.6	66.7	67.4	55.9	60.0	
Other	16.5	18.5	14.1	21.0	13.0	17.3	11.9	13.5	17.8	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.6b. Connecticut individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	71.7	64.4	74.6	57.7	80.6	72.2	81.1	82.3	68.8
Hispanic or Latino origin	28.3	35.6	25.4	42.3	19.4	27.8	18.9	17.7	31.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	2.8	4.0	3.6	3.9	†	2.6	4.9	1.6	4.2
Puerto Rican	62.1	59.7	57.2	77.7	62.0	62.4	66.7	60.0	62.7
Cuban	†	†	†	†	1.2	†	--	1.2	†
Other specific	34.3	35.6	38.3	17.8	36.0	34.3	28.4	37.2	32.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	3.3	4.3	2.1	1.1	2.7	2.8	6.3	4.8	3.0
Not a veteran	96.7	95.7	97.9	98.9	97.3	97.2	93.8	95.2	97.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.7a. Delaware individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	8,800	612	205	254	1,490	1,925	33	1,683	2,598
Gender									
Male	4,766	319	95	184	709	925	13	1,026	1,495
Female	3,887	287	110	70	773	995	20	654	978
Total	8,653	606	205	254	1,482	1,920	33	1,680	2,473
Age²									
0 to 11 years	308	72	11	77	73	22	--	2	51
12 to 14 years	399	59	9	72	79	107	1	2	70
15 to 17 years	461	47	16	55	100	159	1	3	80
18 to 20 years	453	85	14	27	71	146	--	37	73
21 to 24 years	715	58	14	13	113	159	5	124	229
25 to 29 years	1,033	63	23	3	151	177	4	173	439
30 to 34 years	893	38	23	1	131	152	4	161	383
35 to 39 years	916	41	22	1	153	154	4	180	361
40 to 44 years	682	34	13	3	109	137	4	141	241
45 to 49 years	715	31	14	--	119	164	3	170	214
50 to 54 years	690	22	12	1	125	162	1	175	192
55 to 59 years	663	38	17	--	118	157	3	191	139
60 to 64 years	424	6	10	1	79	113	1	147	67
65 years and older	447	18	7	--	69	116	2	176	59
Total	8,799	612	205	254	1,490	1,925	33	1,682	2,598
Race									
American Indian or Alaska Native	60	5	--	1	13	9	--	14	18
Asian ³	54	6	2	--	9	10	--	18	9
Black or African American	2,818	209	53	121	475	583	10	794	573
Native Hawaiian or Other Pacific Islander	46	4	--	--	6	9	1	16	10
White	5,451	356	141	131	937	1,191	20	775	1,900
Other	121	8	2	--	12	35	--	32	32
Total	8,550	588	198	253	1,452	1,837	31	1,649	2,542

Continued. See notes at end of table.

Table 10.7a. Delaware individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	7,708	511	166	229	1,325	1,623	28	1,480	2,346
Hispanic or Latino origin	502	62	18	24	73	131	2	81	111
Total	8,210	573	184	253	1,398	1,754	30	1,561	2,457
Specific Hispanic or Latino origin									
Mexican	57	9	2	--	4	12	--	13	17
Puerto Rican	105	4	2	--	15	29	--	28	27
Cuban	5	1	--	--	2	--	--	1	1
Other specific	132	12	7	--	20	35	1	25	32
Total	299	26	11	--	41	76	1	67	77
Veteran status⁴									
Veteran	252	25	8	--	45	64	--	57	53
Not a veteran	5,412	322	128	11	931	1,266	26	1,181	1,547
Total	5,664	347	136	11	976	1,330	26	1,238	1,600

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.7b. Delaware individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	55.1	52.6	46.3	72.4	47.8	48.2	39.4	61.1	60.5	
Female	44.9	47.4	53.7	27.6	52.2	51.8	60.6	38.9	39.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	3.5	11.8	5.4	30.3	4.9	1.1	--	†	2.0	
12 to 14 years	4.5	9.6	4.4	28.3	5.3	5.6	3.0	†	2.7	
15 to 17 years	5.2	7.7	7.8	21.7	6.7	8.3	3.0	†	3.1	
18 to 20 years	5.1	13.9	6.8	10.6	4.8	7.6	--	2.2	2.8	
21 to 24 years	8.1	9.5	6.8	5.1	7.6	8.3	15.2	7.4	8.8	
25 to 29 years	11.7	10.3	11.2	1.2	10.1	9.2	12.1	10.3	16.9	
30 to 34 years	10.1	6.2	11.2	†	8.8	7.9	12.1	9.6	14.7	
35 to 39 years	10.4	6.7	10.7	†	10.3	8.0	12.1	10.7	13.9	
40 to 44 years	7.8	5.6	6.3	1.2	7.3	7.1	12.1	8.4	9.3	
45 to 49 years	8.1	5.1	6.8	--	8.0	8.5	9.1	10.1	8.2	
50 to 54 years	7.8	3.6	5.9	†	8.4	8.4	3.0	10.4	7.4	
55 to 59 years	7.5	6.2	8.3	--	7.9	8.2	9.1	11.4	5.4	
60 to 64 years	4.8	†	4.9	†	5.3	5.9	3.0	8.7	2.6	
65 years and older	5.1	2.9	3.4	--	4.6	6.0	6.1	10.5	2.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	--	†	†	†	--	†	†	
Asian ³	†	1.0	1.0	--	†	†	--	1.1	†	
Black or African American	33.0	35.5	26.8	47.8	32.7	31.7	32.3	48.2	22.5	
Native Hawaiian or Other Pacific Islander	†	†	--	--	†	†	3.2	†	†	
White	63.8	60.5	71.2	51.8	64.5	64.8	64.5	47.0	74.7	
Other	1.4	1.4	1.0	--	†	1.9	--	1.9	1.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.7b. Delaware individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	93.9	89.2	90.2	90.5	94.8	92.5	93.3	94.8	95.5
Hispanic or Latino origin	6.1	10.8	9.8	9.5	5.2	7.5	6.7	5.2	4.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	19.1	34.6	18.2	--	9.8	15.8	--	19.4	22.1
Puerto Rican	35.1	15.4	18.2	--	36.6	38.2	--	41.8	35.1
Cuban	1.7	3.8	--	--	4.9	--	--	1.5	1.3
Other specific	44.1	46.2	63.6	--	48.8	46.1	100.0	37.3	41.6
Total	100.0	100.0	100.0	--	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	4.4	7.2	5.9	--	4.6	4.8	--	4.6	3.3
Not a veteran	95.6	92.8	94.1	100.0	95.4	95.2	100.0	95.4	96.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.8a. District of Columbia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	35,828	2,880	1,263	2,399	6,155	12,846	38	6,670	3,577
Gender									
Male	18,544	1,390	506	1,658	2,619	6,154	11	4,044	2,162
Female	17,223	1,489	756	740	3,529	6,668	27	2,614	1,400
Total	35,767	2,879	1,262	2,398	6,148	12,822	38	6,658	3,562
Age²									
0 to 11 years	1,484	260	70	703	21	188	--	2	240
12 to 14 years	1,208	220	48	460	26	253	--	8	193
15 to 17 years	1,587	259	61	400	87	411	--	13	356
18 to 20 years	1,584	221	52	308	194	500	1	24	284
21 to 24 years	1,927	165	68	206	376	696	1	196	219
25 to 29 years	3,124	242	117	90	751	1,170	2	446	306
30 to 34 years	3,003	220	125	48	714	1,100	5	531	260
35 to 39 years	2,964	234	120	38	602	1,090	5	612	263
40 to 44 years	2,433	176	95	30	517	895	3	491	226
45 to 49 years	2,787	202	95	36	572	1,111	2	576	193
50 to 54 years	3,991	252	114	20	827	1,612	5	860	301
55 to 59 years	4,289	193	118	30	777	1,809	7	1,037	318
60 to 64 years	3,097	149	114	15	434	1,259	4	875	247
65 years and older	2,350	87	66	15	257	752	3	999	171
Total	35,828	2,880	1,263	2,399	6,155	12,846	38	6,670	3,577
Race									
American Indian or Alaska Native	57	2	2	--	12	21	--	18	2
Asian ³	106	4	11	--	12	30	--	41	8
Black or African American	32,638	2,695	1,019	2,300	5,654	11,944	34	5,715	3,277
Native Hawaiian or Other Pacific Islander	1	--	--	--	--	--	--	1	--
White	1,153	53	75	33	252	291	3	353	93
Other	67	7	7	1	16	20	--	10	6
Total	34,022	2,761	1,114	2,334	5,946	12,306	37	6,138	3,386

Continued. See notes at end of table.

Table 10.8a. District of Columbia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	12,736	978	482	614	2,446	4,222	14	2,476	1,504
Hispanic or Latino origin	675	46	49	22	96	234	1	162	65
Total	13,411	1,024	531	636	2,542	4,456	15	2,638	1,569
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.8b. District of Columbia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	51.8	48.3	40.1	69.1	42.6	48.0	28.9	60.7	60.7
Female	48.2	51.7	59.9	30.9	57.4	52.0	71.1	39.3	39.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	4.1	9.0	5.5	29.3	†	1.5	--	†	6.7
12 to 14 years	3.4	7.6	3.8	19.2	†	2.0	--	†	5.4
15 to 17 years	4.4	9.0	4.8	16.7	1.4	3.2	--	†	10.0
18 to 20 years	4.4	7.7	4.1	12.8	3.2	3.9	2.6	†	7.9
21 to 24 years	5.4	5.7	5.4	8.6	6.1	5.4	2.6	2.9	6.1
25 to 29 years	8.7	8.4	9.3	3.8	12.2	9.1	5.3	6.7	8.6
30 to 34 years	8.4	7.6	9.9	2.0	11.6	8.6	13.2	8.0	7.3
35 to 39 years	8.3	8.1	9.5	1.6	9.8	8.5	13.2	9.2	7.4
40 to 44 years	6.8	6.1	7.5	1.3	8.4	7.0	7.9	7.4	6.3
45 to 49 years	7.8	7.0	7.5	1.5	9.3	8.6	5.3	8.6	5.4
50 to 54 years	11.1	8.8	9.0	†	13.4	12.5	13.2	12.9	8.4
55 to 59 years	12.0	6.7	9.3	1.3	12.6	14.1	18.4	15.5	8.9
60 to 64 years	8.6	5.2	9.0	†	7.1	9.8	10.5	13.1	6.9
65 years and older	6.6	3.0	5.2	†	4.2	5.9	7.9	15.0	4.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	†	†	--	†	†	--	†	†
Asian ³	†	†	†	--	†	†	--	†	†
Black or African American	95.9	97.6	91.5	98.5	95.1	97.1	91.9	93.1	96.8
Native Hawaiian or Other Pacific Islander	†	--	--	--	--	--	--	†	--
White	3.4	1.9	6.7	1.4	4.2	2.4	8.1	5.8	2.7
Other	†	†	†	†	†	†	--	†	†
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.8b. District of Columbia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

<i>Selected characteristics</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Ethnicity									
Not of Hispanic or Latino origin	95.0	95.5	90.8	96.5	96.2	94.7	93.3	93.9	95.9
Hispanic or Latino origin	5.0	4.5	9.2	3.5	3.8	5.3	6.7	6.1	4.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.9a. Florida individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	185,688	26,900	14,027	13,081	26,120	53,057	515	32,965	19,023
Gender									
Male	93,454	12,666	5,517	9,119	10,905	22,761	236	21,308	10,942
Female	92,233	14,234	8,510	3,962	15,215	30,295	279	11,657	8,081
Total	185,687	26,900	14,027	13,081	26,120	53,056	515	32,965	19,023
Age²									
0 to 11 years	12,642	3,843	529	5,095	36	498	2	24	2,615
12 to 14 years	9,142	2,119	461	2,892	131	1,622	3	51	1,863
15 to 17 years	11,206	2,311	705	2,202	341	3,181	9	155	2,302
18 to 20 years	8,606	1,544	740	882	743	2,758	18	570	1,351
21 to 24 years	11,233	1,684	919	389	1,543	3,421	55	2,097	1,125
25 to 29 years	17,868	2,593	1,575	387	2,837	4,936	83	3,843	1,614
30 to 34 years	18,216	2,670	1,674	331	3,046	5,033	79	3,918	1,465
35 to 39 years	16,956	2,357	1,504	253	3,108	4,625	70	3,763	1,276
40 to 44 years	13,980	1,767	1,148	185	2,648	4,048	44	3,152	988
45 to 49 years	14,482	1,571	1,044	158	2,887	4,571	50	3,228	973
50 to 54 years	15,508	1,537	1,089	100	3,005	5,366	39	3,391	981
55 to 59 years	15,650	1,378	1,070	90	2,809	5,556	30	3,748	969
60 to 64 years	10,872	758	815	76	1,753	4,009	21	2,757	683
65 years and older	9,327	768	754	41	1,233	3,433	12	2,268	818
Total	185,688	26,900	14,027	13,081	26,120	53,057	515	32,965	19,023
Race									
American Indian or Alaska Native	636	89	43	34	98	173	1	129	69
Asian ³	1,102	175	72	35	110	319	2	308	81
Black or African American	40,974	5,441	1,558	3,009	4,056	9,447	84	13,152	4,227
Native Hawaiian or Other Pacific Islander	360	47	31	20	53	112	1	47	49
White	123,259	17,881	10,764	8,299	19,635	36,879	394	16,778	12,629
Other	12,667	2,006	911	1,136	1,529	3,822	26	1,811	1,426
Total	178,998	25,639	13,379	12,533	25,481	50,752	508	32,225	18,481

Continued. See notes at end of table.

Table 10.9a. Florida individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	150,482	22,250	11,065	10,378	22,198	41,901	464	27,963	14,263
Hispanic or Latino origin	35,182	4,647	2,960	2,703	3,920	11,153	51	4,993	4,755
Total	185,664	26,897	14,025	13,081	26,118	53,054	515	32,956	19,018
Specific Hispanic or Latino origin									
Mexican	2,803	626	218	360	166	838	5	301	289
Puerto Rican	7,260	821	523	674	1,062	2,282	13	1,140	745
Cuban	6,289	370	497	258	717	2,263	9	1,077	1,098
Other specific	14,461	2,152	1,245	1,109	1,552	4,240	19	1,935	2,209
Total	30,813	3,969	2,483	2,401	3,497	9,623	46	4,453	4,341
Veteran status⁴									
Veteran	7,047	790	523	61	1,119	2,410	13	1,824	307
Not a veteran	143,390	17,549	11,641	2,808	24,181	44,623	469	30,419	11,700
Total	150,437	18,339	12,164	2,869	25,300	47,033	482	32,243	12,007

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting periods. Data as of 07.08.2019.

Table 10.9b. Florida individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	50.3	47.1	39.3	69.7	41.7	42.9	45.8	64.6	57.5	
Female	49.7	52.9	60.7	30.3	58.3	57.1	54.2	35.4	42.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	6.8	14.3	3.8	38.9	†	†	†	†	13.7	
12 to 14 years	4.9	7.9	3.3	22.1	†	3.1	†	†	9.8	
15 to 17 years	6.0	8.6	5.0	16.8	1.3	6.0	1.7	†	12.1	
18 to 20 years	4.6	5.7	5.3	6.7	2.8	5.2	3.5	1.7	7.1	
21 to 24 years	6.0	6.3	6.6	3.0	5.9	6.4	10.7	6.4	5.9	
25 to 29 years	9.6	9.6	11.2	3.0	10.9	9.3	16.1	11.7	8.5	
30 to 34 years	9.8	9.9	11.9	2.5	11.7	9.5	15.3	11.9	7.7	
35 to 39 years	9.1	8.8	10.7	1.9	11.9	8.7	13.6	11.4	6.7	
40 to 44 years	7.5	6.6	8.2	1.4	10.1	7.6	8.5	9.6	5.2	
45 to 49 years	7.8	5.8	7.4	1.2	11.1	8.6	9.7	9.8	5.1	
50 to 54 years	8.4	5.7	7.8	†	11.5	10.1	7.6	10.3	5.2	
55 to 59 years	8.4	5.1	7.6	†	10.8	10.5	5.8	11.4	5.1	
60 to 64 years	5.9	2.8	5.8	†	6.7	7.6	4.1	8.4	3.6	
65 years and older	5.0	2.9	5.4	†	4.7	6.5	2.3	6.9	4.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†	
Asian ³	†	†	†	†	†	†	†	†	†	
Black or African American	22.9	21.2	11.6	24.0	15.9	18.6	16.5	40.8	22.9	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	68.9	69.7	80.5	66.2	77.1	72.7	77.6	52.1	68.3	
Other	7.1	7.8	6.8	9.1	6.0	7.5	5.1	5.6	7.7	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.9b. Florida individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	81.1	82.7	78.9	79.3	85.0	79.0	90.1	84.8	75.0
Hispanic or Latino origin	18.9	17.3	21.1	20.7	15.0	21.0	9.9	15.2	25.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	9.1	15.8	8.8	15.0	4.7	8.7	10.9	6.8	6.7
Puerto Rican	23.6	20.7	21.1	28.1	30.4	23.7	28.3	25.6	17.2
Cuban	20.4	9.3	20.0	10.7	20.5	23.5	19.6	24.2	25.3
Other specific	46.9	54.2	50.1	46.2	44.4	44.1	41.3	43.5	50.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	4.7	4.3	4.3	2.1	4.4	5.1	2.7	5.7	2.6
Not a veteran	95.3	95.7	95.7	97.9	95.6	94.9	97.3	94.3	97.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.10a. Hawaii individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	9,097	2,401	275	395	1,032	1,557	15	2,719	703
Gender									
Male	5,275	1,215	158	307	540	799	10	1,765	481
Female	3,819	1,186	117	88	492	756	5	953	222
Total	9,094	2,401	275	395	1,032	1,555	15	2,718	703
Age²									
0 to 11 years	534	154	24	179	2	18	--	--	157
12 to 14 years	379	78	28	93	2	48	1	1	128
15 to 17 years	578	115	47	62	9	117	1	13	214
18 to 20 years	282	106	11	13	15	54	--	32	51
21 to 24 years	315	144	3	4	25	58	1	73	7
25 to 29 years	576	222	7	3	66	93	2	174	9
30 to 34 years	665	213	10	7	89	113	2	218	13
35 to 39 years	673	209	9	3	88	97	2	256	9
40 to 44 years	624	174	13	3	79	109	2	233	11
45 to 49 years	742	185	24	4	117	107	1	292	12
50 to 54 years	881	192	18	4	143	157	1	344	22
55 to 59 years	1,028	202	30	5	147	215	--	407	22
60 to 64 years	859	180	22	10	110	178	1	339	19
65 years and older	955	224	29	4	140	193	1	336	28
Total	9,091	2,398	275	394	1,032	1,557	15	2,718	702
Race									
American Indian or Alaska Native	65	22	1	--	9	11	--	19	3
Asian ³	1,388	180	45	13	106	167	2	799	76
Black or African American	206	38	3	8	19	34	--	97	7
Native Hawaiian or Other Pacific Islander	968	255	24	31	91	126	2	350	89
White	2,086	488	63	48	394	398	4	615	76
Other	1,493	276	62	131	153	217	3	481	170
Total	6,206	1,259	198	231	772	953	11	2,361	421

Continued. See notes at end of table.

Table 10.10a. Hawaii individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	5,512	1,114	167	135	722	852	9	2,250	263
Hispanic or Latino origin	527	96	16	63	61	81	1	127	82
Total	6,039	1,210	183	198	783	933	10	2,377	345
Specific Hispanic or Latino origin									
Mexican	30	6	--	--	8	5	--	9	2
Puerto Rican	131	22	1	3	22	27	--	52	4
Cuban	4	--	1	--	2	--	--	1	--
Other specific	115	24	--	--	19	19	1	48	4
Total	280	52	2	3	51	51	1	110	10
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.10b. Hawaii individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	58.0	50.6	57.5	77.7	52.3	51.4	66.7	64.9	68.4	
Female	42.0	49.4	42.5	22.3	47.7	48.6	33.3	35.1	31.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	5.9	6.4	8.7	45.4	†	1.2	--	--	22.4	
12 to 14 years	4.2	3.3	10.2	23.6	†	3.1	6.7	†	18.2	
15 to 17 years	6.4	4.8	17.1	15.7	†	7.5	6.7	†	30.5	
18 to 20 years	3.1	4.4	4.0	3.3	1.5	3.5	--	1.2	7.3	
21 to 24 years	3.5	6.0	1.1	1.0	2.4	3.7	6.7	2.7	†	
25 to 29 years	6.3	9.3	2.5	†	6.4	6.0	13.3	6.4	1.3	
30 to 34 years	7.3	8.9	3.6	1.8	8.6	7.3	13.3	8.0	1.9	
35 to 39 years	7.4	8.7	3.3	†	8.5	6.2	13.3	9.4	1.3	
40 to 44 years	6.9	7.3	4.7	†	7.7	7.0	13.3	8.6	1.6	
45 to 49 years	8.2	7.7	8.7	1.0	11.3	6.9	6.7	10.7	1.7	
50 to 54 years	9.7	8.0	6.5	1.0	13.9	10.1	6.7	12.7	3.1	
55 to 59 years	11.3	8.4	10.9	1.3	14.2	13.8	--	15.0	3.1	
60 to 64 years	9.4	7.5	8.0	2.5	10.7	11.4	6.7	12.5	2.7	
65 years and older	10.5	9.3	10.5	1.0	13.6	12.4	6.7	12.4	4.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	1.0	1.7	†	--	1.2	1.2	--	†	†	
Asian ³	22.4	14.3	22.7	5.6	13.7	17.5	18.2	33.8	18.1	
Black or African American	3.3	3.0	1.5	3.5	2.5	3.6	--	4.1	1.7	
Native Hawaiian or Other Pacific Islander	15.6	20.3	12.1	13.4	11.8	13.2	18.2	14.8	21.1	
White	33.6	38.8	31.8	20.8	51.0	41.8	36.4	26.0	18.1	
Other	24.1	21.9	31.3	56.7	19.8	22.8	27.3	20.4	40.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.10b. Hawaii individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	91.3	92.1	91.3	68.2	92.2	91.3	90.0	94.7	76.2
Hispanic or Latino origin	8.7	7.9	8.7	31.8	7.8	8.7	10.0	5.3	23.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	10.7	11.5	--	--	15.7	9.8	--	8.2	20.0
Puerto Rican	46.8	42.3	50.0	100.0	43.1	52.9	--	47.3	40.0
Cuban	1.4	--	50.0	--	3.9	--	--	†	--
Other specific	41.1	46.2	--	--	37.3	37.3	100.0	43.6	40.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.11a. Idaho individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	7,666	291	383	218	1,536	2,271	154	1,985	828
Gender									
Male	4,203	152	185	185	678	1,077	84	1,281	561
Female	3,462	139	197	33	858	1,194	70	704	267
Total	7,665	291	382	218	1,536	2,271	154	1,985	828
Age²									
0 to 11 years	141	14	15	30	3	12	--	1	66
12 to 14 years	458	49	57	61	9	92	1	5	184
15 to 17 years	707	77	99	76	25	204	3	12	211
18 to 20 years	437	19	28	20	55	155	14	75	71
21 to 24 years	588	21	25	8	124	174	16	163	57
25 to 29 years	855	20	37	4	185	268	31	264	46
30 to 34 years	801	28	26	6	203	230	15	254	39
35 to 39 years	818	9	26	6	230	243	19	246	39
40 to 44 years	616	14	16	2	168	183	19	186	28
45 to 49 years	616	16	21	2	164	201	10	180	22
50 to 54 years	533	9	11	1	128	177	11	176	20
55 to 59 years	510	8	13	1	121	155	9	187	16
60 to 64 years	310	2	3	1	69	101	4	118	12
65 years and older	272	5	6	--	49	75	2	118	17
Total	7,662	291	383	218	1,533	2,270	154	1,985	828
Race									
American Indian or Alaska Native	161	7	5	2	26	46	6	48	21
Asian ³	47	2	4	1	4	12	1	18	5
Black or African American	138	9	13	3	14	28	2	56	13
Native Hawaiian or Other Pacific Islander	25	--	2	1	3	6	3	7	3
White	6,082	216	283	172	1,279	1,835	115	1,560	622
Other	594	32	33	19	97	175	8	155	75
Total	7,047	266	340	198	1,423	2,102	135	1,844	739

Continued. See notes at end of table.

Table 10.11a. Idaho individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	6,070	222	285	175	1,267	1,824	121	1,573	603
Hispanic or Latino origin	712	32	44	23	107	202	6	197	101
Total	6,782	254	329	198	1,374	2,026	127	1,770	704
Specific Hispanic or Latino origin									
Mexican	100	5	8	--	18	24	--	34	11
Puerto Rican	14	1	--	1	4	2	1	5	--
Cuban	1	--	--	--	--	--	--	--	1
Other specific	89	1	3	2	21	24	2	28	8
Total	204	7	11	3	43	50	3	67	20
Veteran status⁴									
Veteran	245	6	13	1	39	68	4	101	13
Not a veteran	4,928	115	165	41	1,178	1,569	109	1,459	292
Total	5,173	121	178	42	1,217	1,637	113	1,560	305

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.11b. Idaho individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	54.8	52.2	48.4	84.9	44.1	47.4	54.5	64.5	67.8	
Female	45.2	47.8	51.6	15.1	55.9	52.6	45.5	35.5	32.2	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	1.8	4.8	3.9	13.8	†	†	--	†	8.0	
12 to 14 years	6.0	16.8	14.9	28.0	†	4.1	†	†	22.2	
15 to 17 years	9.2	26.5	25.8	34.9	1.6	9.0	1.9	†	25.5	
18 to 20 years	5.7	6.5	7.3	9.2	3.6	6.8	9.1	3.8	8.6	
21 to 24 years	7.7	7.2	6.5	3.7	8.1	7.7	10.4	8.2	6.9	
25 to 29 years	11.2	6.9	9.7	1.8	12.1	11.8	20.1	13.3	5.6	
30 to 34 years	10.5	9.6	6.8	2.8	13.2	10.1	9.7	12.8	4.7	
35 to 39 years	10.7	3.1	6.8	2.8	15.0	10.7	12.3	12.4	4.7	
40 to 44 years	8.0	4.8	4.2	†	11.0	8.1	12.3	9.4	3.4	
45 to 49 years	8.0	5.5	5.5	†	10.7	8.9	6.5	9.1	2.7	
50 to 54 years	7.0	3.1	2.9	†	8.3	7.8	7.1	8.9	2.4	
55 to 59 years	6.7	2.7	3.4	†	7.9	6.8	5.8	9.4	1.9	
60 to 64 years	4.0	†	†	†	4.5	4.4	2.6	5.9	1.4	
65 years and older	3.5	1.7	1.6	--	3.2	3.3	1.3	5.9	2.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	2.3	2.6	1.5	1.0	1.8	2.2	4.4	2.6	2.8	
Asian ³	†	†	1.2	†	†	†	†	†	†	
Black or African American	2.0	3.4	3.8	1.5	†	1.3	1.5	3.0	1.8	
Native Hawaiian or Other Pacific Islander	†	--	†	†	†	†	2.2	†	†	
White	86.3	81.2	83.2	86.9	89.9	87.3	85.2	84.6	84.2	
Other	8.4	12.0	9.7	9.6	6.8	8.3	5.9	8.4	10.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.11b. Idaho individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	89.5	87.4	86.6	88.4	92.2	90.0	95.3	88.9	85.7
Hispanic or Latino origin	10.5	12.6	13.4	11.6	7.8	10.0	4.7	11.1	14.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	49.0	71.4	72.7	--	41.9	48.0	--	50.7	55.0
Puerto Rican	6.9	14.3	--	33.3	9.3	4.0	33.3	7.5	--
Cuban	†	--	--	--	--	--	--	--	5.0
Other specific	43.6	14.3	27.3	66.7	48.8	48.0	66.7	41.8	40.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	4.7	5.0	7.3	2.4	3.2	4.2	3.5	6.5	4.3
Not a veteran	95.3	95.0	92.7	97.6	96.8	95.8	96.5	93.5	95.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.12a. Illinois individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	66,235	8,809	5,655	5,127	10,132	18,504	241	10,409	7,358
Gender									
Male	31,862	3,764	2,186	3,691	4,135	6,854	86	6,802	4,344
Female	34,373	5,045	3,469	1,436	5,997	11,650	155	3,607	3,014
Total	66,235	8,809	5,655	5,127	10,132	18,504	241	10,409	7,358
Age²									
0 to 11 years	7,725	2,548	505	2,636	44	382	--	17	1,593
12 to 14 years	4,710	977	481	1,185	109	1,020	1	24	913
15 to 17 years	5,014	840	562	698	289	1,736	3	40	846
18 to 20 years	3,191	428	356	238	436	1,125	18	212	378
21 to 24 years	3,931	443	412	76	749	1,221	22	562	446
25 to 29 years	5,816	686	622	81	1,091	1,717	43	955	621
30 to 34 years	5,403	619	558	66	1,100	1,580	38	929	513
35 to 39 years	5,315	540	521	39	1,110	1,507	37	1,101	460
40 to 44 years	4,572	440	341	36	1,055	1,336	21	979	364
45 to 49 years	4,706	343	324	17	1,069	1,510	21	1,094	328
50 to 54 years	4,930	333	333	23	1,021	1,609	11	1,315	285
55 to 59 years	4,943	290	291	16	1,036	1,636	13	1,386	275
60 to 64 years	3,198	160	170	12	589	1,121	7	970	169
65 years and older	2,781	162	179	4	434	1,004	6	825	167
Total	66,235	8,809	5,655	5,127	10,132	18,504	241	10,409	7,358
Race									
American Indian or Alaska Native	213	27	14	11	45	57	1	40	18
Asian ³	676	88	41	22	93	206	1	192	33
Black or African American	11,775	1,158	410	884	1,712	2,678	36	3,474	1,423
Native Hawaiian or Other Pacific Islander	50	2	5	3	7	15	--	12	6
White	48,139	6,529	4,742	3,630	7,685	14,185	188	6,067	5,113
Other	910	207	71	92	110	183	1	139	107
Total	61,763	8,011	5,283	4,642	9,652	17,324	227	9,924	6,700

Continued. See notes at end of table.

Table 10.12a. Illinois individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	57,572	7,381	4,863	4,356	8,994	16,126	219	9,257	6,376
Hispanic or Latino origin	5,241	882	471	436	699	1,478	15	723	537
Total	62,813	8,263	5,334	4,792	9,693	17,604	234	9,980	6,913
Specific Hispanic or Latino origin									
Mexican	2,967	568	285	275	352	861	9	354	263
Puerto Rican	821	113	62	71	139	227	2	133	74
Cuban	55	7	7	4	8	14	1	12	2
Other specific	1,247	189	116	86	166	349	3	158	180
Total	5,090	877	470	436	665	1,451	15	657	519
Veteran status⁴									
Veteran	1,153	133	83	8	224	292	5	335	73
Not a veteran	43,866	3,908	3,814	572	8,696	13,952	185	9,169	3,570
Total	45,019	4,041	3,897	580	8,920	14,244	190	9,504	3,643

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.12b. Illinois individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	48.1	42.7	38.7	72.0	40.8	37.0	35.7	65.3	59.0	
Female	51.9	57.3	61.3	28.0	59.2	63.0	64.3	34.7	41.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	11.7	28.9	8.9	51.4	†	2.1	--	†	21.6	
12 to 14 years	7.1	11.1	8.5	23.1	1.1	5.5	†	†	12.4	
15 to 17 years	7.6	9.5	9.9	13.6	2.9	9.4	1.2	†	11.5	
18 to 20 years	4.8	4.9	6.3	4.6	4.3	6.1	7.5	2.0	5.1	
21 to 24 years	5.9	5.0	7.3	1.5	7.4	6.6	9.1	5.4	6.1	
25 to 29 years	8.8	7.8	11.0	1.6	10.8	9.3	17.8	9.2	8.4	
30 to 34 years	8.2	7.0	9.9	1.3	10.9	8.5	15.8	8.9	7.0	
35 to 39 years	8.0	6.1	9.2	†	11.0	8.1	15.4	10.6	6.3	
40 to 44 years	6.9	5.0	6.0	†	10.4	7.2	8.7	9.4	4.9	
45 to 49 years	7.1	3.9	5.7	†	10.6	8.2	8.7	10.5	4.5	
50 to 54 years	7.4	3.8	5.9	†	10.1	8.7	4.6	12.6	3.9	
55 to 59 years	7.5	3.3	5.1	†	10.2	8.8	5.4	13.3	3.7	
60 to 64 years	4.8	1.8	3.0	†	5.8	6.1	2.9	9.3	2.3	
65 years and older	4.2	1.8	3.2	†	4.3	5.4	2.5	7.9	2.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†	
Asian ³	1.1	1.1	†	†	†	1.2	†	1.9	†	
Black or African American	19.1	14.5	7.8	19.0	17.7	15.5	15.9	35.0	21.2	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†	
White	77.9	81.5	89.8	78.2	79.6	81.9	82.8	61.1	76.3	
Other	1.5	2.6	1.3	2.0	1.1	1.1	†	1.4	1.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.12b. Illinois individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	91.7	89.3	91.2	90.9	92.8	91.6	93.6	92.8	92.2
Hispanic or Latino origin	8.3	10.7	8.8	9.1	7.2	8.4	6.4	7.2	7.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	58.3	64.8	60.6	63.1	52.9	59.3	60.0	53.9	50.7
Puerto Rican	16.1	12.9	13.2	16.3	20.9	15.6	13.3	20.2	14.3
Cuban	1.1	†	1.5	†	1.2	†	6.7	1.8	†
Other specific	24.5	21.6	24.7	19.7	25.0	24.1	20.0	24.0	34.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	2.6	3.3	2.1	1.4	2.5	2.0	2.6	3.5	2.0
Not a veteran	97.4	96.7	97.9	98.6	97.5	98.0	97.4	96.5	98.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.13a. Indiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	137,363	11,718	13,401	14,142	13,813	47,509	1,044	14,899	20,837
Gender									
Male	65,045	4,903	5,792	10,238	5,055	16,834	363	9,084	12,776
Female	72,318	6,815	7,609	3,904	8,758	30,675	681	5,815	8,061
Total	137,363	11,718	13,401	14,142	13,813	47,509	1,044	14,899	20,837
Age²									
0 to 11 years	29,660	4,399	3,572	8,703	120	2,983	--	49	9,834
12 to 14 years	13,682	1,447	1,404	2,963	207	4,174	6	66	3,415
15 to 17 years	12,418	1,016	1,119	1,434	480	5,553	22	123	2,671
18 to 20 years	5,885	411	635	344	554	2,873	70	285	713
21 to 24 years	6,659	516	750	140	891	3,001	103	746	512
25 to 29 years	9,445	739	1,134	179	1,326	3,979	197	1,296	595
30 to 34 years	8,841	701	1,015	110	1,401	3,599	154	1,373	488
35 to 39 years	8,868	654	978	108	1,647	3,342	149	1,478	512
40 to 44 years	7,930	481	635	55	1,492	3,214	111	1,491	451
45 to 49 years	8,041	435	580	37	1,571	3,430	79	1,536	373
50 to 54 years	8,311	334	520	29	1,468	3,659	72	1,820	409
55 to 59 years	7,853	283	481	22	1,270	3,410	41	1,976	370
60 to 64 years	5,216	158	281	15	776	2,245	25	1,471	245
65 years and older	4,554	144	297	3	610	2,047	15	1,189	249
Total	137,363	11,718	13,401	14,142	13,813	47,509	1,044	14,899	20,837
Race									
American Indian or Alaska Native	506	41	46	49	43	178	5	72	72
Asian ³	527	32	55	32	37	192	3	118	58
Black or African American	20,172	1,260	1,329	2,678	1,534	5,557	142	4,120	3,552
Native Hawaiian or Other Pacific Islander	127	7	9	8	20	47	--	14	22
White	106,690	9,415	11,072	10,134	11,530	38,529	854	9,952	15,204
Other	9,340	963	890	1,241	649	3,006	39	623	1,929
Total	137,362	11,718	13,401	14,142	13,813	47,509	1,043	14,899	20,837

Continued. See notes at end of table.

Table 10.13a. Indiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	125,504	10,610	12,214	12,663	12,943	43,402	999	13,981	18,692
Hispanic or Latino origin	9,612	928	1,018	1,203	613	3,459	31	600	1,760
Total	135,116	11,538	13,232	13,866	13,556	46,861	1,030	14,581	20,452
Specific Hispanic or Latino origin									
Mexican	3,410	331	376	446	164	1,310	5	148	630
Puerto Rican	739	48	75	89	81	271	1	74	100
Cuban	69	7	6	8	7	23	--	7	11
Other specific	5,362	541	559	660	358	1,847	25	360	1,012
Total	9,580	927	1,016	1,203	610	3,451	31	589	1,753
Veteran status⁴									
Veteran	2,626	225	178	24	388	985	29	670	127
Not a veteran	653	9	5	1	82	38	39	410	69
Total	3,279	234	183	25	470	1,023	68	1,080	196

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.13b. Indiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	47.4	41.8	43.2	72.4	36.6	35.4	34.8	61.0	61.3
Female	52.6	58.2	56.8	27.6	63.4	64.6	65.2	39.0	38.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	21.6	37.5	26.7	61.5	†	6.3	--	†	47.2
12 to 14 years	10.0	12.3	10.5	21.0	1.5	8.8	†	†	16.4
15 to 17 years	9.0	8.7	8.4	10.1	3.5	11.7	2.1	†	12.8
18 to 20 years	4.3	3.5	4.7	2.4	4.0	6.0	6.7	1.9	3.4
21 to 24 years	4.8	4.4	5.6	†	6.5	6.3	9.9	5.0	2.5
25 to 29 years	6.9	6.3	8.5	1.3	9.6	8.4	18.9	8.7	2.9
30 to 34 years	6.4	6.0	7.6	†	10.1	7.6	14.8	9.2	2.3
35 to 39 years	6.5	5.6	7.3	†	11.9	7.0	14.3	9.9	2.5
40 to 44 years	5.8	4.1	4.7	†	10.8	6.8	10.6	10.0	2.2
45 to 49 years	5.9	3.7	4.3	†	11.4	7.2	7.6	10.3	1.8
50 to 54 years	6.1	2.9	3.9	†	10.6	7.7	6.9	12.2	2.0
55 to 59 years	5.7	2.4	3.6	†	9.2	7.2	3.9	13.3	1.8
60 to 64 years	3.8	1.3	2.1	†	5.6	4.7	2.4	9.9	1.2
65 years and older	3.3	1.2	2.2	†	4.4	4.3	1.4	8.0	1.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†
Asian ³	†	†	†	†	†	†	†	†	†
Black or African American	14.7	10.8	9.9	18.9	11.1	11.7	13.6	27.7	17.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†
White	77.7	80.3	82.6	71.7	83.5	81.1	81.9	66.8	73.0
Other	6.8	8.2	6.6	8.8	4.7	6.3	3.7	4.2	9.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.13b. Indiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	92.9	92.0	92.3	91.3	95.5	92.6	97.0	95.9	91.4
Hispanic or Latino origin	7.1	8.0	7.7	8.7	4.5	7.4	3.0	4.1	8.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	35.6	35.7	37.0	37.1	26.9	38.0	16.1	25.1	35.9
Puerto Rican	7.7	5.2	7.4	7.4	13.3	7.9	3.2	12.6	5.7
Cuban	†	†	†	†	1.1	†	--	1.2	†
Other specific	56.0	58.4	55.0	54.9	58.7	53.5	80.6	61.1	57.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	80.1	96.2	97.3	96.0	82.6	96.3	42.6	62.0	64.8
Not a veteran	19.9	3.8	2.7	4.0	17.4	3.7	57.4	38.0	35.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.14a. Iowa individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	174,011	29,171	35,102	25,669	11,833	43,078	790	10,473	17,895
Gender									
Male	75,300	12,228	11,408	17,459	4,351	13,785	236	6,259	9,574
Female	98,710	16,943	23,694	8,209	7,482	29,293	554	4,214	8,321
Total	174,010	29,171	35,102	25,668	11,833	43,078	790	10,473	17,895
Age²									
0 to 11 years	32,635	11,888	2,665	12,324	36	735	10	17	4,960
12 to 14 years	14,292	3,725	1,797	4,694	84	1,823	5	38	2,126
15 to 17 years	13,768	2,454	2,519	2,942	225	3,322	25	78	2,203
18 to 20 years	8,843	1,010	2,272	1,301	463	2,662	49	213	873
21 to 24 years	10,007	1,122	2,840	854	773	3,017	78	480	843
25 to 29 years	15,382	1,726	4,420	1,130	1,396	4,401	116	956	1,237
30 to 34 years	13,862	1,561	3,950	836	1,472	4,070	102	948	923
35 to 39 years	12,931	1,500	3,429	595	1,526	3,851	96	1,044	890
40 to 44 years	10,169	1,065	2,511	349	1,304	3,237	71	909	723
45 to 49 years	9,634	869	2,202	239	1,181	3,289	61	1,117	676
50 to 54 years	9,727	787	1,938	191	1,198	3,682	65	1,225	641
55 to 59 years	9,586	733	1,883	119	1,039	3,774	46	1,292	700
60 to 64 years	6,415	406	1,286	80	613	2,540	22	972	496
65 years and older	6,757	325	1,390	15	522	2,675	44	1,182	604
Total	174,008	29,171	35,102	25,669	11,832	43,078	790	10,471	17,895
Race									
American Indian or Alaska Native	2,437	550	475	354	156	561	13	105	223
Asian ³	863	126	190	74	34	246	2	101	90
Black or African American	9,667	1,746	1,535	1,353	548	2,280	39	959	1,207
Native Hawaiian or Other Pacific Islander	742	135	224	125	28	148	3	13	66
White	114,332	17,385	24,349	15,285	8,784	29,885	571	7,289	10,784
Other	6,089	1,487	961	1,372	244	1,072	14	153	786
Total	134,130	21,429	27,734	18,563	9,794	34,192	642	8,620	13,156

Continued. See notes at end of table.

Table 10.14a. Iowa individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	78	7	3	10	11	17	4	13	13
Hispanic or Latino origin	37	4	6	1	2	10	--	10	4
Total	115	11	9	11	13	27	4	23	17
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.14b. Iowa individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	43.3	41.9	32.5	68.0	36.8	32.0	29.9	59.8	53.5	
Female	56.7	58.1	67.5	32.0	63.2	68.0	70.1	40.2	46.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	18.8	40.8	7.6	48.0	†	1.7	1.3	†	27.7	
12 to 14 years	8.2	12.8	5.1	18.3	†	4.2	†	†	11.9	
15 to 17 years	7.9	8.4	7.2	11.5	1.9	7.7	3.2	†	12.3	
18 to 20 years	5.1	3.5	6.5	5.1	3.9	6.2	6.2	2.0	4.9	
21 to 24 years	5.8	3.8	8.1	3.3	6.5	7.0	9.9	4.6	4.7	
25 to 29 years	8.8	5.9	12.6	4.4	11.8	10.2	14.7	9.1	6.9	
30 to 34 years	8.0	5.4	11.3	3.3	12.4	9.4	12.9	9.1	5.2	
35 to 39 years	7.4	5.1	9.8	2.3	12.9	8.9	12.2	10.0	5.0	
40 to 44 years	5.8	3.7	7.2	1.4	11.0	7.5	9.0	8.7	4.0	
45 to 49 years	5.5	3.0	6.3	†	10.0	7.6	7.7	10.7	3.8	
50 to 54 years	5.6	2.7	5.5	†	10.1	8.5	8.2	11.7	3.6	
55 to 59 years	5.5	2.5	5.4	†	8.8	8.8	5.8	12.3	3.9	
60 to 64 years	3.7	1.4	3.7	†	5.2	5.9	2.8	9.3	2.8	
65 years and older	3.9	1.1	4.0	†	4.4	6.2	5.6	11.3	3.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	1.8	2.6	1.7	1.9	1.6	1.6	2.0	1.2	1.7	
Asian ³	†	†	†	†	†	†	†	1.2	†	
Black or African American	7.2	8.1	5.5	7.3	5.6	6.7	6.1	11.1	9.2	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	85.2	81.1	87.8	82.3	89.7	87.4	88.9	84.6	82.0	
Other	4.5	6.9	3.5	7.4	2.5	3.1	2.2	1.8	6.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.14b. Iowa individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	67.8	63.6	33.3	90.9	84.6	63.0	100.0	56.5	76.5
Hispanic or Latino origin	32.2	36.4	66.7	9.1	15.4	37.0	--	43.5	23.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.15a. Kentucky individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	155,989	22,995	17,814	13,405	11,021	35,319	941	10,367	44,127
Gender									
Male	74,856	9,677	6,950	9,490	3,856	12,531	438	6,122	25,792
Female	81,103	13,313	10,859	3,915	7,161	22,780	503	4,244	18,328
Total	155,959	22,990	17,809	13,405	11,017	35,311	941	10,366	44,120
Age²									
0 to 11 years	26,377	6,702	2,274	7,215	49	845	4	41	9,247
12 to 14 years	12,610	2,383	1,467	2,692	97	2,039	5	25	3,902
15 to 17 years	12,800	2,063	1,670	1,734	208	3,149	10	82	3,884
18 to 20 years	7,241	945	974	529	391	1,968	57	229	2,148
21 to 24 years	9,049	1,088	1,146	319	697	2,254	110	468	2,967
25 to 29 years	13,154	1,666	1,575	275	1,100	2,950	134	884	4,570
30 to 34 years	12,132	1,581	1,478	179	1,183	2,526	111	906	4,168
35 to 39 years	11,649	1,513	1,320	156	1,251	2,670	118	1,066	3,555
40 to 44 years	9,882	1,274	1,135	97	1,150	2,612	84	1,007	2,523
45 to 49 years	9,808	1,160	1,045	89	1,172	3,032	100	1,114	2,096
50 to 54 years	9,957	1,028	1,019	55	1,267	3,436	69	1,259	1,824
55 to 59 years	8,577	772	920	49	1,065	3,011	71	1,268	1,421
60 to 64 years	5,663	413	674	12	682	2,032	37	990	823
65 years and older	7,064	401	1,111	4	705	2,789	31	1,027	996
Total	155,963	22,989	17,808	13,405	11,017	35,313	941	10,366	44,124
Race									
American Indian or Alaska Native	302	49	29	30	36	54	2	27	75
Asian ³	422	64	42	19	20	79	1	67	130
Black or African American	14,904	2,272	981	1,706	987	2,627	110	1,859	4,362
Native Hawaiian or Other Pacific Islander	148	30	17	11	9	32	1	8	40
White	124,180	17,965	14,756	10,317	9,222	29,187	738	7,565	34,430
Other	12,912	1,969	1,654	1,229	549	2,488	74	638	4,311
Total	152,868	22,349	17,479	13,312	10,823	34,467	926	10,164	43,348

Continued. See notes at end of table.

Table 10.15a. Kentucky individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	148,555	21,660	17,184	12,863	10,528	33,934	721	9,598	42,067
Hispanic or Latino origin	3,427	710	356	445	163	671	10	146	926
Total	151,982	22,370	17,540	13,308	10,691	34,605	731	9,744	42,993
Specific Hispanic or Latino origin									
Mexican	1,377	288	125	224	43	269	2	46	380
Puerto Rican	504	80	47	61	42	111	2	23	138
Cuban	285	54	27	19	18	76	2	23	66
Other specific	1,236	287	154	141	56	211	4	48	335
Total	3,402	709	353	445	159	667	10	140	919
Veteran status⁴									
Veteran	1,147	169	108	16	151	287	8	158	250
Not a veteran	58,746	5,253	6,132	1,213	7,437	15,620	484	7,719	14,888
Total	59,893	5,422	6,240	1,229	7,588	15,907	492	7,877	15,138

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.15b. Kentucky individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	48.0	42.1	39.0	70.8	35.0	35.5	46.5	59.1	58.5	
Female	52.0	57.9	61.0	29.2	65.0	64.5	53.5	40.9	41.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	16.9	29.2	12.8	53.8	†	2.4	†	†	21.0	
12 to 14 years	8.1	10.4	8.2	20.1	†	5.8	†	†	8.8	
15 to 17 years	8.2	9.0	9.4	12.9	1.9	8.9	1.1	†	8.8	
18 to 20 years	4.6	4.1	5.5	3.9	3.5	5.6	6.1	2.2	4.9	
21 to 24 years	5.8	4.7	6.4	2.4	6.3	6.4	11.7	4.5	6.7	
25 to 29 years	8.4	7.2	8.8	2.1	10.0	8.4	14.2	8.5	10.4	
30 to 34 years	7.8	6.9	8.3	1.3	10.7	7.2	11.8	8.7	9.4	
35 to 39 years	7.5	6.6	7.4	1.2	11.4	7.6	12.5	10.3	8.1	
40 to 44 years	6.3	5.5	6.4	†	10.4	7.4	8.9	9.7	5.7	
45 to 49 years	6.3	5.0	5.9	†	10.6	8.6	10.6	10.7	4.8	
50 to 54 years	6.4	4.5	5.7	†	11.5	9.7	7.3	12.1	4.1	
55 to 59 years	5.5	3.4	5.2	†	9.7	8.5	7.5	12.2	3.2	
60 to 64 years	3.6	1.8	3.8	†	6.2	5.8	3.9	9.6	1.9	
65 years and older	4.5	1.7	6.2	†	6.4	7.9	3.3	9.9	2.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†	
Asian ³	†	†	†	†	†	†	†	†	†	
Black or African American	9.7	10.2	5.6	12.8	9.1	7.6	11.9	18.3	10.1	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	81.2	80.4	84.4	77.5	85.2	84.7	79.7	74.4	79.4	
Other	8.4	8.8	9.5	9.2	5.1	7.2	8.0	6.3	9.9	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.15b. Kentucky individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	97.7	96.8	98.0	96.7	98.5	98.1	98.6	98.5	97.8
Hispanic or Latino origin	2.3	3.2	2.0	3.3	1.5	1.9	1.4	1.5	2.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	40.5	40.6	35.4	50.3	27.0	40.3	20.0	32.9	41.3
Puerto Rican	14.8	11.3	13.3	13.7	26.4	16.6	20.0	16.4	15.0
Cuban	8.4	7.6	7.6	4.3	11.3	11.4	20.0	16.4	7.2
Other specific	36.3	40.5	43.6	31.7	35.2	31.6	40.0	34.3	36.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	1.9	3.1	1.7	1.3	2.0	1.8	1.6	2.0	1.7
Not a veteran	98.1	96.9	98.3	98.7	98.0	98.2	98.4	98.0	98.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

† Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes substance use diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.16a. Louisiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	23,000	1,127	1,729	1,993	3,559	8,596	78	4,631	1,287
Gender									
Male	10,421	514	609	1,427	1,205	2,920	28	2,889	829
Female	12,508	613	1,118	561	2,349	5,650	50	1,715	452
Total	22,929	1,127	1,727	1,988	3,554	8,570	78	4,604	1,281
Age²									
0 to 11 years	1,770	233	119	1,008	8	75	1	3	323
12 to 14 years	1,114	135	84	490	13	199	--	5	188
15 to 17 years	1,098	90	119	294	28	360	2	16	189
18 to 20 years	850	52	88	102	97	376	1	57	77
21 to 24 years	1,209	59	135	19	225	484	8	210	69
25 to 29 years	2,007	95	191	19	364	809	12	426	91
30 to 34 years	2,128	101	219	13	443	764	11	512	65
35 to 39 years	2,119	90	192	14	449	785	9	522	58
40 to 44 years	1,796	53	128	9	411	706	7	436	46
45 to 49 years	1,907	65	123	4	404	848	6	410	47
50 to 54 years	2,289	59	116	6	424	1,047	9	588	40
55 to 59 years	2,199	55	103	3	329	1,025	6	640	38
60 to 64 years	1,479	25	67	1	206	679	5	459	37
65 years and older	1,035	15	45	11	158	439	1	347	19
Total	23,000	1,127	1,729	1,993	3,559	8,596	78	4,631	1,287
Race									
American Indian or Alaska Native	276	19	38	19	50	118	1	21	10
Asian ³	70	7	5	1	3	30	--	20	4
Black or African American	9,351	397	481	1,170	969	2,826	23	2,844	641
Native Hawaiian or Other Pacific Islander	20	--	2	--	4	8	--	2	4
White	10,569	557	1,012	640	2,097	4,477	44	1,270	472
Other	1,435	65	110	61	214	673	6	243	63
Total	21,721	1,045	1,648	1,891	3,337	8,132	74	4,400	1,194

Continued. See notes at end of table.

Table 10.16a. Louisiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	21,410	1,050	1,630	1,881	3,297	8,026	72	4,255	1,199
Hispanic or Latino origin	567	49	56	49	83	214	1	91	24
Total	21,977	1,099	1,686	1,930	3,380	8,240	73	4,346	1,223
Specific Hispanic or Latino origin									
Mexican	17	2	3	2	4	5	--	--	1
Puerto Rican	9	--	2	--	2	2	--	3	--
Cuban	8	--	--	--	1	2	--	5	--
Other specific	23	--	3	1	--	10	--	9	--
Total	57	2	8	3	7	19	--	17	1
Veteran status⁴									
Veteran	226	9	16	3	41	80	1	76	--
Not a veteran	16,955	603	1,273	163	3,086	7,212	65	4,052	501
Total	17,181	612	1,289	166	3,127	7,292	66	4,128	501

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it excludes substance use diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.16b. Louisiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	45.4	45.6	35.3	71.8	33.9	34.1	35.9	62.7	64.7	
Female	54.6	54.4	64.7	28.2	66.1	65.9	64.1	37.3	35.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	7.7	20.7	6.9	50.6	†	†	1.3	†	25.1	
12 to 14 years	4.8	12.0	4.9	24.6	†	2.3	--	†	14.6	
15 to 17 years	4.8	8.0	6.9	14.8	†	4.2	2.6	†	14.7	
18 to 20 years	3.7	4.6	5.1	5.1	2.7	4.4	1.3	1.2	6.0	
21 to 24 years	5.3	5.2	7.8	†	6.3	5.6	10.3	4.5	5.4	
25 to 29 years	8.7	8.4	11.0	†	10.2	9.4	15.4	9.2	7.1	
30 to 34 years	9.3	9.0	12.7	†	12.4	8.9	14.1	11.1	5.1	
35 to 39 years	9.2	8.0	11.1	†	12.6	9.1	11.5	11.3	4.5	
40 to 44 years	7.8	4.7	7.4	†	11.5	8.2	9.0	9.4	3.6	
45 to 49 years	8.3	5.8	7.1	†	11.4	9.9	7.7	8.9	3.7	
50 to 54 years	10.0	5.2	6.7	†	11.9	12.2	11.5	12.7	3.1	
55 to 59 years	9.6	4.9	6.0	†	9.2	11.9	7.7	13.8	3.0	
60 to 64 years	6.4	2.2	3.9	†	5.8	7.9	6.4	9.9	2.9	
65 years and older	4.5	1.3	2.6	†	4.4	5.1	1.3	7.5	1.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	1.3	1.8	2.3	1.0	1.5	1.5	1.4	†	†	
Asian ³	†	†	†	†	†	†	--	†	†	
Black or African American	43.1	38.0	29.2	61.9	29.0	34.8	31.1	64.6	53.7	
Native Hawaiian or Other Pacific Islander	†	--	†	--	†	†	--	†	†	
White	48.7	53.3	61.4	33.8	62.8	55.1	59.5	28.9	39.5	
Other	6.6	6.2	6.7	3.2	6.4	8.3	8.1	5.5	5.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.16b. Louisiana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	97.4	95.5	96.7	97.5	97.5	97.4	98.6	97.9	98.0
Hispanic or Latino origin	2.6	4.5	3.3	2.5	2.5	2.6	1.4	2.1	2.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	29.8	100.0	37.5	66.7	57.1	26.3	--	--	100.0
Puerto Rican	15.8	--	25.0	--	28.6	10.5	--	17.6	--
Cuban	14.0	--	--	--	14.3	10.5	--	29.4	--
Other specific	40.4	--	37.5	33.3	--	52.6	--	52.9	--
Total	100.0	100.0	100.0	100.0	100.0	100.0	--	100.0	100.0
Veteran status⁴									
Veteran	1.3	1.5	1.2	1.8	1.3	1.1	1.5	1.8	--
Not a veteran	98.7	98.5	98.8	98.2	98.7	98.9	98.5	98.2	100.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.17a. Maryland individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	211,849	33,567	29,107	28,739	26,331	56,339	498	16,323	20,945
Gender									
Male	97,400	14,699	10,517	19,296	10,264	20,473	153	10,186	11,812
Female	114,432	18,867	18,589	9,442	16,066	35,864	345	6,136	9,123
Total	211,832	33,566	29,106	28,738	26,330	56,337	498	16,322	20,935
Age²									
0 to 11 years	37,472	11,578	3,714	13,867	119	1,321	10	22	6,841
12 to 14 years	17,782	4,303	2,160	5,690	249	2,375	9	33	2,963
15 to 17 years	16,540	3,122	2,345	3,734	660	3,812	10	109	2,748
18 to 20 years	10,455	1,376	1,791	1,317	1,284	3,210	39	362	1,076
21 to 24 years	11,943	1,433	2,086	703	2,030	3,813	57	966	855
25 to 29 years	20,774	2,336	3,638	1,053	3,776	6,741	66	1,841	1,323
30 to 34 years	19,043	2,127	3,209	820	3,675	6,285	79	1,744	1,104
35 to 39 years	15,737	1,735	2,591	571	3,112	5,178	56	1,599	895
40 to 44 years	11,913	1,254	1,865	327	2,362	4,155	39	1,278	633
45 to 49 years	12,918	1,295	1,711	257	2,645	4,798	30	1,491	691
50 to 54 years	14,298	1,297	1,650	187	2,819	5,638	45	1,964	698
55 to 59 years	12,272	937	1,358	131	2,084	5,011	32	2,140	579
60 to 64 years	7,563	615	837	72	1,082	2,971	17	1,585	384
65 years and older	3,135	159	151	10	434	1,030	8	1,189	154
Total	211,845	33,567	29,106	28,739	26,331	56,338	497	16,323	20,944
Race									
American Indian or Alaska Native	936	195	139	103	105	244	3	67	80
Asian ³	2,837	470	463	220	233	737	6	424	284
Black or African American	97,442	14,825	8,741	14,413	11,546	26,808	148	9,879	11,082
Native Hawaiian or Other Pacific Islander	580	118	98	86	37	140	--	29	72
White	99,617	15,571	17,864	12,281	13,758	26,252	320	5,500	8,071
Other	4,458	1,041	596	1,083	315	761	5	142	515
Total	205,870	32,220	27,901	28,186	25,994	54,942	482	16,041	20,104

Continued. See notes at end of table.

Table 10.17a. Maryland individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	179,709	26,252	23,889	25,013	23,900	49,272	420	13,623	17,340
Hispanic or Latino origin	22,534	5,124	2,980	3,045	1,991	4,974	56	2,107	2,257
Total	202,243	31,376	26,869	28,058	25,891	54,246	476	15,730	19,597
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	4,047	415	411	104	797	1,383	17	717	203
Not a veteran	128,372	12,960	18,801	5,128	23,987	45,677	416	14,009	7,394
Total	132,419	13,375	19,212	5,232	24,784	47,060	433	14,726	7,597

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.17b. Maryland individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	46.0	43.8	36.1	67.1	39.0	36.3	30.7	62.4	56.4	
Female	54.0	56.2	63.9	32.9	61.0	63.7	69.3	37.6	43.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	17.7	34.5	12.8	48.3	†	2.3	2.0	†	32.7	
12 to 14 years	8.4	12.8	7.4	19.8	†	4.2	1.8	†	14.1	
15 to 17 years	7.8	9.3	8.1	13.0	2.5	6.8	2.0	†	13.1	
18 to 20 years	4.9	4.1	6.2	4.6	4.9	5.7	7.8	2.2	5.1	
21 to 24 years	5.6	4.3	7.2	2.4	7.7	6.8	11.5	5.9	4.1	
25 to 29 years	9.8	7.0	12.5	3.7	14.3	12.0	13.3	11.3	6.3	
30 to 34 years	9.0	6.3	11.0	2.9	14.0	11.2	15.9	10.7	5.3	
35 to 39 years	7.4	5.2	8.9	2.0	11.8	9.2	11.3	9.8	4.3	
40 to 44 years	5.6	3.7	6.4	1.1	9.0	7.4	7.8	7.8	3.0	
45 to 49 years	6.1	3.9	5.9	†	10.0	8.5	6.0	9.1	3.3	
50 to 54 years	6.7	3.9	5.7	†	10.7	10.0	9.1	12.0	3.3	
55 to 59 years	5.8	2.8	4.7	†	7.9	8.9	6.4	13.1	2.8	
60 to 64 years	3.6	1.8	2.9	†	4.1	5.3	3.4	9.7	1.8	
65 years and older	1.5	†	†	†	1.6	1.8	1.6	7.3	†	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†	
Asian ³	1.4	1.5	1.7	†	†	1.3	1.2	2.6	1.4	
Black or African American	47.3	46.0	31.3	51.1	44.4	48.8	30.7	61.6	55.1	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†	
White	48.4	48.3	64.0	43.6	52.9	47.8	66.4	34.3	40.1	
Other	2.2	3.2	2.1	3.8	1.2	1.4	1.0	†	2.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.17b. Maryland individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	88.9	83.7	88.9	89.1	92.3	90.8	88.2	86.6	88.5
Hispanic or Latino origin	11.1	16.3	11.1	10.9	7.7	9.2	11.8	13.4	11.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	3.1	3.1	2.1	2.0	3.2	2.9	3.9	4.9	2.7
Not a veteran	96.9	96.9	97.9	98.0	96.8	97.1	96.1	95.1	97.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.18a. Massachusetts individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	9,248	712	827	318	1,656	1,861	31	3,322	521
Gender									
Male	4,965	298	351	230	799	755	12	2,258	262
Female	4,283	414	476	88	857	1,106	19	1,064	259
Total	9,248	712	827	318	1,656	1,861	31	3,322	521
Age²									
0 to 11 years	651	132	199	167	9	45	--	2	97
12 to 14 years	439	73	82	85	26	113	--	9	51
15 to 17 years	556	62	66	30	55	221	3	46	73
18 to 20 years	688	67	84	16	121	193	2	147	58
21 to 24 years	735	48	58	5	152	124	3	308	37
25 to 29 years	864	56	54	4	197	120	5	392	36
30 to 34 years	773	44	43	4	173	112	2	358	37
35 to 39 years	697	47	38	3	155	118	3	306	27
40 to 44 years	599	36	36	1	127	106	4	273	16
45 to 49 years	661	36	39	2	144	140	5	275	20
50 to 54 years	785	39	61	1	160	198	--	308	18
55 to 59 years	811	40	33	--	163	172	2	380	21
60 to 64 years	550	24	19	--	101	125	2	261	18
65 years and older	439	8	15	--	73	74	--	257	12
Total	9,248	712	827	318	1,656	1,861	31	3,322	521
Race									
American Indian or Alaska Native	17	1	1	1	5	1	--	8	--
Asian ³	182	9	11	1	19	35	--	92	15
Black or African American	1,282	99	74	54	137	107	3	751	57
Native Hawaiian or Other Pacific Islander	4	--	--	--	--	--	--	3	1
White	5,279	360	371	106	1,101	1,050	22	1,979	290
Other	175	21	23	11	20	29	1	49	21
Total	6,939	490	480	173	1,282	1,222	26	2,882	384

Continued. See notes at end of table.

Table 10.18a. Massachusetts individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	5,308	329	351	110	1,041	893	21	2,256	307
Hispanic or Latino origin	809	97	60	29	108	165	4	312	34
Total	6,117	426	411	139	1,149	1,058	25	2,568	341
Specific Hispanic or Latino origin									
Mexican	7	1	4	--	--	1	--	1	--
Puerto Rican	288	37	17	5	33	57	2	123	14
Cuban	--	--	--	--	--	--	--	--	--
Other specific	170	21	17	11	18	35	1	61	6
Total	465	59	38	16	51	93	3	185	20
Veteran status⁴									
Veteran	159	4	2	--	37	18	--	95	3
Not a veteran	6,423	359	366	27	1,321	1,141	24	2,932	253
Total	6,582	363	368	27	1,358	1,159	24	3,027	256

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.18b. Massachusetts individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	53.7	41.9	42.4	72.3	48.2	40.6	38.7	68.0	50.3	
Female	46.3	58.1	57.6	27.7	51.8	59.4	61.3	32.0	49.7	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	7.0	18.5	24.1	52.5	†	2.4	--	†	18.6	
12 to 14 years	4.7	10.3	9.9	26.7	1.6	6.1	--	†	9.8	
15 to 17 years	6.0	8.7	8.0	9.4	3.3	11.9	9.7	1.4	14.0	
18 to 20 years	7.4	9.4	10.2	5.0	7.3	10.4	6.5	4.4	11.1	
21 to 24 years	7.9	6.7	7.0	1.6	9.2	6.7	9.7	9.3	7.1	
25 to 29 years	9.3	7.9	6.5	1.3	11.9	6.4	16.1	11.8	6.9	
30 to 34 years	8.4	6.2	5.2	1.3	10.4	6.0	6.5	10.8	7.1	
35 to 39 years	7.5	6.6	4.6	†	9.4	6.3	9.7	9.2	5.2	
40 to 44 years	6.5	5.1	4.4	†	7.7	5.7	12.9	8.2	3.1	
45 to 49 years	7.1	5.1	4.7	†	8.7	7.5	16.1	8.3	3.8	
50 to 54 years	8.5	5.5	7.4	†	9.7	10.6	--	9.3	3.5	
55 to 59 years	8.8	5.6	4.0	--	9.8	9.2	6.5	11.4	4.0	
60 to 64 years	5.9	3.4	2.3	--	6.1	6.7	6.5	7.9	3.5	
65 years and older	4.7	1.1	1.8	--	4.4	4.0	--	7.7	2.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	--	†	--	
Asian ³	2.6	1.8	2.3	†	1.5	2.9	--	3.2	3.9	
Black or African American	18.5	20.2	15.4	31.2	10.7	8.8	11.5	26.1	14.8	
Native Hawaiian or Other Pacific Islander	†	--	--	--	--	--	--	†	†	
White	76.1	73.5	77.3	61.3	85.9	85.9	84.6	68.7	75.5	
Other	2.5	4.3	4.8	6.4	1.6	2.4	3.8	1.7	5.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.18b. Massachusetts individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

<i>Selected characteristics</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Ethnicity									
Not of Hispanic or Latino origin	86.8	77.2	85.4	79.1	90.6	84.4	84.0	87.9	90.0
Hispanic or Latino origin	13.2	22.8	14.6	20.9	9.4	15.6	16.0	12.1	10.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	1.5	1.7	10.5	--	--	1.1	--	†	--
Puerto Rican	61.9	62.7	44.7	31.3	64.7	61.3	66.7	66.5	70.0
Cuban	--	--	--	--	--	--	--	--	--
Other specific	36.6	35.6	44.7	68.8	35.3	37.6	33.3	33.0	30.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	2.4	1.1	†	--	2.7	1.6	--	3.1	1.2
Not a veteran	97.6	98.9	99.5	100.0	97.3	98.4	100.0	96.9	98.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.19a. Michigan individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	224,567	18,469	13,515	12,034	28,423	53,417	1,302	30,743	66,664
Gender									
Male	115,470	8,379	5,633	8,795	11,779	20,762	422	18,953	40,747
Female	109,096	10,090	7,882	3,239	16,644	32,655	880	11,790	25,916
Total	224,566	18,469	13,515	12,034	28,423	53,417	1,302	30,743	66,663
Age²									
0 to 11 years	29,110	5,691	1,494	5,756	112	792	2	53	15,210
12 to 14 years	12,793	1,778	813	2,441	261	2,179	4	59	5,258
15 to 17 years	14,236	1,681	1,027	1,544	572	3,846	16	184	5,366
18 to 20 years	10,277	829	736	611	996	2,773	96	529	3,707
21 to 24 years	14,132	962	912	419	1,847	3,744	158	1,407	4,683
25 to 29 years	20,707	1,391	1,474	443	3,281	5,255	194	2,557	6,112
30 to 34 years	18,550	1,229	1,390	314	3,291	4,833	212	2,679	4,602
35 to 39 years	17,144	1,106	1,203	166	3,318	4,632	151	2,761	3,807
40 to 44 years	15,092	829	884	92	2,998	4,414	128	2,763	2,984
45 to 49 years	16,797	873	1,016	90	3,192	5,153	102	3,181	3,190
50 to 54 years	18,041	856	911	67	3,271	5,491	94	3,961	3,390
55 to 59 years	16,713	630	786	41	2,604	4,988	80	4,444	3,140
60 to 64 years	10,953	325	461	29	1,542	3,076	39	3,150	2,331
65 years and older	10,021	288	408	21	1,138	2,241	26	3,015	2,884
Total	224,566	18,468	13,515	12,034	28,423	53,417	1,302	30,743	66,664
Race									
American Indian or Alaska Native	1,370	169	100	54	161	352	11	200	323
Asian ³	940	44	48	21	78	156	2	179	412
Black or African American	45,198	2,981	1,652	2,586	4,772	10,604	117	10,209	12,277
Native Hawaiian or Other Pacific Islander	136	13	7	5	14	44	--	23	30
White	132,606	11,419	9,234	6,431	18,076	31,546	1,020	14,127	40,753
Other	42,752	3,610	2,356	2,895	5,194	10,242	138	5,896	12,421
Total	223,002	18,236	13,397	11,992	28,295	52,944	1,288	30,634	66,216

Continued. See notes at end of table.

Table 10.19a. Michigan individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	192,659	15,174	11,571	10,329	25,297	46,625	1,145	27,712	54,806
Hispanic or Latino origin	13,083	1,655	895	927	1,253	3,277	52	1,085	3,939
Total	205,742	16,829	12,466	11,256	26,550	49,902	1,197	28,797	58,745
Specific Hispanic or Latino origin									
Mexican	3,252	544	262	246	209	826	10	171	984
Puerto Rican	966	146	66	70	88	261	5	73	257
Cuban	142	18	12	7	12	31	1	19	42
Other specific	1,565	190	87	104	158	405	5	133	483
Total	5,925	898	427	427	467	1,523	21	396	1,766
Veteran status⁴									
Veteran	3,731	333	208	21	678	1,098	18	922	453
Not a veteran	155,952	8,256	9,617	2,222	26,188	43,707	1,176	28,838	35,948
Total	159,683	8,589	9,825	2,243	26,866	44,805	1,194	29,760	36,401

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.19b. Michigan individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	51.4	45.4	41.7	73.1	41.4	38.9	32.4	61.6	61.1
Female	48.6	54.6	58.3	26.9	58.6	61.1	67.6	38.4	38.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	13.0	30.8	11.1	47.8	†	1.5	†	†	22.8
12 to 14 years	5.7	9.6	6.0	20.3	†	4.1	†	†	7.9
15 to 17 years	6.3	9.1	7.6	12.8	2.0	7.2	1.2	†	8.0
18 to 20 years	4.6	4.5	5.4	5.1	3.5	5.2	7.4	1.7	5.6
21 to 24 years	6.3	5.2	6.7	3.5	6.5	7.0	12.1	4.6	7.0
25 to 29 years	9.2	7.5	10.9	3.7	11.5	9.8	14.9	8.3	9.2
30 to 34 years	8.3	6.7	10.3	2.6	11.6	9.0	16.3	8.7	6.9
35 to 39 years	7.6	6.0	8.9	1.4	11.7	8.7	11.6	9.0	5.7
40 to 44 years	6.7	4.5	6.5	†	10.5	8.3	9.8	9.0	4.5
45 to 49 years	7.5	4.7	7.5	†	11.2	9.6	7.8	10.3	4.8
50 to 54 years	8.0	4.6	6.7	†	11.5	10.3	7.2	12.9	5.1
55 to 59 years	7.4	3.4	5.8	†	9.2	9.3	6.1	14.5	4.7
60 to 64 years	4.9	1.8	3.4	†	5.4	5.8	3.0	10.2	3.5
65 years and older	4.5	1.6	3.0	†	4.0	4.2	2.0	9.8	4.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†
Asian ³	†	†	†	†	†	†	†	†	†
Black or African American	20.3	16.3	12.3	21.6	16.9	20.0	9.1	33.3	18.5
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†
White	59.5	62.6	68.9	53.6	63.9	59.6	79.2	46.1	61.5
Other	19.2	19.8	17.6	24.1	18.4	19.3	10.7	19.2	18.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.19b. Michigan individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	93.6	90.2	92.8	91.8	95.3	93.4	95.7	96.2	93.3
Hispanic or Latino origin	6.4	9.8	7.2	8.2	4.7	6.6	4.3	3.8	6.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	54.9	60.6	61.4	57.6	44.8	54.2	47.6	43.2	55.7
Puerto Rican	16.3	16.3	15.5	16.4	18.8	17.1	23.8	18.4	14.6
Cuban	2.4	2.0	2.8	1.6	2.6	2.0	4.8	4.8	2.4
Other specific	26.4	21.2	20.4	24.4	33.8	26.6	23.8	33.6	27.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	2.3	3.9	2.1	†	2.5	2.5	1.5	3.1	1.2
Not a veteran	97.7	96.1	97.9	99.1	97.5	97.5	98.5	96.9	98.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.20a. Minnesota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	269,422	59,657	61,843	23,686	15,009	58,737	4,578	17,509	28,403
Gender									
Male	119,452	23,701	22,972	15,561	6,405	21,685	1,401	10,944	16,783
Female	149,970	35,956	38,871	8,125	8,604	37,052	3,177	6,565	11,620
Total	269,422	59,657	61,843	23,686	15,009	58,737	4,578	17,509	28,403
Age²									
0 to 11 years	38,736	14,652	6,897	8,409	67	997	4	11	7,699
12 to 14 years	17,478	4,862	3,518	3,932	128	1,966	7	24	3,041
15 to 17 years	17,841	4,285	3,955	2,866	250	3,311	17	80	3,077
18 to 20 years	11,825	2,227	3,052	1,473	405	2,581	151	288	1,648
21 to 24 years	15,053	2,781	4,303	1,226	830	3,224	424	842	1,423
25 to 29 years	25,493	4,783	7,463	1,727	1,661	5,487	762	1,550	2,060
30 to 34 years	24,423	4,969	6,812	1,261	1,830	5,327	734	1,822	1,668
35 to 39 years	21,883	4,524	5,679	885	1,831	5,084	608	1,831	1,441
40 to 44 years	17,280	3,475	4,344	555	1,522	4,394	445	1,481	1,064
45 to 49 years	17,441	3,329	3,890	468	1,553	5,095	402	1,623	1,081
50 to 54 years	18,352	3,125	3,781	339	1,607	5,997	334	1,991	1,178
55 to 59 years	17,548	2,574	3,479	302	1,473	5,895	312	2,288	1,225
60 to 64 years	12,256	1,766	2,300	181	948	4,318	209	1,669	865
65 years and older	13,794	2,300	2,363	61	903	5,056	169	2,009	933
Total	269,403	59,652	61,836	23,685	15,008	58,732	4,578	17,509	28,403
Race									
American Indian or Alaska Native	10,098	2,687	2,042	746	504	2,446	131	502	1,040
Asian ³	7,513	1,554	1,389	226	140	2,942	46	613	603
Black or African American	32,335	8,489	5,401	2,184	1,717	7,469	403	3,568	3,104
Native Hawaiian or Other Pacific Islander	266	73	51	17	19	53	10	23	20
White	163,948	31,267	39,443	14,142	10,820	36,176	3,335	11,651	17,114
Other	12,002	3,110	2,701	1,524	577	2,056	178	397	1,459
Total	226,162	47,180	51,027	18,839	13,777	51,142	4,103	16,754	23,340

Continued. See notes at end of table.

Table 10.20a. Minnesota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	237,008	50,676	54,015	20,699	13,818	52,512	4,048	16,261	24,979
Hispanic or Latino origin	12,716	3,118	2,486	1,027	707	2,806	282	977	1,313
Total	249,724	53,794	56,501	21,726	14,525	55,318	4,330	17,238	26,292
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.20b. Minnesota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	44.3	39.7	37.1	65.7	42.7	36.9	30.6	62.5	59.1	
Female	55.7	60.3	62.9	34.3	57.3	63.1	69.4	37.5	40.9	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	14.4	24.6	11.2	35.5	†	1.7	†	†	27.1	
12 to 14 years	6.5	8.2	5.7	16.6	†	3.3	†	†	10.7	
15 to 17 years	6.6	7.2	6.4	12.1	1.7	5.6	†	†	10.8	
18 to 20 years	4.4	3.7	4.9	6.2	2.7	4.4	3.3	1.6	5.8	
21 to 24 years	5.6	4.7	7.0	5.2	5.5	5.5	9.3	4.8	5.0	
25 to 29 years	9.5	8.0	12.1	7.3	11.1	9.3	16.6	8.9	7.3	
30 to 34 years	9.1	8.3	11.0	5.3	12.2	9.1	16.0	10.4	5.9	
35 to 39 years	8.1	7.6	9.2	3.7	12.2	8.7	13.3	10.5	5.1	
40 to 44 years	6.4	5.8	7.0	2.3	10.1	7.5	9.7	8.5	3.7	
45 to 49 years	6.5	5.6	6.3	2.0	10.3	8.7	8.8	9.3	3.8	
50 to 54 years	6.8	5.2	6.1	1.4	10.7	10.2	7.3	11.4	4.1	
55 to 59 years	6.5	4.3	5.6	1.3	9.8	10.0	6.8	13.1	4.3	
60 to 64 years	4.5	3.0	3.7	†	6.3	7.4	4.6	9.5	3.0	
65 years and older	5.1	3.9	3.8	†	6.0	8.6	3.7	11.5	3.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	4.5	5.7	4.0	4.0	3.7	4.8	3.2	3.0	4.5	
Asian ³	3.3	3.3	2.7	1.2	1.0	5.8	1.1	3.7	2.6	
Black or African American	14.3	18.0	10.6	11.6	12.5	14.6	9.8	21.3	13.3	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	72.5	66.3	77.3	75.1	78.5	70.7	81.3	69.5	73.3	
Other	5.3	6.6	5.3	8.1	4.2	4.0	4.3	2.4	6.3	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.20b. Minnesota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	94.9	94.2	95.6	95.3	95.1	94.9	93.5	94.3	95.0
Hispanic or Latino origin	5.1	5.8	4.4	4.7	4.9	5.1	6.5	5.7	5.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.21a. Mississippi individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	84,918	7,577	6,237	13,776	8,855	20,203	300	10,365	17,605
Gender									
Male	41,240	3,177	2,109	9,403	3,258	5,922	138	6,077	11,156
Female	43,620	4,397	4,103	4,370	5,596	14,274	162	4,286	6,432
Total	84,860	7,574	6,212	13,773	8,854	20,196	300	10,363	17,588
Age²									
0 to 11 years	15,444	2,633	353	7,517	461	310	4	23	4,143
12 to 14 years	7,544	1,134	266	3,046	461	791	8	34	1,804
15 to 17 years	7,033	988	360	1,861	573	1,218	15	72	1,946
18 to 20 years	3,969	355	312	691	431	921	23	225	1,011
21 to 24 years	3,979	220	383	207	530	1,067	30	486	1,056
25 to 29 years	6,207	327	628	150	805	1,687	42	909	1,659
30 to 34 years	6,182	323	676	98	837	1,633	42	1,033	1,540
35 to 39 years	6,199	350	596	62	845	1,861	39	1,205	1,241
40 to 44 years	5,070	268	477	54	772	1,628	25	954	892
45 to 49 years	5,108	262	484	34	816	1,816	15	986	695
50 to 54 years	5,364	272	478	23	748	2,014	19	1,212	598
55 to 59 years	5,278	207	463	16	693	2,148	10	1,261	480
60 to 64 years	3,714	123	330	14	454	1,530	13	982	268
65 years and older	3,827	115	431	3	429	1,579	15	983	272
Total	84,918	7,577	6,237	13,776	8,855	20,203	300	10,365	17,605
Race									
American Indian or Alaska Native	212	25	17	26	22	43	--	19	60
Asian ³	144	8	17	13	21	33	--	31	21
Black or African American	42,373	2,911	1,831	8,472	2,985	9,672	125	7,361	9,016
Native Hawaiian or Other Pacific Islander	12	2	--	2	1	2	--	2	3
White	40,012	4,243	4,205	4,791	5,626	10,095	164	2,861	8,027
Other	1,212	210	81	309	115	207	3	49	238
Total	83,965	7,399	6,151	13,613	8,770	20,052	292	10,323	17,365

Continued. See notes at end of table.

Table 10.21a. Mississippi individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	79,011	6,724	5,767	12,932	8,498	19,004	286	10,075	15,725
Hispanic or Latino origin	1,332	220	103	232	93	258	3	70	353
Total	80,343	6,944	5,870	13,164	8,591	19,262	289	10,145	16,078
Specific Hispanic or Latino origin									
Mexican	537	113	44	105	38	116	3	21	97
Puerto Rican	81	11	7	14	10	14	--	9	16
Cuban	28	3	3	4	2	3	--	4	9
Other specific	--	--	--	--	--	--	--	--	--
Total	646	127	54	123	50	133	3	34	122
Veteran status⁴									
Veteran	1,155	123	102	19	135	321	5	220	230
Not a veteran	52,049	2,638	5,030	1,316	7,062	17,230	254	9,414	9,105
Total	53,204	2,761	5,132	1,335	7,197	17,551	259	9,634	9,335

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.21b. Mississippi individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	48.6	41.9	34.0	68.3	36.8	29.3	46.0	58.6	63.4	
Female	51.4	58.1	66.0	31.7	63.2	70.7	54.0	41.4	36.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	18.2	34.7	5.7	54.6	5.2	1.5	1.3	†	23.5	
12 to 14 years	8.9	15.0	4.3	22.1	5.2	3.9	2.7	†	10.2	
15 to 17 years	8.3	13.0	5.8	13.5	6.5	6.0	5.0	†	11.1	
18 to 20 years	4.7	4.7	5.0	5.0	4.9	4.6	7.7	2.2	5.7	
21 to 24 years	4.7	2.9	6.1	1.5	6.0	5.3	10.0	4.7	6.0	
25 to 29 years	7.3	4.3	10.1	1.1	9.1	8.4	14.0	8.8	9.4	
30 to 34 years	7.3	4.3	10.8	†	9.5	8.1	14.0	10.0	8.7	
35 to 39 years	7.3	4.6	9.6	†	9.5	9.2	13.0	11.6	7.0	
40 to 44 years	6.0	3.5	7.6	†	8.7	8.1	8.3	9.2	5.1	
45 to 49 years	6.0	3.5	7.8	†	9.2	9.0	5.0	9.5	3.9	
50 to 54 years	6.3	3.6	7.7	†	8.4	10.0	6.3	11.7	3.4	
55 to 59 years	6.2	2.7	7.4	†	7.8	10.6	3.3	12.2	2.7	
60 to 64 years	4.4	1.6	5.3	†	5.1	7.6	4.3	9.5	1.5	
65 years and older	4.5	1.5	6.9	†	4.8	7.8	5.0	9.5	1.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	--	†	†	
Asian ³	†	†	†	†	†	†	--	†	†	
Black or African American	50.5	39.3	29.8	62.2	34.0	48.2	42.8	71.3	51.9	
Native Hawaiian or Other Pacific Islander	†	†	--	†	†	†	--	†	†	
White	47.7	57.3	68.4	35.2	64.2	50.3	56.2	27.7	46.2	
Other	1.4	2.8	1.3	2.3	1.3	1.0	1.0	†	1.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.21b. Mississippi individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	98.3	96.8	98.2	98.2	98.9	98.7	99.0	99.3	97.8
Hispanic or Latino origin	1.7	3.2	1.8	1.8	1.1	1.3	1.0	†	2.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	83.1	89.0	81.5	85.4	76.0	87.2	100.0	61.8	79.5
Puerto Rican	12.5	8.7	13.0	11.4	20.0	10.5	--	26.5	13.1
Cuban	4.3	2.4	5.6	3.3	4.0	2.3	--	11.8	7.4
Other specific	--	--	--	--	--	--	--	--	--
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	2.2	4.5	2.0	1.4	1.9	1.8	1.9	2.3	2.5
Not a veteran	97.8	95.5	98.0	98.6	98.1	98.2	98.1	97.7	97.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.22a. Missouri individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	73,128	3,028	9,068	6,764	12,909	22,356	797	13,187	5,019
Gender									
Male	34,713	1,043	3,589	5,058	4,791	8,169	286	8,546	3,231
Female	38,415	1,985	5,479	1,706	8,118	14,187	511	4,641	1,788
Total	73,128	3,028	9,068	6,764	12,909	22,356	797	13,187	5,019
Age²									
0 to 11 years	7,805	429	958	3,661	125	454	1	16	2,161
12 to 14 years	4,670	271	577	1,575	196	984	4	34	1,029
15 to 17 years	4,156	236	515	903	340	1,331	24	89	718
18 to 20 years	2,524	127	402	255	453	871	42	228	146
21 to 24 years	3,164	147	472	98	692	1,009	67	566	113
25 to 29 years	5,259	256	736	93	1,173	1,596	92	1,166	147
30 to 34 years	5,786	300	896	55	1,377	1,633	92	1,301	132
35 to 39 years	6,190	279	817	45	1,480	1,901	116	1,445	107
40 to 44 years	5,598	254	720	19	1,351	1,792	66	1,297	99
45 to 49 years	6,389	223	785	18	1,460	2,289	82	1,445	87
50 to 54 years	7,191	217	821	20	1,493	2,779	76	1,686	99
55 to 59 years	6,995	166	738	8	1,408	2,757	61	1,772	85
60 to 64 years	4,507	84	403	11	842	1,818	40	1,254	55
65 years and older	2,894	39	228	3	519	1,142	34	888	41
Total	73,128	3,028	9,068	6,764	12,909	22,356	797	13,187	5,019
Race									
American Indian or Alaska Native	165	13	23	10	26	54	2	20	17
Asian ³	214	4	24	5	30	59	4	81	7
Black or African American	15,567	380	1,027	1,875	2,082	3,979	131	5,129	964
Native Hawaiian or Other Pacific Islander	32	2	7	4	4	10	--	2	3
White	52,703	2,371	7,422	4,246	10,153	17,101	613	7,270	3,527
Other	2,761	148	350	362	451	660	31	466	293
Total	71,442	2,918	8,853	6,502	12,746	21,863	781	12,968	4,811

Continued. See notes at end of table.

Table 10.22a. Missouri individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	71,431	2,923	8,839	6,572	12,702	21,829	776	12,958	4,832
Hispanic or Latino origin	1,695	105	228	192	207	527	21	229	186
Total	73,126	3,028	9,067	6,764	12,909	22,356	797	13,187	5,018
Specific Hispanic or Latino origin									
Mexican	890	60	126	107	105	284	15	107	86
Puerto Rican	169	4	23	11	32	59	3	26	11
Cuban	49	5	5	2	6	14	1	13	3
Other specific	571	36	73	72	61	169	1	77	82
Total	1,679	105	227	192	204	526	20	223	182
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.22b. Missouri individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	47.5	34.4	39.6	74.8	37.1	36.5	35.9	64.8	64.4	
Female	52.5	65.6	60.4	25.2	62.9	63.5	64.1	35.2	35.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	10.7	14.2	10.6	54.1	†	2.0	†	†	43.1	
12 to 14 years	6.4	8.9	6.4	23.3	1.5	4.4	†	†	20.5	
15 to 17 years	5.7	7.8	5.7	13.4	2.6	6.0	3.0	†	14.3	
18 to 20 years	3.5	4.2	4.4	3.8	3.5	3.9	5.3	1.7	2.9	
21 to 24 years	4.3	4.9	5.2	1.4	5.4	4.5	8.4	4.3	2.3	
25 to 29 years	7.2	8.5	8.1	1.4	9.1	7.1	11.5	8.8	2.9	
30 to 34 years	7.9	9.9	9.9	†	10.7	7.3	11.5	9.9	2.6	
35 to 39 years	8.5	9.2	9.0	†	11.5	8.5	14.6	11.0	2.1	
40 to 44 years	7.7	8.4	7.9	†	10.5	8.0	8.3	9.8	2.0	
45 to 49 years	8.7	7.4	8.7	†	11.3	10.2	10.3	11.0	1.7	
50 to 54 years	9.8	7.2	9.1	†	11.6	12.4	9.5	12.8	2.0	
55 to 59 years	9.6	5.5	8.1	†	10.9	12.3	7.7	13.4	1.7	
60 to 64 years	6.2	2.8	4.4	†	6.5	8.1	5.0	9.5	1.1	
65 years and older	4.0	1.3	2.5	†	4.0	5.1	4.3	6.7	†	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†	
Asian ³	†	†	†	†	†	†	†	†	†	
Black or African American	21.8	13.0	11.6	28.8	16.3	18.2	16.8	39.6	20.0	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†	
White	73.8	81.3	83.8	65.3	79.7	78.2	78.5	56.1	73.3	
Other	3.9	5.1	4.0	5.6	3.5	3.0	4.0	3.6	6.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.22b. Missouri individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	97.7	96.5	97.5	97.2	98.4	97.6	97.4	98.3	96.3
Hispanic or Latino origin	2.3	3.5	2.5	2.8	1.6	2.4	2.6	1.7	3.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	53.0	57.1	55.5	55.7	51.5	54.0	75.0	48.0	47.3
Puerto Rican	10.1	3.8	10.1	5.7	15.7	11.2	15.0	11.7	6.0
Cuban	2.9	4.8	2.2	1.0	2.9	2.7	5.0	5.8	1.6
Other specific	34.0	34.3	32.2	37.5	29.9	32.1	5.0	34.5	45.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.23a. Montana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	61,189	9,788	13,719	4,220	3,759	14,071	550	3,370	11,712
Gender									
Male	27,027	3,734	4,919	2,813	1,429	4,945	169	2,048	6,970
Female	34,162	6,054	8,800	1,407	2,330	9,126	381	1,322	4,742
Total	61,189	9,788	13,719	4,220	3,759	14,071	550	3,370	11,712
Age²									
0 to 11 years	13,042	2,740	2,687	2,009	27	482	5	16	5,076
12 to 14 years	5,622	911	1,176	790	60	1,022	2	28	1,633
15 to 17 years	5,417	793	1,196	412	140	1,533	4	48	1,291
18 to 20 years	2,976	429	711	143	177	938	22	92	464
21 to 24 years	3,242	438	802	116	248	950	51	192	445
25 to 29 years	5,198	871	1,383	188	417	1,409	92	315	523
30 to 34 years	5,106	848	1,330	194	464	1,404	89	352	425
35 to 39 years	4,484	739	1,102	109	457	1,266	89	352	370
40 to 44 years	3,356	534	819	81	356	931	38	307	290
45 to 49 years	3,062	452	704	59	378	904	48	288	229
50 to 54 years	2,923	410	572	56	334	944	41	346	220
55 to 59 years	2,835	306	554	36	300	970	35	414	220
60 to 64 years	2,045	219	385	17	207	738	19	285	175
65 years and older	1,879	98	298	10	194	579	15	334	351
Total	61,187	9,788	13,719	4,220	3,759	14,070	550	3,369	11,712
Race									
American Indian or Alaska Native	7,054	1,450	1,525	388	288	1,684	43	352	1,324
Asian ³	166	26	30	12	12	31	2	15	38
Black or African American	653	106	115	52	34	123	7	54	162
Native Hawaiian or Other Pacific Islander	50	5	10	4	2	10	1	3	15
White	46,986	7,084	10,583	3,342	3,126	10,806	460	2,732	8,853
Other	1,340	259	282	109	71	254	11	56	298
Total	56,249	8,930	12,545	3,907	3,533	12,908	524	3,212	10,690

Continued. See notes at end of table.

Table 10.23a. Montana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	56,744	8,996	12,686	3,915	3,582	13,090	521	3,189	10,765
Hispanic or Latino origin	2,282	423	516	169	107	467	23	117	460
Total	59,026	9,419	13,202	4,084	3,689	13,557	544	3,306	11,225
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	134	16	15	2	23	39	1	33	5
Not a veteran	7,748	672	1,119	88	1,223	2,219	209	1,690	528
Total	7,882	688	1,134	90	1,246	2,258	210	1,723	533

‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.23b. Montana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	44.2	38.1	35.9	66.7	38.0	35.1	30.7	60.8	59.5	
Female	55.8	61.9	64.1	33.3	62.0	64.9	69.3	39.2	40.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	21.3	28.0	19.6	47.6	†	3.4	†	†	43.3	
12 to 14 years	9.2	9.3	8.6	18.7	1.6	7.3	†	†	13.9	
15 to 17 years	8.9	8.1	8.7	9.8	3.7	10.9	†	1.4	11.0	
18 to 20 years	4.9	4.4	5.2	3.4	4.7	6.7	4.0	2.7	4.0	
21 to 24 years	5.3	4.5	5.8	2.7	6.6	6.8	9.3	5.7	3.8	
25 to 29 years	8.5	8.9	10.1	4.5	11.1	10.0	16.7	9.3	4.5	
30 to 34 years	8.3	8.7	9.7	4.6	12.3	10.0	16.2	10.4	3.6	
35 to 39 years	7.3	7.6	8.0	2.6	12.2	9.0	16.2	10.4	3.2	
40 to 44 years	5.5	5.5	6.0	1.9	9.5	6.6	6.9	9.1	2.5	
45 to 49 years	5.0	4.6	5.1	1.4	10.1	6.4	8.7	8.5	2.0	
50 to 54 years	4.8	4.2	4.2	1.3	8.9	6.7	7.5	10.3	1.9	
55 to 59 years	4.6	3.1	4.0	†	8.0	6.9	6.4	12.3	1.9	
60 to 64 years	3.3	2.2	2.8	†	5.5	5.2	3.5	8.5	1.5	
65 years and older	3.1	1.0	2.2	†	5.2	4.1	2.7	9.9	3.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	12.5	16.2	12.2	9.9	8.2	13.0	8.2	11.0	12.4	
Asian ³	†	†	†	†	†	†	†	†	†	
Black or African American	1.2	1.2	†	1.3	†	†	1.3	1.7	1.5	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	83.5	79.3	84.4	85.5	88.5	83.7	87.8	85.1	82.8	
Other	2.4	2.9	2.2	2.8	2.0	2.0	2.1	1.7	2.8	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.23b. Montana individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	96.1	95.5	96.1	95.9	97.1	96.6	95.8	96.5	95.9
Hispanic or Latino origin	3.9	4.5	3.9	4.1	2.9	3.4	4.2	3.5	4.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	1.7	2.3	1.3	2.2	1.8	1.7	†	1.9	†
Not a veteran	98.3	97.7	98.7	97.8	98.2	98.3	99.5	98.1	99.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.24a. Nebraska individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	20,011	2,462	1,760	721	3,479	5,700	116	2,264	3,509
Gender									
Male	10,061	1,172	793	503	1,525	2,391	56	1,488	2,133
Female	9,856	1,286	946	217	1,934	3,273	60	772	1,368
Total	19,917	2,458	1,739	720	3,459	5,664	116	2,260	3,501
Age²									
0 to 11 years	715	184	31	178	8	25	--	4	285
12 to 14 years	614	92	40	145	13	62	1	1	260
15 to 17 years	798	123	53	108	17	163	3	5	326
18 to 20 years	1,016	163	85	49	110	302	6	59	242
21 to 24 years	1,810	260	188	58	306	491	11	193	303
25 to 29 years	2,581	357	280	58	458	680	23	293	432
30 to 34 years	2,511	307	275	33	518	668	15	305	390
35 to 39 years	2,203	268	225	26	437	623	11	314	299
40 to 44 years	1,720	208	151	23	362	510	12	205	249
45 to 49 years	1,729	171	114	26	387	554	11	239	227
50 to 54 years	1,700	140	135	9	364	599	15	237	201
55 to 59 years	1,369	96	96	4	267	540	2	201	163
60 to 64 years	813	60	53	3	146	323	4	134	90
65 years and older	430	33	34	1	86	159	2	73	42
Total	20,009	2,462	1,760	721	3,479	5,699	116	2,263	3,509
Race									
American Indian or Alaska Native	480	78	33	8	73	149	5	49	85
Asian ³	136	18	8	4	11	36	1	28	30
Black or African American	1,741	153	87	37	244	408	3	409	400
Native Hawaiian or Other Pacific Islander	82	12	5	1	15	22	--	12	15
White	16,002	1,955	1,478	590	2,913	4,672	90	1,610	2,694
Other	992	155	85	68	122	224	7	119	212
Total	19,433	2,371	1,696	708	3,378	5,511	106	2,227	3,436

Continued. See notes at end of table.

Table 10.24a. Nebraska individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	16,128	1,947	1,433	581	2,936	4,680	79	1,772	2,700
Hispanic or Latino origin	1,770	291	151	101	209	488	9	187	334
Total	17,898	2,238	1,584	682	3,145	5,168	88	1,959	3,034
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	370	55	27	1	69	105	2	81	30
Not a veteran	16,444	1,751	1,566	272	3,252	5,168	103	2,041	2,291
Total	16,814	1,806	1,593	273	3,321	5,273	105	2,122	2,321

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.24b. Nebraska individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	50.5	47.7	45.6	69.9	44.1	42.2	48.3	65.8	60.9	
Female	49.5	52.3	54.4	30.1	55.9	57.8	51.7	34.2	39.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	3.6	7.5	1.8	24.7	†	†	--	†	8.1	
12 to 14 years	3.1	3.7	2.3	20.1	†	1.1	†	†	7.4	
15 to 17 years	4.0	5.0	3.0	15.0	†	2.9	2.6	†	9.3	
18 to 20 years	5.1	6.6	4.8	6.8	3.2	5.3	5.2	2.6	6.9	
21 to 24 years	9.0	10.6	10.7	8.0	8.8	8.6	9.5	8.5	8.6	
25 to 29 years	12.9	14.5	15.9	8.0	13.2	11.9	19.8	12.9	12.3	
30 to 34 years	12.5	12.5	15.6	4.6	14.9	11.7	12.9	13.5	11.1	
35 to 39 years	11.0	10.9	12.8	3.6	12.6	10.9	9.5	13.9	8.5	
40 to 44 years	8.6	8.4	8.6	3.2	10.4	8.9	10.3	9.1	7.1	
45 to 49 years	8.6	6.9	6.5	3.6	11.1	9.7	9.5	10.6	6.5	
50 to 54 years	8.5	5.7	7.7	1.2	10.5	10.5	12.9	10.5	5.7	
55 to 59 years	6.8	3.9	5.5	†	7.7	9.5	1.7	8.9	4.6	
60 to 64 years	4.1	2.4	3.0	†	4.2	5.7	3.4	5.9	2.6	
65 years and older	2.1	1.3	1.9	†	2.5	2.8	1.7	3.2	1.2	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	2.5	3.3	1.9	1.1	2.2	2.7	4.7	2.2	2.5	
Asian ³	†	†	†	†	†	†	†	1.3	†	
Black or African American	9.0	6.5	5.1	5.2	7.2	7.4	2.8	18.4	11.6	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†	
White	82.3	82.5	87.1	83.3	86.2	84.8	84.9	72.3	78.4	
Other	5.1	6.5	5.0	9.6	3.6	4.1	6.6	5.3	6.2	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.24b. Nebraska individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	90.1	87.0	90.5	85.2	93.4	90.6	89.8	90.5	89.0
Hispanic or Latino origin	9.9	13.0	9.5	14.8	6.6	9.4	10.2	9.5	11.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	2.2	3.0	1.7	†	2.1	2.0	1.9	3.8	1.3
Not a veteran	97.8	97.0	98.3	99.6	97.9	98.0	98.1	96.2	98.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.25a. Nevada individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	13,250	1,681	1,589	367	1,008	2,022	437	2,003	4,143
Gender									
Male	6,688	649	618	265	401	730	182	1,310	2,533
Female	6,555	1,031	969	102	606	1,291	255	692	1,609
Total	13,243	1,680	1,587	367	1,007	2,021	437	2,002	4,142
Age²									
0 to 11 years	1,467	497	253	170	19	169	--	4	355
12 to 14 years	1,025	254	113	68	23	312	--	7	248
15 to 17 years	1,268	231	128	38	71	394	3	29	374
18 to 20 years	568	71	68	19	32	77	24	67	210
21 to 24 years	793	58	76	10	50	62	44	134	359
25 to 29 years	1,114	73	119	14	85	90	51	193	489
30 to 34 years	1,106	78	127	11	72	93	39	226	460
35 to 39 years	986	70	92	7	96	83	46	212	380
40 to 44 years	879	65	91	9	96	96	50	200	272
45 to 49 years	976	68	124	5	106	121	54	227	271
50 to 54 years	1,091	80	133	2	114	153	54	252	303
55 to 59 years	916	60	106	4	122	148	39	216	221
60 to 64 years	608	35	90	5	70	116	24	143	125
65 years and older	448	38	68	5	51	108	9	93	76
Total	13,245	1,678	1,588	367	1,007	2,022	437	2,003	4,143
Race									
American Indian or Alaska Native	244	37	26	5	12	34	4	47	79
Asian ³	344	34	27	4	27	54	7	122	69
Black or African American	1,718	204	130	38	93	185	53	381	634
Native Hawaiian or Other Pacific Islander	89	9	13	3	4	15	2	18	25
White	7,405	1,004	861	201	646	1,217	251	966	2,259
Other	1,284	72	144	6	99	140	48	322	453
Total	11,084	1,360	1,201	257	881	1,645	365	1,856	3,519

Continued. See notes at end of table.

Table 10.25a. Nevada individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	8,263	962	846	201	723	1,136	290	1,458	2,647
Hispanic or Latino origin	2,218	325	287	35	115	446	45	314	651
Total	10,481	1,287	1,133	236	838	1,582	335	1,772	3,298
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	6	--	--	--	--	1	--	1	4
Not a veteran	3,228	201	378	25	335	339	138	886	926
Total	3,234	201	378	25	335	340	138	887	930

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.25b. Nevada individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	50.5	38.6	38.9	72.2	39.8	36.1	41.6	65.4	61.2	
Female	49.5	61.4	61.1	27.8	60.2	63.9	58.4	34.6	38.8	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	11.1	29.6	15.9	46.3	1.9	8.4	--	†	8.6	
12 to 14 years	7.7	15.1	7.1	18.5	2.3	15.4	--	†	6.0	
15 to 17 years	9.6	13.8	8.1	10.4	7.1	19.5	†	1.4	9.0	
18 to 20 years	4.3	4.2	4.3	5.2	3.2	3.8	5.5	3.3	5.1	
21 to 24 years	6.0	3.5	4.8	2.7	5.0	3.1	10.1	6.7	8.7	
25 to 29 years	8.4	4.4	7.5	3.8	8.4	4.5	11.7	9.6	11.8	
30 to 34 years	8.4	4.6	8.0	3.0	7.1	4.6	8.9	11.3	11.1	
35 to 39 years	7.4	4.2	5.8	1.9	9.5	4.1	10.5	10.6	9.2	
40 to 44 years	6.6	3.9	5.7	2.5	9.5	4.7	11.4	10.0	6.6	
45 to 49 years	7.4	4.1	7.8	1.4	10.5	6.0	12.4	11.3	6.5	
50 to 54 years	8.2	4.8	8.4	†	11.3	7.6	12.4	12.6	7.3	
55 to 59 years	6.9	3.6	6.7	1.1	12.1	7.3	8.9	10.8	5.3	
60 to 64 years	4.6	2.1	5.7	1.4	7.0	5.7	5.5	7.1	3.0	
65 years and older	3.4	2.3	4.3	1.4	5.1	5.3	2.1	4.6	1.8	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	2.2	2.7	2.2	1.9	1.4	2.1	1.1	2.5	2.2	
Asian ³	3.1	2.5	2.2	1.6	3.1	3.3	1.9	6.6	2.0	
Black or African American	15.5	15.0	10.8	14.8	10.6	11.2	14.5	20.5	18.0	
Native Hawaiian or Other Pacific Islander	†	†	1.1	1.2	†	†	†	†	†	
White	66.8	73.8	71.7	78.2	73.3	74.0	68.8	52.0	64.2	
Other	11.6	5.3	12.0	2.3	11.2	8.5	13.2	17.3	12.9	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.25b. Nevada individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	78.8	74.7	74.7	85.2	86.3	71.8	86.6	82.3	80.3
Hispanic or Latino origin	21.2	25.3	25.3	14.8	13.7	28.2	13.4	17.7	19.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	†	--	--	--	--	†	--	†	†
Not a veteran	99.8	100.0	100.0	100.0	100.0	99.7	100.0	99.9	99.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.26a. New Mexico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	149,498	25,568	20,189	7,559	6,705	20,739	230	5,557	62,951
Gender									
Male	73,678	10,226	6,992	5,318	2,466	6,891	89	3,546	38,150
Female	75,602	15,332	13,188	2,237	4,234	13,838	141	2,006	24,626
Total	149,280	25,558	20,180	7,555	6,700	20,729	230	5,552	62,776
Age²									
0 to 11 years	33,849	6,164	1,250	3,550	49	433	8	17	22,378
12 to 14 years	10,108	2,687	881	1,490	61	1,044	5	24	3,916
15 to 17 years	9,859	2,379	1,222	849	172	1,693	8	49	3,487
18 to 20 years	5,739	1,069	1,051	323	251	1,132	6	130	1,777
21 to 24 years	7,393	1,184	1,311	207	376	1,246	12	361	2,696
25 to 29 years	12,839	2,004	2,182	300	669	1,842	27	550	5,265
30 to 34 years	13,043	2,051	2,286	223	739	1,754	30	621	5,339
35 to 39 years	11,868	1,885	2,115	201	782	1,673	19	653	4,540
40 to 44 years	9,116	1,416	1,604	135	743	1,412	30	523	3,253
45 to 49 years	8,445	1,269	1,458	99	721	1,535	20	497	2,846
50 to 54 years	8,341	1,148	1,505	58	685	1,742	20	561	2,622
55 to 59 years	7,678	1,008	1,310	52	636	1,851	17	581	2,223
60 to 64 years	5,449	694	998	53	415	1,387	11	417	1,474
65 years and older	5,770	610	1,016	19	406	1,995	17	573	1,134
Total	149,497	25,568	20,189	7,559	6,705	20,739	230	5,557	62,950
Race									
American Indian or Alaska Native	11,844	1,767	1,046	425	159	1,425	8	269	6,745
Asian ³	1,598	285	183	66	32	190	--	41	801
Black or African American	3,015	587	391	160	159	451	11	226	1,030
Native Hawaiian or Other Pacific Islander	141	15	18	6	2	27	1	13	59
White	104,091	18,651	15,284	5,167	4,987	15,155	172	3,603	41,072
Other	3,360	553	437	191	184	421	12	254	1,308
Total	124,049	21,858	17,359	6,015	5,523	17,669	204	4,406	51,015

Continued. See notes at end of table.

Table 10.26a. New Mexico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	57,560	9,854	8,158	2,681	3,378	8,603	112	2,413	22,361
Hispanic or Latino origin	66,599	11,830	9,338	3,344	2,185	9,230	89	2,145	28,438
Total	124,159	21,684	17,496	6,025	5,563	17,833	201	4,558	50,799
Specific Hispanic or Latino origin									
Mexican	834	131	101	76	55	136	2	83	250
Puerto Rican	64	10	8	9	7	12	--	5	13
Cuban	44	10	2	5	5	7	--	3	12
Other specific	494	72	68	26	39	53	1	51	184
Total	1,436	223	179	116	106	208	3	142	459
Veteran status⁴									
Veteran	699	146	93	13	54	137	2	45	209
Not a veteran	92,372	13,888	16,476	1,618	6,061	16,903	202	5,108	32,116
Total	93,071	14,034	16,569	1,631	6,115	17,040	204	5,153	32,325

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.26b. New Mexico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	49.4	40.0	34.6	70.4	36.8	33.2	38.7	63.9	60.8	
Female	50.6	60.0	65.4	29.6	63.2	66.8	61.3	36.1	39.2	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Age²										
0 to 11 years	22.6	24.1	6.2	47.0	†	2.1	3.5	†	35.5	
12 to 14 years	6.8	10.5	4.4	19.7	†	5.0	2.2	†	6.2	
15 to 17 years	6.6	9.3	6.1	11.2	2.6	8.2	3.5	†	5.5	
18 to 20 years	3.8	4.2	5.2	4.3	3.7	5.5	2.6	2.3	2.8	
21 to 24 years	4.9	4.6	6.5	2.7	5.6	6.0	5.2	6.5	4.3	
25 to 29 years	8.6	7.8	10.8	4.0	10.0	8.9	11.7	9.9	8.4	
30 to 34 years	8.7	8.0	11.3	3.0	11.0	8.5	13.0	11.2	8.5	
35 to 39 years	7.9	7.4	10.5	2.7	11.7	8.1	8.3	11.8	7.2	
40 to 44 years	6.1	5.5	7.9	1.8	11.1	6.8	13.0	9.4	5.2	
45 to 49 years	5.6	5.0	7.2	1.3	10.8	7.4	8.7	8.9	4.5	
50 to 54 years	5.6	4.5	7.5	†	10.2	8.4	8.7	10.1	4.2	
55 to 59 years	5.1	3.9	6.5	†	9.5	8.9	7.4	10.5	3.5	
60 to 64 years	3.6	2.7	4.9	†	6.2	6.7	4.8	7.5	2.3	
65 years and older	3.9	2.4	5.0	†	6.1	9.6	7.4	10.3	1.8	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Race										
American Indian or Alaska Native	9.5	8.1	6.0	7.1	2.9	8.1	3.9	6.1	13.2	
Asian ³	1.3	1.3	1.1	1.1	†	1.1	--	†	1.6	
Black or African American	2.4	2.7	2.3	2.7	2.9	2.6	5.4	5.1	2.0	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	83.9	85.3	88.0	85.9	90.3	85.8	84.3	81.8	80.5	
Other	2.7	2.5	2.5	3.2	3.3	2.4	5.9	5.8	2.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	

Continued. See notes at end of table.

Table 10.26b. New Mexico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	46.4	45.4	46.6	44.5	60.7	48.2	55.7	52.9	44.0
Hispanic or Latino origin	53.6	54.6	53.4	55.5	39.3	51.8	44.3	47.1	56.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	58.1	58.7	56.4	65.5	51.9	65.4	66.7	58.5	54.5
Puerto Rican	4.5	4.5	4.5	7.8	6.6	5.8	--	3.5	2.8
Cuban	3.1	4.5	1.1	4.3	4.7	3.4	--	2.1	2.6
Other specific	34.4	32.3	38.0	22.4	36.8	25.5	33.3	35.9	40.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	†	1.0	†	†	†	†	†	†	†
Not a veteran	99.2	99.0	99.4	99.2	99.1	99.2	99.0	99.1	99.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.27a. New York individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals²	56,509	7,579	3,271	1,870	5,166	7,603	4,586	19,316	7,118
Gender									
Male	36,966	5,253	1,707	1,427	2,503	3,401	4,229	13,297	5,149
Female	19,543	2,326	1,564	443	2,663	4,202	357	6,019	1,969
Total	56,509	7,579	3,271	1,870	5,166	7,603	4,586	19,316	7,118
Age³									
0 to 11 years	2,171	274	169	920	43	89	--	17	659
12 to 14 years	2,308	284	243	423	105	513	3	51	686
15 to 17 years	2,927	320	325	282	270	892	20	173	645
18 to 20 years	1,391	291	101	56	132	267	132	236	176
21 to 24 years	2,895	737	147	31	231	269	430	717	333
25 to 29 years	5,200	1,192	306	54	404	478	730	1,401	635
30 to 34 years	5,294	1,067	325	28	433	482	780	1,522	657
35 to 39 years	5,251	882	308	28	477	511	689	1,688	668
40 to 44 years	4,347	609	233	23	448	494	470	1,581	489
45 to 49 years	4,836	644	256	5	513	607	448	1,822	541
50 to 54 years	5,630	568	260	7	561	846	452	2,359	577
55 to 59 years	5,471	372	216	8	572	756	253	2,815	479
60 to 64 years	3,996	192	144	2	469	580	99	2,236	274
65 years and older	4,788	147	238	3	508	818	80	2,697	297
Total	56,505	7,579	3,271	1,870	5,166	7,602	4,586	19,315	7,116
Race									
American Indian or Alaska Native	372	85	22	9	27	55	47	67	60
Asian ⁴	966	48	28	9	70	131	11	621	48
Black or African American	16,155	2,308	362	392	818	1,272	2,023	7,061	1,919
Native Hawaiian or Other Pacific Islander	51	7	--	--	4	4	2	26	8
White	28,378	3,673	2,309	1,036	3,330	4,431	1,615	8,300	3,684
Other	6,740	994	322	315	522	926	639	2,043	979
Total	52,662	7,115	3,043	1,761	4,771	6,819	4,337	18,118	6,698

Continued. See notes at end of table.

Table 10.27a. New York individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	41,959	5,744	2,533	1,412	3,837	5,350	3,461	14,230	5,392
Hispanic or Latino origin	8,520	1,198	385	245	644	1,266	918	2,752	1,112
Total	50,479	6,942	2,918	1,657	4,481	6,616	4,379	16,982	6,504
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁵									
Veteran	331	20	5	--	31	21	20	202	32
Not a veteran	12,556	777	362	19	1,005	1,056	1,188	7,082	1,067
Total	12,887	797	367	19	1,036	1,077	1,208	7,284	1,099

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² New York reports only a subset of the total individuals served through their SMHAs.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.27b. New York individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	65.4	69.3	52.2	76.3	48.5	44.7	92.2	68.8	72.3
Female	34.6	30.7	47.8	23.7	51.5	55.3	7.8	31.2	27.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age³									
0 to 11 years	3.8	3.6	5.2	49.2	†	1.2	--	†	9.3
12 to 14 years	4.1	3.7	7.4	22.6	2.0	6.7	†	†	9.6
15 to 17 years	5.2	4.2	9.9	15.1	5.2	11.7	†	†	9.1
18 to 20 years	2.5	3.8	3.1	3.0	2.6	3.5	2.9	1.2	2.5
21 to 24 years	5.1	9.7	4.5	1.7	4.5	3.5	9.4	3.7	4.7
25 to 29 years	9.2	15.7	9.4	2.9	7.8	6.3	15.9	7.3	8.9
30 to 34 years	9.4	14.1	9.9	1.5	8.4	6.3	17.0	7.9	9.2
35 to 39 years	9.3	11.6	9.4	1.5	9.2	6.7	15.0	8.7	9.4
40 to 44 years	7.7	8.0	7.1	1.2	8.7	6.5	10.2	8.2	6.9
45 to 49 years	8.6	8.5	7.8	†	9.9	8.0	9.8	9.4	7.6
50 to 54 years	10.0	7.5	7.9	†	10.9	11.1	9.9	12.2	8.1
55 to 59 years	9.7	4.9	6.6	†	11.1	9.9	5.5	14.6	6.7
60 to 64 years	7.1	2.5	4.4	†	9.1	7.6	2.2	11.6	3.9
65 years and older	8.5	1.9	7.3	†	9.8	10.8	1.7	14.0	4.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	1.2	†	†	†	†	1.1	†	†
Asian ⁴	1.8	†	†	†	1.5	1.9	†	3.4	†
Black or African American	30.7	32.4	11.9	22.3	17.1	18.7	46.6	39.0	28.7
Native Hawaiian or Other Pacific Islander	†	†	--	--	†	†	†	†	†
White	53.9	51.6	75.9	58.8	69.8	65.0	37.2	45.8	55.0
Other	12.8	14.0	10.6	17.9	10.9	13.6	14.7	11.3	14.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.27b. New York individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	83.1	82.7	86.8	85.2	85.6	80.9	79.0	83.8	82.9
Hispanic or Latino origin	16.9	17.3	13.2	14.8	14.4	19.1	21.0	16.2	17.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁵									
Veteran	2.6	2.5	1.4	--	3.0	1.9	1.7	2.8	2.9
Not a veteran	97.4	97.5	98.6	100.0	97.0	98.1	98.3	97.2	97.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² New York reports only a subset of the total individuals served through their SMHAs.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.28a. North Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	90,175	11,858	8,114	2,439	14,038	29,505	1,110	13,345	9,766
Gender									
Male	44,058	5,042	3,644	1,766	6,124	12,778	397	8,566	5,741
Female	46,100	6,814	4,468	671	7,910	16,726	713	4,774	4,024
Total	90,158	11,856	8,112	2,437	14,034	29,504	1,110	13,340	9,765
Age²									
0 to 11 years	3,565	890	164	1,007	34	176	1	13	1,280
12 to 14 years	3,454	708	173	587	96	558	--	27	1,305
15 to 17 years	3,950	693	270	490	208	866	15	80	1,328
18 to 20 years	3,479	470	349	150	382	1,207	72	278	571
21 to 24 years	7,117	931	763	66	1,062	2,430	141	1,060	664
25 to 29 years	11,246	1,533	1,245	59	1,905	3,695	210	1,792	807
30 to 34 years	10,220	1,372	1,126	27	1,938	3,307	165	1,631	654
35 to 39 years	9,346	1,212	985	20	1,791	3,002	143	1,588	605
40 to 44 years	8,411	1,099	787	10	1,630	2,964	108	1,319	494
45 to 49 years	8,856	1,005	690	9	1,736	3,425	97	1,354	540
50 to 54 years	8,374	878	679	6	1,499	3,353	71	1,439	449
55 to 59 years	6,428	584	482	5	1,012	2,636	49	1,275	385
60 to 64 years	3,569	276	269	3	495	1,327	29	888	282
65 years and older	2,159	207	132	--	250	559	9	600	402
Total	90,174	11,858	8,114	2,439	14,038	29,505	1,110	13,344	9,766
Race									
American Indian or Alaska Native	1,659	276	126	56	333	503	18	196	151
Asian ³	494	68	37	7	57	160	4	124	37
Black or African American	27,145	3,090	1,408	1,052	3,361	7,848	197	7,053	3,136
Native Hawaiian or Other Pacific Islander	52	10	6	1	5	17	--	9	4
White	56,922	7,537	6,152	1,200	9,911	19,715	849	5,536	6,022
Other	3,598	836	363	113	326	1,171	37	372	380
Total	89,870	11,817	8,092	2,429	13,993	29,414	1,105	13,290	9,730

Continued. See notes at end of table.

Table 10.28a. North Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	85,826	10,837	7,677	2,301	13,687	28,044	1,082	12,892	9,306
Hispanic or Latino origin	4,034	981	404	117	300	1,382	26	391	433
Total	89,860	11,818	8,081	2,418	13,987	29,426	1,108	13,283	9,739
Specific Hispanic or Latino origin									
Mexican	1,147	241	121	41	75	417	5	114	133
Puerto Rican	334	40	30	6	57	121	7	54	19
Cuban	96	14	13	--	11	30	--	13	15
Other specific	2,415	682	239	69	152	798	13	203	259
Total	3,992	977	403	116	295	1,366	25	384	426
Veteran status⁴									
Veteran	1,230	202	86	3	164	398	7	224	146
Not a veteran	77,313	9,327	7,390	346	13,436	27,354	1,072	12,729	5,659
Total	78,543	9,529	7,476	349	13,600	27,752	1,079	12,953	5,805

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.28b. North Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	48.9	42.5	44.9	72.5	43.6	43.3	35.8	64.2	58.8
Female	51.1	57.5	55.1	27.5	56.4	56.7	64.2	35.8	41.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	4.0	7.5	2.0	41.3	†	†	†	†	13.1
12 to 14 years	3.8	6.0	2.1	24.1	†	1.9	--	†	13.4
15 to 17 years	4.4	5.8	3.3	20.1	1.5	2.9	1.4	†	13.6
18 to 20 years	3.9	4.0	4.3	6.2	2.7	4.1	6.5	2.1	5.8
21 to 24 years	7.9	7.9	9.4	2.7	7.6	8.2	12.7	7.9	6.8
25 to 29 years	12.5	12.9	15.3	2.4	13.6	12.5	18.9	13.4	8.3
30 to 34 years	11.3	11.6	13.9	1.1	13.8	11.2	14.9	12.2	6.7
35 to 39 years	10.4	10.2	12.1	†	12.8	10.2	12.9	11.9	6.2
40 to 44 years	9.3	9.3	9.7	†	11.6	10.0	9.7	9.9	5.1
45 to 49 years	9.8	8.5	8.5	†	12.4	11.6	8.7	10.1	5.5
50 to 54 years	9.3	7.4	8.4	†	10.7	11.4	6.4	10.8	4.6
55 to 59 years	7.1	4.9	5.9	†	7.2	8.9	4.4	9.6	3.9
60 to 64 years	4.0	2.3	3.3	†	3.5	4.5	2.6	6.7	2.9
65 years and older	2.4	1.7	1.6	--	1.8	1.9	†	4.5	4.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	1.8	2.3	1.6	2.3	2.4	1.7	1.6	1.5	1.6
Asian ³	†	†	†	†	†	†	†	†	†
Black or African American	30.2	26.1	17.4	43.3	24.0	26.7	17.8	53.1	32.2
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†
White	63.3	63.8	76.0	49.4	70.8	67.0	76.8	41.7	61.9
Other	4.0	7.1	4.5	4.7	2.3	4.0	3.3	2.8	3.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.28b. North Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	95.5	91.7	95.0	95.2	97.9	95.3	97.7	97.1	95.6
Hispanic or Latino origin	4.5	8.3	5.0	4.8	2.1	4.7	2.3	2.9	4.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	28.7	24.7	30.0	35.3	25.4	30.5	20.0	29.7	31.2
Puerto Rican	8.4	4.1	7.4	5.2	19.3	8.9	28.0	14.1	4.5
Cuban	2.4	1.4	3.2	--	3.7	2.2	--	3.4	3.5
Other specific	60.5	69.8	59.3	59.5	51.5	58.4	52.0	52.9	60.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	1.6	2.1	1.2	†	1.2	1.4	†	1.7	2.5
Not a veteran	98.4	97.9	98.8	99.1	98.8	98.6	99.4	98.3	97.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.29a. North Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	7,182	505	776	443	925	1,853	87	1,259	1,334
Gender									
Male	3,530	209	332	296	416	604	58	789	826
Female	3,645	295	444	145	507	1,248	29	470	507
Total	7,175	504	776	441	923	1,852	87	1,259	1,333
Age²									
0 to 11 years	131	36	14	46	2	--	--	--	33
12 to 14 years	191	32	19	74	11	7	--	--	48
15 to 17 years	292	56	20	78	26	55	--	3	54
18 to 20 years	279	28	26	50	26	45	1	4	99
21 to 24 years	428	40	56	44	45	77	--	17	149
25 to 29 years	760	48	80	47	90	156	10	70	259
30 to 34 years	873	67	107	35	107	199	20	108	230
35 to 39 years	724	45	111	28	94	210	17	89	130
40 to 44 years	610	30	74	17	99	180	12	113	85
45 to 49 years	592	33	66	9	101	174	10	119	80
50 to 54 years	631	30	56	3	93	217	5	162	65
55 to 59 years	649	30	46	5	88	200	8	223	49
60 to 64 years	497	22	40	5	80	159	2	161	28
65 years and older	524	8	61	2	63	174	2	190	24
Total	7,181	505	776	443	925	1,853	87	1,259	1,333
Race									
American Indian or Alaska Native	833	82	63	45	74	191	13	95	270
Asian ³	28	2	1	--	4	4	--	13	4
Black or African American	132	13	6	9	23	19	2	36	24
Native Hawaiian or Other Pacific Islander	4	--	--	--	1	--	--	2	1
White	5,962	382	687	368	798	1,588	67	1,088	984
Other	214	25	19	21	24	47	4	25	49
Total	7,173	504	776	443	924	1,849	86	1,259	1,332

Continued. See notes at end of table.

Table 10.29a. North Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	7,013	488	758	428	910	1,815	82	1,240	1,292
Hispanic or Latino origin	150	16	17	14	11	34	5	15	38
Total	7,163	504	775	442	921	1,849	87	1,255	1,330
Specific Hispanic or Latino origin									
Mexican	27	4	3	2	4	7	1	1	5
Puerto Rican	10	1	--	--	1	3	1	--	4
Cuban	--	--	--	--	--	--	--	--	--
Other specific	33	2	5	6	1	5	--	8	6
Total	70	7	8	8	6	15	2	9	15
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.29b. North Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	49.2	41.5	42.8	67.1	45.1	32.6	66.7	62.7	62.0	
Female	50.8	58.5	57.2	32.9	54.9	67.4	33.3	37.3	38.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Age²										
0 to 11 years	1.8	7.1	1.8	10.4	†	--	--	--	2.5	
12 to 14 years	2.7	6.3	2.4	16.7	1.2	†	--	--	3.6	
15 to 17 years	4.1	11.1	2.6	17.6	2.8	3.0	--	†	4.1	
18 to 20 years	3.9	5.5	3.4	11.3	2.8	2.4	1.1	†	7.4	
21 to 24 years	6.0	7.9	7.2	9.9	4.9	4.2	--	1.4	11.2	
25 to 29 years	10.6	9.5	10.3	10.6	9.7	8.4	11.5	5.6	19.4	
30 to 34 years	12.2	13.3	13.8	7.9	11.6	10.7	23.0	8.6	17.3	
35 to 39 years	10.1	8.9	14.3	6.3	10.2	11.3	19.5	7.1	9.8	
40 to 44 years	8.5	5.9	9.5	3.8	10.7	9.7	13.8	9.0	6.4	
45 to 49 years	8.2	6.5	8.5	2.0	10.9	9.4	11.5	9.5	6.0	
50 to 54 years	8.8	5.9	7.2	†	10.1	11.7	5.7	12.9	4.9	
55 to 59 years	9.0	5.9	5.9	1.1	9.5	10.8	9.2	17.7	3.7	
60 to 64 years	6.9	4.4	5.2	1.1	8.6	8.6	2.3	12.8	2.1	
65 years and older	7.3	1.6	7.9	†	6.8	9.4	2.3	15.1	1.8	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
Race										
American Indian or Alaska Native	11.6	16.3	8.1	10.2	8.0	10.3	15.1	7.5	20.3	
Asian ³	†	†	†	--	†	†	--	1.0	†	
Black or African American	1.8	2.6	†	2.0	2.5	1.0	2.3	2.9	1.8	
Native Hawaiian or Other Pacific Islander	†	--	--	--	†	--	--	†	†	
White	83.1	75.8	88.5	83.1	86.4	85.9	77.9	86.4	73.9	
Other	3.0	5.0	2.4	4.7	2.6	2.5	4.7	2.0	3.7	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	

Continued. See notes at end of table.

Table 10.29b. North Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	97.9	96.8	97.8	96.8	98.8	98.2	94.3	98.8	97.1
Hispanic or Latino origin	2.1	3.2	2.2	3.2	1.2	1.8	5.7	1.2	2.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	38.6	57.1	37.5	25.0	66.7	46.7	50.0	11.1	33.3
Puerto Rican	14.3	14.3	--	--	16.7	20.0	50.0	--	26.7
Cuban	--	--	--	--	--	--	--	--	--
Other specific	47.1	28.6	62.5	75.0	16.7	33.3	--	88.9	40.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.30a. Ohio individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	449,140	81,609	56,369	46,037	50,839	109,867	2,835	39,583	62,001
Gender									
Male	205,836	34,676	21,525	31,175	18,670	39,139	1,172	23,906	35,573
Female	234,261	45,065	33,886	13,415	31,228	68,977	1,598	15,212	24,880
Total	440,097	79,741	55,411	44,590	49,898	108,116	2,770	39,118	60,453
Age²									
0 to 11 years	78,132	25,965	7,747	23,331	252	2,361	15	65	18,396
12 to 14 years	37,048	8,985	4,349	9,471	473	5,514	17	100	8,139
15 to 17 years	36,947	7,510	4,805	6,157	1,040	8,884	29	253	8,269
18 to 20 years	20,310	3,333	2,964	2,016	1,791	5,889	131	834	3,352
21 to 24 years	24,045	3,760	3,593	1,081	3,355	7,157	269	1,966	2,864
25 to 29 years	36,614	5,977	5,813	1,182	5,714	10,367	429	3,374	3,758
30 to 34 years	35,708	5,726	5,912	924	6,201	9,891	418	3,550	3,086
35 to 39 years	33,794	5,130	5,055	657	6,481	9,409	341	4,090	2,631
40 to 44 years	27,486	3,879	3,508	416	5,417	8,152	288	3,690	2,136
45 to 49 years	28,362	3,434	3,396	330	5,454	9,113	262	4,149	2,224
50 to 54 years	28,592	2,900	3,060	201	5,315	10,081	241	4,645	2,149
55 to 59 years	27,321	2,407	2,665	155	4,532	9,893	170	5,452	2,047
60 to 64 years	18,525	1,373	1,788	83	2,814	6,772	134	4,097	1,464
65 years and older	16,254	1,230	1,714	33	2,000	6,384	91	3,317	1,485
Total	449,138	81,609	56,369	46,037	50,839	109,867	2,835	39,582	62,000
Race									
American Indian or Alaska Native	1,422	284	175	106	163	379	12	116	187
Asian ³	2,076	364	248	131	157	522	11	321	322
Black or African American	102,146	17,857	7,810	13,444	8,437	21,924	551	14,859	17,264
Native Hawaiian or Other Pacific Islander	386	83	59	30	43	114	--	22	35
White	293,887	53,769	41,666	28,607	36,164	74,046	1,824	19,839	37,972
Other	2,286	434	231	236	213	590	10	248	324
Total	402,203	72,791	50,189	42,554	45,177	97,575	2,408	35,405	56,104

Continued. See notes at end of table.

Table 10.30a. Ohio individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	408,507	75,034	52,343	43,242	45,897	99,680	2,332	32,891	57,088
Hispanic or Latino origin	12,958	2,614	1,527	1,668	938	3,226	39	821	2,125
Total	421,465	77,648	53,870	44,910	46,835	102,906	2,371	33,712	59,213
Specific Hispanic or Latino origin									
Mexican	387	79	57	23	40	120	--	34	34
Puerto Rican	368	65	37	20	41	105	1	53	46
Cuban	20	3	3	2	3	4	--	2	3
Other specific	2,108	471	155	231	163	447	5	166	470
Total	2,883	618	252	276	247	676	6	255	553
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.30b. Ohio individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	46.8	43.5	38.8	69.9	37.4	36.2	42.3	61.1	58.8	
Female	53.2	56.5	61.2	30.1	62.6	63.8	57.7	38.9	41.2	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	17.4	31.8	13.7	50.7	†	2.1	†	†	29.7	
12 to 14 years	8.2	11.0	7.7	20.6	†	5.0	†	†	13.1	
15 to 17 years	8.2	9.2	8.5	13.4	2.0	8.1	1.0	†	13.3	
18 to 20 years	4.5	4.1	5.3	4.4	3.5	5.4	4.6	2.1	5.4	
21 to 24 years	5.4	4.6	6.4	2.3	6.6	6.5	9.5	5.0	4.6	
25 to 29 years	8.2	7.3	10.3	2.6	11.2	9.4	15.1	8.5	6.1	
30 to 34 years	8.0	7.0	10.5	2.0	12.2	9.0	14.7	9.0	5.0	
35 to 39 years	7.5	6.3	9.0	1.4	12.7	8.6	12.0	10.3	4.2	
40 to 44 years	6.1	4.8	6.2	†	10.7	7.4	10.2	9.3	3.4	
45 to 49 years	6.3	4.2	6.0	†	10.7	8.3	9.2	10.5	3.6	
50 to 54 years	6.4	3.6	5.4	†	10.5	9.2	8.5	11.7	3.5	
55 to 59 years	6.1	2.9	4.7	†	8.9	9.0	6.0	13.8	3.3	
60 to 64 years	4.1	1.7	3.2	†	5.5	6.2	4.7	10.4	2.4	
65 years and older	3.6	1.5	3.0	†	3.9	5.8	3.2	8.4	2.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†	
Asian ³	†	†	†	†	†	†	†	†	†	
Black or African American	25.4	24.5	15.6	31.6	18.7	22.5	22.9	42.0	30.8	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†	
White	73.1	73.9	83.0	67.2	80.0	75.9	75.7	56.0	67.7	
Other	†	†	†	†	†	†	†	†	†	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.30b. Ohio individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	96.9	96.6	97.2	96.3	98.0	96.9	98.4	97.6	96.4
Hispanic or Latino origin	3.1	3.4	2.8	3.7	2.0	3.1	1.6	2.4	3.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	13.4	12.8	22.6	8.3	16.2	17.8	--	13.3	6.1
Puerto Rican	12.8	10.5	14.7	7.2	16.6	15.5	16.7	20.8	8.3
Cuban	†	†	1.2	†	1.2	†	--	†	†
Other specific	73.1	76.2	61.5	83.7	66.0	66.1	83.3	65.1	85.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.31a. Oklahoma individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	87,512	14,598	9,960	3,580	10,429	32,048	141	9,690	7,066
Gender									
Male	40,580	6,634	4,114	2,687	3,797	12,970	61	6,173	4,144
Female	46,932	7,964	5,846	893	6,632	19,078	80	3,517	2,922
Total	87,512	14,598	9,960	3,580	10,429	32,048	141	9,690	7,066
Age²									
0 to 11 years	11,189	5,359	1,109	2,002	15	553	1	13	2,137
12 to 14 years	5,450	1,746	668	778	59	1,149	6	18	1,026
15 to 17 years	5,720	1,475	759	464	166	1,853	5	40	958
18 to 20 years	4,129	676	600	153	320	1,795	11	189	385
21 to 24 years	5,338	554	688	43	668	2,501	5	561	318
25 to 29 years	8,374	857	1,177	44	1,142	3,606	17	1,106	425
30 to 34 years	8,232	835	1,080	29	1,313	3,363	16	1,215	381
35 to 39 years	8,040	799	993	18	1,381	3,303	26	1,181	339
40 to 44 years	6,602	602	695	15	1,258	2,718	12	1,071	231
45 to 49 years	6,716	530	630	10	1,213	2,974	17	1,097	245
50 to 54 years	6,439	497	588	12	1,088	2,953	12	1,085	204
55 to 59 years	5,868	349	529	5	991	2,769	5	1,030	190
60 to 64 years	3,382	199	286	3	541	1,559	5	668	121
65 years and older	2,033	120	158	4	274	952	3	416	106
Total	87,512	14,598	9,960	3,580	10,429	32,048	141	9,690	7,066
Race									
American Indian or Alaska Native	7,437	1,423	781	345	728	2,741	13	765	641
Asian ³	484	58	66	14	48	164	--	111	23
Black or African American	11,502	1,707	969	455	1,049	3,986	14	2,399	923
Native Hawaiian or Other Pacific Islander	343	62	37	9	34	127	--	46	28
White	62,856	10,000	7,469	2,413	8,095	23,814	97	6,055	4,913
Other	4,886	1,348	637	344	474	1,216	17	313	537
Total	87,508	14,598	9,959	3,580	10,428	32,048	141	9,689	7,065

Continued. See notes at end of table.

Table 10.31a. Oklahoma individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	80,755	13,059	9,014	3,249	9,951	29,830	132	9,182	6,338
Hispanic or Latino origin	6,745	1,539	945	331	477	2,214	9	504	726
Total	87,500	14,598	9,959	3,580	10,428	32,044	141	9,686	7,064
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	1,528	198	153	3	223	578	2	306	65
Not a veteran	56,600	5,033	6,219	283	9,195	25,045	110	8,587	2,128
Total	58,128	5,231	6,372	286	9,418	25,623	112	8,893	2,193

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.31b. Oklahoma individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	46.4	45.4	41.3	75.1	36.4	40.5	43.3	63.7	58.6	
Female	53.6	54.6	58.7	24.9	63.6	59.5	56.7	36.3	41.4	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	12.8	36.7	11.1	55.9	†	1.7	†	†	30.2	
12 to 14 years	6.2	12.0	6.7	21.7	†	3.6	4.3	†	14.5	
15 to 17 years	6.5	10.1	7.6	13.0	1.6	5.8	3.5	†	13.6	
18 to 20 years	4.7	4.6	6.0	4.3	3.1	5.6	7.8	2.0	5.4	
21 to 24 years	6.1	3.8	6.9	1.2	6.4	7.8	3.5	5.8	4.5	
25 to 29 years	9.6	5.9	11.8	1.2	11.0	11.3	12.1	11.4	6.0	
30 to 34 years	9.4	5.7	10.8	†	12.6	10.5	11.3	12.5	5.4	
35 to 39 years	9.2	5.5	10.0	†	13.2	10.3	18.4	12.2	4.8	
40 to 44 years	7.5	4.1	7.0	†	12.1	8.5	8.5	11.1	3.3	
45 to 49 years	7.7	3.6	6.3	†	11.6	9.3	12.1	11.3	3.5	
50 to 54 years	7.4	3.4	5.9	†	10.4	9.2	8.5	11.2	2.9	
55 to 59 years	6.7	2.4	5.3	†	9.5	8.6	3.5	10.6	2.7	
60 to 64 years	3.9	1.4	2.9	†	5.2	4.9	3.5	6.9	1.7	
65 years and older	2.3	†	1.6	†	2.6	3.0	2.1	4.3	1.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	8.5	9.7	7.8	9.6	7.0	8.6	9.2	7.9	9.1	
Asian ³	†	†	†	†	†	†	--	1.1	†	
Black or African American	13.1	11.7	9.7	12.7	10.1	12.4	9.9	24.8	13.1	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†	
White	71.8	68.5	75.0	67.4	77.6	74.3	68.8	62.5	69.5	
Other	5.6	9.2	6.4	9.6	4.5	3.8	12.1	3.2	7.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.31b. Oklahoma individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	92.3	89.5	90.5	90.8	95.4	93.1	93.6	94.8	89.7
Hispanic or Latino origin	7.7	10.5	9.5	9.2	4.6	6.9	6.4	5.2	10.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	2.6	3.8	2.4	1.0	2.4	2.3	1.8	3.4	3.0
Not a veteran	97.4	96.2	97.6	99.0	97.6	97.7	98.2	96.6	97.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.32a. Oregon individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	134,637	42,073	25,243	7,672	8,850	25,810	691	11,182	13,116
Gender									
Male	57,095	16,348	8,805	5,300	3,202	9,247	200	7,131	6,862
Female	77,540	25,725	16,438	2,372	5,648	16,563	490	4,050	6,254
Total	134,635	42,073	25,243	7,672	8,850	25,810	690	11,181	13,116
Age²									
0 to 11 years	20,989	10,855	2,897	3,609	23	559	--	11	3,035
12 to 14 years	9,180	3,745	1,537	1,173	36	1,299	1	18	1,371
15 to 17 years	11,081	3,483	2,237	757	177	2,705	7	88	1,627
18 to 20 years	6,555	1,726	1,454	270	271	1,691	32	333	778
21 to 24 years	7,113	1,759	1,585	188	529	1,653	56	682	661
25 to 29 years	12,829	3,362	3,021	345	1,122	2,687	118	1,158	1,016
30 to 34 years	12,635	3,494	2,893	373	1,138	2,564	105	1,225	843
35 to 39 years	11,188	3,191	2,331	279	1,050	2,295	101	1,258	683
40 to 44 years	8,743	2,496	1,761	159	916	1,781	74	1,018	538
45 to 49 years	8,568	2,277	1,577	158	907	1,920	62	1,108	559
50 to 54 years	8,186	2,026	1,348	103	883	1,931	53	1,291	551
55 to 59 years	7,720	1,780	1,180	76	820	1,986	41	1,261	576
60 to 64 years	5,460	1,159	818	58	543	1,507	24	921	430
65 years and older	4,390	720	604	124	435	1,232	17	810	448
Total	134,637	42,073	25,243	7,672	8,850	25,810	691	11,182	13,116
Race									
American Indian or Alaska Native	4,390	1,640	840	209	232	769	24	305	371
Asian ³	2,069	440	386	78	92	506	7	367	193
Black or African American	4,923	1,510	804	283	262	902	21	754	387
Native Hawaiian or Other Pacific Islander	461	152	59	31	25	98	1	66	29
White	101,913	31,264	18,908	5,915	7,363	19,257	568	8,729	9,909
Other	5,060	1,822	971	207	238	1,064	21	286	451
Total	118,816	36,828	21,968	6,723	8,212	22,596	642	10,507	11,340

Continued. See notes at end of table.

Table 10.32a. Oregon individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	97,386	28,797	17,869	5,284	7,338	18,730	566	9,685	9,117
Hispanic or Latino origin	14,904	5,344	2,586	930	652	2,643	45	1,026	1,678
Total	112,290	34,141	20,455	6,214	7,990	21,373	611	10,711	10,795
Specific Hispanic or Latino origin									
Mexican	2,521	887	430	114	125	451	5	268	241
Puerto Rican	193	58	45	4	15	31	--	29	11
Cuban	264	125	32	21	6	21	--	30	29
Other specific	1,354	462	236	65	116	242	5	114	114
Total	4,332	1,532	743	204	262	745	10	441	395
Veteran status⁴									
Veteran	2,301	609	287	35	265	496	27	441	141
Not a veteran	61,831	15,981	11,023	1,282	6,375	13,784	520	8,406	4,460
Total	64,132	16,590	11,310	1,317	6,640	14,280	547	8,847	4,601

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.32b. Oregon individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	42.4	38.9	34.9	69.1	36.2	35.8	29.0	63.8	52.3
Female	57.6	61.1	65.1	30.9	63.8	64.2	71.0	36.2	47.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	15.6	25.8	11.5	47.0	†	2.2	--	†	23.1
12 to 14 years	6.8	8.9	6.1	15.3	†	5.0	†	†	10.5
15 to 17 years	8.2	8.3	8.9	9.9	2.0	10.5	1.0	†	12.4
18 to 20 years	4.9	4.1	5.8	3.5	3.1	6.6	4.6	3.0	5.9
21 to 24 years	5.3	4.2	6.3	2.5	6.0	6.4	8.1	6.1	5.0
25 to 29 years	9.5	8.0	12.0	4.5	12.7	10.4	17.1	10.4	7.7
30 to 34 years	9.4	8.3	11.5	4.9	12.9	9.9	15.2	11.0	6.4
35 to 39 years	8.3	7.6	9.2	3.6	11.9	8.9	14.6	11.3	5.2
40 to 44 years	6.5	5.9	7.0	2.1	10.4	6.9	10.7	9.1	4.1
45 to 49 years	6.4	5.4	6.2	2.1	10.2	7.4	9.0	9.9	4.3
50 to 54 years	6.1	4.8	5.3	1.3	10.0	7.5	7.7	11.5	4.2
55 to 59 years	5.7	4.2	4.7	†	9.3	7.7	5.9	11.3	4.4
60 to 64 years	4.1	2.8	3.2	†	6.1	5.8	3.5	8.2	3.3
65 years and older	3.3	1.7	2.4	1.6	4.9	4.8	2.5	7.2	3.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	3.7	4.5	3.8	3.1	2.8	3.4	3.7	2.9	3.3
Asian ³	1.7	1.2	1.8	1.2	1.1	2.2	1.1	3.5	1.7
Black or African American	4.1	4.1	3.7	4.2	3.2	4.0	3.3	7.2	3.4
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	85.8	84.9	86.1	88.0	89.7	85.2	88.5	83.1	87.4
Other	4.3	4.9	4.4	3.1	2.9	4.7	3.3	2.7	4.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.32b. Oregon individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	86.7	84.3	87.4	85.0	91.8	87.6	92.6	90.4	84.5
Hispanic or Latino origin	13.3	15.7	12.6	15.0	8.2	12.4	7.4	9.6	15.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	58.2	57.9	57.9	55.9	47.7	60.5	50.0	60.8	61.0
Puerto Rican	4.5	3.8	6.1	2.0	5.7	4.2	--	6.6	2.8
Cuban	6.1	8.2	4.3	10.3	2.3	2.8	--	6.8	7.3
Other specific	31.3	30.2	31.8	31.9	44.3	32.5	50.0	25.9	28.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	3.6	3.7	2.5	2.7	4.0	3.5	4.9	5.0	3.1
Not a veteran	96.4	96.3	97.5	97.3	96.0	96.5	95.1	95.0	96.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.33a. Pennsylvania individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	487,399	73,952	53,963	60,220	53,537	129,357	1,714	38,221	76,435
Gender									
Male	228,825	31,708	20,032	41,980	20,181	46,085	735	22,287	45,817
Female	258,574	42,244	33,931	18,240	33,356	83,272	979	15,934	30,618
Total	487,399	73,952	53,963	60,220	53,537	129,357	1,714	38,221	76,435
Age²									
0 to 11 years	86,502	22,437	5,237	31,245	456	1,940	8	92	25,087
12 to 14 years	38,186	8,363	3,262	11,591	713	4,461	22	120	9,654
15 to 17 years	37,600	7,102	3,928	7,438	1,608	8,050	35	288	9,151
18 to 20 years	22,217	3,159	2,927	3,042	2,193	6,060	91	649	4,096
21 to 24 years	25,047	3,282	3,646	1,624	3,538	7,881	172	1,652	3,252
25 to 29 years	39,800	5,390	6,115	1,685	6,446	12,760	242	3,079	4,083
30 to 34 years	38,412	4,944	6,018	1,195	6,661	12,638	236	3,248	3,472
35 to 39 years	35,248	4,256	5,127	861	6,542	11,901	169	3,421	2,971
40 to 44 years	29,487	3,364	3,876	548	5,405	10,539	155	3,190	2,410
45 to 49 years	32,188	3,199	3,622	379	5,848	12,548	145	3,911	2,536
50 to 54 years	34,431	3,079	3,616	292	5,722	13,910	133	4,942	2,737
55 to 59 years	30,409	2,482	3,042	200	4,367	12,401	109	5,437	2,371
60 to 64 years	19,653	1,481	1,879	90	2,443	7,877	86	4,141	1,656
65 years and older	18,218	1,414	1,668	30	1,595	6,391	111	4,050	2,959
Total	487,398	73,952	53,963	60,220	53,537	129,357	1,714	38,220	76,435
Race									
American Indian or Alaska Native	1,004	192	110	87	128	247	4	69	167
Asian ³	4,504	729	411	326	271	1,404	10	605	748
Black or African American	102,831	17,882	6,714	13,505	8,973	24,831	298	13,225	17,403
Native Hawaiian or Other Pacific Islander	257	43	38	24	16	86	--	24	26
White	322,268	46,869	40,709	36,697	40,289	85,471	1,266	21,447	49,520
Other	51,198	7,260	5,258	9,089	3,468	15,652	113	2,596	7,762
Total	482,062	72,975	53,240	59,728	53,145	127,691	1,691	37,966	75,626

Continued. See notes at end of table.

Table 10.33a. Pennsylvania individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	425,978	65,148	47,794	49,812	49,157	109,957	1,543	35,003	67,564
Hispanic or Latino origin	60,171	8,703	6,072	10,355	4,289	19,188	114	2,713	8,737
Total	486,149	73,851	53,866	60,167	53,446	129,145	1,657	37,716	76,301
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.33b. Pennsylvania individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	46.9	42.9	37.1	69.7	37.7	35.6	42.9	58.3	59.9
Female	53.1	57.1	62.9	30.3	62.3	64.4	57.1	41.7	40.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	17.7	30.3	9.7	51.9	†	1.5	†	†	32.8
12 to 14 years	7.8	11.3	6.0	19.2	1.3	3.4	1.3	†	12.6
15 to 17 years	7.7	9.6	7.3	12.4	3.0	6.2	2.0	†	12.0
18 to 20 years	4.6	4.3	5.4	5.1	4.1	4.7	5.3	1.7	5.4
21 to 24 years	5.1	4.4	6.8	2.7	6.6	6.1	10.0	4.3	4.3
25 to 29 years	8.2	7.3	11.3	2.8	12.0	9.9	14.1	8.1	5.3
30 to 34 years	7.9	6.7	11.2	2.0	12.4	9.8	13.8	8.5	4.5
35 to 39 years	7.2	5.8	9.5	1.4	12.2	9.2	9.9	9.0	3.9
40 to 44 years	6.0	4.5	7.2	†	10.1	8.1	9.0	8.3	3.2
45 to 49 years	6.6	4.3	6.7	†	10.9	9.7	8.5	10.2	3.3
50 to 54 years	7.1	4.2	6.7	†	10.7	10.8	7.8	12.9	3.6
55 to 59 years	6.2	3.4	5.6	†	8.2	9.6	6.4	14.2	3.1
60 to 64 years	4.0	2.0	3.5	†	4.6	6.1	5.0	10.8	2.2
65 years and older	3.7	1.9	3.1	†	3.0	4.9	6.5	10.6	3.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†
Asian ³	†	†	†	†	†	1.1	†	1.6	†
Black or African American	21.3	24.5	12.6	22.6	16.9	19.4	17.6	34.8	23.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†
White	66.9	64.2	76.5	61.4	75.8	66.9	74.9	56.5	65.5
Other	10.6	9.9	9.9	15.2	6.5	12.3	6.7	6.8	10.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.33b. Pennsylvania individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	87.6	88.2	88.7	82.8	92.0	85.1	93.1	92.8	88.5
Hispanic or Latino origin	12.4	11.8	11.3	17.2	8.0	14.9	6.9	7.2	11.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.34a. Puerto Rico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	2,669	105	101	193	180	718	19	670	683
Gender									
Male	1,568	55	47	148	70	278	14	432	524
Female	1,101	50	54	45	110	440	5	238	159
Total	2,669	105	101	193	180	718	19	670	683
Age²									
0 to 11 years	235	25	10	98	2	29	--	--	71
12 to 14 years	196	22	5	41	3	71	1	1	52
15 to 17 years	209	23	3	29	11	73	1	1	68
18 to 20 years	74	6	2	14	4	9	--	5	34
21 to 24 years	103	3	3	2	7	15	--	33	40
25 to 29 years	173	3	6	3	16	33	3	47	62
30 to 34 years	180	3	7	4	12	28	2	68	56
35 to 39 years	180	1	7	2	19	32	2	59	58
40 to 44 years	170	5	3	--	18	36	--	65	43
45 to 49 years	195	1	5	--	18	48	3	82	38
50 to 54 years	253	2	9	--	24	87	4	86	41
55 to 59 years	243	1	12	--	23	71	1	95	40
60 to 64 years	188	1	9	--	15	69	2	62	30
65 years and older	270	9	20	--	8	117	--	66	50
Total	2,669	105	101	193	180	718	19	670	683
Race									
American Indian or Alaska Native	30	5	1	4	2	4	--	4	10
Asian ³	6	--	--	1	--	1	--	2	2
Black or African American	38	2	--	6	1	7	--	5	17
Native Hawaiian or Other Pacific Islander	3	--	--	--	--	1	--	2	--
White	401	23	8	55	25	64	4	93	129
Other	2,021	66	88	107	139	608	10	514	489
Total	2,499	96	97	173	167	685	14	620	647

Continued. See notes at end of table.

Table 10.34a. Puerto Rico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	18	1	--	--	3	3	--	6	5
Hispanic or Latino origin	2,634	102	101	191	175	711	19	660	675
Total	2,652	103	101	191	178	714	19	666	680
Specific Hispanic or Latino origin									
Mexican	3	--	--	1	--	2	--	--	--
Puerto Rican	2,548	100	99	186	169	698	19	625	652
Cuban	6	--	--	1	1	2	--	1	1
Other specific	40	1	2	2	1	5	--	17	12
Total	2,597	101	101	190	171	707	19	643	665
Veteran status⁴									
Veteran	76	4	1	1	5	17	--	28	20
Not a veteran	1,698	23	80	11	139	492	16	517	420
Total	1,774	27	81	12	144	509	16	545	440

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.34b. Puerto Rico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	58.7	52.4	46.5	76.7	38.9	38.7	73.7	64.5	76.7
Female	41.3	47.6	53.5	23.3	61.1	61.3	26.3	35.5	23.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	8.8	23.8	9.9	50.8	1.1	4.0	--	--	10.4
12 to 14 years	7.3	21.0	5.0	21.2	1.7	9.9	5.3	†	7.6
15 to 17 years	7.8	21.9	3.0	15.0	6.1	10.2	5.3	†	10.0
18 to 20 years	2.8	5.7	2.0	7.3	2.2	1.3	--	†	5.0
21 to 24 years	3.9	2.9	3.0	1.0	3.9	2.1	--	4.9	5.9
25 to 29 years	6.5	2.9	5.9	1.6	8.9	4.6	15.8	7.0	9.1
30 to 34 years	6.7	2.9	6.9	2.1	6.7	3.9	10.5	10.1	8.2
35 to 39 years	6.7	†	6.9	1.0	10.6	4.5	10.5	8.8	8.5
40 to 44 years	6.4	4.8	3.0	--	10.0	5.0	--	9.7	6.3
45 to 49 years	7.3	†	5.0	--	10.0	6.7	15.8	12.2	5.6
50 to 54 years	9.5	1.9	8.9	--	13.3	12.1	21.1	12.8	6.0
55 to 59 years	9.1	†	11.9	--	12.8	9.9	5.3	14.2	5.9
60 to 64 years	7.0	†	8.9	--	8.3	9.6	10.5	9.3	4.4
65 years and older	10.1	8.6	19.8	--	4.4	16.3	--	9.9	7.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	1.2	5.2	1.0	2.3	1.2	†	--	†	1.5
Asian ³	†	--	--	†	--	†	--	†	†
Black or African American	1.5	2.1	--	3.5	†	1.0	--	†	2.6
Native Hawaiian or Other Pacific Islander	†	--	--	--	--	†	--	†	--
White	16.0	24.0	8.2	31.8	15.0	9.3	28.6	15.0	19.9
Other	80.9	68.8	90.7	61.8	83.2	88.8	71.4	82.9	75.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.34b. Puerto Rico individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	†	†	--	--	1.7	†	--	†	†
Hispanic or Latino origin	99.3	99.0	100.0	100.0	98.3	99.6	100.0	99.1	99.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	†	--	--	†	--	†	--	--	--
Puerto Rican	98.1	99.0	98.0	97.9	98.8	98.7	100.0	97.2	98.0
Cuban	†	--	--	†	†	†	--	†	†
Other specific	1.5	†	2.0	1.1	†	†	--	2.6	1.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	4.3	14.8	1.2	8.3	3.5	3.3	--	5.1	4.5
Not a veteran	95.7	85.2	98.8	91.7	96.5	96.7	100.0	94.9	95.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.35a. Rhode Island individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	20,637	2,447	1,671	1,223	2,860	7,445	233	3,414	1,344
Gender									
Male	9,985	1,116	732	951	1,165	2,875	74	2,190	882
Female	10,652	1,331	939	272	1,695	4,570	159	1,224	462
Total	20,637	2,447	1,671	1,223	2,860	7,445	233	3,414	1,344
Age²									
0 to 11 years	1,837	501	242	515	--	80	--	--	499
12 to 14 years	1,070	255	137	280	5	219	--	1	173
15 to 17 years	1,188	242	178	198	23	372	1	14	160
18 to 20 years	877	108	118	114	66	301	5	52	113
21 to 24 years	947	105	89	38	156	319	23	156	61
25 to 29 years	1,441	168	144	24	256	451	32	314	52
30 to 34 years	1,501	167	133	18	266	530	34	319	34
35 to 39 years	1,534	157	100	17	298	584	22	304	52
40 to 44 years	1,369	154	83	2	265	563	23	246	33
45 to 49 years	1,831	165	86	6	359	806	21	357	31
50 to 54 years	2,134	148	101	6	390	1,008	27	413	41
55 to 59 years	2,082	134	101	1	334	1,005	22	442	43
60 to 64 years	1,489	86	75	3	222	673	11	390	29
65 years and older	1,337	57	84	1	220	534	12	406	23
Total	20,637	2,447	1,671	1,223	2,860	7,445	233	3,414	1,344
Race									
American Indian or Alaska Native	178	19	19	8	32	57	2	33	8
Asian ³	238	34	8	7	16	101	--	65	7
Black or African American	2,542	341	151	158	302	895	17	508	170
Native Hawaiian or Other Pacific Islander	43	7	3	2	2	19	1	8	1
White	14,411	1,525	1,218	733	2,227	5,220	189	2,438	861
Other	157	14	7	10	20	52	2	41	11
Total	17,569	1,940	1,406	918	2,599	6,344	211	3,093	1,058

Continued. See notes at end of table.

Table 10.35a. Rhode Island individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	14,858	1,599	1,266	644	2,400	5,158	210	2,724	857
Hispanic or Latino origin	4,746	697	309	461	361	1,997	17	521	383
Total	19,604	2,296	1,575	1,105	2,761	7,155	227	3,245	1,240
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	438	66	34	3	67	146	4	110	8
Not a veteran	14,758	1,287	971	200	2,573	6,075	215	3,006	431
Total	15,196	1,353	1,005	203	2,640	6,221	219	3,116	439

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.35b. Rhode Island individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	48.4	45.6	43.8	77.8	40.7	38.6	31.8	64.1	65.6
Female	51.6	54.4	56.2	22.2	59.3	61.4	68.2	35.9	34.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	8.9	20.5	14.5	42.1	--	1.1	--	--	37.1
12 to 14 years	5.2	10.4	8.2	22.9	†	2.9	--	†	12.9
15 to 17 years	5.8	9.9	10.7	16.2	†	5.0	†	†	11.9
18 to 20 years	4.2	4.4	7.1	9.3	2.3	4.0	2.1	1.5	8.4
21 to 24 years	4.6	4.3	5.3	3.1	5.5	4.3	9.9	4.6	4.5
25 to 29 years	7.0	6.9	8.6	2.0	9.0	6.1	13.7	9.2	3.9
30 to 34 years	7.3	6.8	8.0	1.5	9.3	7.1	14.6	9.3	2.5
35 to 39 years	7.4	6.4	6.0	1.4	10.4	7.8	9.4	8.9	3.9
40 to 44 years	6.6	6.3	5.0	†	9.3	7.6	9.9	7.2	2.5
45 to 49 years	8.9	6.7	5.1	†	12.6	10.8	9.0	10.5	2.3
50 to 54 years	10.3	6.0	6.0	†	13.6	13.5	11.6	12.1	3.1
55 to 59 years	10.1	5.5	6.0	†	11.7	13.5	9.4	12.9	3.2
60 to 64 years	7.2	3.5	4.5	†	7.8	9.0	4.7	11.4	2.2
65 years and older	6.5	2.3	5.0	†	7.7	7.2	5.2	11.9	1.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	1.0	†	1.4	†	1.2	†	†	1.1	†
Asian ³	1.4	1.8	†	†	†	1.6	--	2.1	†
Black or African American	14.5	17.6	10.7	17.2	11.6	14.1	8.1	16.4	16.1
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	82.0	78.6	86.6	79.8	85.7	82.3	89.6	78.8	81.4
Other	†	†	†	1.1	†	†	†	1.3	1.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.35b. Rhode Island individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	75.8	69.6	80.4	58.3	86.9	72.1	92.5	83.9	69.1
Hispanic or Latino origin	24.2	30.4	19.6	41.7	13.1	27.9	7.5	16.1	30.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	2.9	4.9	3.4	1.5	2.5	2.3	1.8	3.5	1.8
Not a veteran	97.1	95.1	96.6	98.5	97.5	97.7	98.2	96.5	98.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.36a. South Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	84,322	10,564	6,100	10,116	10,409	20,051	872	15,733	10,477
Gender									
Male	39,285	4,186	2,277	7,310	3,414	6,271	200	9,197	6,430
Female	45,031	6,378	3,823	2,806	6,995	13,777	672	6,535	4,045
Total	84,316	10,564	6,100	10,116	10,409	20,048	872	15,732	10,475
Age²									
0 to 11 years	12,299	2,699	788	5,848	23	401	--	24	2,516
12 to 14 years	7,458	1,466	628	2,258	69	1,207	3	41	1,786
15 to 17 years	7,925	1,413	853	1,407	196	1,965	21	117	1,953
18 to 20 years	3,371	398	386	296	342	1,085	51	340	473
21 to 24 years	3,849	418	370	73	573	1,129	86	802	398
25 to 29 years	5,861	642	547	74	984	1,629	136	1,313	536
30 to 34 years	5,641	599	478	48	1,013	1,427	122	1,445	509
35 to 39 years	5,620	579	375	31	1,097	1,445	98	1,613	382
40 to 44 years	5,244	509	308	25	1,109	1,461	97	1,376	359
45 to 49 years	5,722	516	291	23	1,233	1,733	79	1,491	356
50 to 54 years	6,316	519	307	12	1,141	2,043	87	1,796	411
55 to 59 years	6,137	403	304	9	1,158	1,928	47	1,957	331
60 to 64 years	4,394	215	221	9	795	1,307	26	1,602	219
65 years and older	4,483	187	244	3	676	1,290	19	1,816	248
Total	84,320	10,563	6,100	10,116	10,409	20,050	872	15,733	10,477
Race									
American Indian or Alaska Native	269	36	18	25	29	72	4	48	37
Asian ³	59	3	1	2	6	14	--	23	10
Black or African American	33,076	3,318	1,203	5,097	2,533	6,515	181	10,022	4,207
Native Hawaiian or Other Pacific Islander	13	2	--	2	5	1	--	1	2
White	44,408	5,860	4,297	3,966	7,396	11,909	645	5,139	5,196
Other	4,158	929	404	644	247	1,032	26	291	585
Total	81,983	10,148	5,923	9,736	10,216	19,543	856	15,524	10,037

Continued. See notes at end of table.

Table 10.36a. South Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	63,507	7,674	4,578	7,398	8,254	15,323	693	11,915	7,672
Hispanic or Latino origin	2,995	592	274	358	250	726	23	373	399
Total	66,502	8,266	4,852	7,756	8,504	16,049	716	12,288	8,071
Specific Hispanic or Latino origin									
Mexican	896	201	90	117	57	218	8	91	114
Puerto Rican	267	49	25	31	34	73	2	27	26
Cuban	363	40	23	15	64	67	2	118	34
Other specific	1,449	300	136	195	91	366	10	129	222
Total	2,975	590	274	358	246	724	22	365	396
Veteran status⁴									
Veteran	813	98	39	4	133	221	6	258	54
Not a veteran	55,777	4,886	3,792	599	9,981	16,245	840	15,270	4,164
Total	56,590	4,984	3,831	603	10,114	16,466	846	15,528	4,218

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.36b. South Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	46.6	39.6	37.3	72.3	32.8	31.3	22.9	58.5	61.4
Female	53.4	60.4	62.7	27.7	67.2	68.7	77.1	41.5	38.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	14.6	25.6	12.9	57.8	†	2.0	--	†	24.0
12 to 14 years	8.8	13.9	10.3	22.3	†	6.0	†	†	17.0
15 to 17 years	9.4	13.4	14.0	13.9	1.9	9.8	2.4	†	18.6
18 to 20 years	4.0	3.8	6.3	2.9	3.3	5.4	5.8	2.2	4.5
21 to 24 years	4.6	4.0	6.1	†	5.5	5.6	9.9	5.1	3.8
25 to 29 years	7.0	6.1	9.0	†	9.5	8.1	15.6	8.3	5.1
30 to 34 years	6.7	5.7	7.8	†	9.7	7.1	14.0	9.2	4.9
35 to 39 years	6.7	5.5	6.1	†	10.5	7.2	11.2	10.3	3.6
40 to 44 years	6.2	4.8	5.0	†	10.7	7.3	11.1	8.7	3.4
45 to 49 years	6.8	4.9	4.8	†	11.8	8.6	9.1	9.5	3.4
50 to 54 years	7.5	4.9	5.0	†	11.0	10.2	10.0	11.4	3.9
55 to 59 years	7.3	3.8	5.0	†	11.1	9.6	5.4	12.4	3.2
60 to 64 years	5.2	2.0	3.6	†	7.6	6.5	3.0	10.2	2.1
65 years and older	5.3	1.8	4.0	†	6.5	6.4	2.2	11.5	2.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†
Asian ³	†	†	†	†	†	†	--	†	†
Black or African American	40.3	32.7	20.3	52.4	24.8	33.3	21.1	64.6	41.9
Native Hawaiian or Other Pacific Islander	†	†	--	†	†	†	--	†	†
White	54.2	57.7	72.5	40.7	72.4	60.9	75.4	33.1	51.8
Other	5.1	9.2	6.8	6.6	2.4	5.3	3.0	1.9	5.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.36b. South Carolina individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	95.5	92.8	94.4	95.4	97.1	95.5	96.8	97.0	95.1
Hispanic or Latino origin	4.5	7.2	5.6	4.6	2.9	4.5	3.2	3.0	4.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	30.1	34.1	32.8	32.7	23.2	30.1	36.4	24.9	28.8
Puerto Rican	9.0	8.3	9.1	8.7	13.8	10.1	9.1	7.4	6.6
Cuban	12.2	6.8	8.4	4.2	26.0	9.3	9.1	32.3	8.6
Other specific	48.7	50.8	49.6	54.5	37.0	50.6	45.5	35.3	56.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	1.4	2.0	1.0	†	1.3	1.3	†	1.7	1.3
Not a veteran	98.6	98.0	99.0	99.3	98.7	98.7	99.3	98.3	98.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.37a. South Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	15,310	3,932	1,561	1,225	1,490	3,800	5	1,442	1,855
Gender									
Male	6,953	1,785	597	876	552	1,243	1	857	1,042
Female	8,357	2,147	964	349	938	2,557	4	585	813
Total	15,310	3,932	1,561	1,225	1,490	3,800	5	1,442	1,855
Age²									
0 to 11 years	2,503	1,452	149	498	18	69	--	--	317
12 to 14 years	1,376	649	87	235	20	193	--	3	189
15 to 17 years	1,307	471	114	171	45	270	--	5	231
18 to 20 years	757	166	92	61	47	237	--	15	139
21 to 24 years	855	144	136	63	72	238	--	51	151
25 to 29 years	1,079	188	165	72	118	321	1	80	134
30 to 34 years	1,081	186	152	41	127	321	--	111	143
35 to 39 years	1,138	154	159	32	181	352	--	142	118
40 to 44 years	841	120	91	14	146	275	--	110	85
45 to 49 years	894	121	87	14	158	289	3	147	75
50 to 54 years	877	97	74	7	160	314	--	161	64
55 to 59 years	970	77	73	11	162	354	--	237	56
60 to 64 years	670	39	71	4	117	235	1	161	42
65 years and older	962	68	111	2	119	332	--	219	111
Total	15,310	3,932	1,561	1,225	1,490	3,800	5	1,442	1,855
Race									
American Indian or Alaska Native	9	--	--	--	--	--	--	3	6
Asian ³	104	41	8	3	5	22	--	12	13
Black or African American	430	192	18	39	21	55	--	56	49
Native Hawaiian or Other Pacific Islander	18	6	1	--	2	6	--	1	2
White	10,959	2,394	1,202	863	1,241	2,846	5	1,118	1,290
Other	3,790	1,299	332	320	221	871	--	252	495
Total	15,310	3,932	1,561	1,225	1,490	3,800	5	1,442	1,855

Continued. See notes at end of table.

Table 10.37a. South Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	14,588	3,692	1,505	1,164	1,437	3,641	5	1,395	1,749
Hispanic or Latino origin	722	240	56	61	53	159	--	47	106
Total	15,310	3,932	1,561	1,225	1,490	3,800	5	1,442	1,855
Specific Hispanic or Latino origin									
Mexican	236	65	20	23	22	53	--	14	39
Puerto Rican	62	19	6	3	3	14	--	4	13
Cuban	15	5	1	1	3	4	--	1	--
Other specific	103	44	6	8	6	18	--	8	13
Total	416	133	33	35	34	89	--	27	65
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.37b. South Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	45.4	45.4	38.2	71.5	37.0	32.7	20.0	59.4	56.2
Female	54.6	54.6	61.8	28.5	63.0	67.3	80.0	40.6	43.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	16.3	36.9	9.5	40.7	1.2	1.8	--	--	17.1
12 to 14 years	9.0	16.5	5.6	19.2	1.3	5.1	--	†	10.2
15 to 17 years	8.5	12.0	7.3	14.0	3.0	7.1	--	†	12.5
18 to 20 years	4.9	4.2	5.9	5.0	3.2	6.2	--	1.0	7.5
21 to 24 years	5.6	3.7	8.7	5.1	4.8	6.3	--	3.5	8.1
25 to 29 years	7.0	4.8	10.6	5.9	7.9	8.4	20.0	5.5	7.2
30 to 34 years	7.1	4.7	9.7	3.3	8.5	8.4	--	7.7	7.7
35 to 39 years	7.4	3.9	10.2	2.6	12.1	9.3	--	9.8	6.4
40 to 44 years	5.5	3.1	5.8	1.1	9.8	7.2	--	7.6	4.6
45 to 49 years	5.8	3.1	5.6	1.1	10.6	7.6	60.0	10.2	4.0
50 to 54 years	5.7	2.5	4.7	†	10.7	8.3	--	11.2	3.5
55 to 59 years	6.3	2.0	4.7	†	10.9	9.3	--	16.4	3.0
60 to 64 years	4.4	†	4.5	†	7.9	6.2	20.0	11.2	2.3
65 years and older	6.3	1.7	7.1	†	8.0	8.7	--	15.2	6.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	--	--	--	--	--	--	†	†
Asian ³	†	1.0	†	†	†	†	--	†	†
Black or African American	2.8	4.9	1.2	3.2	1.4	1.4	--	3.9	2.6
Native Hawaiian or Other Pacific Islander	†	†	†	--	†	†	--	†	†
White	71.6	60.9	77.0	70.4	83.3	74.9	100.0	77.5	69.5
Other	24.8	33.0	21.3	26.1	14.8	22.9	--	17.5	26.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.37b. South Dakota individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	95.3	93.9	96.4	95.0	96.4	95.8	100.0	96.7	94.3
Hispanic or Latino origin	4.7	6.1	3.6	5.0	3.6	4.2	--	3.3	5.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	56.7	48.9	60.6	65.7	64.7	59.6	--	51.9	60.0
Puerto Rican	14.9	14.3	18.2	8.6	8.8	15.7	--	14.8	20.0
Cuban	3.6	3.8	3.0	2.9	8.8	4.5	--	3.7	--
Other specific	24.8	33.1	18.2	22.9	17.6	20.2	--	29.6	20.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	--	100.0	100.0
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.38a. Tennessee individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals²	64,037	6,604	7,804	2,728	12,094	22,288	479	7,150	4,890
Gender									
Male	30,613	2,759	3,208	1,850	5,077	9,851	202	4,806	2,860
Female	33,345	3,841	4,588	876	7,003	12,415	277	2,325	2,020
Total	63,958	6,600	7,796	2,726	12,080	22,266	479	7,131	4,880
Age³									
0 to 11 years	3,670	806	314	1,304	21	134	1	14	1,076
12 to 14 years	2,105	359	249	542	30	362	1	11	551
15 to 17 years	1,994	305	274	291	93	593	2	34	402
18 to 20 years	2,120	227	299	91	352	696	21	208	226
21 to 24 years	4,433	436	598	92	866	1,510	45	590	296
25 to 29 years	7,601	757	1,026	124	1,685	2,522	73	1,000	414
30 to 34 years	7,037	753	867	80	1,664	2,331	75	942	325
35 to 39 years	7,009	715	848	72	1,601	2,488	59	894	332
40 to 44 years	6,042	596	722	41	1,461	2,162	58	753	249
45 to 49 years	6,778	606	787	35	1,535	2,736	57	714	308
50 to 54 years	6,696	521	771	26	1,309	2,981	35	755	298
55 to 59 years	4,986	319	587	13	896	2,301	32	609	229
60 to 64 years	2,602	142	324	14	419	1,132	14	438	119
65 years and older	964	62	138	3	162	340	6	188	65
Total	64,037	6,604	7,804	2,728	12,094	22,288	479	7,150	4,890
Race									
American Indian or Alaska Native	185	22	22	9	40	50	1	28	13
Asian ⁴	6,476	1,044	1,023	556	1,006	1,617	116	361	753
Black or African American	17,863	1,838	1,478	1,328	2,528	4,991	103	3,586	2,011
Native Hawaiian or Other Pacific Islander	32	2	1	4	7	8	--	6	4
White	37,296	3,443	4,976	689	8,193	14,869	250	2,917	1,959
Other	399	40	31	27	76	117	5	72	31
Total	62,251	6,389	7,531	2,613	11,850	21,652	475	6,970	4,771

Continued. See notes at end of table.

Table 10.38a. Tennessee individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	33,646	3,392	3,621	1,784	6,413	9,997	395	4,966	3,078
Hispanic or Latino origin	643	76	71	26	102	204	6	102	56
Total	34,289	3,468	3,692	1,810	6,515	10,201	401	5,068	3,134
Specific Hispanic or Latino origin									
Mexican	22	5	1	--	3	4	1	8	--
Puerto Rican	8	--	--	--	2	2	1	3	--
Cuban	1	--	--	--	--	--	--	1	--
Other specific	37	5	--	--	6	15	1	9	1
Total	68	10	1	--	11	21	3	21	1
Veteran status⁵									
Veteran	254	36	4	--	62	101	7	35	9
Not a veteran	8,603	616	215	15	2,142	2,998	182	1,956	479
Total	8,857	652	219	15	2,204	3,099	189	1,991	488

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Tennessee reports only a subset of the total individuals served through their SMHAs.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.38b. Tennessee individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals²	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	47.9	41.8	41.1	67.9	42.0	44.2	42.2	67.4	58.6
Female	52.1	58.2	58.9	32.1	58.0	55.8	57.8	32.6	41.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age³									
0 to 11 years	5.7	12.2	4.0	47.8	†	†	†	†	22.0
12 to 14 years	3.3	5.4	3.2	19.9	†	1.6	†	†	11.3
15 to 17 years	3.1	4.6	3.5	10.7	†	2.7	†	†	8.2
18 to 20 years	3.3	3.4	3.8	3.3	2.9	3.1	4.4	2.9	4.6
21 to 24 years	6.9	6.6	7.7	3.4	7.2	6.8	9.4	8.3	6.1
25 to 29 years	11.9	11.5	13.1	4.5	13.9	11.3	15.2	14.0	8.5
30 to 34 years	11.0	11.4	11.1	2.9	13.8	10.5	15.7	13.2	6.6
35 to 39 years	10.9	10.8	10.9	2.6	13.2	11.2	12.3	12.5	6.8
40 to 44 years	9.4	9.0	9.3	1.5	12.1	9.7	12.1	10.5	5.1
45 to 49 years	10.6	9.2	10.1	1.3	12.7	12.3	11.9	10.0	6.3
50 to 54 years	10.5	7.9	9.9	†	10.8	13.4	7.3	10.6	6.1
55 to 59 years	7.8	4.8	7.5	†	7.4	10.3	6.7	8.5	4.7
60 to 64 years	4.1	2.2	4.2	†	3.5	5.1	2.9	6.1	2.4
65 years and older	1.5	†	1.8	†	1.3	1.5	1.3	2.6	1.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†
Asian ⁴	10.4	16.3	13.6	21.3	8.5	7.5	24.4	5.2	15.8
Black or African American	28.7	28.8	19.6	50.8	21.3	23.1	21.7	51.4	42.2
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†
White	59.9	53.9	66.1	26.4	69.1	68.7	52.6	41.9	41.1
Other	†	†	†	1.0	†	†	1.1	1.0	†
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.38b. Tennessee individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	98.1	97.8	98.1	98.6	98.4	98.0	98.5	98.0	98.2
Hispanic or Latino origin	1.9	2.2	1.9	1.4	1.6	2.0	1.5	2.0	1.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	32.4	50.0	100.0	--	27.3	19.0	33.3	38.1	--
Puerto Rican	11.8	--	--	--	18.2	9.5	33.3	14.3	--
Cuban	1.5	--	--	--	--	--	--	4.8	--
Other specific	54.4	50.0	--	--	54.5	71.4	33.3	42.9	100.0
Total	100.0	100.0	100.0	--	100.0	100.0	100.0	100.0	100.0
Veteran status⁵									
Veteran	2.9	5.5	1.8	--	2.8	3.3	3.7	1.8	1.8
Not a veteran	97.1	94.5	98.2	100.0	97.2	96.7	96.3	98.2	98.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Tennessee reports only a subset of the total individuals served through their SMHAs.

³ Age is based on the individual's age at the midpoint of the reporting period.

⁴ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁵ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.39a. Texas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	268,741	10,894	5,238	28,366	66,516	92,031	297	49,651	15,748
Gender									
Male	133,139	5,309	2,428	20,605	27,214	35,846	195	31,468	10,074
Female	135,601	5,585	2,810	7,761	39,301	56,185	102	18,183	5,674
Total	268,740	10,894	5,238	28,366	66,515	92,031	297	49,651	15,748
Age²									
0 to 11 years	29,405	3,945	825	17,497	335	1,384	--	108	5,311
12 to 14 years	17,626	1,965	692	6,488	753	4,274	--	174	3,280
15 to 17 years	17,448	1,536	927	3,648	1,621	6,365	3	345	3,003
18 to 20 years	11,362	420	278	386	2,972	5,346	23	1,268	669
21 to 24 years	15,865	378	344	97	4,791	6,578	36	3,048	593
25 to 29 years	24,216	548	494	96	7,973	9,117	59	5,304	625
30 to 34 years	24,333	529	383	63	8,341	8,766	37	5,715	499
35 to 39 years	23,722	423	354	36	8,465	8,124	41	5,878	401
40 to 44 years	19,807	342	194	16	6,917	7,192	29	4,810	307
45 to 49 years	20,900	276	186	17	7,008	8,095	25	5,058	235
50 to 54 years	21,873	195	154	8	6,665	8,897	20	5,708	226
55 to 59 years	20,332	165	149	6	5,636	8,577	13	5,597	189
60 to 64 years	12,779	89	103	4	3,129	5,528	8	3,767	151
65 years and older	9,070	83	155	4	1,910	3,788	3	2,868	259
Total	268,738	10,894	5,238	28,366	66,516	92,031	297	49,648	15,748
Race									
American Indian or Alaska Native	865	37	13	58	226	321	2	161	47
Asian ³	2,294	76	47	89	386	664	1	902	129
Black or African American	60,724	2,048	550	7,932	12,663	16,658	63	17,148	3,662
Native Hawaiian or Other Pacific Islander	210	12	3	11	54	79	--	39	12
White	193,025	7,922	4,335	18,339	51,035	70,519	218	29,727	10,930
Other	11,616	799	290	1,937	2,150	3,789	13	1,670	968
Total	268,734	10,894	5,238	28,366	66,514	92,030	297	49,647	15,748

Continued. See notes at end of table.

Table 10.39a. Texas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	200,057	7,332	3,336	17,925	54,388	66,430	229	39,823	10,594
Hispanic or Latino origin	68,665	3,562	1,902	10,441	12,126	25,598	68	9,820	5,148
Total	268,722	10,894	5,238	28,366	66,514	92,028	297	49,643	15,742
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.39b. Texas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	49.5	48.7	46.4	72.6	40.9	38.9	65.7	63.4	64.0	
Female	50.5	51.3	53.6	27.4	59.1	61.1	34.3	36.6	36.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	10.9	36.2	15.8	61.7	†	1.5	--	†	33.7	
12 to 14 years	6.6	18.0	13.2	22.9	1.1	4.6	--	†	20.8	
15 to 17 years	6.5	14.1	17.7	12.9	2.4	6.9	1.0	†	19.1	
18 to 20 years	4.2	3.9	5.3	1.4	4.5	5.8	7.7	2.6	4.2	
21 to 24 years	5.9	3.5	6.6	†	7.2	7.1	12.1	6.1	3.8	
25 to 29 years	9.0	5.0	9.4	†	12.0	9.9	19.9	10.7	4.0	
30 to 34 years	9.1	4.9	7.3	†	12.5	9.5	12.5	11.5	3.2	
35 to 39 years	8.8	3.9	6.8	†	12.7	8.8	13.8	11.8	2.5	
40 to 44 years	7.4	3.1	3.7	†	10.4	7.8	9.8	9.7	1.9	
45 to 49 years	7.8	2.5	3.6	†	10.5	8.8	8.4	10.2	1.5	
50 to 54 years	8.1	1.8	2.9	†	10.0	9.7	6.7	11.5	1.4	
55 to 59 years	7.6	1.5	2.8	†	8.5	9.3	4.4	11.3	1.2	
60 to 64 years	4.8	†	2.0	†	4.7	6.0	2.7	7.6	†	
65 years and older	3.4	†	3.0	†	2.9	4.1	1.0	5.8	1.6	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†	
Asian ³	†	†	†	†	†	†	†	1.8	†	
Black or African American	22.6	18.8	10.5	28.0	19.0	18.1	21.2	34.5	23.3	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†	
White	71.8	72.7	82.8	64.7	76.7	76.6	73.4	59.9	69.4	
Other	4.3	7.3	5.5	6.8	3.2	4.1	4.4	3.4	6.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.39b. Texas individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	74.4	67.3	63.7	63.2	81.8	72.2	77.1	80.2	67.3
Hispanic or Latino origin	25.6	32.7	36.3	36.8	18.2	27.8	22.9	19.8	32.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.40a. Utah individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	48,319	7,028	8,436	1,859	4,168	14,276	561	5,575	6,416
Gender									
Male	22,574	3,495	3,765	1,407	1,500	4,967	182	3,460	3,798
Female	25,745	3,533	4,671	452	2,668	9,309	379	2,115	2,618
Total	48,319	7,028	8,436	1,859	4,168	14,276	561	5,575	6,416
Age²									
0 to 11 years	11,316	3,566	3,146	1,231	52	1,028	1	35	2,257
12 to 14 years	5,328	928	1,095	285	120	1,959	1	52	888
15 to 17 years	4,225	549	776	117	162	1,918	8	107	588
18 to 20 years	1,616	152	237	39	181	611	17	164	215
21 to 24 years	2,249	197	332	29	282	751	21	335	302
25 to 29 years	3,538	280	538	44	507	1,165	47	562	395
30 to 34 years	3,631	331	543	45	500	1,227	63	580	342
35 to 39 years	3,623	292	495	20	577	1,168	68	686	317
40 to 44 years	2,800	209	330	17	406	924	56	600	258
45 to 49 years	2,498	172	273	9	381	797	67	585	214
50 to 54 years	2,298	126	220	9	342	789	56	575	181
55 to 59 years	2,148	84	203	9	289	780	59	552	172
60 to 64 years	1,460	61	113	3	180	523	53	401	126
65 years and older	1,588	81	135	2	189	636	44	340	161
Total	48,318	7,028	8,436	1,859	4,168	14,276	561	5,574	6,416
Race									
American Indian or Alaska Native	1,018	205	146	34	53	334	8	109	129
Asian ³	369	45	51	8	16	115	3	91	40
Black or African American	1,158	202	174	53	73	282	13	215	146
Native Hawaiian or Other Pacific Islander	969	277	157	42	67	249	4	68	105
White	38,309	5,372	6,802	1,474	3,537	11,547	484	4,227	4,866
Other	5,260	831	1,019	235	287	1,508	35	589	756
Total	47,083	6,932	8,349	1,846	4,033	14,035	547	5,299	6,042

Continued. See notes at end of table.

Table 10.40a. Utah individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	39,590	5,407	6,829	1,471	3,729	11,837	516	4,830	4,971
Hispanic or Latino origin	7,489	1,499	1,457	364	323	2,147	39	587	1,073
Total	47,079	6,906	8,286	1,835	4,052	13,984	555	5,417	6,044
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	668	57	80	4	103	223	21	123	57
Not a veteran	25,728	1,887	3,285	221	3,659	8,941	520	4,957	2,258
Total	26,396	1,944	3,365	225	3,762	9,164	541	5,080	2,315

‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.40b. Utah individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	46.7	49.7	44.6	75.7	36.0	34.8	32.4	62.1	59.2	
Female	53.3	50.3	55.4	24.3	64.0	65.2	67.6	37.9	40.8	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	23.4	50.7	37.3	66.2	1.2	7.2	†	†	35.2	
12 to 14 years	11.0	13.2	13.0	15.3	2.9	13.7	†	†	13.8	
15 to 17 years	8.7	7.8	9.2	6.3	3.9	13.4	1.4	1.9	9.2	
18 to 20 years	3.3	2.2	2.8	2.1	4.3	4.3	3.0	2.9	3.4	
21 to 24 years	4.7	2.8	3.9	1.6	6.8	5.3	3.7	6.0	4.7	
25 to 29 years	7.3	4.0	6.4	2.4	12.2	8.2	8.4	10.1	6.2	
30 to 34 years	7.5	4.7	6.4	2.4	12.0	8.6	11.2	10.4	5.3	
35 to 39 years	7.5	4.2	5.9	1.1	13.8	8.2	12.1	12.3	4.9	
40 to 44 years	5.8	3.0	3.9	†	9.7	6.5	10.0	10.8	4.0	
45 to 49 years	5.2	2.4	3.2	†	9.1	5.6	11.9	10.5	3.3	
50 to 54 years	4.8	1.8	2.6	†	8.2	5.5	10.0	10.3	2.8	
55 to 59 years	4.4	1.2	2.4	†	6.9	5.5	10.5	9.9	2.7	
60 to 64 years	3.0	†	1.3	†	4.3	3.7	9.4	7.2	2.0	
65 years and older	3.3	1.2	1.6	†	4.5	4.5	7.8	6.1	2.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	2.2	3.0	1.7	1.8	1.3	2.4	1.5	2.1	2.1	
Asian ³	†	†	†	†	†	†	†	1.7	†	
Black or African American	2.5	2.9	2.1	2.9	1.8	2.0	2.4	4.1	2.4	
Native Hawaiian or Other Pacific Islander	2.1	4.0	1.9	2.3	1.7	1.8	†	1.3	1.7	
White	81.4	77.5	81.5	79.8	87.7	82.3	88.5	79.8	80.5	
Other	11.2	12.0	12.2	12.7	7.1	10.7	6.4	11.1	12.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.40b. Utah individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	84.1	78.3	82.4	80.2	92.0	84.6	93.0	89.2	82.2
Hispanic or Latino origin	15.9	21.7	17.6	19.8	8.0	15.4	7.0	10.8	17.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	2.5	2.9	2.4	1.8	2.7	2.4	3.9	2.4	2.5
Not a veteran	97.5	97.1	97.6	98.2	97.3	97.6	96.1	97.6	97.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.41a. Vermont individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	29,028	7,648	2,699	1,515	1,085	3,684	355	1,640	10,402
Gender									
Male	15,137	3,501	1,180	1,095	504	1,410	85	1,027	6,335
Female	13,844	4,124	1,512	420	580	2,263	270	613	4,062
Total	28,981	7,625	2,692	1,515	1,084	3,673	355	1,640	10,397
Age²									
0 to 11 years	4,429	2,235	482	629	8	67	--	2	1,006
12 to 14 years	2,183	851	334	330	7	222	--	3	436
15 to 17 years	2,573	890	347	252	25	420	1	11	627
18 to 20 years	1,700	444	193	70	45	300	19	37	592
21 to 24 years	1,905	391	164	34	56	277	24	58	901
25 to 29 years	2,490	429	186	41	86	274	44	113	1,317
30 to 34 years	2,361	377	150	44	77	297	34	131	1,251
35 to 39 years	2,019	360	155	37	94	233	34	151	955
40 to 44 years	1,516	303	105	22	103	195	34	119	635
45 to 49 years	1,627	308	119	13	109	259	46	142	631
50 to 54 years	1,647	305	129	10	101	254	31	174	643
55 to 59 years	1,568	266	98	18	137	261	45	220	523
60 to 64 years	1,172	155	74	8	98	236	23	211	367
65 years and older	1,838	334	163	7	139	389	20	268	518
Total	29,028	7,648	2,699	1,515	1,085	3,684	355	1,640	10,402
Race									
American Indian or Alaska Native	232	95	31	16	7	24	5	16	38
Asian ³	181	65	18	6	6	35	--	21	30
Black or African American	516	217	42	56	11	58	5	45	82
Native Hawaiian or Other Pacific Islander	--	--	--	--	--	--	--	--	--
White	18,689	6,113	2,223	1,239	934	3,045	308	1,367	3,460
Other	538	226	57	50	23	62	5	24	91
Total	20,156	6,716	2,371	1,367	981	3,224	323	1,473	3,701

Continued. See notes at end of table.

Table 10.41a. Vermont individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	17,611	5,785	2,120	1,199	890	2,788	307	1,389	3,133
Hispanic or Latino origin	592	230	59	58	18	85	8	25	109
Total	18,203	6,015	2,179	1,257	908	2,873	315	1,414	3,242
Specific Hispanic or Latino origin									
Mexican	111	42	15	6	2	21	2	2	21
Puerto Rican	119	47	12	9	2	18	2	6	23
Cuban	20	7	2	5	--	2	--	1	3
Other specific	164	74	14	16	6	15	2	6	31
Total	414	170	43	36	10	56	6	15	78
Veteran status⁴									
Veteran	488	145	36	6	46	87	10	88	70
Not a veteran	10,432	2,615	1,199	244	814	2,196	297	1,240	1,827
Total	10,920	2,760	1,235	250	860	2,283	307	1,328	1,897

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.41b. Vermont individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	52.2	45.9	43.8	72.3	46.5	38.4	23.9	62.6	60.9	
Female	47.8	54.1	56.2	27.7	53.5	61.6	76.1	37.4	39.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	15.3	29.2	17.9	41.5	†	1.8	--	†	9.7	
12 to 14 years	7.5	11.1	12.4	21.8	†	6.0	--	†	4.2	
15 to 17 years	8.9	11.6	12.9	16.6	2.3	11.4	†	†	6.0	
18 to 20 years	5.9	5.8	7.2	4.6	4.1	8.1	5.4	2.3	5.7	
21 to 24 years	6.6	5.1	6.1	2.2	5.2	7.5	6.8	3.5	8.7	
25 to 29 years	8.6	5.6	6.9	2.7	7.9	7.4	12.4	6.9	12.7	
30 to 34 years	8.1	4.9	5.6	2.9	7.1	8.1	9.6	8.0	12.0	
35 to 39 years	7.0	4.7	5.7	2.4	8.7	6.3	9.6	9.2	9.2	
40 to 44 years	5.2	4.0	3.9	1.5	9.5	5.3	9.6	7.3	6.1	
45 to 49 years	5.6	4.0	4.4	†	10.0	7.0	13.0	8.7	6.1	
50 to 54 years	5.7	4.0	4.8	†	9.3	6.9	8.7	10.6	6.2	
55 to 59 years	5.4	3.5	3.6	1.2	12.6	7.1	12.7	13.4	5.0	
60 to 64 years	4.0	2.0	2.7	†	9.0	6.4	6.5	12.9	3.5	
65 years and older	6.3	4.4	6.0	†	12.8	10.6	5.6	16.3	5.0	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	1.2	1.4	1.3	1.2	†	†	1.5	1.1	1.0	
Asian ³	†	†	†	†	†	1.1	--	1.4	†	
Black or African American	2.6	3.2	1.8	4.1	1.1	1.8	1.5	3.1	2.2	
Native Hawaiian or Other Pacific Islander	--	--	--	--	--	--	--	--	--	
White	92.7	91.0	93.8	90.6	95.2	94.4	95.4	92.8	93.5	
Other	2.7	3.4	2.4	3.7	2.3	1.9	1.5	1.6	2.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.41b. Vermont individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

<i>Selected characteristics</i>	<i>Total</i>	<i>Mental health diagnosis¹</i>							
		<i>Trauma- and stressor-related disorders</i>	<i>Anxiety disorders</i>	<i>ADHD</i>	<i>Mood disorders</i>		<i>Personality disorders</i>	<i>Schizophrenia, other psychotic disorders</i>	<i>Other mental disorders¹</i>
					<i>Bipolar disorders</i>	<i>Depressive disorders</i>			
Ethnicity									
Not of Hispanic or Latino origin	96.7	96.2	97.3	95.4	98.0	97.0	97.5	98.2	96.6
Hispanic or Latino origin	3.3	3.8	2.7	4.6	2.0	3.0	2.5	1.8	3.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	26.8	24.7	34.9	16.7	20.0	37.5	33.3	13.3	26.9
Puerto Rican	28.7	27.6	27.9	25.0	20.0	32.1	33.3	40.0	29.5
Cuban	4.8	4.1	4.7	13.9	--	3.6	--	6.7	3.8
Other specific	39.6	43.5	32.6	44.4	60.0	26.8	33.3	40.0	39.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	4.5	5.3	2.9	2.4	5.3	3.8	3.3	6.6	3.7
Not a veteran	95.5	94.7	97.1	97.6	94.7	96.2	96.7	93.4	96.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.42a. Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	117,261	15,114	14,530	11,621	13,368	25,589	3,583	17,143	16,313
Gender									
Male	58,304	6,794	5,824	8,158	5,319	10,200	1,679	10,584	9,746
Female	58,937	8,308	8,705	3,462	8,047	15,389	1,904	6,559	6,563
Total	117,241	15,102	14,529	11,620	13,366	25,589	3,583	17,143	16,309
Age²									
0 to 11 years	13,922	3,539	1,459	5,222	103	524	3	41	3,031
12 to 14 years	8,685	1,686	1,131	2,384	168	1,428	5	59	1,824
15 to 17 years	10,220	1,672	1,470	1,750	357	2,503	36	149	2,283
18 to 20 years	5,122	674	771	587	469	1,359	92	343	827
21 to 24 years	6,122	715	850	396	838	1,572	210	900	641
25 to 29 years	9,366	1,043	1,308	361	1,494	2,211	338	1,628	983
30 to 34 years	9,349	1,054	1,197	296	1,514	2,286	392	1,705	905
35 to 39 years	8,894	995	1,167	203	1,556	2,025	436	1,699	813
40 to 44 years	7,573	806	997	114	1,292	1,907	343	1,442	672
45 to 49 years	8,263	820	1,030	103	1,375	2,143	418	1,560	814
50 to 54 years	9,036	798	1,059	84	1,419	2,444	419	1,958	855
55 to 59 years	8,437	633	857	58	1,182	2,280	373	2,181	873
60 to 64 years	5,825	342	613	38	784	1,467	281	1,604	696
65 years and older	6,442	334	621	25	817	1,440	237	1,874	1,094
Total	117,256	15,111	14,530	11,621	13,368	25,589	3,583	17,143	16,311
Race									
American Indian or Alaska Native	373	71	43	22	49	103	8	41	36
Asian ³	1,349	128	133	38	143	292	38	424	153
Black or African American	32,525	3,357	2,248	3,407	3,104	6,190	921	8,046	5,252
Native Hawaiian or Other Pacific Islander	89	9	12	14	8	13	3	23	7
White	71,576	9,404	10,615	6,617	9,222	16,517	2,448	7,497	9,256
Other	10,284	1,886	1,295	1,376	781	2,242	154	1,062	1,488
Total	116,196	14,855	14,346	11,474	13,307	25,357	3,572	17,093	16,192

Continued. See notes at end of table.

Table 10.42a. Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	106,419	13,038	13,021	10,403	12,576	23,100	3,421	16,080	14,780
Hispanic or Latino origin	8,239	1,570	1,098	879	592	1,893	134	855	1,218
Total	114,658	14,608	14,119	11,282	13,168	24,993	3,555	16,935	15,998
Specific Hispanic or Latino origin									
Mexican	966	213	127	135	59	222	11	69	130
Puerto Rican	979	130	145	131	90	236	25	102	120
Cuban	112	13	14	14	19	26	--	18	8
Other specific	4,204	705	557	386	312	954	77	532	681
Total	6,261	1,061	843	666	480	1,438	113	721	939
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.42b. Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	49.7	45.0	40.1	70.2	39.8	39.9	46.9	61.7	59.8	
Female	50.3	55.0	59.9	29.8	60.2	60.1	53.1	38.3	40.2	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	11.9	23.4	10.0	44.9	†	2.0	†	†	18.6	
12 to 14 years	7.4	11.2	7.8	20.5	1.3	5.6	†	†	11.2	
15 to 17 years	8.7	11.1	10.1	15.1	2.7	9.8	1.0	†	14.0	
18 to 20 years	4.4	4.5	5.3	5.1	3.5	5.3	2.6	2.0	5.1	
21 to 24 years	5.2	4.7	5.8	3.4	6.3	6.1	5.9	5.2	3.9	
25 to 29 years	8.0	6.9	9.0	3.1	11.2	8.6	9.4	9.5	6.0	
30 to 34 years	8.0	7.0	8.2	2.5	11.3	8.9	10.9	9.9	5.5	
35 to 39 years	7.6	6.6	8.0	1.7	11.6	7.9	12.2	9.9	5.0	
40 to 44 years	6.5	5.3	6.9	†	9.7	7.5	9.6	8.4	4.1	
45 to 49 years	7.0	5.4	7.1	†	10.3	8.4	11.7	9.1	5.0	
50 to 54 years	7.7	5.3	7.3	†	10.6	9.6	11.7	11.4	5.2	
55 to 59 years	7.2	4.2	5.9	†	8.8	8.9	10.4	12.7	5.4	
60 to 64 years	5.0	2.3	4.2	†	5.9	5.7	7.8	9.4	4.3	
65 years and older	5.5	2.2	4.3	†	6.1	5.6	6.6	10.9	6.7	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	†	†	†	
Asian ³	1.2	†	†	†	1.1	1.2	1.1	2.5	†	
Black or African American	28.0	22.6	15.7	29.7	23.3	24.4	25.8	47.1	32.4	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	61.6	63.3	74.0	57.7	69.3	65.1	68.5	43.9	57.2	
Other	8.9	12.7	9.0	12.0	5.9	8.8	4.3	6.2	9.2	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.42b. Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	92.8	89.3	92.2	92.2	95.5	92.4	96.2	95.0	92.4
Hispanic or Latino origin	7.2	10.7	7.8	7.8	4.5	7.6	3.8	5.0	7.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	15.4	20.1	15.1	20.3	12.3	15.4	9.7	9.6	13.8
Puerto Rican	15.6	12.3	17.2	19.7	18.8	16.4	22.1	14.1	12.8
Cuban	1.8	1.2	1.7	2.1	4.0	1.8	- -	2.5	†
Other specific	67.1	66.4	66.1	58.0	65.0	66.3	68.1	73.8	72.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.43a. Washington individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	269,476	56,177	56,200	20,445	18,161	69,924	4,059	20,155	24,355
Gender									
Male	118,595	21,425	20,649	14,126	7,026	26,226	1,891	12,966	14,286
Female	150,608	34,722	35,523	6,316	11,115	43,631	2,088	7,169	10,044
Total	269,203	56,147	56,172	20,442	18,141	69,857	3,979	20,135	24,330
Age²									
0 to 11 years	32,401	11,747	5,649	7,849	25	1,002	96	12	6,021
12 to 14 years	18,957	5,395	3,841	4,293	57	2,827	176	29	2,339
15 to 17 years	21,369	4,716	4,804	2,597	228	6,173	323	94	2,434
18 to 20 years	16,007	2,852	4,379	1,145	507	5,146	230	307	1,441
21 to 24 years	16,861	2,606	4,671	717	1,137	4,975	305	1,069	1,381
25 to 29 years	26,756	4,521	6,830	1,052	2,243	7,441	413	2,112	2,144
30 to 34 years	26,251	4,878	6,438	902	2,373	7,083	349	2,329	1,899
35 to 39 years	23,087	4,477	5,166	689	2,233	6,249	389	2,403	1,481
40 to 44 years	17,426	3,535	3,479	416	1,891	4,906	293	1,908	998
45 to 49 years	16,289	3,195	2,934	286	1,822	4,989	265	1,875	923
50 to 54 years	15,929	2,934	2,660	204	1,722	5,297	253	2,006	853
55 to 59 years	15,838	2,612	2,401	167	1,700	5,615	255	2,226	862
60 to 64 years	11,279	1,718	1,675	97	1,131	4,157	161	1,709	631
65 years and older	10,878	978	1,235	31	1,084	4,050	493	2,070	937
Total	269,328	56,164	56,162	20,445	18,153	69,910	4,001	20,149	24,344
Race									
American Indian or Alaska Native	7,951	1,919	1,526	621	439	1,876	105	561	904
Asian ³	6,390	1,018	1,141	249	308	2,283	197	699	495
Black or African American	19,662	4,603	3,099	1,211	1,150	4,962	395	2,578	1,664
Native Hawaiian or Other Pacific Islander	3,910	805	744	198	195	1,178	76	411	303
White	195,710	39,520	41,960	14,908	14,765	50,892	2,695	14,111	16,859
Other	31,686	7,510	6,682	2,843	1,199	7,998	357	1,696	3,401
Total	265,309	55,375	55,152	20,030	18,056	69,189	3,825	20,056	23,626

Continued. See notes at end of table.

Table 10.43a. Washington individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	211,150	42,921	43,124	15,113	15,785	55,769	3,172	17,291	17,975
Hispanic or Latino origin	48,215	11,346	10,151	4,433	2,039	11,878	464	2,668	5,236
Total	259,365	54,267	53,275	19,546	17,824	67,647	3,636	19,959	23,211
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	4,052	804	445	47	434	1,384	31	681	226
Not a veteran	100,694	19,212	14,750	1,607	11,020	33,658	953	13,072	6,422
Total	104,746	20,016	15,195	1,654	11,454	35,042	984	13,753	6,648

‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.43b. Washington individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	44.1	38.2	36.8	69.1	38.7	37.5	47.5	64.4	58.7
Female	55.9	61.8	63.2	30.9	61.3	62.5	52.5	35.6	41.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	12.0	20.9	10.1	38.4	†	1.4	2.4	†	24.7
12 to 14 years	7.0	9.6	6.8	21.0	†	4.0	4.4	†	9.6
15 to 17 years	7.9	8.4	8.6	12.7	1.3	8.8	8.1	†	10.0
18 to 20 years	5.9	5.1	7.8	5.6	2.8	7.4	5.7	1.5	5.9
21 to 24 years	6.3	4.6	8.3	3.5	6.3	7.1	7.6	5.3	5.7
25 to 29 years	9.9	8.0	12.2	5.1	12.4	10.6	10.3	10.5	8.8
30 to 34 years	9.7	8.7	11.5	4.4	13.1	10.1	8.7	11.6	7.8
35 to 39 years	8.6	8.0	9.2	3.4	12.3	8.9	9.7	11.9	6.1
40 to 44 years	6.5	6.3	6.2	2.0	10.4	7.0	7.3	9.5	4.1
45 to 49 years	6.0	5.7	5.2	1.4	10.0	7.1	6.6	9.3	3.8
50 to 54 years	5.9	5.2	4.7	†	9.5	7.6	6.3	10.0	3.5
55 to 59 years	5.9	4.7	4.3	†	9.4	8.0	6.4	11.0	3.5
60 to 64 years	4.2	3.1	3.0	†	6.2	5.9	4.0	8.5	2.6
65 years and older	4.0	1.7	2.2	†	6.0	5.8	12.3	10.3	3.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	3.0	3.5	2.8	3.1	2.4	2.7	2.7	2.8	3.8
Asian ³	2.4	1.8	2.1	1.2	1.7	3.3	5.2	3.5	2.1
Black or African American	7.4	8.3	5.6	6.0	6.4	7.2	10.3	12.9	7.0
Native Hawaiian or Other Pacific Islander	1.5	1.5	1.3	†	1.1	1.7	2.0	2.0	1.3
White	73.8	71.4	76.1	74.4	81.8	73.6	70.5	70.4	71.4
Other	11.9	13.6	12.1	14.2	6.6	11.6	9.3	8.5	14.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.43b. Washington individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	81.4	79.1	80.9	77.3	88.6	82.4	87.2	86.6	77.4
Hispanic or Latino origin	18.6	20.9	19.1	22.7	11.4	17.6	12.8	13.4	22.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	3.9	4.0	2.9	2.8	3.8	3.9	3.2	5.0	3.4
Not a veteran	96.1	96.0	97.1	97.2	96.2	96.1	96.8	95.0	96.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.44a. West Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	42,982	3,657	7,412	3,302	5,767	12,457	254	2,262	7,871
Gender									
Male	19,020	1,415	2,842	2,194	2,172	4,386	96	1,331	4,584
Female	23,953	2,242	4,568	1,108	3,594	8,067	158	931	3,285
Total	42,973	3,657	7,410	3,302	5,766	12,453	254	2,262	7,869
Age²									
0 to 11 years	2,780	564	180	1,092	33	79	--	2	830
12 to 14 years	1,983	360	219	617	51	248	--	2	486
15 to 17 years	2,654	380	378	520	147	596	3	14	616
18 to 20 years	2,429	254	407	323	223	695	11	19	497
21 to 24 years	2,671	186	490	185	388	775	26	81	540
25 to 29 years	4,294	293	874	201	643	1,117	40	166	960
30 to 34 years	4,056	257	812	125	675	1,021	26	206	934
35 to 39 years	4,137	288	811	77	732	1,132	38	240	819
40 to 44 years	3,693	253	716	59	645	1,158	30	224	608
45 to 49 years	3,498	218	670	37	651	1,202	25	242	453
50 to 54 years	3,321	203	608	25	538	1,258	23	276	390
55 to 59 years	3,015	175	519	18	443	1,244	15	305	296
60 to 64 years	2,171	126	363	15	308	925	9	228	197
65 years and older	2,280	100	365	8	290	1,007	8	257	245
Total	42,982	3,657	7,412	3,302	5,767	12,457	254	2,262	7,871
Race									
American Indian or Alaska Native	48	2	9	5	4	13	--	6	9
Asian ³	37	4	2	4	1	12	1	5	8
Black or African American	1,625	154	199	155	201	390	7	163	356
Native Hawaiian or Other Pacific Islander	19	2	3	2	5	3	--	2	2
White	36,692	3,170	6,572	2,667	4,836	10,914	207	1,748	6,578
Other	4,410	317	619	468	701	1,091	39	309	866
Total	42,831	3,649	7,404	3,301	5,748	12,423	254	2,233	7,819

Continued. See notes at end of table.

Table 10.44a. West Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	‡	‡	‡	‡	‡	‡	‡	‡	‡
Hispanic or Latino origin	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	891	96	145	18	135	265	7	55	170
Not a veteran	32,385	2,133	6,294	1,017	5,109	10,778	215	1,734	5,105
Total	33,276	2,229	6,439	1,035	5,244	11,043	222	1,789	5,275

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.44b. West Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	44.3	38.7	38.4	66.4	37.7	35.2	37.8	58.8	58.3	
Female	55.7	61.3	61.6	33.6	62.3	64.8	62.2	41.2	41.7	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	6.5	15.4	2.4	33.1	†	†	--	†	10.5	
12 to 14 years	4.6	9.8	3.0	18.7	†	2.0	--	†	6.2	
15 to 17 years	6.2	10.4	5.1	15.7	2.5	4.8	1.2	†	7.8	
18 to 20 years	5.7	6.9	5.5	9.8	3.9	5.6	4.3	†	6.3	
21 to 24 years	6.2	5.1	6.6	5.6	6.7	6.2	10.2	3.6	6.9	
25 to 29 years	10.0	8.0	11.8	6.1	11.1	9.0	15.7	7.3	12.2	
30 to 34 years	9.4	7.0	11.0	3.8	11.7	8.2	10.2	9.1	11.9	
35 to 39 years	9.6	7.9	10.9	2.3	12.7	9.1	15.0	10.6	10.4	
40 to 44 years	8.6	6.9	9.7	1.8	11.2	9.3	11.8	9.9	7.7	
45 to 49 years	8.1	6.0	9.0	1.1	11.3	9.6	9.8	10.7	5.8	
50 to 54 years	7.7	5.6	8.2	†	9.3	10.1	9.1	12.2	5.0	
55 to 59 years	7.0	4.8	7.0	†	7.7	10.0	5.9	13.5	3.8	
60 to 64 years	5.1	3.4	4.9	†	5.3	7.4	3.5	10.1	2.5	
65 years and older	5.3	2.7	4.9	†	5.0	8.1	3.1	11.4	3.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	†	†	†	†	†	†	--	†	†	
Asian ³	†	†	†	†	†	†	†	†	†	
Black or African American	3.8	4.2	2.7	4.7	3.5	3.1	2.8	7.3	4.6	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	--	†	†	
White	85.7	86.9	88.8	80.8	84.1	87.9	81.5	78.3	84.1	
Other	10.3	8.7	8.4	14.2	12.2	8.8	15.4	13.8	11.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.44b. West Virginia individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	‡	‡	‡	‡	‡	‡	‡	‡	‡
Hispanic or Latino origin	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	2.7	4.3	2.3	1.7	2.6	2.4	3.2	3.1	3.2
Not a veteran	97.3	95.7	97.7	98.3	97.4	97.6	96.8	96.9	96.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.45a. Wisconsin individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	63,581	6,785	10,509	2,811	5,714	11,226	6,150	9,134	11,252
Gender									
Male	31,412	2,956	4,902	1,808	2,390	4,533	3,352	5,704	5,767
Female	32,162	3,829	5,606	1,002	3,324	6,693	2,798	3,430	5,480
Total	63,574	6,785	10,508	2,810	5,714	11,226	6,150	9,134	11,247
Age²									
0 to 11 years	3,076	985	445	606	29	88	10	5	908
12 to 14 years	3,251	630	559	433	65	354	38	10	1,162
15 to 17 years	4,743	764	855	416	144	717	90	49	1,708
18 to 20 years	3,456	332	682	162	188	635	423	136	898
21 to 24 years	4,293	361	851	164	315	747	613	411	831
25 to 29 years	5,901	574	1,081	207	596	973	805	693	972
30 to 34 years	5,806	566	1,101	206	624	974	691	818	826
35 to 39 years	5,338	524	936	159	609	925	642	813	730
40 to 44 years	4,476	454	685	92	560	875	490	713	607
45 to 49 years	4,890	390	729	99	584	1,053	554	886	595
50 to 54 years	5,143	396	715	80	612	1,157	559	1,082	542
55 to 59 years	5,069	316	653	79	568	1,060	528	1,331	534
60 to 64 years	3,616	203	474	41	434	763	312	1,027	362
65 years and older	4,523	290	743	67	386	905	395	1,160	577
Total	63,581	6,785	10,509	2,811	5,714	11,226	6,150	9,134	11,252
Race									
American Indian or Alaska Native	1,235	111	244	56	75	204	75	112	358
Asian ³	872	219	90	14	43	234	52	121	99
Black or African American	6,722	527	728	204	398	764	2,026	1,445	630
Native Hawaiian or Other Pacific Islander	74	8	16	4	2	10	6	9	19
White	49,100	5,439	8,445	2,355	4,847	9,037	3,471	6,978	8,528
Other	2,231	252	309	96	246	294	323	361	350
Total	60,234	6,556	9,832	2,729	5,611	10,543	5,953	9,026	9,984

Continued. See notes at end of table.

Table 10.45a. Wisconsin individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	53,764	5,707	8,890	2,468	5,189	9,473	4,986	8,275	8,776
Hispanic or Latino origin	2,788	377	444	130	163	489	440	283	462
Total	56,552	6,084	9,334	2,598	5,352	9,962	5,426	8,558	9,238
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ “Mental health diagnosis” refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. “Other mental disorders” includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, “Asian” includes “Pacific Islanders.” See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.45b. Wisconsin individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹								
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹	
					Bipolar disorders	Depressive disorders				
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender										
Male	49.4	43.6	46.7	64.3	41.8	40.4	54.5	62.4	51.3	
Female	50.6	56.4	53.3	35.7	58.2	59.6	45.5	37.6	48.7	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²										
0 to 11 years	4.8	14.5	4.2	21.6	†	†	†	†	8.1	
12 to 14 years	5.1	9.3	5.3	15.4	1.1	3.2	†	†	10.3	
15 to 17 years	7.5	11.3	8.1	14.8	2.5	6.4	1.5	†	15.2	
18 to 20 years	5.4	4.9	6.5	5.8	3.3	5.7	6.9	1.5	8.0	
21 to 24 years	6.8	5.3	8.1	5.8	5.5	6.7	10.0	4.5	7.4	
25 to 29 years	9.3	8.5	10.3	7.4	10.4	8.7	13.1	7.6	8.6	
30 to 34 years	9.1	8.3	10.5	7.3	10.9	8.7	11.2	9.0	7.3	
35 to 39 years	8.4	7.7	8.9	5.7	10.7	8.2	10.4	8.9	6.5	
40 to 44 years	7.0	6.7	6.5	3.3	9.8	7.8	8.0	7.8	5.4	
45 to 49 years	7.7	5.7	6.9	3.5	10.2	9.4	9.0	9.7	5.3	
50 to 54 years	8.1	5.8	6.8	2.8	10.7	10.3	9.1	11.8	4.8	
55 to 59 years	8.0	4.7	6.2	2.8	9.9	9.4	8.6	14.6	4.7	
60 to 64 years	5.7	3.0	4.5	1.5	7.6	6.8	5.1	11.2	3.2	
65 years and older	7.1	4.3	7.1	2.4	6.8	8.1	6.4	12.7	5.1	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race										
American Indian or Alaska Native	2.1	1.7	2.5	2.1	1.3	1.9	1.3	1.2	3.6	
Asian ³	1.4	3.3	†	†	†	2.2	†	1.3	†	
Black or African American	11.2	8.0	7.4	7.5	7.1	7.2	34.0	16.0	6.3	
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†	
White	81.5	83.0	85.9	86.3	86.4	85.7	58.3	77.3	85.4	
Other	3.7	3.8	3.1	3.5	4.4	2.8	5.4	4.0	3.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.45b. Wisconsin individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	95.1	93.8	95.2	95.0	97.0	95.1	91.9	96.7	95.0
Hispanic or Latino origin	4.9	6.2	4.8	5.0	3.0	4.9	8.1	3.3	5.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Not a veteran	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡

‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.46a. Wyoming individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	15,762	2,971	2,298	862	1,487	4,680	316	939	2,209
Gender									
Male	7,014	1,264	845	599	544	1,699	154	597	1,312
Female	8,748	1,707	1,453	263	943	2,981	162	342	897
Total	15,762	2,971	2,298	862	1,487	4,680	316	939	2,209
Age²									
0 to 11 years	1,813	653	234	305	--	121	--	2	498
12 to 14 years	1,189	293	142	164	4	316	--	6	264
15 to 17 years	1,366	208	162	117	30	514	5	14	316
18 to 20 years	819	118	128	43	52	311	16	25	126
21 to 24 years	995	154	192	31	92	328	28	44	126
25 to 29 years	1,490	269	297	54	156	452	46	89	127
30 to 34 years	1,400	255	240	45	168	419	41	104	128
35 to 39 years	1,385	248	235	29	194	409	40	110	120
40 to 44 years	1,045	177	152	22	157	317	36	83	101
45 to 49 years	1,007	158	135	16	149	329	30	93	97
50 to 54 years	859	121	113	11	139	281	29	95	70
55 to 59 years	854	116	95	13	140	280	17	113	80
60 to 64 years	655	82	82	6	95	237	11	84	58
65 years and older	884	118	91	6	111	366	17	77	98
Total	15,761	2,970	2,298	862	1,487	4,680	316	939	2,209
Race									
American Indian or Alaska Native	313	64	32	15	20	102	6	38	36
Asian ³	69	12	9	3	6	22	1	7	9
Black or African American	246	48	12	15	14	65	8	43	41
Native Hawaiian or Other Pacific Islander	28	9	1	1	1	9	1	2	4
White	13,678	2,569	2,036	747	1,316	4,077	271	759	1,903
Other	598	123	100	24	48	176	12	26	89
Total	14,932	2,825	2,190	805	1,405	4,451	299	875	2,082

Continued. See notes at end of table.

Table 10.46a. Wyoming individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	13,660	2,588	1,993	735	1,313	4,063	279	792	1,897
Hispanic or Latino origin	1,538	289	222	97	96	469	24	107	234
Total	15,198	2,877	2,215	832	1,409	4,532	303	899	2,131
Specific Hispanic or Latino origin									
Mexican	746	152	120	43	38	214	12	57	110
Puerto Rican	64	10	6	5	8	22	--	2	11
Cuban	11	1	--	2	2	5	1	--	--
Other specific	683	122	91	46	45	222	10	43	104
Total	1,504	285	217	96	93	463	23	102	225
Veteran status⁴									
Veteran	210	61	28	2	24	55	7	12	21
Not a veteran	3,705	699	613	91	411	1,244	80	244	323
Total	3,915	760	641	93	435	1,299	87	256	344

-- Quantity is zero.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Total number of client characteristics may not equal the total number of clients due to missing values.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.46b. Wyoming individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Gender									
Male	44.5	42.5	36.8	69.5	36.6	36.3	48.7	63.6	59.4
Female	55.5	57.5	63.2	30.5	63.4	63.7	51.3	36.4	40.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Age²									
0 to 11 years	11.5	22.0	10.2	35.4	--	2.6	--	†	22.5
12 to 14 years	7.5	9.9	6.2	19.0	†	6.8	--	†	12.0
15 to 17 years	8.7	7.0	7.0	13.6	2.0	11.0	1.6	1.5	14.3
18 to 20 years	5.2	4.0	5.6	5.0	3.5	6.6	5.1	2.7	5.7
21 to 24 years	6.3	5.2	8.4	3.6	6.2	7.0	8.9	4.7	5.7
25 to 29 years	9.5	9.1	12.9	6.3	10.5	9.7	14.6	9.5	5.7
30 to 34 years	8.9	8.6	10.4	5.2	11.3	9.0	13.0	11.1	5.8
35 to 39 years	8.8	8.4	10.2	3.4	13.0	8.7	12.7	11.7	5.4
40 to 44 years	6.6	6.0	6.6	2.6	10.6	6.8	11.4	8.8	4.6
45 to 49 years	6.4	5.3	5.9	1.9	10.0	7.0	9.5	9.9	4.4
50 to 54 years	5.5	4.1	4.9	1.3	9.3	6.0	9.2	10.1	3.2
55 to 59 years	5.4	3.9	4.1	1.5	9.4	6.0	5.4	12.0	3.6
60 to 64 years	4.2	2.8	3.6	†	6.4	5.1	3.5	8.9	2.6
65 years and older	5.6	4.0	4.0	†	7.5	7.8	5.4	8.2	4.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Race									
American Indian or Alaska Native	2.1	2.3	1.5	1.9	1.4	2.3	2.0	4.3	1.7
Asian ³	†	†	†	†	†	†	†	†	†
Black or African American	1.6	1.7	†	1.9	†	1.5	2.7	4.9	2.0
Native Hawaiian or Other Pacific Islander	†	†	†	†	†	†	†	†	†
White	91.6	90.9	93.0	92.8	93.7	91.6	90.6	86.7	91.4
Other	4.0	4.4	4.6	3.0	3.4	4.0	4.0	3.0	4.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Continued. See notes at end of table.

Table 10.46b. Wyoming individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018 (continued)

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	89.9	90.0	90.0	88.3	93.2	89.7	92.1	88.1	89.0
Hispanic or Latino origin	10.1	10.0	10.0	11.7	6.8	10.3	7.9	11.9	11.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	49.6	53.3	55.3	44.8	40.9	46.2	52.2	55.9	48.9
Puerto Rican	4.3	3.5	2.8	5.2	8.6	4.8	--	2.0	4.9
Cuban	†	†	--	2.1	2.2	1.1	4.3	--	--
Other specific	45.4	42.8	41.9	47.9	48.4	47.9	43.5	42.2	46.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veteran status⁴									
Veteran	5.4	8.0	4.4	2.2	5.5	4.2	8.0	4.7	6.1
Not a veteran	94.6	92.0	95.6	97.8	94.5	95.8	92.0	95.3	93.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

-- Quantity is zero; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTE: Percentages may not sum to 100 percent because of rounding.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.47a. Other jurisdictions individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	278	8	9	10	4	16	--	112	119
Gender									
Male	198	2	6	9	2	9	--	72	98
Female	80	6	3	1	2	7	--	40	21
Total	278	8	9	10	4	16	--	112	119
Age²									
0 to 11 years	--	--	--	--	--	--	--	--	--
12 to 14 years	3	1	1	--	--	--	--	1	--
15 to 17 years	3	1	--	2	--	--	--	--	--
18 to 20 years	7	1	1	1	--	1	--	1	2
21 to 24 years	15	--	2	1	2	--	--	2	8
25 to 29 years	35	1	--	2	--	1	--	12	19
30 to 34 years	41	1	--	--	--	--	--	19	21
35 to 39 years	38	2	--	--	--	5	--	13	18
40 to 44 years	30	1	2	3	--	2	--	7	15
45 to 49 years	25	--	2	--	--	2	--	13	8
50 to 54 years	35	--	--	--	--	2	--	18	15
55 to 59 years	21	--	1	--	2	3	--	10	5
60 to 64 years	16	--	--	--	--	--	--	14	2
65 years and older	5	--	--	1	--	--	--	2	2
Total	274	8	9	10	4	16	--	112	115
Race									
American Indian or Alaska Native	--	--	--	--	--	--	--	--	--
Asian ³	173	6	7	8	4	6	--	108	34
Black or African American	1	--	--	--	--	--	--	--	1
Native Hawaiian or Other Pacific Islander	--	--	--	--	--	--	--	--	--
White	2	--	--	1	--	--	--	--	1
Other	29	--	--	--	--	3	--	2	24
Total	205	6	7	9	4	9	--	110	60

Continued. See notes at end of table.

Table 10.47a. Other jurisdictions individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Number, 2018 (continued)

Selected characteristics	Mental health diagnosis ¹								
	Total	Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	168	4	6	9	3	6	--	108	32
Hispanic or Latino origin	--	--	--	--	--	--	--	--	--
Total	168	4	6	9	3	6	--	108	32
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	4	1	--	1	--	--	--	--	2
Not a veteran	41	--	1	2	--	4	--	29	5
Total	45	1	1	3	--	4	--	29	7

-- Quantity is zero; ‡ No data submitted.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTES: Total number of client characteristics may not equal the total number of clients due to missing values. Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

Table 10.47b. Other jurisdictions individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Total individuals	100.0	100.0	100.0	100.0	100.0	100.0	--	100.0	100.0
Gender									
Male	71.2	25.0	66.7	90.0	50.0	56.3	--	64.3	82.4
Female	28.8	75.0	33.3	10.0	50.0	43.8	--	35.7	17.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	--	100.0	100.0
Age²									
0 to 11 years	--	--	--	--	--	--	--	--	--
12 to 14 years	1.1	12.5	11.1	--	--	--	--	†	--
15 to 17 years	1.1	12.5	--	20.0	--	--	--	--	--
18 to 20 years	2.6	12.5	11.1	10.0	--	6.3	--	†	1.7
21 to 24 years	5.5	--	22.2	10.0	50.0	--	--	1.8	7.0
25 to 29 years	12.8	12.5	--	20.0	--	6.3	--	10.7	16.5
30 to 34 years	15.0	12.5	--	--	--	--	--	17.0	18.3
35 to 39 years	13.9	25.0	--	--	--	31.3	--	11.6	15.7
40 to 44 years	10.9	12.5	22.2	30.0	--	12.5	--	6.3	13.0
45 to 49 years	9.1	--	22.2	--	--	12.5	--	11.6	7.0
50 to 54 years	12.8	--	--	--	--	12.5	--	16.1	13.0
55 to 59 years	7.7	--	11.1	--	50.0	18.8	--	8.9	4.3
60 to 64 years	5.8	--	--	--	--	--	--	12.5	1.7
65 years and older	1.8	--	--	10.0	--	--	--	1.8	1.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	--	100.0	100.0
Race									
American Indian or Alaska Native	--	--	--	--	--	--	--	--	--
Asian ³	84.4	100.0	100.0	88.9	100.0	66.7	--	98.2	56.7
Black or African American	†	--	--	--	--	--	--	--	1.7
Native Hawaiian or Other Pacific Islander	--	--	--	--	--	--	--	--	--
White	†	--	--	11.1	--	--	--	--	1.7
Other	14.1	--	--	--	--	33.3	--	1.8	40.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	--	100.0	100.0

Continued. See notes at end of table.

**Table 10.47b. Other jurisdictions individuals by mental health diagnosis, gender, age, race, ethnicity, and veteran status: Column percent distribution, 2018
(continued)**

Selected characteristics	Total	Mental health diagnosis ¹							
		Trauma- and stressor-related disorders	Anxiety disorders	ADHD	Mood disorders		Personality disorders	Schizophrenia, other psychotic disorders	Other mental disorders ¹
					Bipolar disorders	Depressive disorders			
Ethnicity									
Not of Hispanic or Latino origin	100.0	100.0	100.0	100.0	100.0	100.0	--	100.0	100.0
Hispanic or Latino origin	--	--	--	--	--	--	--	--	--
Total	100.0	100.0	100.0	100.0	100.0	100.0	--	100.0	100.0
Specific Hispanic or Latino origin									
Mexican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Puerto Rican	‡	‡	‡	‡	‡	‡	‡	‡	‡
Cuban	‡	‡	‡	‡	‡	‡	‡	‡	‡
Other specific	‡	‡	‡	‡	‡	‡	‡	‡	‡
Total	‡	‡	‡	‡	‡	‡	‡	‡	‡
Veteran status⁴									
Veteran	8.9	100.0	--	33.3	--	--	--	--	28.6
Not a veteran	91.1	--	100.0	66.7	--	100.0	--	100.0	71.4
Total	100.0	100.0	100.0	100.0	--	100.0	--	100.0	100.0

-- Quantity is zero; ‡ No data submitted; † Less than 1 percent of individuals.

ADHD = attention deficit/hyperactivity disorder.

¹ "Mental health diagnosis" refers to the first-mention mental health diagnosis, which is the first of up to three diagnoses reported for each individual. "Other mental disorders" includes all other mental health diagnoses and diagnoses not included in another diagnostic category; it also includes alcohol- and substance-related diagnoses. Mental health diagnostic category definitions can be found in Appendix E.

² Age is based on the individual's age at the midpoint of the reporting period.

³ For some states, "Asian" includes "Pacific Islanders." See Appendix C.

⁴ Veteran status is only reported for individuals aged 18 years and older.

NOTES: Percentages may not sum to 100 percent because of rounding. Other jurisdictions include the Commonwealth of the Northern Mariana Islands and the Republic of Palau.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2018 reporting period. Data as of 07.08.2019.

**Appendix A
MH-TEDS/MH-CLD Crosswalk Tables**

MH-TEDS					MH-CLD			
Variable Name	Data Field Reference No.			Categories	Variable Name	Data Field Reference No.		Categories
	Admission Record	Discharge/Update Record				Data Files		
		Value taken from admission record	Value taken at discharge or data update			BCI	SHR	
System Transaction Type	SDS 1		DIS 1	A=Add C=Change D=Delete	Transaction Type	C-00	R-00	A=Add C=Change D=Delete
State Code — Key field	SDS 2		DIS 2	2 alphanumeric	Reporting State Code	H-02	H-01	2 alphanumeric
Reporting Date	SDS 3		DIS 3	MMYYYY	contained in the file name algorithm			MMDDYYYY
CLIENT RECORD					CLIENT RECORD			
State Provider Identifier – Key field	MDS 1	DIS 11	DIS 4	1-15 alphanumeric	Not reported			
Client Identifier – Key field	MDS 2	DIS 12	DIS 5	1-15 alphanumeric	Client Identifier	C-01	R-01	1-15 alphanumeric
Codependent or Collateral – Key field	MDS 3	DIS 13	DIS 6	1=codependent/collateral 2=client	Not reported			
Client Transaction Type – Key field	MDS 4			A=Initial Admission (SA) T=Transfer or change in service (SA) M=Initial Admission (MH) X=Transfer or change in service (MH)	Not reported			
Date of Admission – Key field	MDS 5	DIS 15		MMDDYYYY	Client treatment status at the start of the reporting period	C-02		1=New client; 2=Continuing client;
Date of Birth	MDS 8	DIS 17		MMDDYYYY 01010007=unknown 01010008=not collected	Age	C-05		
Gender	MDS 9	DIS 18		1=male 2=female 7=unknown 8=not collected	Sex	C-04		1=male 2=female 7=unknown 8=not collected
Race	MDS 10	DIS 19		01=Alaska Native 02=American Indian and Alaska Native 13=Asian 23=Native Hawaiian or other Pacific Islander	Race	C-06		02=American Indian and Alaska Native 13=Asian 23=Native Hawaiian or other Pacific Islander

**Appendix A
MH-TEDS/MH-CLD Crosswalk Tables**

Variable Name	MH-TEDS			Categories	Variable Name	MH-CLD		Categories
	Data Field Reference No.					Data Field Reference No.		
	Admission Record	Discharge/Update Record				BCI	SHR	
Value taken from admission record		Value taken at discharge or data update						
				03=Asian or Pacific Islander				03=Asian or Pacific Islander (temporary code)
				04= Black or African American				04= Black or African American
				05=White				05=White
				20=Other single race				20=Some other race alone
				21=Two or more races				21=Two or more races
								22=Hispanic (temporary code)
				97=unknown				97=unknown
				98=not collected				98=not collected
Hispanic or Latino Origin (Ethnicity)	MDS 11	DIS 20		01=Puerto Rico	Ethnicity	C-07		11=Puerto Rican (optional)
				02=Mexican				12=Mexican (optional)
				03=Cuban				13=Cuban (optional)
				04=Other Specific Hispanic or Latino				14=Other Specific Hispanic
				05 Not of Hispanic Origin				02=Not of Hispanic Origin
				06=Hispanic, specific origin not specified				01=Hispanic origin regardless of race
				97=unknown				97=unknown
				98=not collected				98=Not collected
Education	MDS 12		MHD 5 (reported only for ages 3-17 and up to 21 if in special education).	00= less than one grade completed or no schooling	School grade level	C-25		00=no years of schooling
				01-11=grades 1 thru 11th				01-11=grades 1 thru 11th
				12=12th grade or GED				12 = 12th grade
				13=Freshman or first year in a 2-year (Associate) Degree				17=freshman (1st year in College)
				14=Sophomore or second year in a 2-year (Associate) Degree				18=sophomore (2nd year in College)
				15=Junior in college				19=junior (3rd year in College)
				16=Senior in college or Bachelor's Degree				20=senior (4th year in College)
				17=Some Post-Graduate Study-Degree not completed				
				18= Master's Degree completed				
				19-25= Post graduate study				
				70= graduate or professional school (can be used instead of codes 17-25)				21=Graduate or professional school
				71=vocational school				16=vocational school

**Appendix A
MH-TEDS/MH-CLD Crosswalk Tables**

Variable Name	MH-TEDS			Categories	Variable Name	MH-CLD		Categories
	Data Field Reference No.					Data Field Reference No.		
	Admission Record	Discharge/Update Record				Data Files		
Value taken from admission record		Value taken at discharge or data update	BCI	SHR				
				72=nursery school, pre-school 73=kindergarten 74=self-contained special education class 97=unknown 98=not collected				13=nursery school, pre-school (head start) 14=kindergarten 15=self-contained special education class 97 = unknown 98=not collected
Employment Status	MDS 13		DIS 24	01=full time 02=part-time 03=unemployed 04=not in the labor force 05=employed, full/part-time not specified (temporary code) 97=unknown 98=not collected	Competitive Employment Status at admission/start of the reporting period and discharge/end of the reporting period (>= 16 yrs old)	C-09 and C-10		01=full time 02=part time 03=unemployed 84=not in the labor force 05=employed, full/part-time not specified (temporary code) 97=unknown 98=not collected
Type of Treatment Service/ Treatment Setting – Key field	MDS 18	DIS 16	DIS 7	01=detox, 24-hr, hospital 02= detox, 24-hr, residential 03=rehabilitation/residential-hospital (not detox) 04=rehabilitation/residential-short term (<30 days) 05=rehabilitation/ residential-long term (>30 days) 06=ambulatory-intensive outpatient 07=ambulatory-non-intensive outpatient 08=ambulatory-detoxification 72=state psychiatric hospital 73=SMHA funded/operated community- based program 74=residential treatment center 75=other psychiatric inpatient	Service Setting Status Throughout the Reporting Period	C-15		1=state psychiatric hospital 2=SMHA funded/operated community-based program 3=residential treatment center 4=other psychiatric inpatient

**Appendix A
MH-TEDS/MH-CLD Crosswalk Tables**

Variable Name	MH-TEDS			Categories	Variable Name	MH-CLD		Categories
	Data Field Reference No.					Data Field Reference No.		
	Admission Record	Discharge/Update Record				Data Files		
Value taken from admission record		Value taken at discharge or data update	BCI	SHR				
				76=institutions under the justice system 96=Not applicable (use only for codependents or collateral clients (SA) and for MH clients receiving MH assessments, evaluation, or screening only)				5=institutions under the justice system
Date of Last Contact or Data Update - Key field			DIS 8	MMDDYYYY 01010007 = Unknown 01010008 = Not Collected	Information used in determining client status			
Reason for Discharge, Transfer, or Discontinuance of Treatment			DIS 10	01=treatment completed 02= dropped out of treatment (lost contact, administrative discharge, left against medical advise, eloped, failed to return from leave, and client choice) 03=terminated by facility 04=transferred to another treatment program or facility provider 14=transferred to another treatment program/facility but did not report 05=incarcerated or released by or to courts 06=death 07=Other (includes aging out of the children's MH system, extended placement (conditional release), and all other reasons) 08= Unknown - This code will continue to be accepted. However, states are encouraged to use the code 97 Unknown instead 24=transferred to another treatment program or facility that is not in the state reporting system	Discharge reasons	BCI (C-03) 12=treatment completed 22= lost contact or administrative discharge 32=discharged to corrections, jail 43=death of client 52=aged out; 62= other specified reasons	SHR (R-03) 01=completed SH inpatient treatment 07=discharged to other inpatient provider 02=released by or to courts 06=death 05=extended placement; 09=reason not classified elsewhere	

**Appendix A
MH-TEDS/MH-CLD Crosswalk Tables**

Variable Name	MH-TEDS			Categories	Variable Name	MH-CLD		Categories
	Data Field Reference No.					Data Field Reference No.		
	Admission Record	Discharge/Update Record				Data Files		
Value taken from admission record		Value taken at discharge or data update	BCI	SHR				
				34 =discharged temporarily to an acute medical facility for medical services 96 = Not applicable 97=unknown 98=not collected				08=discharged to an acute medical facility for medical services 72=reason unknown 82=reason not collected 97=unknown 98=not collected
Client Discharge Transaction Type - Key field		DIS 14	DIS 28	D=Discharge (SA client) E=Discharge (MH client) U=D ata update (MH client)	Not reported			
Co-occurring Mental and Substance Use Disorders	SuDS 5			1=yes 2=no 7=unknown 8=not collected	Substance use problem	C-21		1=yes 2=no 7=unknown 8=not collected
Date of Discharge - Key field			DIS 9	MMDDYYYY	Information used in determining Client status - at end of the reporting period	C-03		
Living Arrangements	SuDS 8		DIS 23	01=homeless 02=dependent living (under age 18 living with parents, relatives, or guardians) 22=dependent living: residential care and other residential status 32=dependent living: foster home/foster care 42=dependent living: crisis residence 52=dependent living:institutional setting 62=dependent living: jail and other institutions under the justice system 72=dependent living: adults in private residence who needs assistance in daily living	Residential status at start and end of the reporting period	C-12 and C-13		01=homeless 37=private residence (children only) 03=residential care 08=other residential status 02=fosterhome/foster care 04=crisis residence 05=institutional setting 06=jail/correctional facility 27=private residence, dependent living (adult)

**Appendix A
MH-TEDS/MH-CLD Crosswalk Tables**

Variable Name	MH-TEDS			Categories	Variable Name	MH-CLD		Categories
	Data Field Reference No.					Data Field Reference No.		
	Admission Record	Discharge/Update Record				Data Files		
Value taken from admission record		Value taken at discharge or data update	BCI	SHR				
				03=independent living 04=private residence, living arrangement not specified, adults (temporary code) 97=unknown 98=not collected				17=private residence, independent living (adult) 07=private residence, living arrangement not specified, adults (temporary code) 97=unknown 98=not collected
Detailed Not-in-Labor Force	SuDS 12		DIS 25	01=homemaker 02=student 03=retired 04=disabled 05= resident of institution or persons receiving services from institutional facilities such as hospitals, jails, prisons, etc. 06=other 07=sheltered/non-competitive employment 96=not applicable (if employment status is not 04, NILF) 97=unknown 98=not collected	Competitive Employment Status - at start and end of the reporting period	C-09 and C-10		14=homemaker 24=student 34=retired 44=disabled 64=other reported classification 74=sheltered/non-competitive employment 96=not applicable (if in institutional setting or if age <16) 97=unknown 84=NILF, classification not specified 98=not collected
Arrests in Past 30 Days	SuDS 16		DIS 26	00-96=number of arrests 97=unknown 98=not collected	Number of Arrests in Prior 30 days - at start and end of the reporting period	C-22 and C-23		00-96=number of arrests 97=unknown 98=not collected
Diagnostic Code Set Identifier	SuDS 18		MHD 1	1=DSM-IV 2=ICD-9 3=ICD-10 4=DSM-V 5=DSM-IIIIR 7=unknown 8=not collected	Diagnostic Code Identifier	H-08		1=DSM IV 2=ICD-9 3=ICD 10

**Appendix A
MH-TEDS/MH-CLD Crosswalk Tables**

		MH-TEDS				MH-CLD			
Variable Name	Data Field Reference No.			Categories	Variable Name	Data Field Reference No.		Categories	
	Admission Record	Discharge/Update Record				Data Files			
		Value taken from admission record	Value taken at discharge or data update			BCI	SHR		
SH-Legal Status at Admission	MHA 4			01=voluntary-self	State Hospital-Admission Legal Status		R-05	01=voluntary-self	
				02=voluntary-others (parents, guardians, etc)				02=voluntary-others (parents, guardians, etc)	
				03=invuntary-civil				03=invuntary-civil	
				04=invuntary-criminal				04=invuntary-criminal	
				05=invuntary-juvenile justice				05=invuntary-juvenile justice	
				06=invuntary-civil, sexual				06=invuntary-civil, sexual	
				96=not applicable					
				97=unknown				97=unknown	
				98=not collected				98=not collected	
SMI/SED Status	MHA 2		MHD 3	1=SMI	SMI/SED Status	C-08		1=SMI	
				2=SED				2=SED	
				3=at risk for SED (optional)				3=at risk for SED (optional)	
				4=Not SMI/SED				4=Not SMI/SED	
				7=unknown				7=unknown	
				8=not collected				8=not collected	
School Attendance Status	MHA 3		MHD 4	1=Yes, client has attended school at any time in the past 3 months	School Attendance Status (reported only for school age children, 3-17, and up to 21 if in special education).	C-24		1=Yes, client has attended school at any time in the past 3 months	
				2=No, client has not attended school at any time in the past three months				2=No, client has not attended school at any time in the past three months	
				6=not applicable (non-school age)				6=not applicable	
				7=unknown				7=unknown	
				8=not collected				8=not collected	
MH Diagnostic Code - one (ICD-10 Compatible)	MHA 1a		MHD 2a	xxx.xxxx; xxx-----; xxx.---;	Mental Health Diagnosis - one	C-17		xxx.xxxx; xxx.xxx-; xxx.xx--; xxx.x---; xxx--	
				999.9997 unknown				999.9997=unknown	
				999.998=not collected				999.9998=not collected	

**Appendix A
MH-TEDS/MH-CLD Crosswalk Tables**

MH-TEDS				MH-CLD				
Variable Name	Data Field Reference No.			Categories	Variable Name	Data Field Reference No.		Categories
	Admission Record	Discharge/Update Record				Data Files		
		Value taken from admission record	Value taken at discharge or data update			BCI	SHR	
MH Diagnostic Code - two (ICD-10 Compatible)	MHA 1b		MHD 2b	xxx.xxxx; xxx-----; xxx.----; xxx.x- --; xxx.xx--; xxx.xxx;	Mental Health Diagnosis - two	C-18		xxx.xxxx; xxx-----; xxx.----; xxx.x- --; xxx.xx--; xxx.xxx-
				999.9996=no second MH diagnosis				999.9996=no second diagnosis
				999.9997=unknown				999.9997=unknown
				999.9998=not collected				999.9998=not collected
MH Diagnostic Code - three (ICD-10 Compatible)	MHA 1c		MHD 2c	xxx.xxxx; xxx-----; xxx.----; xxx.x- --; xxx.xx--; xxx.xxx-	Mental Health Diagnosis - three	C-19		xxx.xxxx; xxx.xxx-; xxx.xx--; xxx.x- --; xxx.xx--; xxx.xxx-
				999.9996=no third MH diagnosis				999.9996=no third diagnosis
				999.9997=unknown				999.9997=unknown
				999.9998=not collected				999.9998=not collected
Diagnostic Code - ICD 10 Compatible, Substance use	SuDS 19			optional reporting for MH	Substance use Diagnosis	C-20		
				xxx.xxxx; xxx-----; xxx.----; xxx.x- --; xxx.xx--; xxx.xxx-				xxx.xxxx; xxx.xxx-; xxx.xx--; xxx.x- --; xxx.xx--; xxx.xxx-
				999.9997=unknown				999.9996 no substance use diagnosis
				999.9998=not collected				999.9997=unknown
Marital Status	SuDS 14			optional reporting for MH	Marital Status	O-1		optional reporting
				01=never married				01=never married
				02=now married/living as couple or cohabiting				02=married/living as couple
				03=separated				03=separated
				04=divorced				04=divorced
				05=widowed				05=widowed
				97=unknown				97=unknown
				98=not collected				98=not collected
GAF/CGAS Score – Optional	MHA 5		MHD 6	optional reporting	GAF/CGAS Score	O-02		optional reporting
				0-100 GAF/CGAS Score				0-100 GAF/CGAS score
				997 = Unknown				997=unknown
				998 = Not collected				
Prior Treatment Episodes	MDS 6			optional reporting for MH	Not reported			
Referral Source	MDS 7			optional reporting for MH	Not reported			
Substance Use (Primary, Secondary, Tertiary)	MDS 14a MDS 14b MDS 14c		DIS 21a DIS 21b DIS 21c	optional reporting for MH	Not reported			

**Appendix A
MH-TEDS/MH-CLD Crosswalk Tables**

		MH-TEDS				MH-CLD			
Variable Name	Data Field Reference No.			Categories	Variable Name	Data Field Reference No.		Categories	
	Admission Record	Discharge/Update Record				Data Files			
		Value taken from admission record	Value taken at discharge or data update			BCI	SHR		
Route of Administration (Primary, Secondary, Tertiary Substances)	MDS 15a MDS 15b MDS 15c			optional reporting for MH	Not reported				
Frequency of Use (Primary, Secondary, Tertiary Substances)	MDS 16a MDS 16b MDS 16c		DIS 22a DIS 22b DIS 22c	optional reporting for MH	Not reported				
Age at First Use (Primary, Secondary, Tertiary, Substances)	MDS 17a MDS 17b MDS 17c			optional reporting for MH	Not reported				
Medication-Assisted Opioid Therapy	MDS 19			optional reporting for MH	Not reported				
Detailed Drug Code (Primary, Secondary, Tertiary)	SuDS 1 SuDS 2 SuDS 3			optional reporting for MH	Not reported				
Diagnostic Code (DSM/ICD)	SuDS 4			optional reporting	Not reported				
Pregnant at Admission	SuDS 6			optional reporting	Not reported				
Veteran Status	SuDS 7			optional reporting for MH	Veteran Status	O-05		optional reporting for MH	
				1= Veteran				1= Veteran	
				2 = Not a Veteran				2 = Not a Veteran	
				7 = Unknown				7 = Unknown	
				8 = Not Collected				8 = Not Collected	
Source of Income/Support	SuDS 9			optional reporting for MH	Not reported				
Health Insurance	SuDS 10			optional reporting for MH	Not reported				
Payment Source, Primary (Expected or Actual)	SuDS 11			optional reporting for MH	Not reported				
Detailed Criminal Justice Referral	SuDS 13			optional reporting	Not reported				
Days Waiting to Enter Treatment	SuDS 15			optional reporting	Not reported				
Attendance at Self-Help Groups in Past 30 Days	SuDS 17		DIS 27	optional reporting for MH	Not reported				
Not reported				optional reporting for MH	Type of Funding Support	O-03		optional reporting for MH	
				01 = Medicaid Only				01 = Medicaid Only	
				02 = Medicaid and Non-Medicaid Sources				02 = Medicaid and Non-Medicaid Sources	
				03 = Non-Medicaid Only				03 = Non-Medicaid Only	
				97 = Unknown				97 = Unknown	
98 = Not Collected	98 = Not Collected								
Not reported				optional reporting for MH	Mental Health Block Grant funded services	O-04		optional reporting for MH	
				01 = Yes, MHBG Used to pay for services and supports				01 = Yes, MHBG Used to pay for services and supports	
				02 = No, MHBG funds were not used				02 = No, MHBG funds were not used	
				97 = Unknown				97 = Unknown	

**Appendix A
MH-TEDS/MH-CLD Crosswalk Tables**

MH-TEDS				MH-CLD				
Variable Name	Data Field Reference No.			Categories	Variable Name	Data Field Reference No.		Categories
	Admission Record	Discharge/Update Record				Data Files		
		Value taken from admission record	Value taken at discharge or data update			BCI	SHR	
								98 = Not Collected
LEGEND								
Key field = must not be blank. Key fields, taken together, are used to uniquely identify and link records.								
Variable highlighted in green = variable not reported in MH-CLD but required in MH-TEDS.								
Variable highlighted in orange = variable reported in MH-CLD and incorporated in MH-TEDS.								

APPENDIX B

Table B1. MH-TEDS and MH-CLD 2011–2018 reporting methods and 2018 reporting period, by state or jurisdiction

State or jurisdiction	Data reporting method (C: MH-CLD, T: MH-TEDS)								
	2011	2012	2013	2014	2015	2016	2017	2018	2018 reporting period
Alabama	-	-	C	C	C	C	C	C	10/1/2017 to 9/30/2018
Alaska	-	-	-	-	-	-	-	-	-
American Samoa	-	-	-	-	-	-	-	-	-
Arizona	C	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Arkansas	C	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
California	-	-	C	C	C	C	C	C	7/1/2016 to 6/30/2017
Colorado	C	-	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Connecticut	-	C	C	T, C ¹	7/1/2017 to 6/30/2018				
Delaware	-	C	C	C	C	T	T	T	7/1/2017 to 6/30/2018
District of Columbia	-	-	-	T*	T*	T	T	T	10/1/2017 to 9/30/2018
Federated States of Micronesia	-	-	-	-	-	-	-	-	-
Florida	C	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Georgia	-	-	-	-	-	-	-	-	-
Guam	-	-	-	-	-	-	-	-	-
Hawaii	-	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Idaho	-	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Illinois	-	-	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Indiana	C	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Iowa	-	-	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Kansas	C	C	C	C	*	*	*	*	<i>n.a.</i>
Kentucky	C	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Louisiana	C	C	C	T	T	T	T	T	7/1/2017 to 6/30/2018
Maine	-	-	C	*	*	*	C	**	7/1/2017 to 6/30/2018
Marshall Islands	-	-	-	-	-	-	-	-	-
Maryland	-	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Massachusetts	-	-	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Michigan	-	-	-	-	-	-	T	T	10/1/2016 to 9/30/2017
Minnesota	-	-	-	C	C	C	C	C	1/1/2017 to 12/30/2017
Mississippi	-	-	C	T	T	T	T	T	7/1/2016 to 6/30/2017
Missouri	C	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Montana	C	-	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Nebraska	-	C	C	C	C	C	T	T	7/1/2017 to 6/30/2018

Continued. See notes at end of table.

Table B1. MH-TEDS and MH-CLD 2011–2018 reporting methods and 2018 reporting period, by state or jurisdiction (continued)

State or jurisdiction	Data reporting method (C: MH-CLD, T: MH-TEDS)								
	2011	2012	2013	2014	2015	2016	2017	2018	2018 reporting period
Nevada	C	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
New Hampshire	-	-	-	-	-	-	-	-	-
New Jersey	-	C	C	C	C	C	***	***	n.a.
New Mexico	-	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
New York ²	-	C	C	C	C	C	C	C	4/1/2017 to 3/30/2018
North Carolina	-	-	C	C	C	T	T	T	7/1/2017 to 6/30/2018
North Dakota	-	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Commonwealth of the Northern Mariana Islands	-	C	C	C	C	C	C	C	10/1/2017 to 9/30/2018
Ohio	-	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Oklahoma	C	C	C	T	T	T	T	T	7/1/2017 to 6/30/2018
Oregon	-	-	C	C	C	C	C	C	1/1/2018 to 12/30/2018
Republic of Palau	-	-	-	-	-	C	C	C	10/1/2016 to 9/30/2017
Pennsylvania	-	-	C	C	*	T	T	T	7/1/2016 to 6/30/2017
Puerto Rico	-	-	-	T	T	T	T	T	7/1/2017 to 6/30/2018
Rhode Island	C	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
South Carolina	C	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
South Dakota	-	-	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Tennessee ²	-	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Texas	C	C	C	C	C	C	C	C	9/1/2017 to 8/30/2018
Utah	C	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
Vermont	-	C	C	C	C	T	T	T	7/1/2017 to 6/30/2018
Virginia	-	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
U.S. Virgin Islands	-	-	-	-	-	-	-	-	-
Washington	C	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018
West Virginia	-	-	-	-	-	-	-	T	7/1/2017 to 6/30/2018
Wisconsin	C	C	C	C	C	C	C	C	1/1/2017 to 12/30/2018
Wyoming	C	C	C	C	C	C	C	C	7/1/2017 to 6/30/2018

n.a. Not reporting; - State is building capacity for future reporting of client-level data; * State is building capacity to switch to MH-TEDS reporting; ** State switched to building capacity (for MH-CLD reporting) due to staff turnover; *** The state agency that is responsible for data reporting moved to a new location and was not able to submit data.

T* District of Columbia: 2014 and 2015 reporting Period: 7/1–6/30. Reporting period changed in FY 2016.

¹ Connecticut, 2014–2018: MH TEDS for adults, MH CLD for children/adolescents.

² New York and Tennessee report only a subset of the total individuals served through their SMHAs.

APPENDIX C

MH-TEDS AND MH-CLD DATA DICTIONARY OF INCLUDED VARIABLES

Variable Name: Age

Description: This field is calculated from the client's date of birth at midpoint of the state's elected reporting period.

Guidelines: Do not round up age. For example, if the calculated age is 13 and 8 months, report only 13. When the client's calculated age is less than 12 months, report age as zero. If the reported date of birth of a client changes during the reporting period, use the most recent available information in calculating age.

Variable Name: Competitive Employment Status (Aged 16 +)

Description: This data element is reported for all individuals (16 years old and over) who are receiving services in non-institutional settings. Institutional settings include correctional facilities like prison, jail, detention centers, and mental health care facilities like state hospitals, other psychiatric inpatient facilities, nursing homes or other institutions that keep a person, otherwise able, from entering the labor force.

Full Time – use state definition for full time employment; includes members of the Armed Forces, and individuals in full-time Supported Employment.

Part Time – use state definition for part time; includes individuals in part-time Supported Employment.

Unemployed – defined as actively looking for work or laid off from job (and waiting to be recalled) in the past 30 days.

Not in labor force – defined as not employed and not actively looking for work during the past 30 days, includes Homemaker, Student, Retired, Disabled, Other Reported Classification (e.g., volunteers), Sheltered/Not-Competitive Employment, an inmate of an institution.

Guidelines: **Employment Status** is defined by SAMHSA as an outcome measure and is collected at admission and discharge/update to assess change.

Coding of individuals with overlapping employment statuses: When individuals are engaged in two or more activities (have overlapping status) during the period when their status is collected by the SMHA, use DOL's system of priorities to determine the appropriate employment status. The prioritization rule is, labor force activities (such as working or looking for work) take precedence over non-labor force activities (such as student and homemaker); and working or having a job takes precedence over looking for work.

Reporting of a person in an internship program: The Individual with Disabilities Education Act (IDEA) ceases to apply for young individuals (18-21 years old) once they received their high school diploma. The following rules should be observed in making determination whether the adult client in an internship program should be reported as ‘Not in Labor Force’ or ‘Employed’.

1. If the internship is a school requirement, whether paid or not, the person should be considered a “student” (Not In Labor Force).
2. If the internship is not a school requirement, an unpaid position, does not displace regular employees or does not entitle the person for the job at the end of the internship period, then report the person’s status as Not In Labor Force - (Other).
3. If the internship is not a school requirement, paid minimum wage and over- time provision, the employer benefits from the internship through the client’s engagement in actual operations of the business and performing productive work, then the person is employed (Full time or Part time).

Variable Name: Detailed Not In Labor Force

Description: This field provides more detailed information about those individuals who are coded as “Not in labor force” in Employment Status.

Homemaker, Student, Retired, Disabled

Resident of Institution – Individuals receiving services from institutional facilities such as hospitals, jails, prisons, long-term residential care, etc.

Other – includes volunteer, seasonal worker, other categories used by the state not specified.

Guidelines: **Detailed Not in Labor Force** is defined by SAMHSA as an outcome measure, and is collected at admission and at discharge/update to assess change.

This field is reported for individuals aged 16 and older whose **Employment Status** is coded “Not in labor force.”

Variable Name: Co-occurring mental and substance use disorders

Description: This field indicates whether the client has co-occurring substance use and mental health disorders.

Yes – Client has co-occurring substance use and mental health disorders.

No – Client does not have co-occurring substance use and mental health disorders.

Guidelines: The assessment of co-occurring mental and substance use disorders may be based on clinical diagnoses, screening results, claims information, or self-report.

Variable Name: Ethnicity

Description: This field identifies whether or not the client is of Hispanic origin.

Guidelines: If the ethnicity of a client changes during the reporting period, report the most recent available information at the end of the reporting period.

Variable Name: Gender

Description: This field identifies the client’s most recent reported sex at the end of the reporting period.

Guidelines: States that collect transgender as an option must report the data as follows: Transgender man (female sex at birth but identifies as male) – code as female; transgender woman (male sex at birth but identifies as female) – code as male.

If the gender of a client changes during the reporting period, report the most recent available information.

Variable Name: Living Arrangements (Residential Status)

Description: This field specifies client’s residential status at time of admission (new individuals) or start of the reporting period (continuing individuals).

Homeless – person has no fixed address; includes homeless shelters.

Foster Home/Foster Care – individual resides in foster home. A foster home is a home that is licensed by a county or state department to provide foster care to children, adolescents, and/or individuals. This includes therapeutic foster care facilities. Therapeutic foster care is a service that provides treatment for troubled children within private homes of trained families.

Residential Care – individual resides in a residential care facility. This level of care may include a group home, therapeutic group home, board and care residential treatment, rehabilitation center, or agency-operated residential care facilities.

Crisis Residence – a time-limited residential (24 hours/day) stabilization program that delivers services for acute symptom reduction and restores clients to a pre-crisis level of functioning.

Institutional Setting – individual resides in an institutional care facility with care provided on a 24 hour, 7 days a week basis. This level of care may include skilled nursing/intermediate care facility, nursing homes, institute of mental disease (IMD), inpatient psychiatric hospital, psychiatric health facility, veterans’ affairs hospital, state hospital, or intermediate care facility/MR.

Jail/Correctional Facility – individual resides in a jail and/or correctional facility with care provided on a 24 hour, 7 days a week basis. This includes a jail, correctional facility, detention centers, and prison.

Dependent Living: Private Residence – adult individuals living in a house, apartment, or other similar dwelling who are heavily dependent on others for daily living assistance.

Independent Living: Private Residence – individuals living alone or with others in a private residence and capable of self-care. Includes adult children (age 18 and over)

living with parents and adolescents living independently. Also includes individuals who live independently with case management and supported housing support.

Private Residence – for adult with unknown dependency status, or for all children living in a private residence regardless of living arrangement.

Other – Other residential status

Guidelines: **Living Arrangements** is defined by SAMHSA as an outcome measure and is collected at admission and at discharge/update to assess change.

Variable Name: Marital Status (Optional Reporting)

Description: This field identifies the client’s marital status. The following categories are compatible with categories used in the U.S. Census.

Never married – includes individuals who are single or whose only marriage was annulled.

Currently married – includes married couples, those living together as married, living with partners, or cohabitating.

Separated – includes those legally separated or otherwise absent from spouse because of marital discord.

Divorced – legally divorced.

Widowed

Guidelines: Report marital status at time of discharge or the most recent available status at the end of the reporting period.

Variable Name: Mental Health Diagnosis

Description: This field specifies the client’s current mental health diagnoses during the reporting period.

Guidelines: This measure can be reported by using either the *Diagnostic and Statistical Manual of Mental Disorders (DSM)* from the American Psychiatric Association or the *International Classification of Diseases (ICD)* from the World Health Organization.

States should report the most recent mental health diagnoses that are current during the reporting period. Most recent is defined by the date when the diagnosis is reported.

States can report up to three mental health diagnoses by completing the Mental Health Diagnosis 1, 2, and 3 data elements in sequential order.

If the client has more than three most recent mental health diagnoses, use the algorithm below. Use this if the diagnoses are collected through administrative method (i.e., based on the clinician’s evaluation of the person and reported in the client’s case record):

1. Report the primary and secondary diagnoses (if available). If the state does not classify diagnosis into primary/secondary; then
2. Conduct your search for diagnostic codes using both Axis I and II. Report

all diagnoses in Axis I (clinical disorders) first followed by diagnosis in Axis II (personality disorders and mental retardation) unless a personality disorder in Axis II was labeled as primary diagnosis, then it should be reported first (note: this method is applicable if using DSM-IV and other earlier editions or if the state continues to use the DSM axial structure;

If primary/secondary labels and Axis classifications (I and II) are not used by the state, then

3. Report in chronological order starting from the diagnosis that appears on top of the list or first cited in the clinician's report. Report only mental health diagnosis.

If the state is using claims/encounter data to collect a client's diagnosis, use the three most frequently reported mental health diagnoses in the client's service claims/encounters data throughout the reporting period.

Variable Name: Race

Description: This field specifies the client's most recent reported race at the end of the reporting period.

Alaska Native – used in MH-TEDS only

American Indian or Alaska Native – A person having origins in any of the original people of North and South America (including Central America and the original people of Alaska) and who maintain cultural identification through tribal affiliation or community attachment.

Asian – A person having origins in any of the following people of the Far East, the Indian Subcontinent, or Southeast Asia, including Cambodia, China, India, Japan, Korea, Malaysia, Philippine Islands, Thailand, and Vietnam.

Asian or Pacific Islander (temporary code) – older racial category used only by some states that have not yet adopted the OMB recommended racial categories.

Black or African American – A person having origins in any of the Black racial groups of Africa.

Hispanic (temporary code) – used in MH-CLD only

Native Hawaiian or Other Pacific Islander – A person having origins in any of the original peoples of Hawaii, Guam, American Samoa, or Other Pacific Islander.

White – A person having origins in any of the original people of Europe, North Africa or the Middle East.

Some Other Race Alone – use this category for instances in which the client is not identified in any category above or whose origin group, because of area custom, is regarded as racial class distinct from the above categories (do not use this category for individuals indicating multiple or mixed races).

Two or more races – use this code when the state data system allows multiple race selection and more than one race is indicated (see guidelines).

Guidelines: For states that collect “Other Race” or allow individuals to specify a single race different from the OMB racial categories, use “Some Other Race Alone.” Note that this category should not be used if the client is indicating multiple or mixed race.

[MH-TEDS only] States that collect “Alaska Native” as a separate category should use “American Indian/Alaska Native” for other American Indians.

In cases where the method of collecting race information differs between community programs and state hospitals, or where the race reported differs between data systems, the state should resolve the issue in accordance to the state business rule in resolving data quality issues. States may use the following guidelines in the absence of a state business rule or to supplement an existing one:

1. Use the most recent race reported if variation in a person’s race was reported by the same provider across time
2. Use the most frequently reported race if variation in a person’s race was reported by different providers across time

Variable Name: Treatment Status at the End of the Reporting Period

Description: This field indicates client status at the end of the reporting period—continuing to receive services or reason for discontinuance of treatment.

Continuing Client

Discharged with Treatment Completed – All parts of the treatment plan or program were completed.

Discharged due to Lost Contact/Administrative Discharge

Discharged to Corrections, Jail – Individuals whose course of treatment is terminated because the client has been subject to jail, prison, or house confinement, or has been released by or to the courts.

Discharged due Death Discharged due to Aged Out

Discharged due to Other Specified Reasons Discharged, Reason Unknown

Discharged, Reason not Collected

Variable Name: School Grade Level – Education

Description: This field specifies the school grade level of three sub-populations of individuals, as follows:

Current grade level for school-age children (3–17 years old and 18–21 years of age if the client is in special education) who attended school at any time in the past three months.

Highest grade level completed for school-age children who have not attended school at any time within the past three months.

Highest educational attainment for all individuals, whether currently in school or not.

Guidelines: Information for this data element should be collected and reported for all individuals.

“School” includes, but is not limited to, any one or combination of home-schooling, online education, alternative school, vocational school, or regular school (public, private, charter, traditional, military, magnet, independent, parochial, etc.), at which the child is enrolled in any of the following school grade levels: nursery/preschool (including Head Start), kindergarten, elementary/middle school (Grades 1–8), high school (Grades 9–12, including General Equivalency Degree or GED), vocational school (including business, technical, secretarial, trade, or correspondence courses which are not counted as regular school enrollment and are not for recreation or adult education classes), or college/professional degree.

This measure is to be reported only once during the reporting period, either at time of client discharge or at the end of the reporting period if the client is continuing to receive services.

If the information is collected at the time when the school year just ended, report the recent school grade level completed (not the grade level the child is advancing to in the next school year).

For children who are homeschooled or children in Special Education but have been mainstreamed in regular school grades, please report the equivalent grade level.

“A nursery school is defined as a group or class that is organized to provide educational experiences for children during the year or years preceding kindergarten. It includes instruction as an important and integral phase of its program of child care. Private homes in which essentially custodial care is provided are not considered nursery schools. Children attending nursery school are classified as attending during either part of the day or the full day. Part-day attendance refers to those who attend either in the morning or in the afternoon, but not both. Full-day attendance refers to those who attend in both the morning and the afternoon. Children enrolled in Head Start programs or similar programs sponsored by local agencies to provide preschool education to young children are counted under nursery school” (The American Community Survey [ACS]).

A person who had reached the age of 18 prior to collecting this information should report the last grade completed, unless they meet the exception criteria for young individuals.

Report the highest educational attainment completed for all individuals 18 years and older (except the young individuals aged 18–21 in Special Education and receiving mental health services from the Children Mental Health System) whether they are currently in school or not.

Variable Name: Service Setting Status – Mental Health Service Setting

Description: This field indicates the type of mental health treatment setting(s) in which the client received services throughout the reporting period.

State Psychiatric Hospital – all SMHA-funded and SMHA-operated organizations operated as hospitals that provide primary inpatient care to individuals with mental illness from a specific geographical area and/or statewide.

SMHA-Funded/Operated Community-Based Program – include community mental health centers (CMHCs), outpatient clinics, partial care organizations, partial hospitalization programs, PACT programs, consumer run programs (including Club Houses and drop-in centers), and all community support programs (CSP).

Residential Treatment Center – an organization, not licensed as a psychiatric hospital, whose primary purpose is the provision of individually planned programs of mental health treatment services in conjunction with residential care for children and youth, and in some cases, adult care.

Other Psychiatric Inpatient – a private provider or medical provider licensed and/or contracted through the SMHA. This includes public/private hospitals with separate psychiatric wards.

Institutions under the Justice System – mental health services provided in a jail, prison, juvenile detention center, etc.

Guidelines: This data element refers to the place where services were provided and not particularly the residence of the client.

Variable Name: SMI/SED Status

Description: This field indicates whether the client has serious mental illness (SMI) or serious emotional disturbance (SED) using the state’s definition.

Guidelines: Report the most recent available SMI/SED status at the end of the reporting period.

Variable Name: Treatment Status at the Start of the Reporting Period

Description: This field indicates whether the client is already receiving services at the start of the reporting period (continuing) or is a new admission.

New admission – Admission that happened on the first day of the reporting period or thereafter.

Continuing – Already receiving services at the start of the reporting period. The admission date should have occurred at least a day prior to the start of the reporting period.

Variable Name: Veteran Status

Description: This field specifies whether the client has served in the uniform services (Army, Navy, Air Force, Marine Corps, Coast Guard, Public Health Service Commissioned Corps, Coast and Geodetic Survey, etc.).

Guidelines: A veteran is a person 16 years or over who has served (even for a short time), but is not now serving, on active duty in the U.S. Army, Navy, Air Force, Marine Corps, Coast Guard, or Commissioner Corps of the U.S. Public Health Service or the National Oceanic and Atmospheric Administration, or who served as a Merchant Marine seaman during World War II. Individuals who served in the National Guard or Military Reserves are classified as veterans only if they have ever been called or ordered to active duty (excluding the 4-6 months of initial training and yearly summer camps).

Appendix C Table 1. Number of states reporting data, MH-CLD and MH-TEDS 2013–2018

<i>Data item</i>	<i>Number of states</i>					
	<i>2013</i>	<i>2014</i>	<i>2015</i>	<i>2016</i>	<i>2017</i>	<i>2018</i>
Any data	45	47	45	47	48	48
Service settings	45	47	45	47	48	48
Timing of admission	45	47	45	47	48	48
First mental health diagnosis	45	47	45	47	48	48
Gender	45	47	45	47	48	48
Age	45	47	45	47	48	48
Race	45	47	45	47	48	48
Ethnicity	43	45	43	45	47	46
Detailed Hispanic	26	28	29	28	28	27
Marital status	35	38	37	40	41	41
SED/SMI status	44	47	45	47	47	48
Co-occurring substance use	45	46	44	44	45	45
Employment	44	46	44	46	47	47
Detailed not in labor force	44	46	44	46	47	47
Living arrangement	44	47	45	47	47	48
Education	33	36	34	39	41	41
Veteran status	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	<i>n/c</i>	32	34

n/c Not collected.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD), 2013–2018 reporting periods. Data as of 07.08.2019.

Appendix C Table 2. Item percentage response rate, by state or jurisdiction: MH-CLD and MH-TEDS 2018

[Based on administrative data reported to MH-CLD or MH-TEDS by all reporting states and jurisdictions. Comparisons among states should be made with caution.]

State or jurisdiction	Service setting	Timing of admission	First-mentioned Mental health diagnosis ¹	Gender	Age	Race	Ethnicity	Marital status	Co-occurring substance use		Employment status	Residence	Education	Veteran status
									SED/SMI					
Total	100	100	88	100	100	93	88	58	94	89	65	58	44	34
Alabama	100	100	99	100	100	99	100	100	91	100	82	68	99	85
Arizona	100	100	76	100	100	100	100	60	100	100	40	11	†	4
Arkansas	100	100	80	100	100	85	90	†	100	100	85	89	†	†
California	100	100	95	100	100	84	81	2	100	99	51	56	†	†
Colorado	100	100	99	100	100	88	93	100	100	100	100	100	75	100
Commonwealth of the Northern Mariana Islands	100	100	52	100	98	36	†	84	80	96	91	85	†	†
Connecticut	100	100	83	100	100	94	95	55	98	63	77	77	63	52
Delaware	100	100	89	98	100	97	93	89	86	98	90	89	69	66
District of Columbia	100	100	79	100	100	94	40	59	100	10	12	2	†	†
Florida	100	100	85	100	100	97	100	85	100	13	89	85	86	88
Hawaii	100	100	91	100	100	64	62	57	97	89	41	36	44	†
Idaho	100	100	46	100	100	91	87	46	41	46	99	35	7	81
Illinois	100	100	99	100	100	93	95	95	95	87	95	92	84	93
Indiana	100	100	100	100	100	100	98	100	100	100	93	99	96	2
Iowa	100	100	92	100	100	75	†	†	100	100	39	15	1	†
Kentucky	100	100	99	100	100	98	97	99	96	99	98	98	93	58
Louisiana	100	100	58	100	100	94	95	54	52	100	50	45	44	85
Maryland	100	100	99	100	100	97	95	86	100	100	70	86	66	95
Massachusetts	100	100	35	100	100	80	74	89	100	100	66	65	53	84
Michigan	100	100	97	100	100	99	91	97	92	100	99	100	91	95
Minnesota	100	100	94	100	100	83	92	†	100	100	99	99	96	†
Mississippi	100	100	99	100	100	99	95	99	99	†	95	98	89	97
Missouri	100	100	96	100	100	98	100	69	95	100	72	86	55	†
Montana	100	100	97	100	100	91	96	†	98	99	50	28	13	14
Nebraska	100	100	80	96	100	91	82	82	100	100	87	72	71	78
Nevada	100	100	67	100	100	80	76	71	65	95	51	31	4	26
New Mexico	100	100	100	100	100	83	83	32	100	100	38	6	18	98
New York	100	100	98	100	100	93	89	94	100	100	95	85	90	23
North Carolina	100	100	89	100	100	100	100	99	100	†	99	77	98	99

Continued. See notes at end of table.

Appendix C Table 2. Item percentage response rate, by state or jurisdiction: MH-CLD and MH-TEDS 2018 (continued)

[Based on administrative data reported to MH-CLD or MH-TEDS by all reporting states and jurisdictions. Comparisons among states should be made with caution.]

State or jurisdiction	Service setting	Timing of admission	First-mentioned Mental health diagnosis ¹					Co-occurring substance use						Veteran status
				Gender	Age	Race	Ethnicity	Marital status	SED/SMI	Employment status	Residence	Education		
Total	100	100	88	100	100	93	88	58	94	89	65	58	44	34
North Dakota	100	100	52	99	100	99	99	†	99	100	86	90	†	†
Ohio	100	100	83	98	100	88	93	37	100	100	25	2	†	†
Oklahoma	100	100	86	100	100	100	100	86	84	100	86	84	83	83
Oregon	100	100	100	100	100	88	83	76	100	99	71	96	76	71
Palau	100	100	92	100	99	100	100	57	100	100	53	63	8	30
Pennsylvania	100	100	86	100	100	99	100	99	100	100	33	†	†	†
Puerto Rico	100	100	91	100	100	94	99	98	94	98	92	98	96	69
Rhode Island	100	100	78	100	100	84	93	91	80	45	86	83	†	87
South Carolina	100	100	97	100	100	97	79	98	100	100	83	93	91	100
South Dakota	100	100	94	100	100	100	100	100	90	49	55	38	5	†
Tennessee	100	100	54	100	100	93	65	59	54	74	59	70	34	34
Texas	100	100	67	100	100	100	100	78	95	71	72	71	70	†
Utah	100	100	86	100	100	92	91	90	92	100	92	93	83	91
Vermont	100	100	93	100	100	69	63	72	3	†	59	65	2	63
Virginia	100	100	95	100	100	99	97	†	91	100	96	93	88	†
Washington	100	100	93	99	100	97	95	57	58	100	60	54	50	50
West Virginia	100	100	85	100	100	99	†	94	87	95	92	99	89	87
Wisconsin	100	100	89	100	100	94	88	†	100	100	44	40	†	†
Wyoming	100	100	97	100	100	95	96	96	99	100	99	98	96	34

† Less than 1 percent of individuals.

¹ The first-mentioned mental health diagnosis is the first of up to three diagnoses reported for each individual. Mental health diagnostic category definitions can be found in Appendix E.

SOURCE: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Mental Health Treatment Episode Data Set (MH-TEDS) and Mental Health Client-Level Data (MH-CLD). Data received 07.08.2019 as part of 2018 reporting period.

APPENDIX D

SMI/SED FEDERAL DEFINITIONS

Note that identification of individuals with a serious mental illness (SMI) or serious emotional disturbance (SED) is based on each state or jurisdiction's definitions for SMI and SED; some states and jurisdictions may use definitions that vary from the federal definitions that appear in this appendix.

Serious Mental Illness (SMI)

Pursuant to section 1912 (c) of the Public Health Service Act, as amended by Public Law 102-321 "adult[s] with a serious mental illness" are individuals:

1. age 18 years and older;
2. who currently or at any time during the past year, have had a diagnosable mental, behavioral, or emotional disorder of sufficient duration to meet diagnostic criteria specified within DSM-III-R; and
3. this disorder has resulted in functional impairment which substantially interferes with or limits one or more major life activities.

"These disorders include any mental disorders (including those of biological etiology) listed in DSM-III-R or their ICD-9-CM equivalent (and subsequent revisions), with the exception of DSM-III-R "V" codes, substance use disorders, and developmental disorders, which are excluded, unless they co-occur with another diagnosable serious mental illness..." (p. 29425).

Serious Emotional Disturbance (SED)

Pursuant to section 1912 (c) of the Public Health Service Act, as amended by Public Law 102-321 "children with a serious emotional disturbance" are individuals:

1. from birth up to age 18;
2. who currently or at any time during the past year have had a diagnosable mental, behavioral, or emotional disorder of sufficient duration to meet diagnostic criteria specified within DSM-III-R; and
3. this disorder resulted in functional impairment, which substantially interferes with or limits the child's role or functioning in family, school, or community activities (p.29425).

"These disorders include any mental disorder (including those of biological etiology) listed in DSM-III-R or their ICD-9-CM equivalent (and subsequent revisions) with the exception of DSM-III-R 'V' codes, substance use, and developmental disorders, which are excluded, unless they co-occur with another diagnosable serious emotional disturbance..." (p. 29425).

"Functional impairment is defined as difficulties that substantially interfere with or limit a child or adolescent from achieving or maintaining one or more developmentally-appropriate social,

behavioral, cognitive, communicative, or adaptive skills. Functional impairments of episodic, recurrent, and continuous duration are included unless they are temporary and expected responses to stressful events in their environment. Children who would have met functional impairment criteria during the referenced year without the benefit of treatment or other support services are included in this definition....” (p. 29425) [Source: Federal Register 58(96): 29422–29425. May 20, 1993; <http://www.samhsa.gov/sites/default/files/federal-register-notice-58-96-definitions.pdf>].

APPENDIX E

MENTAL HEALTH AND SUBSTANCE USE DIAGNOSIS CODES

Mental Health Diagnostic Groups With Corresponding ICD-9 Diagnostic Codes

The listings below group ICD-9 codes into a modified form of Clinical Classification Software (CCS) mental health “multi-level” diagnostic category codes (5.1, 5.2, and so on) and corresponding labels (trauma- and stressor-related disorders, anxiety disorders, and so on). Notes:

- **Bold** diagnostic groups are displayed in one or more tables in this report; records with ICD-9 codes in other diagnostic groups are grouped together as “other” mental health diagnoses.
- *Italicized codes* in parentheses are nonstandard ICD-9 codes in the data that were recognized to belong to the same CCS mental health diagnostic group.
- Entries in brackets ([650], [651], and so on) are single-level diagnosis categories that are otherwise ignored in this report but are noted here for convenience.
- The ICD-9 diagnostic codes shown below have omitted the dot separator after the first three digits.
- The groupings of certain DSM-IV/ICD-9 codes have been reorganized according to DSM-5 nomenclature.

5.1 **Trauma- and stressor-related disorders** [650]^{15, 16}

3083	3090	3091	30922	30923	30924	30928	30929	3093
3094	30981	30982	30983	30989	3099			
<i>(30830</i>	<i>309</i>	<i>30900</i>	<i>30910</i>	<i>30920</i>	<i>30930</i>	<i>30940</i>	<i>30980</i>	<i>3098</i>
<i>30990)</i>								

5.2 **Anxiety disorders** [651]¹⁷

29384	30000	30001	30002	30009	30010	30020	30021	30022
30023	30029	3003	3005	30089	3009	3080	3081	3082
3084	3089	30921	3130	3131	31321	31322	3133	31382
31383								
<i>(3000</i>	<i>300</i>	<i>3001</i>	<i>3002</i>	<i>30030</i>	<i>30050</i>	<i>30090</i>	<i>30800</i>	<i>308</i>
<i>30810</i>	<i>30820</i>	<i>30840</i>	<i>30890</i>	<i>313</i>	<i>31300</i>	<i>3138</i>	<i>31380)</i>	

¹⁵ For these and other bolded (i.e., tabulated) CCS diagnostic groups, table column and row headers drop the word “disorders” from the description.

¹⁶ Codes for posttraumatic stress disorder and acute stress disorder diagnoses have been grouped under *trauma- and stressor-related disorders*, consistent with DSM-5. Codes for adjustment disorders have also been subsumed into this diagnostic group.

¹⁷ The code for separation anxiety disorder has been grouped with anxiety disorders, consistent with DSM-5.

5.3 Attention deficit, conduct, and disruptive behavior disorders [652]

5.3.1 Conduct disorder [6521]

31200	31201	31202	31203	31210	31211	31212	31213	31220
31221	31222	31223	3124	3128	31281	31282	31289	3129
(312	3120	3121	3122	31240	31280	31290)		

5.3.2 Oppositional defiant disorder [6522]

31381

5.3.3 Attention-deficit/hyperactivity disorder [6523]

31400	31401	3141	3142	3148	3149			
(314	3140	31410	31480	31490)				

5.4 Delirium, dementia, and amnestic and other cognitive disorders [653]¹⁸

2900	29010	29011	29012	29013	29020	29021	2903	29040
29041	29042	29043	2908	2909	2930	2931	2940	2941
29410	29411	29420	29421	2948	2949	3100	3102	3108
31081	31089	3109	3310	3311	33111	33119	3312	33182
797								
(29000	290	2901	2902	29030	2904	29080	29090	293
29300	29400	294	2942	29480	29490	310	31000	31020
31090	331	33100	33110)					

5.5 Developmental disorders [654]

5.5.1 Communication disorders [6541]

3070	3079	31531	31534	31535	31539	V401		
(30700	307	30790)						

5.5.2 Developmental disabilities [6542]

31501	31502	31509	31532	3155	3158			
(31550	1580)							

5.5.3 Intellectual disabilities [6543]

317	3180	3181	3182	319				
(3170	31700	318	31800	31810	31820	31900)		

5.5.4 Learning disorders [6544]

31500	3151	3152	3159	V400				
(3150	315	31510	31520	31540)				

5.5.5 Motor skill disorders [6545]

3154
(31540)

¹⁸ Referred to as “delirium/dementia” in tables and text.

5.6 Disorders usually diagnosed in infancy, childhood, or adolescence [655]

5.6.1 Elimination disorders [6551]

3076 3077
(30760 30770)

5.6.2 Other disorders of infancy, childhood, or adolescence [6552]

3073 31323 31389 3139
(31390)

5.6.3 **Pervasive developmental disorders** [6553]

29900 29901 29910 29911 29980 29981 29990 29991
(299 2991 2998 2999)

5.6.4 Tic disorders [6554]

30720 30721 30722 30723

5.7 Impulse control disorders, not elsewhere classified [656]

31230 31231 31232 31233 31234 31235 31239
(3123)

5.8 Mood disorders [657]

5.8.1 **Bipolar disorders** [6571]

29600 29601 29602 29603 29604 29605 29606 29610 29611
29612 29613 29614 29615 29616 29640 29641 29642 29643
29644 29645 29646 29650 29651 29652 29653 29654 29655
29656 29660 29661 29662 29663 29664 29665 29666 2967
29680 29681 29682 29689 29690 29699
(296 2960 2961 2964 2965 2966 29670 2968 2969)

5.8.2 **Depressive disorders** [6572]

29383 29620 29621 29622 29623 29624 29625 29626 29630
29631 29632 29633 29634 29635 29636 3004 311
(2962 2963 30040 31100 3110)

5.9 **Personality disorders** [658]

3010 30110 30111 30112 30113 30120 30121 30122 3013
3014 30150 0151 30159 3016 3017 30181 30182 30183
30184 30189 3019
(30100 301 3011 3012 30130 30140 3015 30160 30170
30190)

5.10 Schizophrenia and other psychotic disorders [659]¹⁹

29381	29382	29500	29501	29502	29503	29504	29505	29510
29511	29512	29513	29514	29515	29520	29521	29522	29523
29524	29525	29530	29531	29532	29533	29534	29535	29540
29541	29542	29543	29544	29545	29550	29551	29552	29553
29554	29555	29560	29561	29562	29563	29564	29565	29570
29571	29572	29573	29574	29575	29580	29581	29582	29583
29584	29585	29590	29591	29592	29593	29594	29595	2970
2971	2972	2973	2978	2979	2980	2981	2982	2983
2984	2988	2989						
(295	2950	2951	2952	2953	2954	2955	2956	2957
2958	2959	29700	297	29710	29720	20730	20780	20790
29800	298	29810	29820	29830	29840	29880	29890)	

5.11 Alcohol-related disorders [660]²⁰

2910	2911	2912	2913	2914	2915	2918	29181	29182
29189	2919	30300	30301	30302	30303	30390	30391	30392
30393	30500	30501	30502	30503	3575	4255	53530	53531
5710	5711	5712	5713	76071	9800			
(29100	291	29110	29120	29130	29140	29150	29180	20190
303	3030	3039	305	3050)				

5.12 Substance-related disorders [661]²¹

2920	29211	29212	2922	29281	29282	29283	29284	29285
29289	2929	30400	30401	30402	30403	30410	30411	30412
30413	30420	30421	30422	30423	30430	30431	30432	30433
30440	30441	30442	30443	30450	30451	30452	30453	30460
30461	30462	30463	30470	30471	30472	30473	30480	30481
30482	30483	30490	30491	30492	30493	30520	30521	30522
30523	30530	30531	30532	30533	30540	30541	30542	30543
30550	30551	30552	30553	30560	30561	30562	30563	30570
30571	30572	30573	30580	30581	30582	30583	30590	30591
30592	30593	64830	64831	64832	64833	64834	65550	65551
65553	76072	76073	76075	7795	96500	96501	96502	96509
V6542								
(292	29200	20220	20290	304	3040	3041	3042	3043
3044	3045	3046	3047	3048	3049	3052	3053	3054
3055	3057	3058	3059)					

¹⁹ Referred to as “Schizophrenia, other psychotic” in tables.

²⁰ Regardless of their “co-occurring” flag in the data, individuals reported to be diagnosed with only alcohol-related disorders and/or substance-related disorders are not counted as having co-occurring mental health and substance use disorders for the purpose of Chapter 6; some other, non-alcohol and non-substance-related disorder diagnosis is necessary.

²¹ See footnote 20.

5.13 Suicide and intentional self-inflicted injury [662]

E9500	E9501	E9502	E9503	E9504	E9505	E9506	E9507	E9508
E9509	E9510	E9511	E9518	E9520	E9521	E9528	E9529	E9530
E9531	E9538	E9539	E954	E9550	E9551	E9552	E9553	E9554
E9555	E9556	E9557	E9559	E956	E9570	E9571	E9572	E9579
E9580	E9581	E9582	E9583	E9584	E9585	E9586	E9587	E9588
E9589	E959	V6284						

5.14 Screening and history of mental health and substance use codes [663]

33392	V110	V111	V112	V114	V118	V119	V154	V1541
V1542	V1549	V1582	V6285	V663	V701	V702	V7101	V7102
V7109	V790	V792	V793	7903	V113	V798	V791	V799
3051	30510	30511	30512	30513	5353			

5.15 Miscellaneous mental disorders [670]

30012	30013	30014	30015	3006	3071	30750	30751	30752
30753	30754	30759	30016	30019	3060	3061	3062	3063
3064	30650	30652	30653	30659	3066	3067	3068	3069
3021	3022	3023	3024	30250	30251	30252	30253	3026
30270	30271	30272	30273	30274	30275	30276	30279	30281
30282	30283	30284	30285	30289	3029	30651	30740	30741
30742	30743	30744	30745	30746	30747	30748	30749	30011
3007	30081	30082	30780	30781	30789	29389	2939	3101
316	64840	64841	64842	64843	64844	V402	V403	V4031
V4039	V409	V673						
<i>(30060</i>	<i>30710</i>	<i>3075</i>	<i>306</i>	<i>30600</i>	<i>30610</i>	<i>30630</i>	<i>30640</i>	<i>30680</i>
<i>30690</i>	<i>30210</i>	<i>30220</i>	<i>30230</i>	<i>30240</i>	<i>3025</i>	<i>30260</i>	<i>3027</i>	<i>30290</i>
<i>30070</i>	<i>3078</i>	<i>29390</i>	<i>31010</i>	<i>31600</i>	<i>316</i>	<i>3160</i>	<i>6484</i>	<i>V4020</i>
<i>V4030</i>	<i>V4090)</i>							

Mental Health Diagnostic Groups With Corresponding ICD-10 Diagnostic Codes

The listings below group ICD-10 codes into a modified form of Clinical Classification Software (CCS) mental health “multi-level” diagnostic category codes (5.1, 5.2, and so on) and corresponding labels (trauma- and stressor-related disorders, anxiety disorders, and so on). Notes:

- **Bold** diagnostic groups are displayed in one or more tables in this report; records with ICD-10 codes in other diagnostic groups are grouped together as “other” mental health diagnoses.
- Entries in brackets ([650], [651], and so on) are single-level diagnosis categories that are otherwise ignored in this report but are noted here for convenience.
- The ICD-10 diagnostic codes shown below have omitted the dot separator after the first three digits.
- Some diagnostic groups implemented by CCS for ICD-9 codes were not implemented for ICD-10 codes (conduct disorders, oppositional defiant disorders, attention deficit and attention deficit hyperactivity disorders, pervasive developmental disorders, bipolar disorders, and depressive disorders). ICD-10 codes from broader CCS diagnostic categories were assigned to these more specific diagnostic groups, based on the ICD-10 code’s detailed meaning.
- The groupings of certain ICD-10 codes have been reorganized according to DSM-5 nomenclature.

5.1 **Trauma- and stressor-related disorders** [650]^{22, 23}

F4310	F4320	F4321	F4322	F4323	F4324	F4325	F4329	F438
F439								

5.2 **Anxiety disorders** [651]²⁴

F064	F4000	F4001	F4002	F4010	F4011	F40210	F40218	F40220
F40228	F40230	F40231	F40232	F40233	F40240	F40241	F40242	F40243
F40248	F40290	F40291	F40298	F408	F409	F410	F411	F413
F418	F419	F42	F430	F4311	F4312	F488	F489	R452
R453	R454	R455	R456	R457	R4581	R4582	R4583	R4584
F930								

5.3.1 **Conduct disorder** [6521]²⁵

F910	F911	F912	F918	F919
------	------	------	------	------

5.3.2 **Oppositional defiant disorder** [6522]

F913

²² For these and other bolded (i.e., tabulated) CCS diagnostic groups, table column and row headers drop the word “disorders” from the description.

²³ Codes for posttraumatic stress disorder and acute stress disorder diagnoses have been grouped under *trauma- and stressor-related disorders*, consistent with DSM-5. Codes for adjustment disorders have also been subsumed into this diagnostic group.

²⁴ The code for separation anxiety disorder has been grouped with anxiety disorders, consistent with DSM-5.

²⁵ Codes in this category were assigned from a more general CCS diagnostic group to this one based on the code’s detailed meaning.

5.3.3 Attention-deficit/hyperactivity disorder [6523]²⁵

F900	F901	F902	F908	F909	R460	R461	R462	R463
R464	R465	R466	R467	R4681	R4689			

5.4 Delirium, dementia, and amnestic and other cognitive disorders [653]²⁶

F0150	F0151	F0280	F0281	F0390	F0391	F04	F05	F070
F0781	F0789	F079	F09	F482	G300	G301	G308	G309
G3101	G3109	G311	G3183	R4181	R54			

5.5 Developmental disorders [654]

F70	F71	F72	F73	F78	F79	F800	F801	F802
F804	F8081	F8089	F809	F810	F812	F8181	F8189	F819
F82	F88	F89	F985	R4183	R480			

5.6 Disorders usually diagnosed in infancy, childhood, or adolescence [655]

F642	F938	F939	F940	F941	F942	F948	F949	F950
F951	F952	F958	F959	F980	F981	F9821	F9829	F983
F984	F988	F989						

5.6.3 Pervasive developmental disorders [6553]²⁷

F840	F842	F843	F845	F848	F849			
------	------	------	------	------	------	--	--	--

5.7 Impulse control disorders not elsewhere classified [656]

F630	F631	F632	F633	F6381	F6389	F639	R45850	
------	------	------	------	-------	-------	------	--------	--

5.8 Mood disorders [657]²⁷

F0630	F0631	F0632	F0633	F0634	F3010	F3011	F3012	F3013
F302	F303	F304	F308	F309	F340	F341	F348	F349
F39	R4586							

5.8.1 Bipolar disorders [6571]

F310	F3110	F3111	F3112	F3113	F312	F3130	F3131	F3132
F314	F315	F3160	F3161	F3162	F3163	F3164	F3170	F3171
F3172	F3173	F3174	F3175	F3176	F3177	F3178	F3181	F3189
F319								

5.8.2 Depressive disorders [6572]

F320	F321	F322	F323	F324	F325	F328	F329	F330
F331	F332	F333	F3340	F3341	F3342	F338	F339	

5.9 Personality disorders [658]

F600	F601	F602	F603	F604	F605	F606	F607	F6081
F6089	F609	F69						

²⁶ Referred to as “delirium/dementia” in tables and text.

²⁷ Mood disorder codes that could not be assigned to bipolar or depressive disorders.

5.10 Schizophrenia and other psychotic disorders [659]²⁸

F060	F062	F200	F201	F202	F203	F205	F2081	F2089
F209	F21	F22	F23	F24	F250	F251	F258	F259
F28	F29							

5.11 Alcohol-related disorders [660]²⁹

F1010	F10120	F10121	F10129	F1014	F10150	F10151	F10159	F10180
F10181	F10182	F10188	F1019	F1020	F1021	F10220	F10221	F10229
F10230	F10231	F10232	F10239	F1024	F10250	F10251	F10259	F1026
F1027	F10280	F10281	F10282	F10288	F1029	F10920	F10921	F10929
F1094	F10950	F10951	F10959	F1096	F1097	F10980	F10981	F10982
F10988	F1099	G621	I426	K2920	K2921	K700	K7010	K7011
K702	K7030	K7031	K7040	K709	O99310	O99311	O99312	O99313
O99314	O99315	P043	Q860					

5.12 Substance-related disorders [661]³⁰

F1110	F11120	F11121	F11122	F11129	F1114	F11150	F11151	F11159
F11181	F11182	F11188	F1119	F1120	F1121	F11220	F11221	F11222
F11229	F1123	F1124	F11250	F11251	F11259	F11281	F11282	F11288
F1129	F1190	F11920	F11921	F11922	F11929	F1193	F1194	F11950
F11951	F11959	F11981	F11982	F11988	F1199	F1210	F12120	F12121
F12122	F12129	F12150	F12151	F12159	F12180	F12188	F1219	F1220
F1221	F12220	F12221	F12222	F12229	F12250	F12251	F12259	F12280
F12288	F1229	F1290	F12920	F12921	F12922	F12929	F12950	F12951
F12959	F12980	F12988	F1299	F1310	F13120	F13121	F13129	F1314
F13150	F13151	F13159	F13180	F13181	F13182	F13188	F1319	F1320
F1321	F13220	F13221	F13229	F13230	F13231	F13232	F13239	F1324
F13250	F13251	F13259	F1326	F1327	F13280	F13281	F13282	F13288
F1329	F1390	F13920	F13921	F13929	F13930	F13931	F13932	F13939
F1394	F13950	F13951	F13959	F1396	F1397	F13980	F13981	F13982
F13988	F1399	F1410	F14120	F14121	F14122	F14129	F1414	F14150
F14151	F14159	F14180	F14181	F14182	F14188	F1419	F1420	F1421
F14220	F14221	F14222	F14229	F1423	F1424	F14250	F14251	F14259
F14280	F14281	F14282	F14288	F1429	F1490	F14920	F14921	F14922
F14929	F1494	F14950	F14951	F14959	F14980	F14981	F14982	F14988
F1499	F1510	F15120	F15121	F15122	F15129	F1514	F15150	F15151
F15159	F15180	F15181	F15182	F15188	F1519	F1520	F1521	F15220
F15221	F15222	F15229	F1523	F1524	F15250	F15251	F15259	F15280
F15281	F15282	F15288	F1529	F1590	F15920	F15921	F15922	F15929
F1593	F1594	F15950	F15951	F15959	F15980	F15981	F15982	F15988
F1599	F1610	F16120	F16121	F16122	F16129	F1614	F16150	F16151
F16159	F16180	F16183	F16188	F1619	F1620	F1621	F16220	F16221
F16229	F1624	F16250	F16251	F16259	F16280	F16283	F16288	F1629
F1690	F16920	F16921	F16929	F1694	F16950	F16951	F16959	F16980
F16983	F16988	F1699	F17200	F17201	F17203	F17208	F17209	F17210

²⁸ Referred to as “Schizophrenia, other psychotic” in tables.

²⁹ Regardless of their “co-occurring” flag in the data, individuals reported to be diagnosed with only alcohol-related disorders and/or substance-related disorders are not counted as having co-occurring mental health and substance use disorders for the purpose of Chapter 6; some other, non-alcohol and non-substance-related disorder diagnosis is necessary.

³⁰ See footnote 25.

F17211	F17213	F17218	F17219	F17220	F17221	F17223	F17228	F17229
F17290	F17291	F17293	F17298	F17299	F1810	F18120	F18121	F18129
F1814	F18150	F18151	F18159	F1817	F18180	F18188	F1819	F1820
F1821	F18220	F18221	F18229	F1824	F18250	F18251	F18259	F1827
F18280	F18288	F1829	F1890	F18920	F18921	F18929	F1894	F18950
F18951	F18959	F1897	F18980	F18988	F1899	F1910	F19120	F19121
F19122	F19129	F1914	F19150	F19151	F19159	F1916	F1917	F19180
F19181	F19182	F19188	F1919	F1920	F1921	F19220	F19221	F19222
F19229	F19230	F19231	F19232	F19239	F1924	F19250	F19251	F19259
F1926	F1927	F19280	F19281	F19282	F19288	F1929	F1990	F19920
F19921	F19922	F19929	F19930	F19931	F19932	F19939	F1994	F19950
F19951	F19959	F1996	F1997	F19980	F19981	F19982	F19988	F1999
F550	F551	F552	F553	F554	F558	O355XX0	O355XX1	O355XX2
O355XX3	O355XX4	O355XX5	O355XX9	O99320	O99321	O99322	O99323	O99324
O99325	P0441	P0449	P961	P962	T400X1A	T400X1D	T400X1S	T400X3A
T400X3D	T400X3S	T400X4A	T400X4D	T400X4S	T400X5A	T400X5D	T400X5S	T400X6A
T400X6D	T400X6S	T401X1A	T401X1D	T401X1S	T401X3A	T401X3D	T401X3S	T401X4A
T401X4D	T401X4S	T401X5A	T401X5D	T401X5S	T405X1A	T405X1D	T405X1S	T405X3A
T405X3D	T405X3S	T405X4A	T405X4D	T405X4S	T405X5A	T405X5D	T405X5S	T405X6A
T405X6D	T405X6S	T407X1A	T407X1D	T407X1S	T407X3A	T407X3D	T407X3S	T407X4A
T407X4D	T407X4S	T407X5A	T407X5D	T407X5S	T407X6A	T407X6D	T407X6S	T408X1A
T408X1D	T408X1S	T408X3A	T408X3D	T408X3S	T408X4A	T408X4D	T408X4S	T408X5A
T408X5D	T408X5S	T40901A	T40901D	T40901S	T40903A	T40903D	T40903S	T40904A
T40904D	T40904S	T40905A	T40905D	T40905S	T40906A	T40906D	T40906S	T40991A
T40991D	T40991S	T40993A	T40993D	T40993S	T40994A	T40994D	T40994S	T40995A
T40995D	T40995S	T40996A	T40996D	T40996S				

5.13 Suicide and intentional self-inflicted injury [662]

R45851	T1491	T360X2A	T360X2D	T360X2S	T361X2A	T361X2D	T361X2S	T362X2A
T362X2D	T362X2S	T363X2A	T363X2D	T363X2S	T364X2A	T364X2D	T364X2S	T365X2A
T365X2D	T365X2S	T366X2A	T366X2D	T366X2S	T367X2A	T367X2D	T367X2S	T368X2A
T368X2D	T368X2S	T3692XA	T3692XD	T3692XS	T370X2A	T370X2D	T370X2S	T371X2A
T371X2D	T371X2S	T372X2A	T372X2D	T372X2S	T373X2A	T373X2D	T373X2S	T374X2A
T374X2D	T374X2S	T375X2A	T375X2D	T375X2S	T378X2A	T378X2D	T378X2S	T3792XA
T3792XD	T3792XS	T380X2A	T380X2D	T380X2S	T381X2A	T381X2D	T381X2S	T382X2A
T382X2D	T382X2S	T383X2A	T383X2D	T383X2S	T384X2A	T384X2D	T384X2S	T385X2A
T385X2D	T385X2S	T386X2A	T386X2D	T386X2S	T387X2A	T387X2D	T387X2S	T38802A
T38802D	T38802S	T38812A	T38812D	T38812S	T38892A	T38892D	T38892S	T38902A
T38902D	T38902S	T38992A	T38992D	T38992S	T39012A	T39012D	T39012S	T39092A
T39092D	T39092S	T391X2A	T391X2D	T391X2S	T392X2A	T392X2D	T392X2S	T39312A
T39312D	T39312S	T39392A	T39392D	T39392S	T394X2A	T394X2D	T394X2S	T398X2A
T398X2D	T398X2S	T3992XA	T3992XD	T3992XS	T400X2A	T400X2D	T400X2S	T401X2A
T401X2D	T401X2S	T402X2A	T402X2D	T402X2S	T403X2A	T403X2D	T403X2S	T404X2A
T404X2D	T404X2S	T405X2A	T405X2D	T405X2S	T40602A	T40602D	T40602S	T40692A
T40692D	T40692S	T407X2A	T407X2D	T407X2S	T408X2A	T408X2D	T408X2S	T40902A
T40902D	T40902S	T40992A	T40992D	T40992S	T410X2A	T410X2D	T410X2S	T411X2A
T411X2D	T411X2S	T41202A	T41202D	T41202S	T41292A	T41292D	T41292S	T413X2A
T413X2D	T413X2S	T4142XA	T4142XD	T4142XS	T415X2A	T415X2D	T415X2S	T420X2A
T420X2D	T420X2S	T421X2A	T421X2D	T421X2S	T422X2A	T422X2D	T422X2S	T423X2A
T423X2D	T423X2S	T424X2A	T424X2D	T424X2S	T425X2A	T425X2D	T425X2S	T426X2A
T426X2D	T426X2S	T4272XA	T4272XD	T4272XS	T428X2A	T428X2D	T428X2S	T43012A
T43012D	T43012S	T43022A	T43022D	T43022S	T431X2A	T431X2D	T431X2S	T43202A
T43202D	T43202S	T43212A	T43212D	T43212S	T43222A	T43222D	T43222S	T43292A

T43292D T43292S T433X2A T433X2D T433X2S T434X2A T434X2D T434X2S T43502A
T43502D T43502S T43592A T43592D T43592S T43602A T43602D T43602S T43612A
T43612D T43612S T43622A T43622D T43622S T43632A T43632D T43632S T43692A
T43692D T43692S T438X2A T438X2D T438X2S T4392XA T4392XD T4392XS T440X2A
T440X2D T440X2S T441X2A T441X2D T441X2S T442X2A T442X2D T442X2S T443X2A
T443X2D T443X2S T444X2A T444X2D T444X2S T445X2A T445X2D T445X2S T446X2A
T446X2D T446X2S T447X2A T447X2D T447X2S T448X2A T448X2D T448X2S T44902A
T44902D T44902S T44992A T44992D T44992S T450X2A T450X2D T450X2S T451X2A
T451X2D T451X2S T452X2A T452X2D T452X2S T453X2A T453X2D T453X2S T454X2A
T454X2D T454X2S T45512A T45512D T45512S T45522A T45522D T45522S T45602A
T45602D T45602S T45612A T45612D T45612S T45622A T45622D T45622S T45692A
T45692D T45692S T457X2A T457X2D T457X2S T458X2A T458X2D T458X2S T4592XA
T4592XD T4592XS T460X2A T460X2D T460X2S T461X2A T461X2D T461X2S T462X2A
T462X2D T462X2S T463X2A T463X2D T463X2S T464X2A T464X2D T464X2S T465X2A
T465X2D T465X2S T466X2A T466X2D T466X2S T467X2A T467X2D T467X2S T468X2A
T468X2D T468X2S T46902A T46902D T46902S T46992A T46992D T46992S T470X2A
T470X2D T470X2S T471X2A T471X2D T471X2S T472X2A T472X2D T472X2S T473X2A
T473X2D T473X2S T474X2A T474X2D T474X2S T475X2A T475X2D T475X2S T476X2A
T476X2D T476X2S T477X2A T477X2D T477X2S T478X2A T478X2D T478X2S T4792XA
T4792XD T4792XS T480X2A T480X2D T480X2S T481X2A T481X2D T481X2S T48202A
T48202D T48202S T48292A T48292D T48292S T483X2A T483X2D T483X2S T484X2A
T484X2D T484X2S T485X2A T485X2D T485X2S T486X2A T486X2D T486X2S T48902A
T48902D T48902S T48992A T48992D T48992S T490X2A T490X2D T490X2S T491X2A
T491X2D T491X2S T492X2A T492X2D T492X2S T493X2A T493X2D T493X2S T494X2A
T494X2D T494X2S T495X2A T495X2D T495X2S T496X2A T496X2D T496X2S T497X2A
T497X2D T497X2S T498X2A T498X2D T498X2S T4992XA T4992XD T4992XS T500X2A
T500X2D T500X2S T501X2A T501X2D T501X2S T502X2A T502X2D T502X2S T503X2A
T503X2D T503X2S T504X2A T504X2D T504X2S T505X2A T505X2D T505X2S T506X2A
T506X2D T506X2S T507X2A T507X2D T507X2S T508X2A T508X2D T508X2S T50902A
T50902D T50902S T50992A T50992D T50992S T50A12A T50A12D T50A12S T50A22A
T50A22D T50A22S T50A92A T50A92D T50A92S T50B12A T50B12D T50B12S T50B92A
T50B92D T50B92S T50Z12A T50Z12D T50Z12S T50Z92A T50Z92D T50Z92S T510X2A
T510X2D T510X2S T511X2A T511X2D T511X2S T512X2A T512X2D T512X2S T513X2A
T513X2D T513X2S T518X2A T518X2D T518X2S T5192XA T5192XD T5192XS T520X2A
T520X2D T520X2S T521X2A T521X2D T521X2S T522X2A T522X2D T522X2S T523X2A
T523X2D T523X2S T524X2A T524X2D T524X2S T528X2A T528X2D T528X2S T5292XA
T5292XD T5292XS T530X2A T530X2D T530X2S T531X2A T531X2D T531X2S T532X2A
T532X2D T532X2S T533X2A T533X2D T533X2S T534X2A T534X2D T534X2S T535X2A
T535X2D T535X2S T536X2A T536X2D T536X2S T537X2A T537X2D T537X2S T5392XA
T5392XD T5392XS T540X2A T540X2D T540X2S T541X2A T541X2D T541X2S T542X2A
T542X2D T542X2S T543X2A T543X2D T543X2S T5492XA T5492XD T5492XS T550X2A
T550X2D T550X2S T551X2A T551X2D T551X2S T560X2A T560X2D T560X2S T561X2A
T561X2D T561X2S T562X2A T562X2D T562X2S T563X2A T563X2D T563X2S T564X2A
T564X2D T564X2S T565X2A T565X2D T565X2S T566X2A T566X2D T566X2S T567X2A
T567X2D T567X2S T56812A T56812D T56812S T56892A T56892D T56892S T5692XA
T5692XD T5692XS T570X2A T570X2D T570X2S T571X2A T571X2D T571X2S T572X2A
T572X2D T572X2S T573X2A T573X2D T573X2S T578X2A T578X2D T578X2S T5792XA
T5792XD T5792XS T5802XA T5802XD T5802XS T5812XA T5812XD T5812XS T582X2A
T582X2D T582X2S T588X2A T588X2D T588X2S T5892XA T5892XD T5892XS T590X2A
T590X2D T590X2S T591X2A T591X2D T591X2S T592X2A T592X2D T592X2S T593X2A
T593X2D T593X2S T594X2A T594X2D T594X2S T595X2A T595X2D T595X2S T596X2A
T596X2D T596X2S T597X2A T597X2D T597X2S T59812A T59812D T59812S T59892A
T59892D T59892S T5992XA T5992XD T5992XS T600X2A T600X2D T600X2S T601X2A
T601X2D T601X2S T602X2A T602X2D T602X2S T603X2A T603X2D T603X2S T604X2A

T604X2D T604X2S T608X2A T608X2D T608X2S T6092XA T6092XD T6092XS T6102XA
T6102XD T6102XS T6112XA T6112XD T6112XS T61772A T61772D T61772S T61782A
T61782D T61782S T618X2A T618X2D T618X2S T6192XA T6192XD T6192XS T620X2A
T620X2D T620X2S T621X2A T621X2D T621X2S T622X2A T622X2D T622X2S T628X2A
T628X2D T628X2S T6292XA T6292XD T6292XS T63002A T63002D T63002S T63012A
T63012D T63012S T63022A T63022D T63022S T63032A T63032D T63032S T63042A
T63042D T63042S T63062A T63062D T63062S T63072A T63072D T63072S T63082A
T63082D T63082S T63092A T63092D T63092S T63112A T63112D T63112S T63122A
T63122D T63122S T63192A T63192D T63192S T632X2A T632X2D T632X2S T63302A
T63302D T63302S T63312A T63312D T63312S T63322A T63322D T63322S T63332A
T63332D T63332S T63392A T63392D T63392S T63412A T63412D T63412S T63422A
T63422D T63422S T63432A T63432D T63432S T63442A T63442D T63442S T63452A
T63452D T63452S T63462A T63462D T63462S T63482A T63482D T63482S T63512A
T63512D T63512S T63592A T63592D T63592S T63612A T63612D T63612S T63622A
T63622D T63622S T63632A T63632D T63632S T63692A T63692D T63692S T63712A
T63712D T63712S T63792A T63792D T63792S T63812A T63812D T63812S T63822A
T63822D T63822S T63832A T63832D T63832S T63892A T63892D T63892S T6392XA
T6392XD T6392XS T6402XA T6402XD T6402XS T6482XA T6482XD T6482XS T650X2A
T650X2D T650X2S T651X2A T651X2D T651X2S T65212A T65212D T65212S T65222A
T65222D T65222S T65292A T65292D T65292S T653X2A T653X2D T653X2S T654X2A
T654X2D T654X2S T655X2A T655X2D T655X2S T656X2A T656X2D T656X2S T65812A
T65812D T65812S T65822A T65822D T65822S T65832A T65832D T65832S T65892A
T65892D T65892S T6592XA T6592XD T6592XS T71112A T71112D T71112S T71122A
T71122D T71122S T71132A T71132D T71132S T71152A T71152D T71152S T71162A
T71162D T71162S T71192A T71192D T71192S T71222A T71222D T71222S T71232A
T71232D T71232S X710XXA X710XXD X710XXS X711XXA X711XXD X711XXS X712XXA
X712XXD X712XXS X713XXA X713XXD X713XXS X718XXA X718XXD X718XXS X719XXA
X719XXD X719XXS X72XXXA X72XXXD X72XXXS X730XXA X730XXD X730XXS X731XXA
X731XXD X731XXS X732XXA X732XXD X732XXS X738XXA X738XXD X738XXS X739XXA
X739XXD X739XXS X7401XA X7401XD X7401XS X7402XA X7402XD X7402XS X7409XA
X7409XD X7409XS X748XXA X748XXD X748XXS X749XXA X749XXD X749XXS X75XXXA
X75XXXD X75XXXS X76XXXA X76XXXD X76XXXS X770XXA X770XXD X770XXS X771XXA
X771XXD X771XXS X772XXA X772XXD X772XXS X773XXA X773XXD X773XXS X778XXA
X778XXD X778XXS X779XXA X779XXD X779XXS X780XXA X780XXD X780XXS X781XXA
X781XXD X781XXS X782XXA X782XXD X782XXS X788XXA X788XXD X788XXS X789XXA
X789XXD X789XXS X79XXXA X79XXXD X79XXXS X80XXXA X80XXXD X80XXXS X810XXA
X810XXD X810XXS X811XXA X811XXD X811XXS X818XXA X818XXD X818XXS X820XXA
X820XXD X820XXS X821XXA X821XXD X821XXS X822XXA X822XXD X822XXS X828XXA
X828XXD X828XXS X830XXA X830XXD X830XXS X831XXA X831XXD X831XXS X832XXA
X832XXD X832XXS X838XXA X838XXD X838XXS Z915

5.14 Screening and history of mental health and substance use codes [663]

R780 R781 R782 R783 R784 R785 R786 Z046 Z134
Z72810 Z72811 Z8651 Z8659 Z87891 Z91410 Z91411 Z91412 Z91419
Z9149

5.15 Miscellaneous mental disorders [670]

F061	F068	F440	F441	F442	F444	F445	F446	F447
F4481	F4489	F449	F450	F451	F4520	F4521	F4522	F4529
F4541	F4542	F458	F459	F481	F5000	F5001	F5002	F502
F508	F509	F5101	F5102	F5103	F5104	F5105	F5109	F5111
F5112	F5113	F5119	F513	F514	F515	F518	F519	F520
F521	F5221	F5222	F5231	F5232	F524	F525	F526	F528
F529	F53	F54	F59	F641	F648	F649	F650	F651
F652	F653	F654	F6550	F6551	F6552	F6581	F6589	F659
F66	F6810	F6811	F6812	F6813	F688	F99	O906	R37
R4589	R630	R632	Z87890	Z9183				

TEDS substance use diagnosis codes

291.00–291.99: Alcoholic Psychoses

292.00–292.99: Drug Psychoses

303.00–303.89: Alcohol Intoxication

303.90–303.99: Alcohol Dependence

304.00–304.09: Opioid Dependence

304.20–304.29: Cocaine Dependence

304.30–304.39: Cannabis Dependence

304.10–304.19,

304.40–304.99,

305.10–305.19: Other Substance Dependence

305.00–305.09: Alcohol Use

305.60–305.69: Cocaine Use

305.20–305.29: Cannabis Use

305.50–305.59: Opioid Use

305.30–305.49,

305.70–305.99: Other Substance Use

APPENDIX F

TERMS AND ACRONYMS

Admission

Admission indicates the beginning of mental health service provision to a person with mental illness through programs under the auspices of the state mental health agency (SMHA). This includes new admissions (someone who has never received any services from the SMHA) and readmissions (someone who had previously received services from the SMHA, had been discharged, and started receiving services during the reporting period).

ADHD – Attention deficit/hyperactivity disorder

ADHD is a neurodevelopmental disorder defined by impairing levels of inattention, disorganization, and/or hyperactivity-impulsivity.

CBHSQ – Center for Behavioral Health Statistics and Quality

A center within the Substance Use and Mental Health Services Administration (SAMHSA) under the U.S. Department of Health and Human Services (HHS), CBHSQ is the lead government agency for behavioral health statistics, as designed by the Office of Management and Budget (OMB).

CD – Conduct disorder

CD is a serious behavioral and emotional disorder that can occur in children and teens. A child with this disorder may display a pattern of disruptive and violent behavior and have problems following rules.

DSM-5 – Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition

DSM-5 is the standard classification of mental disorders used by mental health professionals in the United States. DSM consists of three major components: the diagnostic classification, the diagnostic criteria sets, and the descriptive text [Source: <https://www.psychiatry.org/psychiatrists/practice/dsm>].

GED[®] – General Educational Development

GED is designed for people who, for various reasons, did not graduate from high school but want a certificate equivalent to the traditional high school diploma.

ICD-10-CM/ICD-9-CM – International Classification of Diseases, Tenth Revision, Clinical Modification/International Classification of Diseases, Ninth Revision, Clinical Modification

ICD-10-CM/ICD-9-CM is based on the World Health Organization's International Classification of Diseases (ICD-10/ICD-9). ICD-10-CM and ICD-9-CM are used as the official system of assigning codes to diagnoses and procedures associated with hospital utilization in the United States.

ODD – Oppositional defiant disorder

ODD is a childhood disorder that is characterized by negative, defiant, disobedient, and often hostile behavior toward individuals and authority figures primarily.

PDD – Pervasive developmental disorders

PDD refers to a group of conditions that involve delays in the development of many basic skills. Most notable among them are the ability to socialize with others, to communicate, and to use imagination.

Sources: Mental Health Client-Level Data (MH-CLD State Instruction Manual ([version 2.8](#)) and Combined Substance Use and Mental Health Treatment Episode Data Set (TEDS) State Instruction Manual With Data Submission System (DSS) Guide ([version 4.3.1](#))

APPENDIX G

LIST OF CONTRIBUTORS

This report was prepared for the Center for Behavioral Health Statistics and Quality (CBHSQ), Substance Abuse and Mental Health Services Administration, U.S. Department of Health and Human Services (HHS), by Eagle Technologies, Inc., under Contract No. HHSS283201600001C.

Contributors and reviewers at Eagle, listed alphabetically, include Anand Borse, Erin Doherty, Tsegereda Kifle, Thomas Nephew, David Peabody, Parth Thakore, Doren Walker (Project Director), Yang Yuan, and John Zuwasti Curran. Contributors and reviewers at NRI, listed alphabetically, include Azeb Berhane and Ted Lutterman. Production of the report at SAMHSA was managed by Nichele Waller (COR). Contributors and reviewers at SAMHSA, listed alphabetically, include Herman Alvarado, Heydy Juarez, Sharon Liu, and Nichele Waller.