

**Table 1.103C Any Use and Misuse of Prescription Pain Reliever Subtypes in Past Year: Among People Aged 12 or Older; Misuse of Prescription Pain Reliever Subtypes in Past Year: Among Past Year Any Users of Prescription Pain Reliever Subtypes Aged 12 or Older; Standard Errors of Numbers in Thousands, 2019 and 2020**

Pain Reliever Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse	Misuse	Misuse	Misuse
			in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Any Users of Pain Reliever Subtypes (2019)	in Past Year among Any Users of Pain Reliever Subtypes (2020)
<b>ANY PRESCRIPTION PAIN RELIEVER</b>	790	<i>1,281</i>	289	<i>474</i>	298	<i>483</i>
Hydrocodone Products	651	<i>1,033</i>	209	<i>336</i>	212	<i>340</i>
Zohydro <sup>®</sup> ER	76	<i>105</i>	13	<i>21</i>	*	*
Oxycodone Products	495	<i>780</i>	151	<i>283</i>	153	<i>283</i>
OxyContin <sup>®</sup>	300	<i>460</i>	89	<i>188</i>	89	<i>188</i>
Tramadol Products	434	<i>690</i>	110	<i>188</i>	110	<i>189</i>
Codeine Products	500	<i>727</i>	156	<i>267</i>	156	<i>268</i>
Morphine Products	246	<i>443</i>	59	<i>95</i>	60	<i>95</i>
Fentanyl Products	166	<i>275</i>	44	<i>84</i>	44	<i>85</i>
Buprenorphine Products	135	<i>237</i>	67	<i>105</i>	67	<i>105</i>
Oxymorphone Products	93	<i>144</i>	41	<i>45</i>	41	*
Demerol <sup>®</sup>	95	<i>96</i>	16	<i>18</i>	16	*
Hydromorphone Products	134	<i>172</i>	34	<i>63</i>	34	<i>63</i>
Methadone	101	<i>189</i>	50	<i>72</i>	50	*
Any Other Prescription Pain Reliever <sup>1</sup>	443	<i>705</i>	81	<i>162</i>	81	<i>163</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

<sup>1</sup> Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of “any other prescription pain reliever” that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 1.103D Any Use and Misuse of Prescription Pain Reliever Subtypes in Past Year: Among People Aged 12 or Older; Misuse of Prescription Pain Reliever Subtypes in Past Year: Among Past Year Any Users of Prescription Pain Reliever Subtypes Aged 12 or Older; Standard Errors of Percentages, 2019 and 2020**

Pain Reliever Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse	Misuse	Misuse	Misuse
			in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Any Users of Pain Reliever Subtypes (2019)	in Past Year among Any Users of Pain Reliever Subtypes (2020)
<b>ANY PRESCRIPTION PAIN RELIEVER</b>	0.29	<i>0.46</i>	0.11	<i>0.17</i>	0.34	<i>0.61</i>
Hydrocodone Products	0.24	<i>0.37</i>	0.08	<i>0.12</i>	0.47	<i>0.86</i>
Zohydro <sup>®</sup> ER	0.03	<i>0.04</i>	0.00	<i>0.01</i>	*	*
Oxycodone Products	0.18	<i>0.28</i>	0.05	<i>0.10</i>	0.61	<i>1.21</i>
OxyContin <sup>®</sup>	0.11	<i>0.17</i>	0.03	<i>0.07</i>	1.00	<i>2.39</i>
Tramadol Products	0.16	<i>0.25</i>	0.04	<i>0.07</i>	0.63	<i>1.08</i>
Codeine Products	0.18	<i>0.26</i>	0.06	<i>0.10</i>	0.62	<i>1.26</i>
Morphine Products	0.09	<i>0.16</i>	0.02	<i>0.03</i>	0.99	<i>1.71</i>
Fentanyl Products	0.06	<i>0.10</i>	0.02	<i>0.03</i>	1.73	<i>3.37</i>
Buprenorphine Products	0.05	<i>0.09</i>	0.02	<i>0.04</i>	2.47	<i>3.60</i>
Oxymorphone Products	0.03	<i>0.05</i>	0.01	<i>0.02</i>	4.52	*
Demerol <sup>®</sup>	0.03	<i>0.03</i>	0.01	<i>0.01</i>	1.77	*
Hydromorphone Products	0.05	<i>0.06</i>	0.01	<i>0.02</i>	2.06	<i>4.46</i>
Methadone	0.04	<i>0.07</i>	0.02	<i>0.03</i>	4.10	*
Any Other Prescription Pain Reliever <sup>1</sup>	0.16	<i>0.25</i>	0.03	<i>0.06</i>	0.35	<i>0.76</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

<sup>1</sup> Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of “any other prescription pain reliever” that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 1.104C Any Use and Misuse of Prescription Pain Reliever Subtypes in Past Year: Among People Aged 12 to 17; Misuse of Prescription Pain Reliever Subtypes in Past Year: Among Past Year Any Users of Prescription Pain Reliever Subtypes Aged 12 to 17; Standard Errors of Numbers in Thousands, 2019 and 2020**

Pain Reliever Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Total Population (2020)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)	Misuse in Past Year among Past Year Any Users of Pain Reliever Subtypes (2020)
<b>ANY PRESCRIPTION PAIN RELIEVER</b>	94	<i>177</i>	39	<i>61</i>	40	<i>62</i>
Hydrocodone Products	43	<i>89</i>	22	<i>25</i>	22	<i>25</i>
Zohydro <sup>®</sup> ER	9	<i>8</i>	6	*	*	*
Oxycodone Products	41	<i>63</i>	19	<i>20</i>	19	<i>20</i>
OxyContin <sup>®</sup>	27	<i>49</i>	14	<i>19</i>	14	*
Tramadol Products	29	<i>35</i>	13	<i>17</i>	13	*
Codeine Products	60	<i>121</i>	26	<i>25</i>	26	<i>25</i>
Morphine Products	29	<i>30</i>	11	<i>10</i>	11	*
Fentanyl Products	12	<i>10</i>	3	*	*	*
Buprenorphine Products	14	<i>17</i>	5	<i>5</i>	*	*
Oxymorphone Products	10	<i>12</i>	4	<i>7</i>	*	*
Demerol <sup>®</sup>	9	<i>4</i>	3	*	*	*
Hydromorphone Products	5	<i>16</i>	4	<i>5</i>	*	*
Methadone	11	<i>10</i>	7	*	*	*
Any Other Prescription Pain Reliever <sup>1</sup>	69	<i>107</i>	16	<i>50</i>	16	<i>51</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

<sup>1</sup> Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of “any other prescription pain reliever” that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 1.104D Any Use and Misuse of Prescription Pain Reliever Subtypes in Past Year: Among People Aged 12 to 17; Misuse of Prescription Pain Reliever Subtypes in Past Year: Among Past Year Any Users of Prescription Pain Reliever Subtypes Aged 12 to 17; Standard Errors of Percentages, 2019 and 2020**

Pain Reliever Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse	Misuse	Misuse	Misuse
			in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Past Year Any Users of Pain Reliever Subtypes (2019)	in Past Year among Past Year Any Users of Pain Reliever Subtypes (2020)
<b>ANY PRESCRIPTION PAIN RELIEVER</b>	0.38	<i>0.71</i>	0.16	<i>0.24</i>	0.91	<i>1.92</i>
Hydrocodone Products	0.17	<i>0.35</i>	0.09	<i>0.10</i>	2.50	<i>3.56</i>
Zohydro <sup>®</sup> ER	0.04	<i>0.03</i>	0.03	*	*	*
Oxycodone Products	0.16	<i>0.25</i>	0.08	<i>0.08</i>	2.43	<i>3.51</i>
OxyContin <sup>®</sup>	0.11	<i>0.20</i>	0.05	<i>0.07</i>	4.08	*
Tramadol Products	0.12	<i>0.14</i>	0.05	<i>0.07</i>	3.40	*
Codeine Products	0.24	<i>0.49</i>	0.10	<i>0.10</i>	1.75	<i>2.33</i>
Morphine Products	0.12	<i>0.12</i>	0.05	<i>0.04</i>	3.26	*
Fentanyl Products	0.05	<i>0.04</i>	0.01	*	*	*
Buprenorphine Products	0.05	<i>0.07</i>	0.02	<i>0.02</i>	*	*
Oxymorphone Products	0.04	<i>0.05</i>	0.02	<i>0.03</i>	*	*
Demerol <sup>®</sup>	0.03	<i>0.02</i>	0.01	*	*	*
Hydromorphone Products	0.02	<i>0.07</i>	0.02	<i>0.02</i>	*	*
Methadone	0.04	<i>0.04</i>	0.03	*	*	*
Any Other Prescription Pain Reliever <sup>1</sup>	0.28	<i>0.43</i>	0.06	<i>0.20</i>	0.82	<i>3.55</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

<sup>1</sup> Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of “any other prescription pain reliever” that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 1.105C Any Use and Misuse of Prescription Pain Reliever Subtypes in Past Year: Among People Aged 18 or Older; Misuse of Prescription Pain Reliever Subtypes in Past Year: Among Past Year Any Users of Prescription Pain Reliever Subtypes Aged 18 or Older; Standard Errors of Numbers in Thousands, 2019 and 2020**

Pain Reliever Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse	Misuse	Misuse	Misuse
			in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Any Users of Pain Reliever Subtypes (2019)	in Past Year among Any Users of Pain Reliever Subtypes (2020)
<b>ANY PRESCRIPTION PAIN RELIEVER</b>	784	<i>1,264</i>	285	<i>466</i>	294	<i>475</i>
Hydrocodone Products	648	<i>1,021</i>	208	<i>336</i>	211	<i>339</i>
Zohydro <sup>®</sup> ER	75	<i>104</i>	12	<i>21</i>	*	*
Oxycodone Products	493	<i>777</i>	150	<i>281</i>	152	<i>282</i>
OxyContin <sup>®</sup>	296	<i>454</i>	88	<i>187</i>	88	<i>187</i>
Tramadol Products	430	<i>687</i>	109	<i>187</i>	110	<i>188</i>
Codeine Products	498	<i>717</i>	152	<i>267</i>	153	<i>268</i>
Morphine Products	245	<i>442</i>	58	<i>95</i>	58	<i>95</i>
Fentanyl Products	166	<i>274</i>	44	<i>85</i>	44	<i>84</i>
Buprenorphine Products	134	<i>237</i>	67	<i>104</i>	67	<i>105</i>
Oxymorphone Products	93	<i>143</i>	41	<i>45</i>	41	*
Demerol <sup>®</sup>	95	<i>96</i>	15	<i>18</i>	15	*
Hydromorphone Products	134	<i>171</i>	34	<i>62</i>	34	*
Methadone	100	<i>188</i>	50	<i>72</i>	50	*
Any Other Prescription Pain Reliever <sup>1</sup>	439	<i>703</i>	79	<i>154</i>	79	<i>155</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

<sup>1</sup> Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of “any other prescription pain reliever” that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 1.105D Any Use and Misuse of Prescription Pain Reliever Subtypes in Past Year: Among People Aged 18 or Older; Misuse of Prescription Pain Reliever Subtypes in Past Year: Among Past Year Any Users of Prescription Pain Reliever Subtypes Aged 18 or Older; Standard Errors of Percentages, 2019 and 2020**

Pain Reliever Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse	Misuse	Misuse	Misuse
			in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Any Users of Pain Reliever Subtypes (2019)	in Past Year among Any Users of Pain Reliever Subtypes (2020)
<b>ANY PRESCRIPTION PAIN RELIEVER</b>	0.31	<i>0.50</i>	0.11	<i>0.18</i>	0.35	<i>0.62</i>
Hydrocodone Products	0.26	<i>0.41</i>	0.08	<i>0.13</i>	0.47	<i>0.88</i>
Zohydro <sup>®</sup> ER	0.03	<i>0.04</i>	0.00	<i>0.01</i>	*	*
Oxycodone Products	0.20	<i>0.31</i>	0.06	<i>0.11</i>	0.63	<i>1.23</i>
OxyContin <sup>®</sup>	0.12	<i>0.18</i>	0.04	<i>0.07</i>	1.03	<i>2.48</i>
Tramadol Products	0.17	<i>0.27</i>	0.04	<i>0.07</i>	0.63	<i>1.09</i>
Codeine Products	0.20	<i>0.28</i>	0.06	<i>0.11</i>	0.64	<i>1.33</i>
Morphine Products	0.10	<i>0.18</i>	0.02	<i>0.04</i>	1.02	<i>1.76</i>
Fentanyl Products	0.07	<i>0.11</i>	0.02	<i>0.03</i>	1.78	<i>3.42</i>
Buprenorphine Products	0.05	<i>0.09</i>	0.03	<i>0.04</i>	2.54	<i>3.67</i>
Oxymorphone Products	0.04	<i>0.06</i>	0.02	<i>0.02</i>	4.79	*
Demerol <sup>®</sup>	0.04	<i>0.04</i>	0.01	<i>0.01</i>	1.79	*
Hydromorphone Products	0.05	<i>0.07</i>	0.01	<i>0.02</i>	2.06	*
Methadone	0.04	<i>0.07</i>	0.02	<i>0.03</i>	4.25	*
Any Other Prescription Pain Reliever <sup>1</sup>	0.18	<i>0.28</i>	0.03	<i>0.06</i>	0.37	<i>0.77</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

<sup>1</sup> Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of “any other prescription pain reliever” that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 1.106C Any Use and Misuse of Prescription Pain Reliever Subtypes in Past Year: Among People Aged 18 to 25; Misuse of Prescription Pain Reliever Subtypes in Past Year: Among Past Year Any Users of Prescription Pain Reliever Subtypes Aged 18 to 25; Standard Errors of Numbers in Thousands, 2019 and 2020**

Pain Reliever Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse	Misuse	Misuse	Misuse
			in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Any Users of Pain Reliever Subtypes (2019)	in Past Year among Any Users of Pain Reliever Subtypes (2020)
<b>ANY PRESCRIPTION PAIN RELIEVER</b>	142	239	75	114	80	117
Hydrocodone Products	101	174	52	86	54	87
Zohydro <sup>®</sup> ER	12	11	2	*	*	*
Oxycodone Products	87	131	42	74	42	74
OxyContin <sup>®</sup>	53	80	29	40	29	40
Tramadol Products	61	125	27	46	27	46
Codeine Products	86	136	41	46	42	46
Morphine Products	50	61	15	18	15	*
Fentanyl Products	28	60	11	16	11	*
Buprenorphine Products	29	54	17	18	17	*
Oxymorphone Products	23	46	15	18	*	*
Demerol <sup>®</sup>	13	7	7	3	*	*
Hydromorphone Products	18	16	10	3	*	*
Methadone	15	44	9	5	*	*
Any Other Prescription Pain Reliever <sup>1</sup>	95	141	30	32	30	32

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

<sup>1</sup> Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of “any other prescription pain reliever” that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 1.106D Any Use and Misuse of Prescription Pain Reliever Subtypes in Past Year: Among People Aged 18 to 25; Misuse of Prescription Pain Reliever Subtypes in Past Year: Among Past Year Any Users of Prescription Pain Reliever Subtypes Aged 18 to 25; Standard Errors of Percentages, 2019 and 2020**

Pain Reliever Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse	Misuse	Misuse	Misuse
			in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Any Users of Pain Reliever Subtypes (2019)	in Past Year among Any Users of Pain Reliever Subtypes (2020)
<b>ANY PRESCRIPTION PAIN RELIEVER</b>	0.42	<i>0.71</i>	0.22	<i>0.34</i>	0.83	<i>1.56</i>
Hydrocodone Products	0.30	<i>0.52</i>	0.15	<i>0.26</i>	1.31	<i>2.45</i>
Zohydro <sup>®</sup> ER	0.04	<i>0.03</i>	0.01	*	*	*
Oxycodone Products	0.26	<i>0.39</i>	0.12	<i>0.22</i>	1.48	<i>3.16</i>
OxyContin <sup>®</sup>	0.16	<i>0.24</i>	0.09	<i>0.12</i>	2.69	<i>5.30</i>
Tramadol Products	0.18	<i>0.37</i>	0.08	<i>0.14</i>	1.99	<i>3.77</i>
Codeine Products	0.25	<i>0.41</i>	0.12	<i>0.14</i>	1.43	<i>2.13</i>
Morphine Products	0.15	<i>0.18</i>	0.05	<i>0.05</i>	1.98	*
Fentanyl Products	0.08	<i>0.18</i>	0.03	<i>0.05</i>	4.10	*
Buprenorphine Products	0.09	<i>0.16</i>	0.05	<i>0.05</i>	4.70	*
Oxymorphone Products	0.07	<i>0.14</i>	0.04	<i>0.05</i>	*	*
Demerol <sup>®</sup>	0.04	<i>0.02</i>	0.02	<i>0.01</i>	*	*
Hydromorphone Products	0.05	<i>0.05</i>	0.03	<i>0.01</i>	*	*
Methadone	0.04	<i>0.13</i>	0.03	<i>0.02</i>	*	*
Any Other Prescription Pain Reliever <sup>1</sup>	0.28	<i>0.42</i>	0.09	<i>0.10</i>	1.04	<i>1.50</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

<sup>1</sup> Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of “any other prescription pain reliever” that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.



**Table 1.107C Any Use and Misuse of Prescription Pain Reliever Subtypes in Past Year: Among People Aged 26 or Older; Misuse of Prescription Pain Reliever Subtypes in Past Year: Among Past Year Any Users of Prescription Pain Reliever Subtypes Aged 26 or Older; Standard Errors of Numbers in Thousands, 2019 and 2020**

Pain Reliever Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse	Misuse	Misuse	Misuse
			in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Any Users of Pain Reliever Subtypes (2019)	in Past Year among Any Users of Pain Reliever Subtypes (2020)
<b>ANY PRESCRIPTION PAIN RELIEVER</b>	770	<i>1,243</i>	276	<i>447</i>	285	<i>456</i>
Hydrocodone Products	636	<i>1,004</i>	203	<i>325</i>	207	<i>327</i>
Zohydro <sup>®</sup> ER	74	<i>103</i>	12	<i>21</i>	*	*
Oxycodone Products	481	<i>765</i>	144	<i>265</i>	146	<i>266</i>
OxyContin <sup>®</sup>	292	<i>445</i>	83	<i>179</i>	83	<i>179</i>
Tramadol Products	423	<i>684</i>	106	<i>179</i>	106	<i>180</i>
Codeine Products	491	<i>707</i>	147	<i>261</i>	147	<i>262</i>
Morphine Products	238	<i>434</i>	57	<i>93</i>	57	<i>93</i>
Fentanyl Products	160	<i>265</i>	42	<i>83</i>	42	<i>83</i>
Buprenorphine Products	131	<i>230</i>	63	<i>103</i>	63	<i>103</i>
Oxymorphone Products	91	<i>135</i>	39	<i>41</i>	*	*
Demerol <sup>®</sup>	95	<i>96</i>	14	<i>18</i>	14	*
Hydromorphone Products	131	<i>171</i>	33	<i>62</i>	33	*
Methadone	99	<i>183</i>	49	<i>72</i>	49	*
Any Other Prescription Pain Reliever <sup>1</sup>	430	<i>689</i>	71	<i>151</i>	71	<i>151</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

<sup>1</sup> Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of “any other prescription pain reliever” that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 1.107D Any Use and Misuse of Prescription Pain Reliever Subtypes in Past Year: Among People Aged 26 or Older; Misuse of Prescription Pain Reliever Subtypes in Past Year: Among Past Year Any Users of Prescription Pain Reliever Subtypes Aged 26 or Older; Standard Errors of Percentages, 2019 and 2020**

Pain Reliever Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse	Misuse	Misuse	Misuse
			in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Any Users of Pain Reliever Subtypes (2019)	in Past Year among Any Users of Pain Reliever Subtypes (2020)
<b>ANY PRESCRIPTION PAIN RELIEVER</b>	0.36	<i>0.57</i>	0.13	<i>0.20</i>	0.39	<i>0.66</i>
Hydrocodone Products	0.29	<i>0.46</i>	0.09	<i>0.15</i>	0.51	<i>0.92</i>
Zohydro <sup>®</sup> ER	0.03	<i>0.05</i>	0.01	<i>0.01</i>	*	*
Oxycodone Products	0.22	<i>0.35</i>	0.07	<i>0.12</i>	0.67	<i>1.30</i>
OxyContin <sup>®</sup>	0.13	<i>0.20</i>	0.04	<i>0.08</i>	1.09	<i>2.64</i>
Tramadol Products	0.20	<i>0.31</i>	0.05	<i>0.08</i>	0.67	<i>1.12</i>
Codeine Products	0.23	<i>0.32</i>	0.07	<i>0.12</i>	0.70	<i>1.46</i>
Morphine Products	0.11	<i>0.20</i>	0.03	<i>0.04</i>	1.16	<i>1.88</i>
Fentanyl Products	0.07	<i>0.12</i>	0.02	<i>0.04</i>	1.88	<i>3.85</i>
Buprenorphine Products	0.06	<i>0.11</i>	0.03	<i>0.05</i>	2.74	<i>4.02</i>
Oxymorphone Products	0.04	<i>0.06</i>	0.02	<i>0.02</i>	*	*
Demerol <sup>®</sup>	0.04	<i>0.04</i>	0.01	<i>0.01</i>	1.71	*
Hydromorphone Products	0.06	<i>0.08</i>	0.02	<i>0.03</i>	2.11	*
Methadone	0.05	<i>0.08</i>	0.02	<i>0.03</i>	4.55	*
Any Other Prescription Pain Reliever <sup>1</sup>	0.20	<i>0.32</i>	0.03	<i>0.07</i>	0.38	<i>0.84</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

<sup>1</sup> Includes use or misuse of pain relievers containing other active ingredients. Reports of misuse of “any other prescription pain reliever” that correspond only to the specific pain reliever categories shown in the table are excluded from estimates for Any Other Prescription Pain Reliever and are included instead in the relevant pain reliever category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.