

Table 1.113A Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 12 or Older; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 12 or Older; Numbers in Thousands, 2019 and 2020

Tranquilizer Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Total Population (2020)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	35,601	<i>35,018</i>	5,220	<i>5,460</i>	5,220	<i>5,460</i>
Benzodiazepine Tranquilizers	28,316	<i>26,095</i>	4,819	<i>4,739</i>	4,819	<i>4,739</i>
Alprazolam Products	16,748	<i>14,507</i>	3,668	<i>3,419</i>	3,668	<i>3,419</i>
Lorazepam Products	7,787	<i>7,189</i>	795	<i>723</i>	795	<i>723</i>
Clonazepam Products	5,998	<i>6,357</i>	1,029	<i>1,234</i>	1,029	<i>1,234</i>
Diazepam Products	5,672	<i>5,706</i>	758	<i>971</i>	758	<i>971</i>
Muscle Relaxants	6,049	<i>8,127</i>	552	<i>845</i>	552	<i>845</i>
Cyclobenzaprine	4,955	<i>7,289</i>	361	<i>738</i>	361	<i>738</i>
Soma [®]	1,182	<i>1,215</i>	223	<i>206</i>	223	<i>*</i>
Any Other Prescription Tranquilizer ¹	5,738	<i>5,428</i>	97	<i>228</i>	97	<i>228</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.113B Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 12 or Older; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 12 or Older; Percentages, 2019 and 2020

Tranquilizer Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Total Population (2020)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	12.9	<i>12.6</i>	1.9	<i>2.0</i>	14.7	<i>15.6</i>
Benzodiazepine Tranquilizers	10.3	<i>9.4</i>	1.8	<i>1.7</i>	17.0	<i>18.2</i>
Alprazolam Products	6.1	<i>5.2</i>	1.3	<i>1.2</i>	21.9	<i>23.6</i>
Lorazepam Products	2.8	<i>2.6</i>	0.3	<i>0.3</i>	10.2	<i>10.1</i>
Clonazepam Products	2.2	<i>2.3</i>	0.4	<i>0.4</i>	17.2	<i>19.4</i>
Diazepam Products	2.1	<i>2.1</i>	0.3	<i>0.4</i>	13.4	<i>17.0</i>
Muscle Relaxants	2.2	<i>2.9</i>	0.2	<i>0.3</i>	9.1	<i>10.4</i>
Cyclobenzaprine	1.8	<i>2.6</i>	0.1	<i>0.3</i>	7.3	<i>10.1</i>
Soma [®]	0.4	<i>0.4</i>	0.1	<i>0.1</i>	18.8	*
Any Other Prescription Tranquilizer ¹	2.1	<i>2.0</i>	0.0	<i>0.1</i>	1.7	<i>4.2</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.114A Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 12 to 17; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 12 to 17; Numbers in Thousands, 2019 and 2020

Tranquilizer Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Total Population (2020)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	1,010	588	392	<i>196</i>	392	<i>196</i>
Benzodiazepine Tranquilizers	700	398	371	<i>152</i>	371	*
Alprazolam Products	613	308	359	<i>151</i>	359	*
Lorazepam Products	84	48	9	9	*	*
Clonazepam Products	83	41	33	2	*	*
Diazepam Products	80	44	16	7	*	*
Muscle Relaxants	45	27	15	7	*	*
Cyclobenzaprine	32	27	4	7	*	*
Soma®	11	*	10	*	*	*
Any Other Prescription Tranquilizer ¹	367	<i>184</i>	16	23	16	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.114B Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 12 to 17; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 12 to 17; Percentages, 2019 and 2020

Tranquilizer Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Total Population (2020)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	4.1	<i>2.4</i>	1.6	<i>0.8</i>	38.8	33.3
Benzodiazepine Tranquilizers	2.8	<i>1.6</i>	1.5	<i>0.6</i>	52.9	*
Alprazolam Products	2.5	<i>1.2</i>	1.4	<i>0.6</i>	58.6	*
Lorazepam Products	0.3	<i>0.2</i>	0.0	<i>0.0</i>	*	*
Clonazepam Products	0.3	<i>0.2</i>	0.1	<i>0.0</i>	*	*
Diazepam Products	0.3	<i>0.2</i>	0.1	<i>0.0</i>	*	*
Muscle Relaxants	0.2	<i>0.1</i>	0.1	<i>0.0</i>	*	*
Cyclobenzaprine	0.1	<i>0.1</i>	0.0	<i>0.0</i>	*	*
Soma [®]	0.0	*	0.0	*	*	*
Any Other Prescription Tranquilizer ¹	1.5	<i>0.7</i>	0.1	<i>0.1</i>	4.3	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.115A Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 18 or Older; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 18 or Older; Numbers in Thousands, 2019 and 2020

Tranquilizer Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Total Population (2020)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	34,591	<i>34,430</i>	4,829	<i>5,264</i>	4,829	<i>5,264</i>
Benzodiazepine Tranquilizers	27,615	<i>25,697</i>	4,448	<i>4,587</i>	4,448	<i>4,587</i>
Alprazolam Products	16,134	<i>14,199</i>	3,308	<i>3,269</i>	3,308	<i>3,269</i>
Lorazepam Products	7,703	<i>7,140</i>	786	<i>714</i>	786	<i>714</i>
Clonazepam Products	5,916	<i>6,316</i>	996	<i>1,232</i>	996	<i>1,232</i>
Diazepam Products	5,592	<i>5,662</i>	742	<i>965</i>	742	<i>965</i>
Muscle Relaxants	6,005	<i>8,099</i>	537	<i>838</i>	537	<i>838</i>
Cyclobenzaprine	4,923	<i>7,262</i>	358	<i>731</i>	358	<i>731</i>
Soma®	1,170	<i>1,214</i>	213	<i>206</i>	213	*
Any Other Prescription Tranquilizer ¹	5,371	<i>5,244</i>	81	<i>205</i>	81	<i>205</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.115B Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 18 or Older; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 18 or Older; Percentages, 2019 and 2020

Tranquilizer Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Total Population (2020)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	13.8	<i>13.7</i>	1.9	<i>2.1</i>	14.0	<i>15.3</i>
Benzodiazepine Tranquilizers	11.0	<i>10.2</i>	1.8	<i>1.8</i>	16.1	<i>17.9</i>
Alprazolam Products	6.4	<i>5.6</i>	1.3	<i>1.3</i>	20.5	<i>23.0</i>
Lorazepam Products	3.1	<i>2.8</i>	0.3	<i>0.3</i>	10.2	<i>10.0</i>
Clonazepam Products	2.4	<i>2.5</i>	0.4	<i>0.5</i>	16.8	<i>19.5</i>
Diazepam Products	2.2	<i>2.2</i>	0.3	<i>0.4</i>	13.3	<i>17.0</i>
Muscle Relaxants	2.4	<i>3.2</i>	0.2	<i>0.3</i>	8.9	<i>10.3</i>
Cyclobenzaprine	2.0	<i>2.9</i>	0.1	<i>0.3</i>	7.3	<i>10.1</i>
Soma [®]	0.5	<i>0.5</i>	0.1	<i>0.1</i>	18.2	*
Any Other Prescription Tranquilizer ¹	2.1	<i>2.1</i>	0.0	<i>0.1</i>	1.5	<i>3.9</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.116A Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 18 to 25; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 18 to 25; Numbers in Thousands, 2019 and 2020

Tranquilizer Subtype	Any Use		Misuse		Misuse	
	in Past Year (2019)	in Past Year (2020)	in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	3,419	3,338	1,313	<i>1,108</i>	1,313	<i>1,108</i>
Benzodiazepine Tranquilizers	2,922	2,822	1,260	<i>1,087</i>	1,260	<i>1,087</i>
Alprazolam Products	2,229	<i>1,992</i>	1,091	<i>932</i>	1,091	<i>932</i>
Lorazepam Products	630	<i>701</i>	132	<i>169</i>	132	<i>169</i>
Clonazepam Products	599	<i>641</i>	214	<i>216</i>	214	*
Diazepam Products	451	<i>561</i>	138	<i>117</i>	138	*
Muscle Relaxants	314	<i>396</i>	59	<i>64</i>	59	*
Cyclobenzaprine	258	<i>351</i>	35	<i>41</i>	35	*
Soma [®]	70	<i>73</i>	33	<i>42</i>	*	*
Any Other Prescription Tranquilizer ¹	515	<i>466</i>	33	<i>3</i>	33	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.116B Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 18 to 25; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 18 to 25; Percentages, 2019 and 2020

Tranquilizer Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Total Population (2020)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	10.1	<i>10.0</i>	3.9	3.3	38.4	33.2
Benzodiazepine Tranquilizers	8.7	<i>8.4</i>	3.7	3.2	43.1	38.5
Alprazolam Products	6.6	<i>5.9</i>	3.2	2.8	49.0	46.8
Lorazepam Products	1.9	<i>2.1</i>	0.4	<i>0.5</i>	21.0	<i>24.1</i>
Clonazepam Products	1.8	<i>1.9</i>	0.6	<i>0.6</i>	35.7	*
Diazepam Products	1.3	<i>1.7</i>	0.4	<i>0.4</i>	30.6	*
Muscle Relaxants	0.9	<i>1.2</i>	0.2	<i>0.2</i>	18.9	*
Cyclobenzaprine	0.8	<i>1.0</i>	0.1	<i>0.1</i>	13.5	*
Soma [®]	0.2	<i>0.2</i>	0.1	<i>0.1</i>	*	*
Any Other Prescription Tranquilizer ¹	1.5	<i>1.4</i>	0.1	<i>0.0</i>	6.4	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.117A Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 26 or Older; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 26 or Older; Numbers in Thousands, 2019 and 2020

Tranquilizer Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Total Population (2020)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	31,172	<i>31,092</i>	3,516	<i>4,156</i>	3,516	<i>4,156</i>
Benzodiazepine Tranquilizers	24,693	<i>22,875</i>	3,188	<i>3,500</i>	3,188	<i>3,500</i>
Alprazolam Products	13,906	<i>12,207</i>	2,217	<i>2,336</i>	2,217	<i>2,336</i>
Lorazepam Products	7,074	<i>6,439</i>	654	<i>545</i>	654	<i>545</i>
Clonazepam Products	5,317	<i>5,675</i>	782	<i>1,016</i>	782	<i>1,016</i>
Diazepam Products	5,141	<i>5,101</i>	604	<i>847</i>	604	<i>847</i>
Muscle Relaxants	5,690	<i>7,703</i>	477	<i>774</i>	477	<i>774</i>
Cyclobenzaprine	4,665	<i>6,911</i>	323	<i>690</i>	323	<i>690</i>
Soma®	1,100	<i>1,141</i>	179	<i>164</i>	179	*
Any Other Prescription Tranquilizer ¹	4,856	<i>4,778</i>	48	<i>202</i>	48	<i>202</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.117B Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 26 or Older; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 26 or Older; Percentages, 2019 and 2020

Tranquilizer Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Total Population (2020)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	14.4	<i>14.2</i>	1.6	<i>1.9</i>	11.3	<i>13.4</i>
Benzodiazepine Tranquilizers	11.4	<i>10.5</i>	1.5	<i>1.6</i>	12.9	<i>15.3</i>
Alprazolam Products	6.4	<i>5.6</i>	1.0	<i>1.1</i>	15.9	<i>19.1</i>
Lorazepam Products	3.3	<i>2.9</i>	0.3	<i>0.2</i>	9.2	<i>8.5</i>
Clonazepam Products	2.5	<i>2.6</i>	0.4	<i>0.5</i>	14.7	<i>17.9</i>
Diazepam Products	2.4	<i>2.3</i>	0.3	<i>0.4</i>	11.7	<i>16.6</i>
Muscle Relaxants	2.6	<i>3.5</i>	0.2	<i>0.4</i>	8.4	<i>10.0</i>
Cyclobenzaprine	2.2	<i>3.2</i>	0.1	<i>0.3</i>	6.9	<i>10.0</i>
Soma [®]	0.5	<i>0.5</i>	0.1	<i>0.1</i>	16.3	*
Any Other Prescription Tranquilizer ¹	2.2	<i>2.2</i>	0.0	<i>0.1</i>	1.0	<i>4.2</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.