

Table 1.113C Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 12 or Older; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 12 or Older; Standard Errors of Numbers in Thousands, 2019 and 2020

Tranquilizer Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Total Population (2020)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	589	871	224	309	228	313
Benzodiazepine Tranquilizers	527	787	212	282	216	287
Alprazolam Products	398	579	157	234	159	237
Lorazepam Products	307	436	102	106	102	106
Clonazepam Products	251	382	95	162	96	163
Diazepam Products	257	392	88	146	88	145
Muscle Relaxants	254	449	75	134	75	135
Cyclobenzaprine	220	424	59	129	59	129
Soma®	121	199	46	71	46	*
Any Other Prescription Tranquilizer ¹	243	374	22	107	22	107

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.113D Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 12 or Older; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 12 or Older; Standard Errors of Percentages, 2019 and 2020

Tranquilizer Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Total Population (2020)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	0.21	<i>0.31</i>	0.08	<i>0.11</i>	0.58	<i>0.86</i>
Benzodiazepine Tranquilizers	0.19	<i>0.28</i>	0.08	<i>0.10</i>	0.70	<i>1.04</i>
Alprazolam Products	0.14	<i>0.21</i>	0.06	<i>0.08</i>	0.90	<i>1.51</i>
Lorazepam Products	0.11	<i>0.16</i>	0.04	<i>0.04</i>	1.23	<i>1.46</i>
Clonazepam Products	0.09	<i>0.14</i>	0.03	<i>0.06</i>	1.46	<i>2.34</i>
Diazepam Products	0.09	<i>0.14</i>	0.03	<i>0.05</i>	1.44	<i>2.38</i>
Muscle Relaxants	0.09	<i>0.16</i>	0.03	<i>0.05</i>	1.13	<i>1.60</i>
Cyclobenzaprine	0.08	<i>0.15</i>	0.02	<i>0.05</i>	1.11	<i>1.70</i>
Soma [®]	0.04	<i>0.07</i>	0.02	<i>0.03</i>	3.60	*
Any Other Prescription Tranquilizer ¹	0.09	<i>0.13</i>	0.01	<i>0.04</i>	0.38	<i>1.92</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.114C Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 12 to 17; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 12 to 17; Standard Errors of Numbers in Thousands, 2019 and 2020

Tranquilizer Subtype	Any Use in Past Year		Misuse in Past Year among Total Population		Misuse in Past Year among Any Users of Tranquilizer Subtypes	
	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)
ANY PRESCRIPTION TRANQUILIZER	51	<i>61</i>	35	38	35	38
Benzodiazepine Tranquilizers	44	<i>54</i>	34	36	34	*
Alprazolam Products	41	<i>49</i>	33	36	33	*
Lorazepam Products	16	<i>16</i>	4	5	*	*
Clonazepam Products	16	<i>15</i>	10	2	*	*
Diazepam Products	16	<i>16</i>	7	5	*	*
Muscle Relaxants	13	<i>10</i>	8	5	*	*
Cyclobenzaprine	10	<i>10</i>	3	5	*	*
Soma®	7	*	7	*	*	*
Any Other Prescription Tranquilizer ¹	32	<i>31</i>	5	16	5	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.114D Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 12 to 17; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 12 to 17; Standard Errors of Percentages, 2019 and 2020

Tranquilizer Subtype	Any Use		Misuse		Misuse	
	in Past Year (2019)	in Past Year (2020)	in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	0.21	<i>0.24</i>	0.14	<i>0.15</i>	2.70	<i>5.09</i>
Benzodiazepine Tranquilizers	0.17	<i>0.22</i>	0.13	<i>0.14</i>	3.23	*
Alprazolam Products	0.16	<i>0.20</i>	0.13	<i>0.14</i>	3.32	*
Lorazepam Products	0.06	<i>0.06</i>	0.02	<i>0.02</i>	*	*
Clonazepam Products	0.06	<i>0.06</i>	0.04	<i>0.01</i>	*	*
Diazepam Products	0.06	<i>0.06</i>	0.03	<i>0.02</i>	*	*
Muscle Relaxants	0.05	<i>0.04</i>	0.03	<i>0.02</i>	*	*
Cyclobenzaprine	0.04	<i>0.04</i>	0.01	<i>0.02</i>	*	*
Soma®	0.03	*	0.03	*	*	*
Any Other Prescription Tranquilizer ¹	0.13	<i>0.13</i>	0.02	<i>0.06</i>	1.41	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.115C Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 18 or Older; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 18 or Older; Standard Errors of Numbers in Thousands, 2019 and 2020

Tranquilizer Subtype	Any Use		Misuse		Misuse	
	in Past Year (2019)	in Past Year (2020)	in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	581	866	219	308	223	311
Benzodiazepine Tranquilizers	520	781	207	282	212	285
Alprazolam Products	392	573	153	233	155	235
Lorazepam Products	307	435	102	106	102	106
Clonazepam Products	250	381	95	162	96	163
Diazepam Products	255	390	87	146	87	145
Muscle Relaxants	252	449	74	134	74	135
Cyclobenzaprine	220	424	59	129	59	129
Soma [®]	119	199	46	71	46	*
Any Other Prescription Tranquilizer ¹	244	372	21	106	21	106

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.115D Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 18 or Older; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 18 or Older; Standard Errors of Percentages, 2019 and 2020

Tranquilizer Subtype	Any Use		Misuse		Misuse	
	in Past Year (2019)	in Past Year (2020)	in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	0.23	<i>0.34</i>	0.09	<i>0.12</i>	0.59	<i>0.86</i>
Benzodiazepine Tranquilizers	0.21	<i>0.31</i>	0.08	<i>0.11</i>	0.70	<i>1.05</i>
Alprazolam Products	0.16	<i>0.23</i>	0.06	<i>0.09</i>	0.91	<i>1.54</i>
Lorazepam Products	0.12	<i>0.17</i>	0.04	<i>0.04</i>	1.25	<i>1.47</i>
Clonazepam Products	0.10	<i>0.15</i>	0.04	<i>0.06</i>	1.47	<i>2.35</i>
Diazepam Products	0.10	<i>0.15</i>	0.03	<i>0.06</i>	1.45	<i>2.39</i>
Muscle Relaxants	0.10	<i>0.18</i>	0.03	<i>0.05</i>	1.14	<i>1.61</i>
Cyclobenzaprine	0.09	<i>0.17</i>	0.02	<i>0.05</i>	1.11	<i>1.71</i>
Soma®	0.05	<i>0.08</i>	0.02	<i>0.03</i>	3.68	*
Any Other Prescription Tranquilizer ¹	0.10	<i>0.15</i>	0.01	<i>0.04</i>	0.40	<i>1.97</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.116C Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 18 to 25; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 18 to 25; Standard Errors of Numbers in Thousands, 2019 and 2020

Tranquilizer Subtype	Any Use		Misuse		Misuse	
	in Past Year (2019)	in Past Year (2020)	in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	103	<i>182</i>	65	<i>104</i>	69	<i>106</i>
Benzodiazepine Tranquilizers	95	<i>173</i>	64	<i>104</i>	68	<i>105</i>
Alprazolam Products	84	<i>145</i>	58	<i>96</i>	61	<i>97</i>
Lorazepam Products	44	<i>91</i>	20	<i>36</i>	20	<i>36</i>
Clonazepam Products	43	<i>89</i>	25	<i>46</i>	26	*
Diazepam Products	37	<i>83</i>	21	<i>34</i>	21	*
Muscle Relaxants	31	<i>62</i>	13	<i>27</i>	13	*
Cyclobenzaprine	29	<i>57</i>	10	<i>17</i>	10	*
Soma [®]	15	<i>30</i>	10	<i>26</i>	*	*
Any Other Prescription Tranquilizer ¹	44	<i>70</i>	12	<i>3</i>	12	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.116D Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 18 to 25; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 18 to 25; Standard Errors of Percentages, 2019 and 2020

Tranquilizer Subtype	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Total Population (2020)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	Misuse in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	0.30	<i>0.54</i>	0.19	<i>0.31</i>	1.55	<i>2.60</i>
Benzodiazepine Tranquilizers	0.28	<i>0.52</i>	0.19	<i>0.31</i>	1.71	<i>2.99</i>
Alprazolam Products	0.25	<i>0.43</i>	0.17	<i>0.29</i>	1.91	<i>3.57</i>
Lorazepam Products	0.13	<i>0.27</i>	0.06	<i>0.11</i>	3.04	<i>4.77</i>
Clonazepam Products	0.13	<i>0.27</i>	0.08	<i>0.14</i>	3.47	*
Diazepam Products	0.11	<i>0.25</i>	0.06	<i>0.10</i>	3.77	*
Muscle Relaxants	0.09	<i>0.19</i>	0.04	<i>0.08</i>	3.71	*
Cyclobenzaprine	0.09	<i>0.17</i>	0.03	<i>0.05</i>	3.73	*
Soma [®]	0.04	<i>0.09</i>	0.03	<i>0.08</i>	*	*
Any Other Prescription Tranquilizer ¹	0.13	<i>0.21</i>	0.03	<i>0.01</i>	2.18	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.117C Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 26 or Older; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 26 or Older; Standard Errors of Numbers in Thousands, 2019 and 2020

Tranquilizer Subtype	Any Use		Misuse		Misuse	
	in Past Year (2019)	in Past Year (2020)	in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	568	<i>846</i>	205	<i>291</i>	208	<i>294</i>
Benzodiazepine Tranquilizers	508	<i>754</i>	194	<i>265</i>	197	<i>268</i>
Alprazolam Products	388	<i>554</i>	138	<i>214</i>	140	<i>215</i>
Lorazepam Products	301	<i>419</i>	98	<i>100</i>	99	<i>100</i>
Clonazepam Products	243	<i>371</i>	91	<i>156</i>	92	<i>157</i>
Diazepam Products	249	<i>378</i>	84	<i>144</i>	84	<i>143</i>
Muscle Relaxants	252	<i>441</i>	73	<i>131</i>	73	<i>131</i>
Cyclobenzaprine	220	<i>418</i>	58	<i>127</i>	58	<i>127</i>
Soma®	118	<i>196</i>	45	<i>66</i>	45	*
Any Other Prescription Tranquilizer ¹	238	<i>366</i>	18	<i>106</i>	18	<i>106</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.117D Any Use and Misuse of Prescription Tranquilizer Subtypes in Past Year: Among People Aged 26 or Older; Misuse of Prescription Tranquilizer Subtypes in Past Year: Among Past Year Any Users of Prescription Tranquilizer Subtypes Aged 26 or Older; Standard Errors of Percentages, 2019 and 2020

Tranquilizer Subtype	Any Use		Misuse		Misuse	
	in Past Year (2019)	in Past Year (2020)	in Past Year among Total Population (2019)	in Past Year among Total Population (2020)	in Past Year among Past Year Any Users of Tranquilizer Subtypes (2019)	in Past Year among Past Year Any Users of Tranquilizer Subtypes (2020)
ANY PRESCRIPTION TRANQUILIZER	0.26	<i>0.39</i>	0.09	<i>0.13</i>	0.61	<i>0.91</i>
Benzodiazepine Tranquilizers	0.23	<i>0.35</i>	0.09	<i>0.12</i>	0.73	<i>1.11</i>
Alprazolam Products	0.18	<i>0.25</i>	0.06	<i>0.10</i>	0.95	<i>1.66</i>
Lorazepam Products	0.14	<i>0.19</i>	0.05	<i>0.05</i>	1.31	<i>1.55</i>
Clonazepam Products	0.11	<i>0.17</i>	0.04	<i>0.07</i>	1.59	<i>2.52</i>
Diazepam Products	0.11	<i>0.17</i>	0.04	<i>0.07</i>	1.54	<i>2.61</i>
Muscle Relaxants	0.12	<i>0.20</i>	0.03	<i>0.06</i>	1.19	<i>1.65</i>
Cyclobenzaprine	0.10	<i>0.19</i>	0.03	<i>0.06</i>	1.16	<i>1.77</i>
Soma [®]	0.05	<i>0.09</i>	0.02	<i>0.03</i>	3.81	*
Any Other Prescription Tranquilizer ¹	0.11	<i>0.17</i>	0.01	<i>0.05</i>	0.37	<i>2.15</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Includes use or misuse of tranquilizers containing other active ingredients. Reports of misuse of “any other prescription tranquilizer” that correspond only to the specific tranquilizer categories shown in the table are excluded from estimates for Any Other Prescription Tranquilizer and are included instead in the relevant tranquilizer category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.