

Table 1.6A Illicit Drug Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	138,027	<i>138,543</i>	57,203	<i>59,277</i>	35,803	<i>37,309</i>
12-20	12,323	<i>10,691</i>	9,116	<i>7,486</i>	5,137	<i>4,208</i>
21 or Older	125,704	<i>127,852</i>	48,087	<i>51,791</i>	30,666	<i>33,101</i>
12-17	6,090	<i>5,229</i>	4,279	<i>3,436</i>	2,167	<i>1,679</i>
12-13	1,041	<i>941</i>	582	<i>442</i>	214	<i>76</i>
14-15	1,970	<i>1,780</i>	1,313	<i>1,209</i>	622	<i>599</i>
16-17	3,080	<i>2,507</i>	2,384	<i>1,785</i>	1,331	<i>1,004</i>
18 or Older	131,937	<i>133,314</i>	52,924	<i>55,841</i>	33,636	<i>35,630</i>
18-25	18,832	<i>17,879</i>	13,193	<i>12,398</i>	8,406	<i>8,016</i>
18-20	6,233	<i>5,463</i>	4,837	<i>4,051</i>	2,970	<i>2,529</i>
21-25	12,599	<i>12,416</i>	8,356	<i>8,347</i>	5,436	<i>5,487</i>
26 or Older	113,104	<i>115,435</i>	39,731	<i>43,443</i>	25,231	<i>27,614</i>
26-49	59,083	<i>58,674</i>	26,092	<i>27,784</i>	16,479	<i>18,165</i>
26-29	11,523	<i>10,746</i>	6,690	<i>6,234</i>	4,419	<i>4,177</i>
30-34	13,677	<i>13,638</i>	6,520	<i>6,967</i>	4,056	<i>4,508</i>
35-39	12,847	<i>13,127</i>	5,338	<i>6,053</i>	3,337	<i>3,914</i>
40-44	10,694	<i>10,448</i>	3,887	<i>4,323</i>	2,337	<i>2,926</i>
45-49	10,342	<i>10,714</i>	3,657	<i>4,207</i>	2,330	<i>2,641</i>
50 or Older	54,021	<i>56,762</i>	13,639	<i>15,660</i>	8,752	<i>9,449</i>
50-54	11,391	<i>11,192</i>	3,108	<i>3,624</i>	2,024	<i>2,302</i>
55-59	12,209	<i>12,248</i>	3,498	<i>3,797</i>	2,203	<i>2,187</i>
60-64	12,176	<i>12,056</i>	3,303	<i>3,955</i>	2,281	<i>2,419</i>
65 or Older	18,245	<i>21,266</i>	3,730	<i>4,284</i>	2,243	<i>2,541</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Illicit Drug Use in Lifetime includes the misuse of prescription psychotherapeutics even though those estimates are not reported due to potential underreporting in lifetime prescription psychotherapeutics estimates.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.6B Illicit Drug Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	50.2	<i>50.0</i>	20.8	<i>21.4</i>	13.0	<i>13.5</i>
12-20	32.4	<i>28.6</i>	23.9	<i>20.0</i>	13.5	<i>11.2</i>
21 or Older	53.0	<i>53.4</i>	20.3	<i>21.6</i>	12.9	<i>13.8</i>
12-17	24.5	<i>20.9</i>	17.2	<i>13.8</i>	8.7	<i>6.7</i>
12-13	12.7	<i>11.1</i>	7.1	<i>5.2</i>	2.6	<i>0.9</i>
14-15	23.2	<i>20.6</i>	15.5	<i>14.0</i>	7.3	<i>6.9</i>
16-17	37.3	<i>31.8</i>	28.9	<i>22.6</i>	16.1	<i>12.7</i>
18 or Older	52.7	<i>52.9</i>	21.1	<i>22.2</i>	13.4	<i>14.1</i>
18-25	55.8	<i>53.4</i>	39.1	<i>37.0</i>	24.9	<i>23.9</i>
18-20	47.3	<i>43.9</i>	36.7	<i>32.6</i>	22.6	<i>20.3</i>
21-25	61.3	<i>59.0</i>	40.6	<i>39.6</i>	26.4	<i>26.1</i>
26 or Older	52.2	<i>52.8</i>	18.3	<i>19.9</i>	11.6	<i>12.6</i>
26-49	58.3	<i>57.8</i>	25.8	<i>27.4</i>	16.3	<i>17.9</i>
26-29	63.8	<i>60.3</i>	37.1	<i>35.0</i>	24.5	<i>23.5</i>
30-34	61.4	<i>60.0</i>	29.3	<i>30.7</i>	18.2	<i>19.8</i>
35-39	60.0	<i>58.8</i>	24.9	<i>27.1</i>	15.6	<i>17.5</i>
40-44	55.5	<i>53.9</i>	20.2	<i>22.3</i>	12.1	<i>15.1</i>
45-49	51.1	<i>55.5</i>	18.1	<i>21.8</i>	11.5	<i>13.7</i>
50 or Older	46.9	<i>48.6</i>	11.8	<i>13.4</i>	7.6	<i>8.1</i>
50-54	55.3	<i>56.2</i>	15.1	<i>18.2</i>	9.8	<i>11.6</i>
55-59	57.4	<i>56.9</i>	16.4	<i>17.6</i>	10.4	<i>10.2</i>
60-64	59.2	<i>57.7</i>	16.1	<i>18.9</i>	11.1	<i>11.6</i>
65 or Older	34.5	<i>39.0</i>	7.1	<i>7.9</i>	4.2	<i>4.7</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Illicit Drug Use in Lifetime includes the misuse of prescription psychotherapeutics even though those estimates are not reported due to potential underreporting in lifetime prescription psychotherapeutics estimates.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.7A Marijuana Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	127,139	<i>126,504</i>	48,242	<i>49,634</i>	31,606	<i>32,784</i>
12-20	9,580	<i>8,003</i>	7,696	<i>6,325</i>	4,626	<i>3,952</i>
21 or Older	117,559	<i>118,501</i>	40,546	<i>43,310</i>	26,981	<i>28,832</i>
12-17	3,947	<i>3,103</i>	3,284	<i>2,531</i>	1,837	<i>1,472</i>
12-13	253	<i>158</i>	201	<i>111</i>	84	<i>32</i>
14-15	1,165	<i>1,031</i>	992	<i>897</i>	527	<i>517</i>
16-17	2,528	<i>1,914</i>	2,091	<i>1,523</i>	1,227	<i>923</i>
18 or Older	123,192	<i>123,401</i>	44,958	<i>47,103</i>	29,769	<i>31,312</i>
18-25	17,450	<i>16,604</i>	11,952	<i>11,566</i>	7,749	<i>7,727</i>
18-20	5,633	<i>4,900</i>	4,412	<i>3,794</i>	2,788	<i>2,480</i>
21-25	11,817	<i>11,704</i>	7,539	<i>7,773</i>	4,961	<i>5,247</i>
26 or Older	105,742	<i>106,797</i>	33,007	<i>35,537</i>	22,020	<i>23,585</i>
26-49	55,315	<i>54,599</i>	22,016	<i>23,445</i>	14,559	<i>15,950</i>
26-29	10,906	<i>10,105</i>	5,958	<i>5,420</i>	4,026	<i>3,787</i>
30-34	12,842	<i>12,750</i>	5,562	<i>5,924</i>	3,627	<i>4,028</i>
35-39	12,077	<i>12,385</i>	4,451	<i>5,214</i>	2,932	<i>3,561</i>
40-44	9,922	<i>9,864</i>	3,134	<i>3,611</i>	2,023	<i>2,478</i>
45-49	9,568	<i>9,496</i>	2,911	<i>3,276</i>	1,951	<i>2,097</i>
50 or Older	50,427	<i>52,198</i>	10,990	<i>12,091</i>	7,461	<i>7,635</i>
50-54	10,707	<i>9,961</i>	2,562	<i>2,789</i>	1,695	<i>1,995</i>
55-59	11,394	<i>11,409</i>	2,867	<i>2,870</i>	1,906	<i>1,580</i>
60-64	11,664	<i>11,474</i>	2,891	<i>3,140</i>	2,024	<i>2,024</i>
65 or Older	16,662	<i>19,354</i>	2,670	<i>3,292</i>	1,836	<i>2,035</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.7B Marijuana Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	46.2	<i>45.7</i>	17.5	<i>17.9</i>	11.5	<i>11.8</i>
12-20	25.2	<i>21.4</i>	20.2	<i>16.9</i>	12.1	<i>10.6</i>
21 or Older	49.6	<i>49.5</i>	17.1	<i>18.1</i>	11.4	<i>12.0</i>
12-17	15.8	<i>12.4</i>	13.2	<i>10.1</i>	7.4	<i>5.9</i>
12-13	3.1	<i>1.9</i>	2.5	<i>1.3</i>	1.0	<i>0.4</i>
14-15	13.7	<i>11.9</i>	11.7	<i>10.4</i>	6.2	<i>6.0</i>
16-17	30.6	<i>24.3</i>	25.3	<i>19.3</i>	14.9	<i>11.7</i>
18 or Older	49.2	<i>49.0</i>	18.0	<i>18.7</i>	11.9	<i>12.4</i>
18-25	51.7	<i>49.6</i>	35.4	<i>34.5</i>	23.0	<i>23.1</i>
18-20	42.8	<i>39.4</i>	33.5	<i>30.5</i>	21.2	<i>19.9</i>
21-25	57.5	<i>55.6</i>	36.7	<i>36.9</i>	24.1	<i>24.9</i>
26 or Older	48.8	<i>48.9</i>	15.2	<i>16.3</i>	10.2	<i>10.8</i>
26-49	54.6	<i>53.8</i>	21.7	<i>23.1</i>	14.4	<i>15.7</i>
26-29	60.4	<i>56.7</i>	33.0	<i>30.4</i>	22.3	<i>21.3</i>
30-34	57.7	<i>56.1</i>	25.0	<i>26.1</i>	16.3	<i>17.7</i>
35-39	56.4	<i>55.5</i>	20.8	<i>23.4</i>	13.7	<i>16.0</i>
40-44	51.5	<i>50.9</i>	16.3	<i>18.6</i>	10.5	<i>12.8</i>
45-49	47.2	<i>49.2</i>	14.4	<i>17.0</i>	9.6	<i>10.9</i>
50 or Older	43.7	<i>44.7</i>	9.5	<i>10.3</i>	6.5	<i>6.5</i>
50-54	51.9	<i>50.0</i>	12.4	<i>14.0</i>	8.2	<i>10.0</i>
55-59	53.6	<i>53.0</i>	13.5	<i>13.3</i>	9.0	<i>7.3</i>
60-64	56.7	<i>54.9</i>	14.0	<i>15.0</i>	9.8	<i>9.7</i>
65 or Older	31.5	<i>35.5</i>	5.1	<i>6.0</i>	3.5	<i>3.7</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.8A Cocaine Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	41,445	39,261	5,468	5,172	1,998	1,831
12-20	925	558	529	357	161	95
21 or Older	40,520	38,703	4,939	4,815	1,838	1,736
12-17	154	105	97	84	34	11
12-13	12	8	12	4	*	*
14-15	30	21	20	19	11	1
16-17	112	76	65	62	23	10
18 or Older	41,291	39,155	5,371	5,088	1,965	1,821
18-25	3,809	3,065	1,789	1,441	540	424
18-20	771	452	432	273	127	84
21-25	3,038	2,612	1,357	1,168	413	340
26 or Older	37,483	36,091	3,582	3,646	1,425	1,396
26-49	17,622	17,641	2,863	3,002	1,051	1,156
26-29	3,298	2,744	957	924	296	286
30-34	4,173	4,021	811	535	309	175
35-39	3,870	4,272	565	676	200	287
40-44	3,346	3,122	284	437	149	226
45-49	2,936	3,482	246	431	98	183
50 or Older	19,861	18,450	720	644	374	240
50-54	4,307	3,548	181	218	118	127
55-59	5,339	4,611	219	150	96	70
60-64	5,323	5,172	220	120	129	16
65 or Older	4,892	5,118	100	157	32	27

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.8B Cocaine Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	15.1	<i>14.2</i>	2.0	<i>1.9</i>	0.7	<i>0.7</i>
12-20	2.4	<i>1.5</i>	1.4	<i>1.0</i>	0.4	<i>0.3</i>
21 or Older	17.1	<i>16.2</i>	2.1	<i>2.0</i>	0.8	<i>0.7</i>
12-17	0.6	<i>0.4</i>	0.4	<i>0.3</i>	0.1	<i>0.0</i>
12-13	0.1	<i>0.1</i>	0.1	<i>0.0</i>	*	*
14-15	0.4	<i>0.2</i>	0.2	<i>0.2</i>	0.1	<i>0.0</i>
16-17	1.4	<i>1.0</i>	0.8	<i>0.8</i>	0.3	<i>0.1</i>
18 or Older	16.5	<i>15.5</i>	2.1	<i>2.0</i>	0.8	<i>0.7</i>
18-25	11.3	<i>9.1</i>	5.3	<i>4.3</i>	1.6	<i>1.3</i>
18-20	5.9	<i>3.6</i>	3.3	<i>2.2</i>	1.0	<i>0.7</i>
21-25	14.8	<i>12.4</i>	6.6	<i>5.5</i>	2.0	<i>1.6</i>
26 or Older	17.3	<i>16.5</i>	1.7	<i>1.7</i>	0.7	<i>0.6</i>
26-49	17.4	<i>17.4</i>	2.8	<i>3.0</i>	1.0	<i>1.1</i>
26-29	18.3	<i>15.4</i>	5.3	<i>5.2</i>	1.6	<i>1.6</i>
30-34	18.7	<i>17.7</i>	3.6	<i>2.4</i>	1.4	<i>0.8</i>
35-39	18.1	<i>19.1</i>	2.6	<i>3.0</i>	0.9	<i>1.3</i>
40-44	17.4	<i>16.1</i>	1.5	<i>2.3</i>	0.8	<i>1.2</i>
45-49	14.5	<i>18.0</i>	1.2	<i>2.2</i>	0.5	<i>0.9</i>
50 or Older	17.2	<i>15.8</i>	0.6	<i>0.6</i>	0.3	<i>0.2</i>
50-54	20.9	<i>17.8</i>	0.9	<i>1.1</i>	0.6	<i>0.6</i>
55-59	25.1	<i>21.4</i>	1.0	<i>0.7</i>	0.5	<i>0.3</i>
60-64	25.9	<i>24.7</i>	1.1	<i>0.6</i>	0.6	<i>0.1</i>
65 or Older	9.3	<i>9.4</i>	0.2	<i>0.3</i>	0.1	<i>0.0</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.9A Crack Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	9,375	9,356	778	657	378	335
12-20	71	46	27	25	10	25
21 or Older	9,304	9,311	751	632	368	310
12-17	20	3	11	2	5	2
12-13	*	*	*	*	*	*
14-15	3	*	*	*	*	*
16-17	16	3	11	2	5	2
18 or Older	9,355	9,354	766	655	372	333
18-25	339	247	61	49	18	35
18-20	51	43	15	23	4	23
21-25	287	204	46	26	13	12
26 or Older	9,016	9,107	706	606	355	298
26-49	4,334	4,463	343	371	168	244
26-29	469	389	38	40	22	19
30-34	949	854	69	52	31	17
35-39	945	1,259	110	118	49	81
40-44	1,024	829	63	83	32	78
45-49	947	1,132	63	79	34	50
50 or Older	4,682	4,644	363	234	187	53
50-54	1,275	1,116	94	138	67	29
55-59	1,266	1,523	96	14	28	6
60-64	1,376	852	99	*	75	*
65 or Older	765	1,153	75	82	18	19

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.9B Crack Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	3.4	<i>3.4</i>	0.3	<i>0.2</i>	0.1	<i>0.1</i>
12-20	0.2	<i>0.1</i>	0.1	<i>0.1</i>	0.0	<i>0.1</i>
21 or Older	3.9	<i>3.9</i>	0.3	<i>0.3</i>	0.2	<i>0.1</i>
12-17	0.1	<i>0.0</i>	0.0	<i>0.0</i>	0.0	<i>0.0</i>
12-13	*	<i>*</i>	*	<i>*</i>	*	<i>*</i>
14-15	0.0	<i>*</i>	*	<i>*</i>	*	<i>*</i>
16-17	0.2	<i>0.0</i>	0.1	<i>0.0</i>	0.1	<i>0.0</i>
18 or Older	3.7	<i>3.7</i>	0.3	<i>0.3</i>	0.1	<i>0.1</i>
18-25	1.0	<i>0.7</i>	0.2	<i>0.1</i>	0.1	<i>0.1</i>
18-20	0.4	<i>0.3</i>	0.1	<i>0.2</i>	0.0	<i>0.2</i>
21-25	1.4	<i>1.0</i>	0.2	<i>0.1</i>	0.1	<i>0.1</i>
26 or Older	4.2	<i>4.2</i>	0.3	<i>0.3</i>	0.2	<i>0.1</i>
26-49	4.3	<i>4.4</i>	0.3	<i>0.4</i>	0.2	<i>0.2</i>
26-29	2.6	<i>2.2</i>	0.2	<i>0.2</i>	0.1	<i>0.1</i>
30-34	4.3	<i>3.8</i>	0.3	<i>0.2</i>	0.1	<i>0.1</i>
35-39	4.4	<i>5.6</i>	0.5	<i>0.5</i>	0.2	<i>0.4</i>
40-44	5.3	<i>4.3</i>	0.3	<i>0.4</i>	0.2	<i>0.4</i>
45-49	4.7	<i>5.9</i>	0.3	<i>0.4</i>	0.2	<i>0.3</i>
50 or Older	4.1	<i>4.0</i>	0.3	<i>0.2</i>	0.2	<i>0.0</i>
50-54	6.2	<i>5.6</i>	0.5	<i>0.7</i>	0.3	<i>0.1</i>
55-59	6.0	<i>7.1</i>	0.5	<i>0.1</i>	0.1	<i>0.0</i>
60-64	6.7	<i>4.1</i>	0.5	<i>*</i>	0.4	<i>*</i>
65 or Older	1.4	<i>2.1</i>	0.1	<i>0.2</i>	0.0	<i>0.0</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.10A Hallucinogen Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	44,087	<i>43,949</i>	6,010	<i>7,133</i>	1,915	<i>1,761</i>
12-20	1,999	<i>1,823</i>	1,271	<i>1,202</i>	360	<i>358</i>
21 or Older	42,088	<i>42,126</i>	4,739	<i>5,931</i>	1,554	<i>1,404</i>
12-17	615	<i>563</i>	440	<i>370</i>	152	<i>80</i>
12-13	50	<i>24</i>	29	<i>14</i>	17	<i>6</i>
14-15	175	<i>161</i>	125	<i>130</i>	50	<i>19</i>
16-17	390	<i>378</i>	285	<i>227</i>	85	<i>55</i>
18 or Older	43,472	<i>43,386</i>	5,571	<i>6,763</i>	1,763	<i>1,681</i>
18-25	5,597	<i>5,269</i>	2,425	<i>2,446</i>	625	<i>645</i>
18-20	1,385	<i>1,260</i>	831	<i>831</i>	208	<i>278</i>
21-25	4,212	<i>4,010</i>	1,593	<i>1,614</i>	416	<i>368</i>
26 or Older	37,876	<i>38,117</i>	3,146	<i>4,317</i>	1,138	<i>1,036</i>
26-49	21,719	<i>21,938</i>	2,742	<i>3,757</i>	1,002	<i>979</i>
26-29	4,398	<i>3,815</i>	1,023	<i>1,142</i>	358	<i>308</i>
30-34	5,101	<i>5,020</i>	878	<i>988</i>	326	<i>159</i>
35-39	4,857	<i>4,825</i>	396	<i>770</i>	157	<i>298</i>
40-44	4,086	<i>3,957</i>	268	<i>460</i>	89	<i>124</i>
45-49	3,277	<i>4,321</i>	178	<i>397</i>	71	<i>90</i>
50 or Older	16,157	<i>16,178</i>	404	<i>560</i>	136	<i>57</i>
50-54	3,312	<i>3,274</i>	114	<i>303</i>	29	<i>11</i>
55-59	3,927	<i>3,353</i>	129	<i>17</i>	51	<i>17</i>
60-64	4,240	<i>4,545</i>	52	<i>81</i>	6	<i>*</i>
65 or Older	4,678	<i>5,006</i>	109	<i>160</i>	51	<i>29</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.10B Hallucinogen Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	16.0	<i>15.9</i>	2.2	<i>2.6</i>	0.7	<i>0.6</i>
12-20	5.3	<i>4.9</i>	3.3	<i>3.2</i>	0.9	<i>1.0</i>
21 or Older	17.7	<i>17.6</i>	2.0	<i>2.5</i>	0.7	<i>0.6</i>
12-17	2.5	<i>2.3</i>	1.8	<i>1.5</i>	0.6	<i>0.3</i>
12-13	0.6	<i>0.3</i>	0.4	<i>0.2</i>	0.2	<i>0.1</i>
14-15	2.1	<i>1.9</i>	1.5	<i>1.5</i>	0.6	<i>0.2</i>
16-17	4.7	<i>4.8</i>	3.4	<i>2.9</i>	1.0	<i>0.7</i>
18 or Older	17.4	<i>17.2</i>	2.2	<i>2.7</i>	0.7	<i>0.7</i>
18-25	16.6	<i>15.7</i>	7.2	<i>7.3</i>	1.9	<i>1.9</i>
18-20	10.5	<i>10.1</i>	6.3	<i>6.7</i>	1.6	<i>2.2</i>
21-25	20.5	<i>19.0</i>	7.7	<i>7.7</i>	2.0	<i>1.7</i>
26 or Older	17.5	<i>17.5</i>	1.5	<i>2.0</i>	0.5	<i>0.5</i>
26-49	21.4	<i>21.6</i>	2.7	<i>3.7</i>	1.0	<i>1.0</i>
26-29	24.4	<i>21.4</i>	5.7	<i>6.4</i>	2.0	<i>1.7</i>
30-34	22.9	<i>22.1</i>	3.9	<i>4.4</i>	1.5	<i>0.7</i>
35-39	22.7	<i>21.6</i>	1.8	<i>3.4</i>	0.7	<i>1.3</i>
40-44	21.2	<i>20.4</i>	1.4	<i>2.4</i>	0.5	<i>0.6</i>
45-49	16.2	<i>22.4</i>	0.9	<i>2.1</i>	0.3	<i>0.5</i>
50 or Older	14.0	<i>13.8</i>	0.3	<i>0.5</i>	0.1	<i>0.0</i>
50-54	16.1	<i>16.4</i>	0.6	<i>1.5</i>	0.1	<i>0.1</i>
55-59	18.5	<i>15.6</i>	0.6	<i>0.1</i>	0.2	<i>0.1</i>
60-64	20.6	<i>21.7</i>	0.3	<i>0.4</i>	0.0	<i>*</i>
65 or Older	8.9	<i>9.2</i>	0.2	<i>0.3</i>	0.1	<i>0.1</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.11A Inhalant Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	25,090	26,749	2,142	2,390	807	904
12-20	3,124	3,254	967	848	259	158
21 or Older	21,966	23,495	1,175	1,541	548	746
12-17	2,202	2,327	743	683	216	106
12-13	675	760	285	247	85	24
14-15	827	874	266	232	69	51
16-17	699	693	192	204	63	31
18 or Older	22,889	24,422	1,399	1,707	591	797
18-25	3,099	2,929	577	507	157	160
18-20	922	927	224	165	43	51
21-25	2,176	2,002	353	341	114	109
26 or Older	19,790	21,493	822	1,200	434	637
26-49	11,893	12,957	623	1,062	332	563
26-29	1,957	2,059	184	325	73	164
30-34	2,671	2,802	167	179	101	71
35-39	2,574	2,723	133	268	56	143
40-44	2,440	2,639	81	97	47	34
45-49	2,252	2,733	59	192	54	151
50 or Older	7,897	8,536	199	139	102	75
50-54	2,035	2,057	63	15	15	1
55-59	2,059	2,052	47	41	37	27
60-64	1,963	2,338	*	52	*	16
65 or Older	1,839	2,089	89	30	50	30

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.11B Inhalant Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	9.1	9.7	0.8	0.9	0.3	0.3
12-20	8.2	8.7	2.5	2.3	0.7	0.4
21 or Older	9.3	9.8	0.5	0.6	0.2	0.3
12-17	8.8	9.3	3.0	2.7	0.9	0.4
12-13	8.3	9.0	3.5	2.9	1.0	0.3
14-15	9.8	10.1	3.1	2.7	0.8	0.6
16-17	8.5	8.8	2.3	2.6	0.8	0.4
18 or Older	9.1	9.7	0.6	0.7	0.2	0.3
18-25	9.2	8.7	1.7	1.5	0.5	0.5
18-20	7.0	7.4	1.7	1.3	0.3	0.4
21-25	10.6	9.5	1.7	1.6	0.6	0.5
26 or Older	9.1	9.8	0.4	0.5	0.2	0.3
26-49	11.7	12.8	0.6	1.0	0.3	0.6
26-29	10.8	11.6	1.0	1.8	0.4	0.9
30-34	12.0	12.3	0.7	0.8	0.5	0.3
35-39	12.0	12.2	0.6	1.2	0.3	0.6
40-44	12.7	13.6	0.4	0.5	0.2	0.2
45-49	11.1	14.1	0.3	1.0	0.3	0.8
50 or Older	6.8	7.3	0.2	0.1	0.1	0.1
50-54	9.9	10.3	0.3	0.1	0.1	0.0
55-59	9.7	9.5	0.2	0.2	0.2	0.1
60-64	9.5	11.2	*	0.2	*	0.1
65 or Older	3.5	3.8	0.2	0.1	0.1	0.1

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.12A Methamphetamine Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	16,013	<i>15,397</i>	1,999	<i>2,550</i>	1,173	<i>1,722</i>
12-20	242	<i>125</i>	115	<i>60</i>	34	<i>34</i>
21 or Older	15,771	<i>15,273</i>	1,884	<i>2,490</i>	1,139	<i>1,688</i>
12-17	75	<i>48</i>	41	<i>21</i>	11	<i>7</i>
12-13	12	<i>1</i>	9	<i>1</i>	3	<i>1</i>
14-15	16	<i>6</i>	9	<i>4</i>	5	<i>4</i>
16-17	47	<i>40</i>	24	<i>16</i>	3	<i>2</i>
18 or Older	15,939	<i>15,349</i>	1,958	<i>2,528</i>	1,161	<i>1,715</i>
18-25	904	<i>538</i>	275	<i>171</i>	110	<i>74</i>
18-20	167	<i>77</i>	74	<i>38</i>	23	<i>27</i>
21-25	737	<i>462</i>	202	<i>133</i>	88	<i>47</i>
26 or Older	15,034	<i>14,811</i>	1,682	<i>2,358</i>	1,051	<i>1,641</i>
26-49	7,727	<i>8,215</i>	1,262	<i>1,549</i>	800	<i>1,107</i>
26-29	1,001	<i>1,002</i>	238	<i>260</i>	139	<i>151</i>
30-34	1,619	<i>1,679</i>	312	<i>233</i>	164	<i>179</i>
35-39	1,779	<i>2,102</i>	318	<i>512</i>	211	<i>334</i>
40-44	1,800	<i>1,588</i>	228	<i>273</i>	162	<i>238</i>
45-49	1,528	<i>1,844</i>	165	<i>270</i>	124	<i>204</i>
50 or Older	7,307	<i>6,597</i>	420	<i>809</i>	251	<i>534</i>
50-54	1,549	<i>1,175</i>	159	<i>202</i>	81	<i>134</i>
55-59	1,687	<i>1,577</i>	102	<i>227</i>	83	<i>90</i>
60-64	1,934	<i>1,966</i>	99	<i>102</i>	73	<i>79</i>
65 or Older	2,137	<i>1,879</i>	60	<i>278</i>	14	<i>231</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.12B Methamphetamine Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	5.8	5.6	0.7	0.9	0.4	0.6
12-20	0.6	0.3	0.3	0.2	0.1	0.1
21 or Older	6.7	6.4	0.8	1.0	0.5	0.7
12-17	0.3	0.2	0.2	0.1	0.0	0.0
12-13	0.1	0.0	0.1	0.0	0.0	0.0
14-15	0.2	0.1	0.1	0.1	0.1	0.1
16-17	0.6	0.5	0.3	0.2	0.0	0.0
18 or Older	6.4	6.1	0.8	1.0	0.5	0.7
18-25	2.7	1.6	0.8	0.5	0.3	0.2
18-20	1.3	0.6	0.6	0.3	0.2	0.2
21-25	3.6	2.2	1.0	0.6	0.4	0.2
26 or Older	6.9	6.8	0.8	1.1	0.5	0.8
26-49	7.6	8.1	1.2	1.5	0.8	1.1
26-29	5.5	5.6	1.3	1.5	0.8	0.8
30-34	7.3	7.4	1.4	1.0	0.7	0.8
35-39	8.3	9.4	1.5	2.3	1.0	1.5
40-44	9.3	8.2	1.2	1.4	0.8	1.2
45-49	7.5	9.5	0.8	1.4	0.6	1.1
50 or Older	6.3	5.6	0.4	0.7	0.2	0.5
50-54	7.5	5.9	0.8	1.0	0.4	0.7
55-59	7.9	7.3	0.5	1.1	0.4	0.4
60-64	9.4	9.4	0.5	0.5	0.4	0.4
65 or Older	4.0	3.4	0.1	0.5	0.0	0.4

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.13A Any Use of Prescription Pain Relievers in Past Year and Misuse of Prescription Pain Relievers in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	82,559	75,198	9,724	9,254	2,819	2,536
12-20	7,054	5,543	1,143	781	301	162
21 or Older	75,505	69,655	8,582	8,474	2,518	2,374
12-17	3,980	3,079	567	396	165	96
12-13	950	710	112	84	36	16
14-15	1,400	1,166	176	130	51	42
16-17	1,630	1,203	278	183	77	38
18 or Older	78,579	72,118	9,158	8,858	2,654	2,440
18-25	8,357	6,858	1,754	1,361	420	288
18-20	3,074	2,463	576	385	136	66
21-25	5,284	4,395	1,178	976	284	222
26 or Older	70,222	65,261	7,404	7,498	2,234	2,152
26-49	30,585	26,783	4,797	4,535	1,370	1,264
26-29	5,089	3,898	1,023	795	322	231
30-34	6,752	5,650	1,205	943	325	191
35-39	6,459	5,927	1,074	946	277	236
40-44	5,992	4,880	802	727	205	218
45-49	6,292	6,429	693	1,125	241	388
50 or Older	39,637	38,477	2,607	2,962	864	888
50-54	6,674	6,084	646	567	227	138
55-59	7,169	7,373	638	682	208	194
60-64	7,325	7,285	422	682	197	266
65 or Older	18,470	17,735	900	1,032	232	289

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.13B Any Use of Prescription Pain Relievers in Past Year and Misuse of Prescription Pain Relievers in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	30.0	<i>27.2</i>	3.5	<i>3.3</i>	1.0	<i>0.9</i>
12-20	18.5	<i>14.8</i>	3.0	<i>2.1</i>	0.8	<i>0.4</i>
21 or Older	31.8	<i>29.1</i>	3.6	<i>3.5</i>	1.1	<i>1.0</i>
12-17	16.0	<i>12.3</i>	2.3	<i>1.6</i>	0.7	<i>0.4</i>
12-13	11.6	<i>8.4</i>	1.4	<i>1.0</i>	0.4	<i>0.2</i>
14-15	16.5	<i>13.5</i>	2.1	<i>1.5</i>	0.6	<i>0.5</i>
16-17	19.7	<i>15.3</i>	3.4	<i>2.3</i>	0.9	<i>0.5</i>
18 or Older	31.4	<i>28.6</i>	3.7	<i>3.5</i>	1.1	<i>1.0</i>
18-25	24.8	<i>20.5</i>	5.2	<i>4.1</i>	1.2	<i>0.9</i>
18-20	23.3	<i>19.8</i>	4.4	<i>3.1</i>	1.0	<i>0.5</i>
21-25	25.7	<i>20.9</i>	5.7	<i>4.6</i>	1.4	<i>1.1</i>
26 or Older	32.4	<i>29.9</i>	3.4	<i>3.4</i>	1.0	<i>1.0</i>
26-49	30.2	<i>26.4</i>	4.7	<i>4.5</i>	1.4	<i>1.2</i>
26-29	28.2	<i>21.9</i>	5.7	<i>4.5</i>	1.8	<i>1.3</i>
30-34	30.3	<i>24.9</i>	5.4	<i>4.2</i>	1.5	<i>0.8</i>
35-39	30.1	<i>26.6</i>	5.0	<i>4.2</i>	1.3	<i>1.1</i>
40-44	31.1	<i>25.2</i>	4.2	<i>3.8</i>	1.1	<i>1.1</i>
45-49	31.1	<i>33.3</i>	3.4	<i>5.8</i>	1.2	<i>2.0</i>
50 or Older	34.4	<i>32.9</i>	2.3	<i>2.5</i>	0.7	<i>0.8</i>
50-54	32.4	<i>30.6</i>	3.1	<i>2.8</i>	1.1	<i>0.7</i>
55-59	33.7	<i>34.2</i>	3.0	<i>3.2</i>	1.0	<i>0.9</i>
60-64	35.6	<i>34.8</i>	2.1	<i>3.3</i>	1.0	<i>1.3</i>
65 or Older	35.0	<i>32.5</i>	1.7	<i>1.9</i>	0.4	<i>0.5</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.14A Any Use of Prescription Stimulants in Past Year and Misuse of Prescription Stimulants in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	18,290	<i>18,332</i>	4,929	<i>5,092</i>	1,566	<i>1,493</i>
12-20	3,339	<i>2,793</i>	1,093	<i>769</i>	301	<i>131</i>
21 or Older	14,950	<i>15,539</i>	3,835	<i>4,323</i>	1,265	<i>1,363</i>
12-17	1,837	<i>1,593</i>	430	<i>288</i>	123	<i>72</i>
12-13	378	<i>389</i>	24	<i>29</i>	3	<i>3</i>
14-15	619	<i>526</i>	119	<i>103</i>	38	<i>19</i>
16-17	841	<i>678</i>	287	<i>157</i>	82	<i>51</i>
18 or Older	16,453	<i>16,739</i>	4,499	<i>4,804</i>	1,443	<i>1,421</i>
18-25	4,320	<i>3,773</i>	1,951	<i>1,616</i>	587	<i>267</i>
18-20	1,502	<i>1,199</i>	664	<i>481</i>	178	<i>58</i>
21-25	2,818	<i>2,574</i>	1,287	<i>1,135</i>	409	<i>209</i>
26 or Older	12,133	<i>12,966</i>	2,548	<i>3,188</i>	856	<i>1,154</i>
26-49	8,105	<i>8,086</i>	2,247	<i>2,640</i>	754	<i>1,072</i>
26-29	1,998	<i>1,841</i>	695	<i>718</i>	198	<i>193</i>
30-34	2,134	<i>2,195</i>	748	<i>835</i>	253	<i>327</i>
35-39	1,588	<i>1,634</i>	410	<i>529</i>	147	<i>260</i>
40-44	1,230	<i>1,159</i>	263	<i>332</i>	114	<i>141</i>
45-49	1,156	<i>1,257</i>	130	<i>226</i>	43	<i>152</i>
50 or Older	4,028	<i>4,880</i>	301	<i>548</i>	102	<i>82</i>
50-54	865	<i>1,464</i>	72	<i>295</i>	5	<i>65</i>
55-59	1,096	<i>962</i>	128	<i>117</i>	38	<i>5</i>
60-64	727	<i>754</i>	66	<i>78</i>	59	<i>13</i>
65 or Older	1,340	<i>1,700</i>	34	<i>58</i>	*	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.14B Any Use of Prescription Stimulants in Past Year and Misuse of Prescription Stimulants in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	6.6	6.6	1.8	1.8	0.6	0.5
12-20	8.8	7.5	2.9	2.1	0.8	0.3
21 or Older	6.3	6.5	1.6	1.8	0.5	0.6
12-17	7.4	6.4	1.7	1.2	0.5	0.3
12-13	4.6	4.6	0.3	0.3	0.0	0.0
14-15	7.3	6.1	1.4	1.2	0.5	0.2
16-17	10.2	8.6	3.5	2.0	1.0	0.6
18 or Older	6.6	6.6	1.8	1.9	0.6	0.6
18-25	12.8	11.3	5.8	4.8	1.7	0.8
18-20	11.4	9.6	5.0	3.9	1.3	0.5
21-25	13.7	12.2	6.3	5.4	2.0	1.0
26 or Older	5.6	5.9	1.2	1.5	0.4	0.5
26-49	8.0	8.0	2.2	2.6	0.7	1.1
26-29	11.1	10.3	3.9	4.0	1.1	1.1
30-34	9.6	9.7	3.4	3.7	1.1	1.4
35-39	7.4	7.3	1.9	2.4	0.7	1.2
40-44	6.4	6.0	1.4	1.7	0.6	0.7
45-49	5.7	6.5	0.6	1.2	0.2	0.8
50 or Older	3.5	4.2	0.3	0.5	0.1	0.1
50-54	4.2	7.4	0.3	1.5	0.0	0.3
55-59	5.2	4.5	0.6	0.5	0.2	0.0
60-64	3.5	3.6	0.3	0.4	0.3	0.1
65 or Older	2.5	3.1	0.1	0.1	*	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.15A Any Use of Prescription Tranquilizers or Prescription Sedatives in Past Year and Misuse of Prescription Tranquilizers or Prescription Sedatives in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	44,051	<i>43,001</i>	5,895	<i>6,205</i>	1,958	<i>2,196</i>
12-20	2,564	<i>2,102</i>	909	<i>641</i>	231	<i>109</i>
21 or Older	41,488	<i>40,899</i>	4,986	<i>5,564</i>	1,727	<i>2,088</i>
12-17	1,370	<i>936</i>	436	<i>226</i>	121	<i>26</i>
12-13	276	<i>234</i>	24	<i>33</i>	10	<i>2</i>
14-15	435	<i>340</i>	124	<i>70</i>	34	<i>12</i>
16-17	659	<i>362</i>	287	<i>122</i>	78	<i>13</i>
18 or Older	42,682	<i>42,065</i>	5,459	<i>5,979</i>	1,837	<i>2,170</i>
18-25	3,995	<i>3,805</i>	1,409	<i>1,237</i>	365	<i>371</i>
18-20	1,194	<i>1,166</i>	473	<i>415</i>	110	<i>82</i>
21-25	2,801	<i>2,638</i>	936	<i>822</i>	254	<i>289</i>
26 or Older	38,687	<i>38,261</i>	4,050	<i>4,742</i>	1,472	<i>1,799</i>
26-49	16,129	<i>16,192</i>	2,582	<i>3,175</i>	1,006	<i>1,053</i>
26-29	2,519	<i>2,301</i>	600	<i>662</i>	230	<i>171</i>
30-34	3,428	<i>2,957</i>	743	<i>648</i>	318	<i>177</i>
35-39	3,305	<i>3,509</i>	514	<i>721</i>	180	<i>294</i>
40-44	3,363	<i>3,603</i>	413	<i>552</i>	181	<i>233</i>
45-49	3,514	<i>3,823</i>	312	<i>593</i>	97	<i>179</i>
50 or Older	22,558	<i>22,068</i>	1,468	<i>1,567</i>	467	<i>746</i>
50-54	3,575	<i>3,642</i>	358	<i>325</i>	106	<i>114</i>
55-59	4,234	<i>3,911</i>	408	<i>479</i>	151	<i>250</i>
60-64	4,133	<i>4,209</i>	378	<i>350</i>	103	<i>119</i>
65 or Older	10,615	<i>10,306</i>	325	<i>412</i>	106	<i>262</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.15B Any Use of Prescription Tranquilizers or Prescription Sedatives in Past Year and Misuse of Prescription Tranquilizers or Prescription Sedatives in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	16.0	<i>15.5</i>	2.1	<i>2.2</i>	0.7	<i>0.8</i>
12-20	6.7	<i>5.6</i>	2.4	<i>1.7</i>	0.6	<i>0.3</i>
21 or Older	17.5	<i>17.1</i>	2.1	<i>2.3</i>	0.7	<i>0.9</i>
12-17	5.5	<i>3.7</i>	1.8	<i>0.9</i>	0.5	<i>0.1</i>
12-13	3.4	<i>2.8</i>	0.3	<i>0.4</i>	0.1	<i>0.0</i>
14-15	5.1	<i>3.9</i>	1.5	<i>0.8</i>	0.4	<i>0.1</i>
16-17	8.0	<i>4.6</i>	3.5	<i>1.6</i>	0.9	<i>0.2</i>
18 or Older	17.1	<i>16.7</i>	2.2	<i>2.4</i>	0.7	<i>0.9</i>
18-25	11.8	<i>11.4</i>	4.2	<i>3.7</i>	1.1	<i>1.1</i>
18-20	9.1	<i>9.4</i>	3.6	<i>3.3</i>	0.8	<i>0.7</i>
21-25	13.6	<i>12.5</i>	4.6	<i>3.9</i>	1.2	<i>1.4</i>
26 or Older	17.9	<i>17.5</i>	1.9	<i>2.2</i>	0.7	<i>0.8</i>
26-49	15.9	<i>15.9</i>	2.5	<i>3.1</i>	1.0	<i>1.0</i>
26-29	14.0	<i>12.9</i>	3.3	<i>3.7</i>	1.3	<i>1.0</i>
30-34	15.4	<i>13.0</i>	3.3	<i>2.9</i>	1.4	<i>0.8</i>
35-39	15.4	<i>15.7</i>	2.4	<i>3.2</i>	0.8	<i>1.3</i>
40-44	17.4	<i>18.6</i>	2.1	<i>2.8</i>	0.9	<i>1.2</i>
45-49	17.3	<i>19.8</i>	1.5	<i>3.1</i>	0.5	<i>0.9</i>
50 or Older	19.6	<i>18.9</i>	1.3	<i>1.3</i>	0.4	<i>0.6</i>
50-54	17.3	<i>18.3</i>	1.7	<i>1.6</i>	0.5	<i>0.6</i>
55-59	19.9	<i>18.2</i>	1.9	<i>2.2</i>	0.7	<i>1.2</i>
60-64	20.1	<i>20.1</i>	1.8	<i>1.7</i>	0.5	<i>0.6</i>
65 or Older	20.1	<i>18.9</i>	0.6	<i>0.8</i>	0.2	<i>0.5</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.16A Any Use of Prescription Tranquilizers in Past Year and Misuse of Prescription Tranquilizers in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	35,601	<i>35,018</i>	5,220	<i>5,460</i>	1,781	<i>1,891</i>
12-20	1,995	<i>1,552</i>	840	<i>547</i>	216	<i>106</i>
21 or Older	33,606	<i>33,466</i>	4,381	<i>4,912</i>	1,565	<i>1,785</i>
12-17	1,010	<i>588</i>	392	<i>196</i>	112	<i>25</i>
12-13	140	<i>125</i>	16	<i>32</i>	5	<i>2</i>
14-15	308	<i>216</i>	105	<i>59</i>	32	<i>12</i>
16-17	562	<i>246</i>	271	<i>105</i>	75	<i>12</i>
18 or Older	34,591	<i>34,430</i>	4,829	<i>5,264</i>	1,669	<i>1,866</i>
18-25	3,419	<i>3,338</i>	1,313	<i>1,108</i>	337	<i>357</i>
18-20	985	<i>965</i>	448	<i>352</i>	104	<i>81</i>
21-25	2,434	<i>2,374</i>	865	<i>756</i>	233	<i>276</i>
26 or Older	31,172	<i>31,092</i>	3,516	<i>4,156</i>	1,332	<i>1,509</i>
26-49	13,668	<i>13,962</i>	2,342	<i>2,901</i>	952	<i>924</i>
26-29	2,183	<i>1,981</i>	570	<i>598</i>	227	<i>154</i>
30-34	2,986	<i>2,624</i>	711	<i>624</i>	312	<i>169</i>
35-39	2,835	<i>3,029</i>	448	<i>648</i>	161	<i>261</i>
40-44	2,886	<i>3,136</i>	373	<i>516</i>	169	<i>199</i>
45-49	2,778	<i>3,192</i>	240	<i>515</i>	83	<i>141</i>
50 or Older	17,504	<i>17,130</i>	1,174	<i>1,255</i>	380	<i>585</i>
50-54	2,905	<i>2,930</i>	300	<i>264</i>	94	<i>100</i>
55-59	3,335	<i>3,136</i>	300	<i>402</i>	124	<i>231</i>
60-64	3,234	<i>3,453</i>	336	<i>333</i>	97	<i>119</i>
65 or Older	8,030	<i>7,611</i>	238	<i>256</i>	65	<i>135</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.16B Any Use of Prescription Tranquilizers in Past Year and Misuse of Prescription Tranquilizers in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	12.9	<i>12.6</i>	1.9	<i>2.0</i>	0.6	<i>0.7</i>
12-20	5.2	<i>4.1</i>	2.2	<i>1.5</i>	0.6	<i>0.3</i>
21 or Older	14.2	<i>14.0</i>	1.8	<i>2.1</i>	0.7	<i>0.7</i>
12-17	4.1	<i>2.4</i>	1.6	<i>0.8</i>	0.5	<i>0.1</i>
12-13	1.7	<i>1.5</i>	0.2	<i>0.4</i>	0.1	<i>0.0</i>
14-15	3.6	<i>2.5</i>	1.2	<i>0.7</i>	0.4	<i>0.1</i>
16-17	6.8	<i>3.1</i>	3.3	<i>1.3</i>	0.9	<i>0.1</i>
18 or Older	13.8	<i>13.7</i>	1.9	<i>2.1</i>	0.7	<i>0.7</i>
18-25	10.1	<i>10.0</i>	3.9	<i>3.3</i>	1.0	<i>1.1</i>
18-20	7.5	<i>7.8</i>	3.4	<i>2.8</i>	0.8	<i>0.7</i>
21-25	11.8	<i>11.3</i>	4.2	<i>3.6</i>	1.1	<i>1.3</i>
26 or Older	14.4	<i>14.2</i>	1.6	<i>1.9</i>	0.6	<i>0.7</i>
26-49	13.5	<i>13.8</i>	2.3	<i>2.9</i>	0.9	<i>0.9</i>
26-29	12.1	<i>11.1</i>	3.2	<i>3.4</i>	1.3	<i>0.9</i>
30-34	13.4	<i>11.6</i>	3.2	<i>2.7</i>	1.4	<i>0.7</i>
35-39	13.2	<i>13.6</i>	2.1	<i>2.9</i>	0.8	<i>1.2</i>
40-44	15.0	<i>16.2</i>	1.9	<i>2.7</i>	0.9	<i>1.0</i>
45-49	13.7	<i>16.5</i>	1.2	<i>2.7</i>	0.4	<i>0.7</i>
50 or Older	15.2	<i>14.7</i>	1.0	<i>1.1</i>	0.3	<i>0.5</i>
50-54	14.1	<i>14.7</i>	1.5	<i>1.3</i>	0.5	<i>0.5</i>
55-59	15.7	<i>14.6</i>	1.4	<i>1.9</i>	0.6	<i>1.1</i>
60-64	15.7	<i>16.5</i>	1.6	<i>1.6</i>	0.5	<i>0.6</i>
65 or Older	15.2	<i>14.0</i>	0.5	<i>0.5</i>	0.1	<i>0.2</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.17A Any Use of Prescription Sedatives in Past Year and Misuse of Prescription Sedatives in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	15,636	<i>15,364</i>	1,098	<i>1,240</i>	304	<i>481</i>
12-20	883	<i>799</i>	138	<i>139</i>	43	<i>23</i>
21 or Older	14,752	<i>14,565</i>	960	<i>1,101</i>	261	<i>458</i>
12-17	545	<i>429</i>	82	<i>35</i>	29	<i>1</i>
12-13	166	<i>131</i>	10	<i>2</i>	5	<i>*</i>
14-15	187	<i>160</i>	35	<i>15</i>	14	<i>*</i>
16-17	192	<i>139</i>	37	<i>18</i>	10	<i>1</i>
18 or Older	15,091	<i>14,935</i>	1,016	<i>1,205</i>	275	<i>480</i>
18-25	1,051	<i>1,066</i>	175	<i>251</i>	44	<i>78</i>
18-20	339	<i>370</i>	56	<i>104</i>	15	<i>21</i>
21-25	712	<i>695</i>	119	<i>147</i>	30	<i>56</i>
26 or Older	14,040	<i>13,869</i>	841	<i>954</i>	231	<i>402</i>
26-49	4,862	<i>4,751</i>	423	<i>483</i>	112	<i>179</i>
26-29	615	<i>626</i>	71	<i>90</i>	11	<i>23</i>
30-34	967	<i>759</i>	85	<i>41</i>	43	<i>9</i>
35-39	978	<i>1,040</i>	104	<i>127</i>	31	<i>54</i>
40-44	1,008	<i>940</i>	72	<i>125</i>	12	<i>55</i>
45-49	1,293	<i>1,385</i>	91	<i>99</i>	14	<i>38</i>
50 or Older	9,178	<i>9,119</i>	417	<i>472</i>	119	<i>223</i>
50-54	1,320	<i>1,574</i>	93	<i>96</i>	35	<i>49</i>
55-59	1,752	<i>1,525</i>	145	<i>171</i>	38	<i>43</i>
60-64	1,701	<i>1,509</i>	72	<i>47</i>	6	<i>4</i>
65 or Older	4,405	<i>4,510</i>	107	<i>158</i>	41	<i>127</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.17B Any Use of Prescription Sedatives in Past Year and Misuse of Prescription Sedatives in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	5.7	5.5	0.4	<i>0.4</i>	0.1	<i>0.2</i>
12-20	2.3	<i>2.1</i>	0.4	<i>0.4</i>	0.1	<i>0.1</i>
21 or Older	6.2	<i>6.1</i>	0.4	<i>0.5</i>	0.1	<i>0.2</i>
12-17	2.2	<i>1.7</i>	0.3	<i>0.1</i>	0.1	<i>0.0</i>
12-13	2.0	<i>1.6</i>	0.1	<i>0.0</i>	0.1	*
14-15	2.2	<i>1.8</i>	0.4	<i>0.2</i>	0.2	*
16-17	2.3	<i>1.8</i>	0.4	<i>0.2</i>	0.1	<i>0.0</i>
18 or Older	6.0	<i>5.9</i>	0.4	<i>0.5</i>	0.1	<i>0.2</i>
18-25	3.1	<i>3.2</i>	0.5	<i>0.7</i>	0.1	<i>0.2</i>
18-20	2.6	<i>3.0</i>	0.4	<i>0.8</i>	0.1	<i>0.2</i>
21-25	3.5	<i>3.3</i>	0.6	<i>0.7</i>	0.1	<i>0.3</i>
26 or Older	6.5	<i>6.3</i>	0.4	<i>0.4</i>	0.1	<i>0.2</i>
26-49	4.8	<i>4.7</i>	0.4	<i>0.5</i>	0.1	<i>0.2</i>
26-29	3.4	<i>3.5</i>	0.4	<i>0.5</i>	0.1	<i>0.1</i>
30-34	4.3	<i>3.3</i>	0.4	<i>0.2</i>	0.2	<i>0.0</i>
35-39	4.6	<i>4.7</i>	0.5	<i>0.6</i>	0.1	<i>0.2</i>
40-44	5.2	<i>4.9</i>	0.4	<i>0.6</i>	0.1	<i>0.3</i>
45-49	6.4	<i>7.2</i>	0.5	<i>0.5</i>	0.1	<i>0.2</i>
50 or Older	8.0	<i>7.8</i>	0.4	<i>0.4</i>	0.1	<i>0.2</i>
50-54	6.4	<i>7.9</i>	0.4	<i>0.5</i>	0.2	<i>0.2</i>
55-59	8.2	<i>7.1</i>	0.7	<i>0.8</i>	0.2	<i>0.2</i>
60-64	8.3	<i>7.2</i>	0.4	<i>0.2</i>	0.0	<i>0.0</i>
65 or Older	8.3	<i>8.3</i>	0.2	<i>0.3</i>	0.1	<i>0.2</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.18A Any Use and Misuse of Prescription Benzodiazepines in Past Year: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)
TOTAL	29,408	<i>27,026</i>	4,841	<i>4,779</i>
12-20	1,619	<i>1,259</i>	817	<i>504</i>
21 or Older	27,789	<i>25,766</i>	4,024	<i>4,275</i>
12-17	749	<i>418</i>	381	<i>157</i>
12-13	78	<i>52</i>	14	<i>10</i>
14-15	215	<i>163</i>	107	<i>59</i>
16-17	455	<i>203</i>	259	<i>88</i>
18 or Older	28,659	<i>26,607</i>	4,460	<i>4,622</i>
18-25	2,983	<i>2,861</i>	1,266	<i>1,097</i>
18-20	870	<i>841</i>	436	<i>347</i>
21-25	2,113	<i>2,019</i>	830	<i>749</i>
26 or Older	25,676	<i>23,747</i>	3,194	<i>3,526</i>
26-49	11,285	<i>10,816</i>	2,169	<i>2,608</i>
26-29	1,850	<i>1,628</i>	550	<i>542</i>
30-34	2,617	<i>2,160</i>	683	<i>538</i>
35-39	2,268	<i>2,339</i>	399	<i>573</i>
40-44	2,343	<i>2,407</i>	312	<i>496</i>
45-49	2,207	<i>2,282</i>	225	<i>457</i>
50 or Older	14,391	<i>12,931</i>	1,026	<i>918</i>
50-54	2,384	<i>2,003</i>	251	<i>199</i>
55-59	2,705	<i>2,428</i>	269	<i>290</i>
60-64	2,629	<i>2,654</i>	268	<i>255</i>
65 or Older	6,673	<i>5,846</i>	238	<i>174</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.18B Any Use and Misuse of Prescription Benzodiazepines in Past Year: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)
TOTAL	10.7	9.8	1.8	1.7
12-20	4.3	3.4	2.1	1.3
21 or Older	11.7	10.8	1.7	1.8
12-17	3.0	1.7	1.5	0.6
12-13	1.0	0.6	0.2	0.1
14-15	2.5	1.9	1.3	0.7
16-17	5.5	2.6	3.1	1.1
18 or Older	11.4	10.6	1.8	1.8
18-25	8.8	8.5	3.8	3.3
18-20	6.6	6.8	3.3	2.8
21-25	10.3	9.6	4.0	3.6
26 or Older	11.9	10.9	1.5	1.6
26-49	11.1	10.7	2.1	2.6
26-29	10.2	9.1	3.0	3.0
30-34	11.7	9.5	3.1	2.4
35-39	10.6	10.5	1.9	2.6
40-44	12.2	12.4	1.6	2.6
45-49	10.9	11.8	1.1	2.4
50 or Older	12.5	11.1	0.9	0.8
50-54	11.6	10.1	1.2	1.0
55-59	12.7	11.3	1.3	1.3
60-64	12.8	12.7	1.3	1.2
65 or Older	12.6	10.7	0.5	0.3

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.19A Misuse of Opioids in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	10,065	<i>9,490</i>	3,101	<i>2,885</i>
12-20	1,145	<i>781</i>	305	<i>162</i>
21 or Older	8,920	<i>8,709</i>	2,796	<i>2,723</i>
12-17	567	<i>396</i>	165	<i>96</i>
12-13	112	<i>84</i>	36	<i>16</i>
14-15	176	<i>130</i>	51	<i>42</i>
16-17	278	<i>183</i>	77	<i>38</i>
18 or Older	9,499	<i>9,094</i>	2,936	<i>2,789</i>
18-25	1,776	<i>1,376</i>	439	<i>296</i>
18-20	578	<i>385</i>	140	<i>66</i>
21-25	1,197	<i>992</i>	300	<i>229</i>
26 or Older	7,723	<i>7,717</i>	2,497	<i>2,493</i>
26-49	4,971	<i>4,704</i>	1,514	<i>1,419</i>
26-29	1,041	<i>841</i>	351	<i>294</i>
30-34	1,271	<i>984</i>	351	<i>221</i>
35-39	1,126	<i>982</i>	328	<i>263</i>
40-44	821	<i>740</i>	222	<i>245</i>
45-49	712	<i>1,157</i>	262	<i>397</i>
50 or Older	2,752	<i>3,013</i>	983	<i>1,075</i>
50-54	657	<i>568</i>	237	<i>188</i>
55-59	680	<i>695</i>	248	<i>295</i>
60-64	436	<i>682</i>	209	<i>276</i>
65 or Older	979	<i>1,068</i>	290	<i>316</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.19B Misuse of Opioids in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	3.7	<i>3.4</i>	1.1	<i>1.0</i>
12-20	3.0	<i>2.1</i>	0.8	<i>0.4</i>
21 or Older	3.8	<i>3.6</i>	1.2	<i>1.1</i>
12-17	2.3	<i>1.6</i>	0.7	<i>0.4</i>
12-13	1.4	<i>1.0</i>	0.4	<i>0.2</i>
14-15	2.1	<i>1.5</i>	0.6	<i>0.5</i>
16-17	3.4	<i>2.3</i>	0.9	<i>0.5</i>
18 or Older	3.8	<i>3.6</i>	1.2	<i>1.1</i>
18-25	5.3	<i>4.1</i>	1.3	<i>0.9</i>
18-20	4.4	<i>3.1</i>	1.1	<i>0.5</i>
21-25	5.8	<i>4.7</i>	1.5	<i>1.1</i>
26 or Older	3.6	<i>3.5</i>	1.2	<i>1.1</i>
26-49	4.9	<i>4.6</i>	1.5	<i>1.4</i>
26-29	5.8	<i>4.7</i>	1.9	<i>1.7</i>
30-34	5.7	<i>4.3</i>	1.6	<i>1.0</i>
35-39	5.3	<i>4.4</i>	1.5	<i>1.2</i>
40-44	4.3	<i>3.8</i>	1.2	<i>1.3</i>
45-49	3.5	<i>6.0</i>	1.3	<i>2.1</i>
50 or Older	2.4	<i>2.6</i>	0.9	<i>0.9</i>
50-54	3.2	<i>2.9</i>	1.2	<i>0.9</i>
55-59	3.2	<i>3.2</i>	1.2	<i>1.4</i>
60-64	2.1	<i>3.3</i>	1.0	<i>1.3</i>
65 or Older	1.9	<i>2.0</i>	0.5	<i>0.6</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.20A Misuse of Central Nervous System Stimulants in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	10,136	<i>10,306</i>	4,264	<i>4,483</i>
12-20	1,444	<i>1,021</i>	465	<i>237</i>
21 or Older	8,693	<i>9,285</i>	3,799	<i>4,246</i>
12-17	499	<i>346</i>	160	<i>87</i>
12-13	39	<i>34</i>	7	<i>4</i>
14-15	144	<i>123</i>	54	<i>24</i>
16-17	316	<i>189</i>	100	<i>59</i>
18 or Older	9,638	<i>9,960</i>	4,103	<i>4,396</i>
18-25	3,205	<i>2,567</i>	1,127	<i>682</i>
18-20	945	<i>675</i>	305	<i>150</i>
21-25	2,260	<i>1,892</i>	823	<i>532</i>
26 or Older	6,433	<i>7,393</i>	2,976	<i>3,714</i>
26-49	5,196	<i>5,742</i>	2,311	<i>2,889</i>
26-29	1,591	<i>1,531</i>	570	<i>593</i>
30-34	1,467	<i>1,305</i>	637	<i>584</i>
35-39	1,039	<i>1,334</i>	489	<i>797</i>
40-44	617	<i>884</i>	370	<i>538</i>
45-49	483	<i>688</i>	245	<i>376</i>
50 or Older	1,236	<i>1,651</i>	665	<i>825</i>
50-54	367	<i>644</i>	202	<i>326</i>
55-59	379	<i>468</i>	208	<i>164</i>
60-64	316	<i>221</i>	223	<i>95</i>
65 or Older	174	<i>318</i>	33	<i>239</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.20B Misuse of Central Nervous System Stimulants in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	3.7	3.7	1.5	1.6
12-20	3.8	2.7	1.2	0.6
21 or Older	3.7	3.9	1.6	1.8
12-17	2.0	1.4	0.6	0.3
12-13	0.5	0.4	0.1	0.0
14-15	1.7	1.4	0.6	0.3
16-17	3.8	2.4	1.2	0.7
18 or Older	3.9	4.0	1.6	1.7
18-25	9.5	7.7	3.3	2.0
18-20	7.2	5.4	2.3	1.2
21-25	11.0	9.0	4.0	2.5
26 or Older	3.0	3.4	1.4	1.7
26-49	5.1	5.7	2.3	2.8
26-29	8.8	8.6	3.2	3.3
30-34	6.6	5.7	2.9	2.6
35-39	4.9	6.0	2.3	3.6
40-44	3.2	4.6	1.9	2.8
45-49	2.4	3.6	1.2	1.9
50 or Older	1.1	1.4	0.6	0.7
50-54	1.8	3.2	1.0	1.6
55-59	1.8	2.2	1.0	0.8
60-64	1.5	1.1	1.1	0.5
65 or Older	0.3	0.6	0.1	0.4

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.21A Use of Illicit Drugs Other Than Marijuana in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	23,588	<i>23,962</i>	9,333	<i>9,305</i>
12-20	3,788	<i>3,077</i>	1,224	<i>799</i>
21 or Older	19,800	<i>20,885</i>	8,110	<i>8,506</i>
12-17	1,862	<i>1,507</i>	598	<i>305</i>
12-13	425	<i>359</i>	141	<i>48</i>
14-15	590	<i>513</i>	165	<i>113</i>
16-17	848	<i>635</i>	292	<i>144</i>
18 or Older	21,727	<i>22,455</i>	8,735	<i>9,000</i>
18-25	5,791	<i>5,169</i>	2,059	<i>1,600</i>
18-20	1,926	<i>1,570</i>	626	<i>494</i>
21-25	3,865	<i>3,598</i>	1,433	<i>1,106</i>
26 or Older	15,936	<i>17,286</i>	6,676	<i>7,400</i>
26-49	11,025	<i>11,693</i>	4,650	<i>5,034</i>
26-29	2,888	<i>2,743</i>	1,132	<i>1,144</i>
30-34	2,966	<i>2,861</i>	1,266	<i>965</i>
35-39	2,169	<i>2,419</i>	926	<i>1,138</i>
40-44	1,625	<i>1,628</i>	685	<i>847</i>
45-49	1,377	<i>2,042</i>	641	<i>940</i>
50 or Older	4,911	<i>5,593</i>	2,026	<i>2,366</i>
50-54	1,257	<i>1,468</i>	504	<i>597</i>
55-59	1,224	<i>1,434</i>	512	<i>689</i>
60-64	996	<i>1,133</i>	518	<i>416</i>
65 or Older	1,433	<i>1,557</i>	493	<i>665</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.21B Use of Illicit Drugs Other Than Marijuana in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Percentages, 2019 and 2020

Age Category	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	8.6	8.7	3.4	3.4
12-20	9.9	8.2	3.2	2.1
21 or Older	8.3	8.7	3.4	3.6
12-17	7.5	6.0	2.4	1.2
12-13	5.2	4.3	1.7	0.6
14-15	6.9	5.9	1.9	1.3
16-17	10.3	8.1	3.5	1.8
18 or Older	8.7	8.9	3.5	3.6
18-25	17.2	15.4	6.1	4.8
18-20	14.6	12.6	4.8	4.0
21-25	18.8	17.1	7.0	5.3
26 or Older	7.4	7.9	3.1	3.4
26-49	10.9	11.5	4.6	5.0
26-29	16.0	15.4	6.3	6.4
30-34	13.3	12.6	5.7	4.2
35-39	10.1	10.8	4.3	5.1
40-44	8.4	8.4	3.6	4.4
45-49	6.8	10.6	3.2	4.9
50 or Older	4.3	4.8	1.8	2.0
50-54	6.1	7.4	2.4	3.0
55-59	5.8	6.7	2.4	3.2
60-64	4.8	5.4	2.5	2.0
65 or Older	2.7	2.9	0.9	1.2

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.