

Table 1.6C Illicit Drug Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	959	<i>1,380</i>	734	<i>1,089</i>	556	<i>890</i>
12-20	247	<i>348</i>	214	<i>306</i>	148	<i>216</i>
21 or Older	1,457	<i>1,982</i>	812	<i>1,202</i>	599	<i>931</i>
12-17	103	<i>206</i>	89	<i>167</i>	69	<i>113</i>
12-13	50	<i>114</i>	35	<i>66</i>	22	<i>16</i>
14-15	75	<i>128</i>	61	<i>108</i>	42	<i>71</i>
16-17	89	<i>150</i>	79	<i>127</i>	57	<i>95</i>
18 or Older	927	<i>1,359</i>	716	<i>1,076</i>	547	<i>880</i>
18-25	184	<i>342</i>	181	<i>325</i>	153	<i>280</i>
18-20	198	<i>273</i>	177	<i>243</i>	121	<i>179</i>
21-25	264	<i>404</i>	207	<i>328</i>	159	<i>256</i>
26 or Older	898	<i>1,310</i>	669	<i>989</i>	506	<i>804</i>
26-49	455	<i>704</i>	392	<i>640</i>	314	<i>554</i>
26-29	312	<i>436</i>	239	<i>343</i>	184	<i>267</i>
30-34	323	<i>540</i>	218	<i>375</i>	166	<i>305</i>
35-39	313	<i>481</i>	196	<i>310</i>	149	<i>256</i>
40-44	309	<i>422</i>	173	<i>277</i>	136	<i>226</i>
45-49	272	<i>417</i>	162	<i>271</i>	123	<i>229</i>
50 or Older	724	<i>1,086</i>	484	<i>750</i>	372	<i>598</i>
50-54	442	<i>658</i>	232	<i>396</i>	182	<i>306</i>
55-59	448	<i>674</i>	242	<i>389</i>	189	<i>297</i>
60-64	465	<i>703</i>	247	<i>453</i>	197	<i>342</i>
65 or Older	450	<i>653</i>	245	<i>373</i>	186	<i>289</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Illicit Drug Use in Lifetime includes the misuse of prescription psychotherapeutics even though those estimates are not reported due to potential underreporting in lifetime prescription psychotherapeutics estimates.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.6D Illicit Drug Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	0.35	<i>0.50</i>	0.27	<i>0.39</i>	0.20	<i>0.32</i>
12-20	0.45	<i>0.77</i>	0.43	<i>0.73</i>	0.33	<i>0.54</i>
21 or Older	0.39	<i>0.56</i>	0.30	<i>0.44</i>	0.23	<i>0.36</i>
12-17	0.41	<i>0.82</i>	0.36	<i>0.67</i>	0.28	<i>0.45</i>
12-13	0.57	<i>1.25</i>	0.42	<i>0.76</i>	0.27	<i>0.19</i>
14-15	0.75	<i>1.34</i>	0.64	<i>1.17</i>	0.48	<i>0.77</i>
16-17	0.82	<i>1.66</i>	0.78	<i>1.48</i>	0.65	<i>1.13</i>
18 or Older	0.37	<i>0.54</i>	0.29	<i>0.43</i>	0.22	<i>0.35</i>
18-25	0.55	<i>1.02</i>	0.54	<i>0.97</i>	0.45	<i>0.84</i>
18-20	0.82	<i>1.59</i>	0.82	<i>1.58</i>	0.68	<i>1.28</i>
21-25	0.67	<i>1.23</i>	0.69	<i>1.18</i>	0.59	<i>1.04</i>
26 or Older	0.41	<i>0.60</i>	0.31	<i>0.45</i>	0.23	<i>0.37</i>
26-49	0.45	<i>0.69</i>	0.39	<i>0.63</i>	0.31	<i>0.55</i>
26-29	0.93	<i>1.43</i>	0.95	<i>1.49</i>	0.82	<i>1.30</i>
30-34	0.87	<i>1.40</i>	0.81	<i>1.31</i>	0.67	<i>1.17</i>
35-39	0.90	<i>1.39</i>	0.78	<i>1.19</i>	0.65	<i>1.07</i>
40-44	1.04	<i>1.51</i>	0.77	<i>1.29</i>	0.66	<i>1.09</i>
45-49	0.95	<i>1.50</i>	0.73	<i>1.21</i>	0.57	<i>1.09</i>
50 or Older	0.63	<i>0.93</i>	0.42	<i>0.64</i>	0.32	<i>0.51</i>
50-54	1.35	<i>2.27</i>	1.02	<i>1.78</i>	0.83	<i>1.43</i>
55-59	1.40	<i>2.21</i>	1.04	<i>1.68</i>	0.84	<i>1.31</i>
60-64	1.29	<i>2.12</i>	1.06	<i>1.90</i>	0.88	<i>1.52</i>
65 or Older	0.85	<i>1.20</i>	0.46	<i>0.68</i>	0.35	<i>0.53</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Illicit Drug Use in Lifetime includes the misuse of prescription psychotherapeutics even though those estimates are not reported due to potential underreporting in lifetime prescription psychotherapeutics estimates.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.7C Marijuana Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	958	<i>1,374</i>	669	<i>1,035</i>	523	825
12-20	222	<i>318</i>	197	<i>285</i>	143	214
21 or Older	1,399	<i>1,898</i>	727	<i>1,096</i>	550	852
12-17	89	<i>162</i>	84	<i>142</i>	65	<i>111</i>
12-13	22	<i>34</i>	20	<i>25</i>	13	<i>10</i>
14-15	61	<i>98</i>	53	<i>94</i>	40	<i>69</i>
16-17	80	<i>130</i>	75	<i>115</i>	55	<i>93</i>
18 or Older	931	<i>1,347</i>	653	<i>1,024</i>	513	814
18-25	191	<i>353</i>	176	<i>324</i>	148	278
18-20	186	<i>269</i>	163	<i>239</i>	115	<i>177</i>
21-25	253	<i>395</i>	197	<i>319</i>	151	<i>252</i>
26 or Older	900	<i>1,301</i>	610	<i>919</i>	478	746
26-49	452	<i>706</i>	373	<i>596</i>	304	520
26-29	301	<i>424</i>	231	<i>310</i>	174	256
30-34	307	<i>517</i>	202	<i>331</i>	156	286
35-39	302	<i>468</i>	175	<i>293</i>	139	247
40-44	291	<i>407</i>	157	<i>261</i>	125	209
45-49	261	<i>395</i>	144	<i>249</i>	113	190
50 or Older	716	<i>1,079</i>	431	<i>683</i>	343	534
50-54	424	<i>613</i>	209	<i>346</i>	163	285
55-59	432	<i>651</i>	211	<i>338</i>	175	239
60-64	452	<i>679</i>	235	<i>431</i>	184	323
65 or Older	437	<i>634</i>	204	<i>316</i>	171	250

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.7D Marijuana Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	0.35	<i>0.50</i>	0.24	<i>0.37</i>	0.19	<i>0.30</i>
12-20	0.44	<i>0.77</i>	0.41	<i>0.69</i>	0.33	<i>0.54</i>
21 or Older	0.39	<i>0.55</i>	0.27	<i>0.41</i>	0.21	<i>0.33</i>
12-17	0.36	<i>0.65</i>	0.34	<i>0.57</i>	0.26	<i>0.45</i>
12-13	0.27	<i>0.40</i>	0.25	<i>0.29</i>	0.15	<i>0.12</i>
14-15	0.66	<i>1.07</i>	0.58	<i>1.03</i>	0.45	<i>0.76</i>
16-17	0.80	<i>1.51</i>	0.77	<i>1.36</i>	0.63	<i>1.11</i>
18 or Older	0.37	<i>0.53</i>	0.26	<i>0.41</i>	0.21	<i>0.32</i>
18-25	0.57	<i>1.06</i>	0.52	<i>0.97</i>	0.44	<i>0.83</i>
18-20	0.83	<i>1.64</i>	0.79	<i>1.56</i>	0.65	<i>1.27</i>
21-25	0.69	<i>1.27</i>	0.66	<i>1.19</i>	0.57	<i>1.03</i>
26 or Older	0.42	<i>0.60</i>	0.28	<i>0.42</i>	0.22	<i>0.34</i>
26-49	0.45	<i>0.69</i>	0.37	<i>0.59</i>	0.30	<i>0.51</i>
26-29	0.96	<i>1.50</i>	0.94	<i>1.41</i>	0.79	<i>1.27</i>
30-34	0.88	<i>1.39</i>	0.78	<i>1.22</i>	0.63	<i>1.12</i>
35-39	0.89	<i>1.38</i>	0.72	<i>1.17</i>	0.62	<i>1.05</i>
40-44	1.02	<i>1.50</i>	0.74	<i>1.23</i>	0.62	<i>1.01</i>
45-49	0.94	<i>1.52</i>	0.65	<i>1.17</i>	0.53	<i>0.92</i>
50 or Older	0.62	<i>0.92</i>	0.37	<i>0.58</i>	0.30	<i>0.46</i>
50-54	1.35	<i>2.28</i>	0.92	<i>1.59</i>	0.75	<i>1.35</i>
55-59	1.40	<i>2.21</i>	0.93	<i>1.50</i>	0.79	<i>1.08</i>
60-64	1.30	<i>2.09</i>	1.02	<i>1.84</i>	0.83	<i>1.43</i>
65 or Older	0.83	<i>1.16</i>	0.39	<i>0.58</i>	0.32	<i>0.46</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.8C Cocaine Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	671	948	215	306	130	192
12-20	59	63	47	53	25	31
21 or Older	750	1,036	212	300	128	189
12-17	21	27	17	26	12	9
12-13	6	6	6	4	*	*
14-15	10	15	9	15	8	1
16-17	17	22	14	21	8	9
18 or Older	665	936	213	305	129	192
18-25	118	171	83	124	44	68
18-20	53	57	42	46	22	30
21-25	118	168	80	116	37	59
26 or Older	647	915	188	271	122	175
26-49	334	520	144	232	87	151
26-29	167	207	81	136	45	66
30-34	182	279	78	77	46	46
35-39	168	265	65	106	39	69
40-44	162	230	47	86	38	67
45-49	136	255	42	108	28	74
50 or Older	524	771	115	136	82	86
50-54	267	371	52	76	42	63
55-59	287	407	55	65	37	54
60-64	314	464	62	66	52	16
65 or Older	265	387	61	67	20	20

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.8D Cocaine Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	0.24	<i>0.34</i>	0.08	<i>0.11</i>	0.05	<i>0.07</i>
12-20	0.15	<i>0.17</i>	0.12	<i>0.14</i>	0.06	<i>0.08</i>
21 or Older	0.28	<i>0.39</i>	0.09	<i>0.12</i>	0.05	<i>0.08</i>
12-17	0.08	<i>0.11</i>	0.07	<i>0.10</i>	0.05	<i>0.03</i>
12-13	0.07	<i>0.07</i>	0.07	<i>0.05</i>	*	*
14-15	0.12	<i>0.17</i>	0.10	<i>0.17</i>	0.10	<i>0.01</i>
16-17	0.20	<i>0.28</i>	0.16	<i>0.27</i>	0.10	<i>0.11</i>
18 or Older	0.27	<i>0.37</i>	0.09	<i>0.12</i>	0.05	<i>0.08</i>
18-25	0.35	<i>0.51</i>	0.24	<i>0.37</i>	0.13	<i>0.20</i>
18-20	0.38	<i>0.46</i>	0.30	<i>0.36</i>	0.16	<i>0.24</i>
21-25	0.49	<i>0.75</i>	0.35	<i>0.54</i>	0.18	<i>0.28</i>
26 or Older	0.30	<i>0.42</i>	0.09	<i>0.12</i>	0.06	<i>0.08</i>
26-49	0.33	<i>0.51</i>	0.14	<i>0.23</i>	0.09	<i>0.15</i>
26-29	0.81	<i>1.08</i>	0.43	<i>0.74</i>	0.25	<i>0.37</i>
30-34	0.71	<i>1.08</i>	0.34	<i>0.33</i>	0.21	<i>0.20</i>
35-39	0.70	<i>1.06</i>	0.30	<i>0.47</i>	0.18	<i>0.31</i>
40-44	0.74	<i>1.08</i>	0.24	<i>0.44</i>	0.20	<i>0.35</i>
45-49	0.63	<i>1.21</i>	0.21	<i>0.55</i>	0.14	<i>0.38</i>
50 or Older	0.45	<i>0.66</i>	0.10	<i>0.12</i>	0.07	<i>0.07</i>
50-54	1.14	<i>1.69</i>	0.25	<i>0.38</i>	0.20	<i>0.32</i>
55-59	1.19	<i>1.74</i>	0.26	<i>0.30</i>	0.17	<i>0.25</i>
60-64	1.26	<i>1.90</i>	0.30	<i>0.31</i>	0.25	<i>0.07</i>
65 or Older	0.50	<i>0.71</i>	0.12	<i>0.12</i>	0.04	<i>0.04</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.9C Crack Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	328	535	100	<i>120</i>	62	83
12-20	14	<i>17</i>	9	<i>14</i>	7	<i>14</i>
21 or Older	331	547	101	<i>118</i>	62	79
12-17	8	2	7	2	5	2
12-13	*	*	*	*	*	*
14-15	3	*	*	*	*	*
16-17	8	2	7	2	5	2
18 or Older	329	532	100	<i>120</i>	62	83
18-25	33	41	12	21	6	16
18-20	12	17	6	14	4	14
21-25	31	37	11	15	5	9
26 or Older	326	527	100	<i>117</i>	63	79
26-49	172	281	46	86	32	73
26-29	52	71	13	18	10	12
30-34	74	129	18	18	12	9
35-39	78	155	27	47	16	32
40-44	86	120	18	46	13	46
45-49	78	142	23	37	18	33
50 or Older	279	455	87	79	54	28
50-54	149	200	35	58	29	21
55-59	144	279	36	10	14	6
60-64	161	194	38	*	34	*
65 or Older	118	223	59	52	15	19

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.9D Crack Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	0.12	<i>0.19</i>	0.04	<i>0.04</i>	0.02	<i>0.03</i>
12-20	0.04	<i>0.05</i>	0.02	<i>0.04</i>	0.02	<i>0.04</i>
21 or Older	0.14	<i>0.22</i>	0.04	<i>0.05</i>	0.03	<i>0.03</i>
12-17	0.03	<i>0.01</i>	0.03	<i>0.01</i>	0.02	<i>0.01</i>
12-13	*	*	*	*	*	*
14-15	0.03	*	*	*	*	*
16-17	0.09	<i>0.02</i>	0.08	<i>0.02</i>	0.06	<i>0.02</i>
18 or Older	0.13	<i>0.21</i>	0.04	<i>0.05</i>	0.02	<i>0.03</i>
18-25	0.10	<i>0.12</i>	0.04	<i>0.06</i>	0.02	<i>0.05</i>
18-20	0.09	<i>0.14</i>	0.05	<i>0.11</i>	0.03	<i>0.11</i>
21-25	0.15	<i>0.18</i>	0.05	<i>0.07</i>	0.02	<i>0.04</i>
26 or Older	0.15	<i>0.24</i>	0.05	<i>0.05</i>	0.03	<i>0.04</i>
26-49	0.17	<i>0.28</i>	0.05	<i>0.08</i>	0.03	<i>0.07</i>
26-29	0.29	<i>0.40</i>	0.07	<i>0.10</i>	0.06	<i>0.07</i>
30-34	0.33	<i>0.55</i>	0.08	<i>0.08</i>	0.05	<i>0.04</i>
35-39	0.37	<i>0.68</i>	0.13	<i>0.21</i>	0.07	<i>0.14</i>
40-44	0.43	<i>0.61</i>	0.09	<i>0.24</i>	0.07	<i>0.24</i>
45-49	0.38	<i>0.73</i>	0.11	<i>0.19</i>	0.09	<i>0.17</i>
50 or Older	0.24	<i>0.39</i>	0.08	<i>0.07</i>	0.05	<i>0.02</i>
50-54	0.70	<i>0.96</i>	0.17	<i>0.29</i>	0.14	<i>0.10</i>
55-59	0.65	<i>1.23</i>	0.17	<i>0.05</i>	0.07	<i>0.03</i>
60-64	0.75	<i>0.91</i>	0.18	*	0.17	*
65 or Older	0.22	<i>0.41</i>	0.11	<i>0.10</i>	0.03	<i>0.03</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.10C Hallucinogen Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	699	972	212	390	115	149
12-20	85	138	69	118	35	71
21 or Older	775	1,047	197	361	110	131
12-17	39	68	33	52	21	19
12-13	10	8	7	6	5	3
14-15	22	38	18	35	15	10
16-17	32	55	27	38	14	16
18 or Older	687	967	205	380	111	147
18-25	134	231	89	166	47	85
18-20	72	122	60	106	27	67
21-25	135	206	78	135	39	55
26 or Older	659	937	170	317	101	123
26-49	362	576	145	272	89	120
26-29	186	263	90	141	52	75
30-34	197	294	79	131	49	45
35-39	190	269	57	114	36	68
40-44	176	255	48	89	24	39
45-49	145	271	32	98	18	33
50 or Older	509	726	86	150	45	25
50-54	242	347	54	107	19	7
55-59	247	310	45	11	27	11
60-64	284	456	28	54	6	*
65 or Older	264	371	41	91	30	21

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.10D Hallucinogen Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	0.25	<i>0.35</i>	0.08	<i>0.14</i>	0.04	<i>0.05</i>
12-20	0.21	<i>0.36</i>	0.17	<i>0.31</i>	0.09	<i>0.19</i>
21 or Older	0.29	<i>0.40</i>	0.08	<i>0.15</i>	0.05	<i>0.05</i>
12-17	0.16	<i>0.27</i>	0.13	<i>0.21</i>	0.09	<i>0.07</i>
12-13	0.12	<i>0.09</i>	0.09	<i>0.07</i>	0.07	<i>0.04</i>
14-15	0.26	<i>0.44</i>	0.21	<i>0.40</i>	0.17	<i>0.11</i>
16-17	0.38	<i>0.69</i>	0.32	<i>0.49</i>	0.17	<i>0.20</i>
18 or Older	0.27	<i>0.38</i>	0.08	<i>0.15</i>	0.04	<i>0.06</i>
18-25	0.40	<i>0.69</i>	0.26	<i>0.50</i>	0.14	<i>0.25</i>
18-20	0.47	<i>0.92</i>	0.40	<i>0.82</i>	0.20	<i>0.53</i>
21-25	0.54	<i>0.89</i>	0.34	<i>0.61</i>	0.19	<i>0.26</i>
26 or Older	0.30	<i>0.43</i>	0.08	<i>0.15</i>	0.05	<i>0.06</i>
26-49	0.36	<i>0.57</i>	0.14	<i>0.27</i>	0.09	<i>0.12</i>
26-29	0.84	<i>1.32</i>	0.47	<i>0.77</i>	0.28	<i>0.42</i>
30-34	0.76	<i>1.13</i>	0.34	<i>0.56</i>	0.22	<i>0.20</i>
35-39	0.77	<i>1.08</i>	0.27	<i>0.51</i>	0.17	<i>0.30</i>
40-44	0.79	<i>1.21</i>	0.25	<i>0.45</i>	0.12	<i>0.20</i>
45-49	0.67	<i>1.25</i>	0.16	<i>0.50</i>	0.09	<i>0.17</i>
50 or Older	0.44	<i>0.62</i>	0.07	<i>0.13</i>	0.04	<i>0.02</i>
50-54	1.05	<i>1.64</i>	0.26	<i>0.53</i>	0.09	<i>0.04</i>
55-59	1.08	<i>1.36</i>	0.21	<i>0.05</i>	0.13	<i>0.05</i>
60-64	1.18	<i>1.91</i>	0.13	<i>0.26</i>	0.03	*
65 or Older	0.50	<i>0.68</i>	0.08	<i>0.17</i>	0.06	<i>0.04</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.11C Inhalant Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	464	735	120	<i>193</i>	84	<i>129</i>
12-20	97	<i>189</i>	51	<i>91</i>	27	<i>32</i>
21 or Older	490	<i>768</i>	105	<i>174</i>	80	<i>126</i>
12-17	70	<i>156</i>	40	<i>81</i>	24	<i>22</i>
12-13	41	<i>108</i>	26	<i>47</i>	15	<i>10</i>
14-15	42	<i>87</i>	22	<i>45</i>	12	<i>14</i>
16-17	45	<i>85</i>	23	<i>47</i>	14	<i>14</i>
18 or Older	455	<i>724</i>	111	<i>176</i>	80	<i>127</i>
18-25	100	<i>159</i>	50	<i>61</i>	26	<i>41</i>
18-20	60	<i>91</i>	29	<i>35</i>	11	<i>23</i>
21-25	94	<i>142</i>	39	<i>49</i>	23	<i>33</i>
26 or Older	443	<i>698</i>	96	<i>169</i>	75	<i>125</i>
26-49	266	<i>461</i>	64	<i>163</i>	52	<i>121</i>
26-29	116	<i>212</i>	37	<i>74</i>	28	<i>52</i>
30-34	140	<i>223</i>	36	<i>48</i>	30	<i>33</i>
35-39	129	<i>210</i>	32	<i>85</i>	20	<i>58</i>
40-44	139	<i>215</i>	24	<i>42</i>	18	<i>16</i>
45-49	124	<i>227</i>	21	<i>86</i>	21	<i>82</i>
50 or Older	344	<i>534</i>	67	<i>45</i>	48	<i>30</i>
50-54	179	<i>260</i>	38	<i>13</i>	15	<i>1</i>
55-59	176	<i>295</i>	26	<i>24</i>	24	<i>19</i>
60-64	186	<i>294</i>	*	<i>30</i>	*	<i>11</i>
65 or Older	164	<i>256</i>	49	<i>21</i>	39	<i>21</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.11D Inhalant Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	0.17	<i>0.27</i>	0.04	<i>0.07</i>	0.03	<i>0.05</i>
12-20	0.23	<i>0.48</i>	0.13	<i>0.24</i>	0.07	<i>0.09</i>
21 or Older	0.19	<i>0.30</i>	0.04	<i>0.07</i>	0.03	<i>0.05</i>
12-17	0.28	<i>0.62</i>	0.16	<i>0.32</i>	0.10	<i>0.09</i>
12-13	0.47	<i>1.20</i>	0.31	<i>0.54</i>	0.19	<i>0.11</i>
14-15	0.47	<i>0.96</i>	0.26	<i>0.52</i>	0.14	<i>0.16</i>
16-17	0.51	<i>1.04</i>	0.27	<i>0.58</i>	0.17	<i>0.17</i>
18 or Older	0.18	<i>0.29</i>	0.04	<i>0.07</i>	0.03	<i>0.05</i>
18-25	0.30	<i>0.48</i>	0.15	<i>0.18</i>	0.08	<i>0.12</i>
18-20	0.42	<i>0.72</i>	0.21	<i>0.28</i>	0.09	<i>0.19</i>
21-25	0.41	<i>0.64</i>	0.19	<i>0.24</i>	0.11	<i>0.16</i>
26 or Older	0.20	<i>0.32</i>	0.04	<i>0.08</i>	0.03	<i>0.06</i>
26-49	0.26	<i>0.45</i>	0.06	<i>0.16</i>	0.05	<i>0.12</i>
26-29	0.58	<i>1.12</i>	0.20	<i>0.41</i>	0.15	<i>0.29</i>
30-34	0.59	<i>0.90</i>	0.16	<i>0.21</i>	0.14	<i>0.14</i>
35-39	0.57	<i>0.89</i>	0.15	<i>0.38</i>	0.09	<i>0.26</i>
40-44	0.66	<i>1.03</i>	0.13	<i>0.22</i>	0.09	<i>0.08</i>
45-49	0.58	<i>1.12</i>	0.11	<i>0.44</i>	0.10	<i>0.42</i>
50 or Older	0.30	<i>0.46</i>	0.06	<i>0.04</i>	0.04	<i>0.03</i>
50-54	0.80	<i>1.25</i>	0.18	<i>0.07</i>	0.07	<i>0.01</i>
55-59	0.79	<i>1.27</i>	0.12	<i>0.11</i>	0.11	<i>0.09</i>
60-64	0.84	<i>1.34</i>	*	<i>0.14</i>	*	<i>0.05</i>
65 or Older	0.31	<i>0.47</i>	0.09	<i>0.04</i>	0.07	<i>0.04</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.12C Methamphetamine Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	443	<i>641</i>	133	<i>265</i>	108	<i>215</i>
12-20	28	<i>32</i>	19	<i>20</i>	11	<i>15</i>
21 or Older	459	<i>661</i>	133	<i>265</i>	107	<i>214</i>
12-17	14	<i>22</i>	10	<i>12</i>	5	<i>5</i>
12-13	6	<i>1</i>	5	<i>1</i>	2	<i>1</i>
14-15	6	<i>5</i>	4	<i>4</i>	3	<i>4</i>
16-17	11	<i>21</i>	7	<i>11</i>	2	<i>2</i>
18 or Older	443	<i>636</i>	134	<i>265</i>	108	<i>214</i>
18-25	56	<i>67</i>	31	<i>31</i>	19	<i>20</i>
18-20	24	<i>21</i>	16	<i>16</i>	10	<i>14</i>
21-25	50	<i>62</i>	26	<i>27</i>	16	<i>14</i>
26 or Older	434	<i>638</i>	129	<i>263</i>	106	<i>213</i>
26-49	230	<i>376</i>	95	<i>178</i>	79	<i>157</i>
26-29	83	<i>126</i>	36	<i>64</i>	32	<i>51</i>
30-34	104	<i>192</i>	46	<i>57</i>	34	<i>54</i>
35-39	115	<i>179</i>	48	<i>101</i>	39	<i>78</i>
40-44	117	<i>170</i>	46	<i>79</i>	38	<i>76</i>
45-49	98	<i>187</i>	34	<i>89</i>	29	<i>84</i>
50 or Older	347	<i>523</i>	89	<i>194</i>	70	<i>143</i>
50-54	154	<i>200</i>	50	<i>98</i>	34	<i>70</i>
55-59	159	<i>235</i>	43	<i>125</i>	41	<i>66</i>
60-64	183	<i>351</i>	46	<i>53</i>	43	<i>48</i>
65 or Older	192	<i>242</i>	28	<i>95</i>	14	<i>89</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.12D Methamphetamine Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	0.16	<i>0.23</i>	0.05	<i>0.10</i>	0.04	<i>0.08</i>
12-20	0.07	<i>0.09</i>	0.05	<i>0.05</i>	0.03	<i>0.04</i>
21 or Older	0.19	<i>0.26</i>	0.06	<i>0.11</i>	0.05	<i>0.09</i>
12-17	0.06	<i>0.09</i>	0.04	<i>0.05</i>	0.02	<i>0.02</i>
12-13	0.07	<i>0.01</i>	0.06	<i>0.01</i>	0.03	<i>0.01</i>
14-15	0.06	<i>0.06</i>	0.05	<i>0.05</i>	0.04	<i>0.05</i>
16-17	0.14	<i>0.27</i>	0.08	<i>0.14</i>	0.02	<i>0.02</i>
18 or Older	0.18	<i>0.25</i>	0.05	<i>0.11</i>	0.04	<i>0.08</i>
18-25	0.16	<i>0.20</i>	0.09	<i>0.09</i>	0.06	<i>0.06</i>
18-20	0.18	<i>0.17</i>	0.12	<i>0.13</i>	0.08	<i>0.11</i>
21-25	0.24	<i>0.29</i>	0.13	<i>0.13</i>	0.08	<i>0.07</i>
26 or Older	0.20	<i>0.29</i>	0.06	<i>0.12</i>	0.05	<i>0.10</i>
26-49	0.23	<i>0.37</i>	0.09	<i>0.18</i>	0.08	<i>0.15</i>
26-29	0.45	<i>0.70</i>	0.20	<i>0.36</i>	0.18	<i>0.29</i>
30-34	0.45	<i>0.80</i>	0.20	<i>0.25</i>	0.15	<i>0.24</i>
35-39	0.52	<i>0.76</i>	0.22	<i>0.45</i>	0.18	<i>0.35</i>
40-44	0.58	<i>0.85</i>	0.24	<i>0.40</i>	0.20	<i>0.39</i>
45-49	0.47	<i>0.93</i>	0.17	<i>0.46</i>	0.14	<i>0.43</i>
50 or Older	0.30	<i>0.45</i>	0.08	<i>0.17</i>	0.06	<i>0.12</i>
50-54	0.70	<i>0.97</i>	0.24	<i>0.49</i>	0.16	<i>0.35</i>
55-59	0.72	<i>1.05</i>	0.20	<i>0.58</i>	0.19	<i>0.31</i>
60-64	0.82	<i>1.57</i>	0.22	<i>0.25</i>	0.21	<i>0.23</i>
65 or Older	0.36	<i>0.44</i>	0.05	<i>0.17</i>	0.03	<i>0.16</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.13C Any Use of Prescription Pain Relievers in Past Year and Misuse of Prescription Pain Relievers in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	790	<i>1,281</i>	289	<i>474</i>	163	<i>245</i>
12-20	166	<i>248</i>	60	<i>86</i>	31	<i>27</i>
21 or Older	1,035	<i>1,585</i>	292	<i>474</i>	163	<i>244</i>
12-17	94	<i>177</i>	39	<i>61</i>	23	<i>20</i>
12-13	49	<i>81</i>	16	<i>41</i>	9	<i>7</i>
14-15	65	<i>121</i>	20	<i>26</i>	11	<i>13</i>
16-17	69	<i>120</i>	29	<i>35</i>	18	<i>12</i>
18 or Older	784	<i>1,264</i>	285	<i>466</i>	162	<i>243</i>
18-25	142	<i>239</i>	75	<i>114</i>	35	<i>47</i>
18-20	123	<i>162</i>	46	<i>63</i>	22	<i>19</i>
21-25	149	<i>219</i>	68	<i>99</i>	29	<i>43</i>
26 or Older	770	<i>1,243</i>	276	<i>447</i>	161	<i>237</i>
26-49	387	<i>608</i>	176	<i>301</i>	102	<i>164</i>
26-29	188	<i>268</i>	82	<i>119</i>	52	<i>55</i>
30-34	206	<i>330</i>	92	<i>140</i>	49	<i>56</i>
35-39	202	<i>320</i>	88	<i>134</i>	48	<i>69</i>
40-44	216	<i>282</i>	76	<i>106</i>	36	<i>69</i>
45-49	214	<i>346</i>	71	<i>166</i>	40	<i>107</i>
50 or Older	648	<i>1,072</i>	206	<i>332</i>	116	<i>174</i>
50-54	328	<i>496</i>	109	<i>145</i>	67	<i>56</i>
55-59	337	<i>580</i>	99	<i>182</i>	52	<i>75</i>
60-64	346	<i>558</i>	74	<i>156</i>	52	<i>98</i>
65 or Older	458	<i>727</i>	123	<i>193</i>	59	<i>111</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.13D Any Use of Prescription Pain Relievers in Past Year and Misuse of Prescription Pain Relievers in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	0.29	<i>0.46</i>	0.11	<i>0.17</i>	0.06	<i>0.09</i>
12-20	0.34	<i>0.62</i>	0.15	<i>0.23</i>	0.08	<i>0.07</i>
21 or Older	0.33	<i>0.53</i>	0.12	<i>0.19</i>	0.07	<i>0.10</i>
12-17	0.38	<i>0.71</i>	0.16	<i>0.24</i>	0.09	<i>0.08</i>
12-13	0.57	<i>0.92</i>	0.20	<i>0.49</i>	0.12	<i>0.09</i>
14-15	0.69	<i>1.31</i>	0.23	<i>0.30</i>	0.13	<i>0.16</i>
16-17	0.72	<i>1.35</i>	0.34	<i>0.45</i>	0.21	<i>0.16</i>
18 or Older	0.31	<i>0.50</i>	0.11	<i>0.18</i>	0.06	<i>0.10</i>
18-25	0.42	<i>0.71</i>	0.22	<i>0.34</i>	0.10	<i>0.14</i>
18-20	0.69	<i>1.21</i>	0.33	<i>0.50</i>	0.16	<i>0.15</i>
21-25	0.54	<i>0.90</i>	0.31	<i>0.46</i>	0.14	<i>0.20</i>
26 or Older	0.36	<i>0.57</i>	0.13	<i>0.20</i>	0.07	<i>0.11</i>
26-49	0.38	<i>0.60</i>	0.17	<i>0.30</i>	0.10	<i>0.16</i>
26-29	0.87	<i>1.34</i>	0.44	<i>0.65</i>	0.28	<i>0.30</i>
30-34	0.79	<i>1.22</i>	0.40	<i>0.60</i>	0.22	<i>0.25</i>
35-39	0.80	<i>1.25</i>	0.40	<i>0.59</i>	0.22	<i>0.31</i>
40-44	0.92	<i>1.31</i>	0.38	<i>0.55</i>	0.19	<i>0.36</i>
45-49	0.90	<i>1.41</i>	0.34	<i>0.82</i>	0.20	<i>0.55</i>
50 or Older	0.56	<i>0.92</i>	0.18	<i>0.28</i>	0.10	<i>0.15</i>
50-54	1.29	<i>2.01</i>	0.52	<i>0.71</i>	0.32	<i>0.28</i>
55-59	1.25	<i>2.18</i>	0.46	<i>0.83</i>	0.24	<i>0.34</i>
60-64	1.30	<i>2.10</i>	0.36	<i>0.75</i>	0.25	<i>0.47</i>
65 or Older	0.87	<i>1.33</i>	0.23	<i>0.35</i>	0.11	<i>0.20</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.14C Any Use of Prescription Stimulants in Past Year and Misuse of Prescription Stimulants in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	387	603	178	292	99	177
12-20	124	183	70	94	34	26
21 or Older	392	594	163	276	94	174
12-17	67	138	37	50	23	19
12-13	30	73	7	12	2	3
14-15	44	75	20	31	14	8
16-17	50	86	31	33	19	18
18 or Older	379	588	173	288	96	177
18-25	134	198	93	121	48	42
18-20	96	112	58	79	25	17
21-25	113	166	72	97	41	36
26 or Older	344	541	138	255	85	168
26-49	229	372	122	224	75	154
26-29	118	188	72	103	38	55
30-34	121	204	68	141	40	95
35-39	108	157	50	105	35	80
40-44	95	150	44	70	31	43
45-49	95	141	29	72	17	68
50 or Older	250	420	64	158	40	66
50-54	127	242	29	123	4	65
55-59	129	185	41	75	22	4
60-64	101	152	34	44	33	13
65 or Older	165	264	20	43	*	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.14D Any Use of Prescription Stimulants in Past Year and Misuse of Prescription Stimulants in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	0.14	<i>0.22</i>	0.06	<i>0.11</i>	0.04	<i>0.06</i>
12-20	0.29	<i>0.47</i>	0.18	<i>0.25</i>	0.09	<i>0.07</i>
21 or Older	0.15	<i>0.24</i>	0.07	<i>0.12</i>	0.04	<i>0.07</i>
12-17	0.27	<i>0.55</i>	0.15	<i>0.20</i>	0.09	<i>0.08</i>
12-13	0.36	<i>0.83</i>	0.09	<i>0.14</i>	0.03	<i>0.03</i>
14-15	0.48	<i>0.85</i>	0.23	<i>0.35</i>	0.16	<i>0.09</i>
16-17	0.55	<i>1.03</i>	0.36	<i>0.42</i>	0.22	<i>0.22</i>
18 or Older	0.15	<i>0.23</i>	0.07	<i>0.11</i>	0.04	<i>0.07</i>
18-25	0.40	<i>0.59</i>	0.27	<i>0.36</i>	0.14	<i>0.12</i>
18-20	0.59	<i>0.87</i>	0.40	<i>0.61</i>	0.18	<i>0.14</i>
21-25	0.47	<i>0.73</i>	0.32	<i>0.45</i>	0.19	<i>0.17</i>
26 or Older	0.16	<i>0.25</i>	0.06	<i>0.12</i>	0.04	<i>0.08</i>
26-49	0.23	<i>0.37</i>	0.12	<i>0.22</i>	0.07	<i>0.15</i>
26-29	0.62	<i>1.00</i>	0.39	<i>0.57</i>	0.21	<i>0.31</i>
30-34	0.51	<i>0.85</i>	0.30	<i>0.61</i>	0.18	<i>0.41</i>
35-39	0.49	<i>0.67</i>	0.23	<i>0.46</i>	0.16	<i>0.36</i>
40-44	0.47	<i>0.75</i>	0.22	<i>0.36</i>	0.16	<i>0.22</i>
45-49	0.45	<i>0.70</i>	0.14	<i>0.37</i>	0.08	<i>0.35</i>
50 or Older	0.22	<i>0.36</i>	0.06	<i>0.13</i>	0.03	<i>0.06</i>
50-54	0.60	<i>1.15</i>	0.14	<i>0.61</i>	0.02	<i>0.32</i>
55-59	0.60	<i>0.85</i>	0.19	<i>0.35</i>	0.10	<i>0.02</i>
60-64	0.48	<i>0.72</i>	0.16	<i>0.21</i>	0.16	<i>0.06</i>
65 or Older	0.31	<i>0.48</i>	0.04	<i>0.08</i>	*	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.15C Any Use of Prescription Tranquilizers or Prescription Sedatives in Past Year and Misuse of Prescription Tranquilizers or Prescription Sedatives in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	656	961	238	336	137	226
12-20	92	141	57	80	29	31
21 or Older	791	1,058	234	333	135	223
12-17	57	77	35	40	21	11
12-13	25	36	8	17	5	1
14-15	34	48	19	24	13	7
16-17	42	47	29	27	17	9
18 or Older	650	958	233	336	136	225
18-25	112	194	67	113	36	59
18-20	67	115	42	69	20	28
21-25	106	171	58	89	31	51
26 or Older	635	936	219	318	129	218
26-49	303	479	133	239	89	138
26-29	131	201	65	104	43	52
30-34	147	225	75	103	53	55
35-39	150	233	56	111	32	77
40-44	158	251	54	97	36	73
45-49	145	254	45	101	22	59
50 or Older	554	809	167	216	88	168
50-54	221	365	79	95	50	51
55-59	255	389	91	139	45	120
60-64	264	386	77	89	38	55
65 or Older	406	559	76	127	43	120

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.15D Any Use of Prescription Tranquilizers or Prescription Sedatives in Past Year and Misuse of Prescription Tranquilizers or Prescription Sedatives in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	0.24	<i>0.35</i>	0.09	<i>0.12</i>	0.05	<i>0.08</i>
12-20	0.22	<i>0.37</i>	0.15	<i>0.21</i>	0.08	<i>0.08</i>
21 or Older	0.27	<i>0.40</i>	0.10	<i>0.14</i>	0.06	<i>0.09</i>
12-17	0.23	<i>0.31</i>	0.14	<i>0.16</i>	0.08	<i>0.04</i>
12-13	0.30	<i>0.43</i>	0.09	<i>0.20</i>	0.06	<i>0.02</i>
14-15	0.38	<i>0.56</i>	0.22	<i>0.28</i>	0.15	<i>0.08</i>
16-17	0.49	<i>0.59</i>	0.34	<i>0.35</i>	0.20	<i>0.11</i>
18 or Older	0.26	<i>0.38</i>	0.09	<i>0.13</i>	0.05	<i>0.09</i>
18-25	0.33	<i>0.58</i>	0.20	<i>0.34</i>	0.11	<i>0.17</i>
18-20	0.45	<i>0.89</i>	0.30	<i>0.55</i>	0.15	<i>0.23</i>
21-25	0.46	<i>0.74</i>	0.27	<i>0.41</i>	0.15	<i>0.24</i>
26 or Older	0.29	<i>0.43</i>	0.10	<i>0.15</i>	0.06	<i>0.10</i>
26-49	0.30	<i>0.47</i>	0.13	<i>0.24</i>	0.09	<i>0.14</i>
26-29	0.67	<i>1.05</i>	0.35	<i>0.57</i>	0.23	<i>0.29</i>
30-34	0.60	<i>0.91</i>	0.33	<i>0.45</i>	0.24	<i>0.24</i>
35-39	0.62	<i>0.96</i>	0.26	<i>0.49</i>	0.15	<i>0.34</i>
40-44	0.73	<i>1.17</i>	0.27	<i>0.50</i>	0.18	<i>0.37</i>
45-49	0.64	<i>1.16</i>	0.22	<i>0.52</i>	0.11	<i>0.30</i>
50 or Older	0.48	<i>0.69</i>	0.15	<i>0.18</i>	0.08	<i>0.14</i>
50-54	0.96	<i>1.68</i>	0.38	<i>0.47</i>	0.24	<i>0.25</i>
55-59	1.08	<i>1.66</i>	0.42	<i>0.64</i>	0.21	<i>0.55</i>
60-64	1.08	<i>1.71</i>	0.37	<i>0.43</i>	0.18	<i>0.26</i>
65 or Older	0.77	<i>1.02</i>	0.14	<i>0.23</i>	0.08	<i>0.22</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.16C Any Use of Prescription Tranquilizers in Past Year and Misuse of Prescription Tranquilizers in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	589	871	224	309	131	200
12-20	82	123	55	71	28	30
21 or Older	684	933	219	306	130	197
12-17	51	61	35	38	20	11
12-13	18	25	6	16	3	1
14-15	30	38	18	23	13	7
16-17	39	39	28	25	16	9
18 or Older	581	866	219	308	130	199
18-25	103	182	65	104	35	58
18-20	58	103	40	59	19	28
21-25	97	164	56	86	30	51
26 or Older	568	846	205	291	125	191
26-49	283	453	131	230	88	134
26-29	122	182	64	98	43	50
30-34	137	216	74	102	53	55
35-39	138	221	54	107	30	73
40-44	144	244	51	96	35	70
45-49	136	236	37	97	21	54
50 or Older	486	727	152	185	82	135
50-54	206	314	69	89	48	48
55-59	228	361	84	136	41	120
60-64	234	354	75	88	37	55
65 or Older	342	489	67	81	37	70

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.16D Any Use of Prescription Tranquilizers in Past Year and Misuse of Prescription Tranquilizers in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	0.21	<i>0.31</i>	0.08	<i>0.11</i>	0.05	<i>0.07</i>
12-20	0.20	<i>0.32</i>	0.14	<i>0.19</i>	0.07	<i>0.08</i>
21 or Older	0.25	<i>0.36</i>	0.09	<i>0.13</i>	0.05	<i>0.08</i>
12-17	0.21	<i>0.24</i>	0.14	<i>0.15</i>	0.08	<i>0.04</i>
12-13	0.22	<i>0.29</i>	0.07	<i>0.19</i>	0.04	<i>0.02</i>
14-15	0.34	<i>0.43</i>	0.21	<i>0.27</i>	0.15	<i>0.08</i>
16-17	0.46	<i>0.50</i>	0.34	<i>0.32</i>	0.20	<i>0.11</i>
18 or Older	0.23	<i>0.34</i>	0.09	<i>0.12</i>	0.05	<i>0.08</i>
18-25	0.30	<i>0.54</i>	0.19	<i>0.31</i>	0.10	<i>0.17</i>
18-20	0.41	<i>0.81</i>	0.29	<i>0.47</i>	0.14	<i>0.23</i>
21-25	0.43	<i>0.72</i>	0.26	<i>0.40</i>	0.14	<i>0.24</i>
26 or Older	0.26	<i>0.39</i>	0.09	<i>0.13</i>	0.06	<i>0.09</i>
26-49	0.28	<i>0.45</i>	0.13	<i>0.23</i>	0.09	<i>0.13</i>
26-29	0.63	<i>0.96</i>	0.35	<i>0.54</i>	0.23	<i>0.28</i>
30-34	0.56	<i>0.88</i>	0.32	<i>0.44</i>	0.23	<i>0.24</i>
35-39	0.59	<i>0.92</i>	0.25	<i>0.47</i>	0.14	<i>0.32</i>
40-44	0.68	<i>1.16</i>	0.26	<i>0.49</i>	0.18	<i>0.36</i>
45-49	0.61	<i>1.10</i>	0.18	<i>0.50</i>	0.10	<i>0.28</i>
50 or Older	0.42	<i>0.62</i>	0.13	<i>0.16</i>	0.07	<i>0.12</i>
50-54	0.92	<i>1.48</i>	0.33	<i>0.44</i>	0.24	<i>0.24</i>
55-59	0.99	<i>1.57</i>	0.39	<i>0.62</i>	0.19	<i>0.55</i>
60-64	1.00	<i>1.59</i>	0.36	<i>0.42</i>	0.18	<i>0.26</i>
65 or Older	0.65	<i>0.90</i>	0.13	<i>0.15</i>	0.07	<i>0.13</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.17C Any Use of Prescription Sedatives in Past Year and Misuse of Prescription Sedatives in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	427	618	102	161	52	123
12-20	51	79	22	46	15	20
21 or Older	457	633	99	154	50	121
12-17	37	51	17	14	13	1
12-13	19	27	5	2	4	*
14-15	24	32	13	8	11	*
16-17	23	29	10	12	6	1
18 or Older	427	614	100	161	51	123
18-25	59	108	24	57	13	32
18-20	35	64	14	43	7	20
21-25	50	87	21	37	11	24
26 or Older	423	607	98	152	49	119
26-49	177	272	55	77	27	49
26-29	61	117	24	37	7	17
30-34	78	118	27	17	20	5
35-39	81	125	25	43	13	30
40-44	85	120	22	34	8	24
45-49	89	163	26	33	8	24
50 or Older	375	532	81	131	41	108
50-54	139	234	45	50	26	38
55-59	155	214	43	67	22	26
60-64	171	211	34	22	6	4
65 or Older	279	385	40	99	22	98

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.17D Any Use of Prescription Sedatives in Past Year and Misuse of Prescription Sedatives in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	0.16	<i>0.22</i>	0.04	<i>0.06</i>	0.02	<i>0.04</i>
12-20	0.13	<i>0.21</i>	0.06	<i>0.12</i>	0.04	<i>0.05</i>
21 or Older	0.18	<i>0.26</i>	0.04	<i>0.06</i>	0.02	<i>0.05</i>
12-17	0.15	<i>0.21</i>	0.07	<i>0.06</i>	0.05	<i>0.01</i>
12-13	0.23	<i>0.32</i>	0.06	<i>0.02</i>	0.05	*
14-15	0.28	<i>0.37</i>	0.15	<i>0.09</i>	0.13	*
16-17	0.27	<i>0.37</i>	0.12	<i>0.15</i>	0.07	<i>0.02</i>
18 or Older	0.17	<i>0.24</i>	0.04	<i>0.06</i>	0.02	<i>0.05</i>
18-25	0.17	<i>0.32</i>	0.07	<i>0.17</i>	0.04	<i>0.09</i>
18-20	0.25	<i>0.51</i>	0.10	<i>0.35</i>	0.06	<i>0.16</i>
21-25	0.23	<i>0.39</i>	0.10	<i>0.17</i>	0.05	<i>0.11</i>
26 or Older	0.20	<i>0.28</i>	0.05	<i>0.07</i>	0.02	<i>0.05</i>
26-49	0.18	<i>0.27</i>	0.05	<i>0.08</i>	0.03	<i>0.05</i>
26-29	0.33	<i>0.65</i>	0.13	<i>0.20</i>	0.04	<i>0.10</i>
30-34	0.34	<i>0.50</i>	0.12	<i>0.08</i>	0.09	<i>0.02</i>
35-39	0.36	<i>0.55</i>	0.11	<i>0.19</i>	0.06	<i>0.13</i>
40-44	0.43	<i>0.62</i>	0.11	<i>0.18</i>	0.04	<i>0.13</i>
45-49	0.43	<i>0.81</i>	0.13	<i>0.17</i>	0.04	<i>0.12</i>
50 or Older	0.33	<i>0.46</i>	0.07	<i>0.11</i>	0.04	<i>0.09</i>
50-54	0.65	<i>1.11</i>	0.22	<i>0.25</i>	0.12	<i>0.19</i>
55-59	0.70	<i>0.97</i>	0.20	<i>0.31</i>	0.10	<i>0.12</i>
60-64	0.78	<i>0.99</i>	0.17	<i>0.10</i>	0.03	<i>0.02</i>
65 or Older	0.53	<i>0.71</i>	0.07	<i>0.18</i>	0.04	<i>0.18</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.18C Any Use and Misuse of Prescription Benzodiazepines in Past Year: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)
TOTAL	537	806	212	283
12-20	73	116	55	68
21 or Older	617	840	207	281
12-17	45	55	34	36
12-13	14	15	6	8
14-15	25	35	18	23
16-17	36	35	28	23
18 or Older	530	800	207	283
18-25	96	176	64	105
18-20	54	99	40	59
21-25	90	154	55	86
26 or Older	519	773	194	266
26-49	257	397	125	220
26-29	113	171	63	94
30-34	128	190	72	91
35-39	121	194	51	101
40-44	128	210	46	95
45-49	117	184	36	92
50 or Older	457	671	144	163
50-54	192	261	65	78
55-59	206	315	79	97
60-64	212	298	67	84
65 or Older	328	454	67	62

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.18D Any Use and Misuse of Prescription Benzodiazepines in Past Year: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year (2019)	Misuse in Past Year (2020)
TOTAL	0.20	<i>0.29</i>	0.08	<i>0.10</i>
12-20	0.18	<i>0.31</i>	0.14	<i>0.18</i>
21 or Older	0.22	<i>0.33</i>	0.09	<i>0.12</i>
12-17	0.18	<i>0.22</i>	0.14	<i>0.15</i>
12-13	0.17	<i>0.18</i>	0.07	<i>0.09</i>
14-15	0.29	<i>0.40</i>	0.22	<i>0.27</i>
16-17	0.43	<i>0.45</i>	0.34	<i>0.29</i>
18 or Older	0.21	<i>0.32</i>	0.08	<i>0.11</i>
18-25	0.28	<i>0.52</i>	0.19	<i>0.31</i>
18-20	0.39	<i>0.77</i>	0.29	<i>0.47</i>
21-25	0.40	<i>0.69</i>	0.26	<i>0.40</i>
26 or Older	0.24	<i>0.35</i>	0.09	<i>0.12</i>
26-49	0.25	<i>0.39</i>	0.12	<i>0.22</i>
26-29	0.59	<i>0.92</i>	0.35	<i>0.52</i>
30-34	0.54	<i>0.79</i>	0.32	<i>0.39</i>
35-39	0.52	<i>0.82</i>	0.24	<i>0.45</i>
40-44	0.61	<i>1.05</i>	0.23	<i>0.49</i>
45-49	0.54	<i>0.90</i>	0.18	<i>0.47</i>
50 or Older	0.40	<i>0.57</i>	0.12	<i>0.14</i>
50-54	0.87	<i>1.25</i>	0.31	<i>0.39</i>
55-59	0.90	<i>1.41</i>	0.37	<i>0.45</i>
60-64	0.92	<i>1.37</i>	0.32	<i>0.41</i>
65 or Older	0.62	<i>0.83</i>	0.13	<i>0.11</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.19C Misuse of Opioids in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	303	<i>478</i>	178	<i>278</i>
12-20	60	<i>86</i>	31	<i>27</i>
21 or Older	308	<i>478</i>	179	<i>277</i>
12-17	39	<i>61</i>	23	<i>20</i>
12-13	16	<i>41</i>	9	<i>7</i>
14-15	20	<i>26</i>	11	<i>13</i>
16-17	29	<i>35</i>	18	<i>12</i>
18 or Older	299	<i>470</i>	177	<i>276</i>
18-25	75	<i>115</i>	36	<i>47</i>
18-20	46	<i>63</i>	22	<i>19</i>
21-25	68	<i>99</i>	30	<i>44</i>
26 or Older	293	<i>452</i>	176	<i>267</i>
26-49	183	<i>304</i>	107	<i>170</i>
26-29	82	<i>122</i>	54	<i>62</i>
30-34	98	<i>142</i>	52	<i>59</i>
35-39	91	<i>137</i>	52	<i>70</i>
40-44	77	<i>106</i>	36	<i>70</i>
45-49	71	<i>168</i>	41	<i>107</i>
50 or Older	223	<i>333</i>	134	<i>205</i>
50-54	110	<i>145</i>	68	<i>75</i>
55-59	105	<i>182</i>	62	<i>119</i>
60-64	75	<i>156</i>	53	<i>99</i>
65 or Older	141	<i>195</i>	81	<i>113</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.19D Misuse of Opioids in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	0.11	<i>0.17</i>	0.06	<i>0.10</i>
12-20	0.15	<i>0.23</i>	0.08	<i>0.07</i>
21 or Older	0.13	<i>0.20</i>	0.07	<i>0.12</i>
12-17	0.16	<i>0.24</i>	0.09	<i>0.08</i>
12-13	0.20	<i>0.49</i>	0.12	<i>0.09</i>
14-15	0.23	<i>0.30</i>	0.13	<i>0.16</i>
16-17	0.34	<i>0.45</i>	0.21	<i>0.16</i>
18 or Older	0.12	<i>0.19</i>	0.07	<i>0.11</i>
18-25	0.22	<i>0.34</i>	0.11	<i>0.14</i>
18-20	0.33	<i>0.50</i>	0.16	<i>0.15</i>
21-25	0.31	<i>0.46</i>	0.15	<i>0.21</i>
26 or Older	0.14	<i>0.21</i>	0.08	<i>0.12</i>
26-49	0.18	<i>0.30</i>	0.11	<i>0.17</i>
26-29	0.44	<i>0.66</i>	0.30	<i>0.35</i>
30-34	0.43	<i>0.61</i>	0.23	<i>0.26</i>
35-39	0.41	<i>0.60</i>	0.24	<i>0.31</i>
40-44	0.38	<i>0.55</i>	0.19	<i>0.36</i>
45-49	0.34	<i>0.83</i>	0.20	<i>0.55</i>
50 or Older	0.19	<i>0.29</i>	0.12	<i>0.18</i>
50-54	0.53	<i>0.71</i>	0.33	<i>0.38</i>
55-59	0.49	<i>0.83</i>	0.29	<i>0.55</i>
60-64	0.36	<i>0.75</i>	0.26	<i>0.47</i>
65 or Older	0.27	<i>0.36</i>	0.15	<i>0.21</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.20C Misuse of Central Nervous System Stimulants in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	276	<i>459</i>	184	<i>316</i>
12-20	86	<i>109</i>	45	<i>41</i>
21 or Older	270	<i>446</i>	182	<i>313</i>
12-17	39	<i>55</i>	26	<i>22</i>
12-13	9	<i>13</i>	3	<i>3</i>
14-15	21	<i>34</i>	16	<i>9</i>
16-17	32	<i>37</i>	20	<i>19</i>
18 or Older	270	<i>454</i>	182	<i>314</i>
18-25	116	<i>159</i>	66	<i>81</i>
18-20	73	<i>89</i>	36	<i>35</i>
21-25	101	<i>140</i>	55	<i>73</i>
26 or Older	240	<i>417</i>	169	<i>296</i>
26-49	194	<i>332</i>	134	<i>236</i>
26-29	107	<i>170</i>	65	<i>95</i>
30-34	102	<i>154</i>	67	<i>108</i>
35-39	85	<i>156</i>	62	<i>117</i>
40-44	72	<i>127</i>	57	<i>107</i>
45-49	61	<i>130</i>	43	<i>100</i>
50 or Older	147	<i>265</i>	108	<i>178</i>
50-54	74	<i>169</i>	54	<i>114</i>
55-59	74	<i>159</i>	59	<i>86</i>
60-64	73	<i>82</i>	65	<i>50</i>
65 or Older	68	<i>97</i>	20	<i>90</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.20D Misuse of Central Nervous System Stimulants in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Misuse in Past Year (2019)	Misuse in Past Year (2020)	Misuse in Past Month (2019)	Misuse in Past Month (2020)
TOTAL	0.10	<i>0.17</i>	0.07	<i>0.11</i>
12-20	0.21	<i>0.29</i>	0.12	<i>0.11</i>
21 or Older	0.11	<i>0.18</i>	0.08	<i>0.13</i>
12-17	0.16	<i>0.22</i>	0.10	<i>0.09</i>
12-13	0.12	<i>0.15</i>	0.04	<i>0.04</i>
14-15	0.25	<i>0.39</i>	0.19	<i>0.10</i>
16-17	0.38	<i>0.47</i>	0.24	<i>0.25</i>
18 or Older	0.11	<i>0.18</i>	0.07	<i>0.12</i>
18-25	0.34	<i>0.48</i>	0.20	<i>0.24</i>
18-20	0.49	<i>0.68</i>	0.26	<i>0.28</i>
21-25	0.42	<i>0.64</i>	0.26	<i>0.34</i>
26 or Older	0.11	<i>0.19</i>	0.08	<i>0.14</i>
26-49	0.19	<i>0.33</i>	0.13	<i>0.23</i>
26-29	0.56	<i>0.92</i>	0.35	<i>0.53</i>
30-34	0.44	<i>0.66</i>	0.29	<i>0.47</i>
35-39	0.39	<i>0.68</i>	0.29	<i>0.51</i>
40-44	0.36	<i>0.64</i>	0.29	<i>0.55</i>
45-49	0.30	<i>0.66</i>	0.21	<i>0.51</i>
50 or Older	0.13	<i>0.23</i>	0.09	<i>0.15</i>
50-54	0.36	<i>0.84</i>	0.26	<i>0.57</i>
55-59	0.34	<i>0.73</i>	0.27	<i>0.40</i>
60-64	0.36	<i>0.39</i>	0.31	<i>0.24</i>
65 or Older	0.13	<i>0.18</i>	0.04	<i>0.16</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.21C Use of Illicit Drugs Other Than Marijuana in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	448	<i>680</i>	286	<i>466</i>
12-20	128	<i>186</i>	66	<i>90</i>
21 or Older	460	<i>682</i>	291	<i>457</i>
12-17	63	<i>117</i>	40	<i>38</i>
12-13	31	<i>64</i>	18	<i>13</i>
14-15	39	<i>66</i>	22	<i>21</i>
16-17	46	<i>69</i>	30	<i>26</i>
18 or Older	440	<i>667</i>	283	<i>462</i>
18-25	136	<i>230</i>	85	<i>129</i>
18-20	106	<i>141</i>	51	<i>79</i>
21-25	129	<i>197</i>	73	<i>103</i>
26 or Older	395	<i>619</i>	263	<i>429</i>
26-49	271	<i>432</i>	180	<i>309</i>
26-29	147	<i>220</i>	91	<i>136</i>
30-34	147	<i>243</i>	96	<i>132</i>
35-39	124	<i>200</i>	86	<i>137</i>
40-44	113	<i>153</i>	75	<i>125</i>
45-49	96	<i>205</i>	62	<i>156</i>
50 or Older	291	<i>450</i>	185	<i>297</i>
50-54	152	<i>243</i>	93	<i>144</i>
55-59	146	<i>256</i>	87	<i>186</i>
60-64	119	<i>181</i>	91	<i>113</i>
65 or Older	167	<i>235</i>	103	<i>170</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.21D Use of Illicit Drugs Other Than Marijuana in Past Year and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
TOTAL	0.16	<i>0.25</i>	0.10	<i>0.17</i>
12-20	0.29	<i>0.48</i>	0.17	<i>0.24</i>
21 or Older	0.18	<i>0.27</i>	0.12	<i>0.19</i>
12-17	0.25	<i>0.47</i>	0.16	<i>0.15</i>
12-13	0.36	<i>0.74</i>	0.22	<i>0.15</i>
14-15	0.43	<i>0.73</i>	0.25	<i>0.25</i>
16-17	0.52	<i>0.86</i>	0.35	<i>0.33</i>
18 or Older	0.18	<i>0.26</i>	0.11	<i>0.18</i>
18-25	0.40	<i>0.69</i>	0.25	<i>0.38</i>
18-20	0.63	<i>1.05</i>	0.36	<i>0.62</i>
21-25	0.51	<i>0.85</i>	0.33	<i>0.47</i>
26 or Older	0.18	<i>0.28</i>	0.12	<i>0.20</i>
26-49	0.27	<i>0.43</i>	0.18	<i>0.30</i>
26-29	0.72	<i>1.13</i>	0.48	<i>0.74</i>
30-34	0.60	<i>0.98</i>	0.42	<i>0.56</i>
35-39	0.54	<i>0.85</i>	0.39	<i>0.59</i>
40-44	0.54	<i>0.77</i>	0.38	<i>0.64</i>
45-49	0.46	<i>0.99</i>	0.30	<i>0.78</i>
50 or Older	0.25	<i>0.38</i>	0.16	<i>0.25</i>
50-54	0.71	<i>1.17</i>	0.45	<i>0.72</i>
55-59	0.66	<i>1.15</i>	0.40	<i>0.85</i>
60-64	0.57	<i>0.87</i>	0.44	<i>0.55</i>
65 or Older	0.32	<i>0.43</i>	0.20	<i>0.31</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.