

Table 1.75A Inhalant Use in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Numbers in Thousands, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	2,142	2,390	743	683	1,399	1,707	577	507	822	1,200
GEOGRAPHIC REGION										
Northeast	356	590	111	87	245	502	107	87	138	415
Midwest	412	492	153	166	260	326	108	93	151	233
South	729	726	292	259	437	468	188	147	248	321
West	645	582	187	171	458	410	173	179	285	231
COUNTY TYPE										
Large Metro	1,354	1,621	392	362	962	1,259	376	361	586	899
Small Metro	578	533	254	211	324	321	139	86	184	236
Nonmetro	211	236	97	110	114	126	62	60	52	66
Urbanized	103	104	40	50	63	54	34	33	29	21
Less Urbanized	81	118	45	48	37	70	20	25	17	45
Completely Rural	27	15	12	*	14	2	8	2	6	*
POVERTY LEVEL¹										
Less Than 100%	408	467	162	140	246	327	143	174	103	153
100-199%	376	357	147	96	229	261	108	112	121	149
200% or More	1,326	1,565	434	447	892	1,118	294	220	599	898
HEALTH INSURANCE²										
Private	1,342	1,251	418	394	923	857	388	273	535	583
Medicaid/CHIP	488	713	282	282	207	431	99	86	107	345
Other ³	248	165	36	66	212	98	46	15	166	83
No Coverage	162	381	25	12	137	368	64	135	73	233

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.75B Inhalant Use in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Percentages, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	0.8	<i>0.9</i>	3.0	<i>2.7</i>	0.6	<i>0.7</i>	1.7	<i>1.5</i>	0.4	<i>0.5</i>
GEOGRAPHIC REGION										
Northeast	0.7	<i>1.2</i>	2.8	<i>2.2</i>	0.6	<i>1.1</i>	1.9	<i>1.5</i>	0.4	<i>1.1</i>
Midwest	0.7	<i>0.9</i>	2.9	<i>3.1</i>	0.5	<i>0.6</i>	1.5	<i>1.3</i>	0.3	<i>0.5</i>
South	0.7	<i>0.7</i>	3.0	<i>2.6</i>	0.5	<i>0.5</i>	1.5	<i>1.2</i>	0.3	<i>0.4</i>
West	1.0	<i>0.9</i>	3.1	<i>2.8</i>	0.8	<i>0.7</i>	2.1	<i>2.2</i>	0.6	<i>0.4</i>
COUNTY TYPE										
Large Metro	0.9	<i>1.1</i>	2.9	<i>2.6</i>	0.7	<i>0.9</i>	2.0	<i>2.0</i>	0.5	<i>0.8</i>
Small Metro	0.7	<i>0.6</i>	3.3	<i>2.8</i>	0.4	<i>0.4</i>	1.3	<i>0.8</i>	0.3	<i>0.4</i>
Nonmetro	0.6	<i>0.6</i>	2.7	<i>3.4</i>	0.3	<i>0.3</i>	1.4	<i>1.3</i>	0.2	<i>0.2</i>
Urbanized	0.6	<i>0.6</i>	2.6	<i>3.8</i>	0.4	<i>0.4</i>	1.7	<i>1.5</i>	0.2	<i>0.2</i>
Less Urbanized	0.5	<i>0.6</i>	2.7	<i>3.0</i>	0.2	<i>0.4</i>	1.0	<i>1.1</i>	0.1	<i>0.3</i>
Completely Rural	0.6	<i>0.3</i>	3.0	<i>*</i>	0.4	<i>0.1</i>	2.2	<i>0.7</i>	0.2	<i>*</i>
POVERTY LEVEL¹										
Less Than 100%	1.1	<i>1.1</i>	3.2	<i>3.0</i>	0.8	<i>0.9</i>	1.8	<i>2.1</i>	0.4	<i>0.5</i>
100-199%	0.7	<i>0.7</i>	2.8	<i>2.1</i>	0.5	<i>0.6</i>	1.4	<i>1.5</i>	0.3	<i>0.4</i>
200% or More	0.7	<i>0.8</i>	3.0	<i>2.8</i>	0.5	<i>0.7</i>	1.7	<i>1.3</i>	0.4	<i>0.6</i>
HEALTH INSURANCE²										
Private	0.7	<i>0.7</i>	2.9	<i>2.6</i>	0.6	<i>0.5</i>	1.9	<i>1.4</i>	0.4	<i>0.4</i>
Medicaid/CHIP	1.1	<i>1.5</i>	3.0	<i>3.4</i>	0.6	<i>1.1</i>	1.4	<i>1.1</i>	0.4	<i>1.1</i>
Other ³	0.4	<i>0.2</i>	2.9	<i>4.0</i>	0.3	<i>0.1</i>	1.8	<i>0.6</i>	0.3	<i>0.1</i>
No Coverage	0.6	<i>1.3</i>	2.1	<i>0.8</i>	0.5	<i>1.4</i>	1.3	<i>2.6</i>	0.3	<i>1.1</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.76A Inhalant Use in Past Month: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Numbers in Thousands, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	807	<i>904</i>	216	<i>106</i>	591	<i>797</i>	157	<i>160</i>	434	<i>637</i>
GEOGRAPHIC REGION										
Northeast	151	<i>289</i>	37	<i>24</i>	114	<i>265</i>	32	<i>29</i>	82	<i>235</i>
Midwest	155	<i>129</i>	49	<i>28</i>	106	<i>101</i>	31	<i>14</i>	76	<i>87</i>
South	259	<i>294</i>	78	<i>43</i>	180	<i>251</i>	37	<i>77</i>	143	<i>174</i>
West	242	<i>192</i>	52	<i>11</i>	190	<i>181</i>	57	<i>40</i>	133	<i>141</i>
COUNTY TYPE										
Large Metro	592	<i>672</i>	125	<i>68</i>	467	<i>604</i>	111	<i>95</i>	356	<i>508</i>
Small Metro	162	<i>153</i>	70	<i>27</i>	92	<i>126</i>	36	<i>38</i>	56	<i>88</i>
Nonmetro	53	<i>80</i>	21	<i>12</i>	32	<i>68</i>	10	<i>26</i>	22	<i>41</i>
Urbanized	26	<i>33</i>	9	<i>5</i>	17	<i>28</i>	8	<i>*</i>	9	<i>10</i>
Less Urbanized	25	<i>39</i>	10	<i>1</i>	15	<i>38</i>	1	<i>7</i>	13	<i>31</i>
Completely Rural	2	<i>7</i>	2	<i>*</i>	<i>*</i>	<i>1</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>
POVERTY LEVEL¹										
Less Than 100%	106	<i>102</i>	40	<i>20</i>	66	<i>82</i>	38	<i>50</i>	29	<i>32</i>
100-199%	174	<i>163</i>	71	<i>10</i>	103	<i>153</i>	39	<i>42</i>	64	<i>111</i>
200% or More	522	<i>639</i>	106	<i>76</i>	417	<i>562</i>	76	<i>67</i>	341	<i>495</i>
HEALTH INSURANCE²										
Private	488	<i>512</i>	116	<i>73</i>	372	<i>439</i>	108	<i>94</i>	264	<i>345</i>
Medicaid/CHIP	174	<i>230</i>	90	<i>45</i>	84	<i>185</i>	25	<i>23</i>	59	<i>162</i>
Other ³	124	<i>65</i>	9	<i>*</i>	115	<i>65</i>	8	<i>1</i>	107	<i>64</i>
No Coverage	60	<i>142</i>	5	<i>*</i>	54	<i>142</i>	17	<i>43</i>	37	<i>100</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.76B Inhalant Use in Past Month: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Percentages, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	0.3	<i>0.3</i>	0.9	<i>0.4</i>	0.2	<i>0.3</i>	0.5	<i>0.5</i>	0.2	<i>0.3</i>
GEOGRAPHIC REGION										
Northeast	0.3	<i>0.6</i>	0.9	<i>0.6</i>	0.3	<i>0.6</i>	0.6	<i>0.5</i>	0.2	<i>0.6</i>
Midwest	0.3	<i>0.2</i>	0.9	<i>0.5</i>	0.2	<i>0.2</i>	0.4	<i>0.2</i>	0.2	<i>0.2</i>
South	0.2	<i>0.3</i>	0.8	<i>0.4</i>	0.2	<i>0.3</i>	0.3	<i>0.6</i>	0.2	<i>0.2</i>
West	0.4	<i>0.3</i>	0.9	<i>0.2</i>	0.3	<i>0.3</i>	0.7	<i>0.5</i>	0.3	<i>0.3</i>
COUNTY TYPE										
Large Metro	0.4	<i>0.4</i>	0.9	<i>0.5</i>	0.3	<i>0.4</i>	0.6	<i>0.5</i>	0.3	<i>0.4</i>
Small Metro	0.2	<i>0.2</i>	0.9	<i>0.3</i>	0.1	<i>0.2</i>	0.3	<i>0.3</i>	0.1	<i>0.1</i>
Nonmetro	0.1	<i>0.2</i>	0.6	<i>0.4</i>	0.1	<i>0.2</i>	0.2	<i>0.6</i>	0.1	<i>0.1</i>
Urbanized	0.2	<i>0.2</i>	0.6	<i>0.4</i>	0.1	<i>0.2</i>	0.4	*	0.1	<i>0.1</i>
Less Urbanized	0.1	<i>0.2</i>	0.6	<i>0.1</i>	0.1	<i>0.2</i>	0.1	<i>0.3</i>	0.1	<i>0.2</i>
Completely Rural	0.1	<i>0.1</i>	0.6	*	*	<i>0.0</i>	*	*	*	*
POVERTY LEVEL¹										
Less Than 100%	0.3	<i>0.2</i>	0.8	<i>0.4</i>	0.2	<i>0.2</i>	0.5	<i>0.6</i>	0.1	<i>0.1</i>
100-199%	0.3	<i>0.3</i>	1.3	<i>0.2</i>	0.2	<i>0.3</i>	0.5	<i>0.5</i>	0.2	<i>0.3</i>
200% or More	0.3	<i>0.3</i>	0.7	<i>0.5</i>	0.2	<i>0.3</i>	0.4	<i>0.4</i>	0.2	<i>0.3</i>
HEALTH INSURANCE²										
Private	0.3	<i>0.3</i>	0.8	<i>0.5</i>	0.2	<i>0.3</i>	0.5	<i>0.5</i>	0.2	<i>0.2</i>
Medicaid/CHIP	0.4	<i>0.5</i>	1.0	<i>0.5</i>	0.2	<i>0.5</i>	0.3	<i>0.3</i>	0.2	<i>0.5</i>
Other ³	0.2	<i>0.1</i>	0.7	*	0.2	<i>0.1</i>	0.3	<i>0.1</i>	0.2	<i>0.1</i>
No Coverage	0.2	<i>0.5</i>	0.5	*	0.2	<i>0.5</i>	0.4	<i>0.8</i>	0.2	<i>0.5</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.