

Table 1.75C Inhalant Use in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Standard Errors of Numbers in Thousands, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	120	<i>193</i>	40	<i>81</i>	111	<i>176</i>	50	<i>61</i>	96	<i>169</i>
GEOGRAPHIC REGION										
Northeast	46	<i>107</i>	14	<i>21</i>	42	<i>106</i>	20	<i>27</i>	36	<i>106</i>
Midwest	39	<i>82</i>	17	<i>37</i>	37	<i>73</i>	18	<i>23</i>	30	<i>69</i>
South	66	<i>100</i>	27	<i>52</i>	59	<i>89</i>	28	<i>36</i>	52	<i>82</i>
West	79	<i>92</i>	22	<i>44</i>	75	<i>82</i>	30	<i>36</i>	67	<i>73</i>
COUNTY TYPE										
Large Metro	107	<i>163</i>	33	<i>57</i>	99	<i>155</i>	46	<i>52</i>	87	<i>151</i>
Small Metro	50	<i>89</i>	23	<i>47</i>	45	<i>76</i>	20	<i>24</i>	39	<i>72</i>
Nonmetro	27	<i>49</i>	14	<i>38</i>	22	<i>31</i>	13	<i>21</i>	18	<i>23</i>
Urbanized	20	<i>34</i>	9	<i>28</i>	18	<i>20</i>	10	<i>18</i>	16	<i>9</i>
Less Urbanized	15	<i>35</i>	11	<i>24</i>	11	<i>24</i>	7	<i>11</i>	8	<i>22</i>
Completely Rural	8	<i>7</i>	4	<i>*</i>	7	<i>2</i>	5	<i>2</i>	5	<i>*</i>
POVERTY LEVEL¹										
Less Than 100%	48	<i>78</i>	19	<i>43</i>	43	<i>65</i>	28	<i>41</i>	33	<i>51</i>
100-199%	42	<i>58</i>	19	<i>27</i>	37	<i>52</i>	21	<i>29</i>	30	<i>43</i>
200% or More	105	<i>170</i>	34	<i>65</i>	97	<i>156</i>	32	<i>35</i>	90	<i>153</i>
HEALTH INSURANCE²										
Private	96	<i>140</i>	33	<i>61</i>	89	<i>122</i>	41	<i>44</i>	78	<i>115</i>
Medicaid/CHIP	44	<i>114</i>	24	<i>55</i>	37	<i>100</i>	20	<i>23</i>	31	<i>97</i>
Other ³	60	<i>41</i>	9	<i>30</i>	59	<i>28</i>	13	<i>7</i>	57	<i>27</i>
No Coverage	26	<i>74</i>	7	<i>7</i>	25	<i>74</i>	14	<i>36</i>	21	<i>64</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.75D Inhalant Use in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Standard Errors of Percentages, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	0.04	<i>0.07</i>	0.16	<i>0.32</i>	0.04	<i>0.07</i>	0.15	<i>0.18</i>	0.04	<i>0.08</i>
GEOGRAPHIC REGION										
Northeast	0.10	<i>0.22</i>	0.34	<i>0.54</i>	0.10	<i>0.24</i>	0.35	<i>0.47</i>	0.09	<i>0.28</i>
Midwest	0.07	<i>0.14</i>	0.32	<i>0.71</i>	0.07	<i>0.14</i>	0.26	<i>0.32</i>	0.07	<i>0.15</i>
South	0.06	<i>0.10</i>	0.28	<i>0.53</i>	0.06	<i>0.09</i>	0.22	<i>0.28</i>	0.06	<i>0.10</i>
West	0.12	<i>0.14</i>	0.36	<i>0.73</i>	0.13	<i>0.14</i>	0.37	<i>0.45</i>	0.13	<i>0.14</i>
COUNTY TYPE										
Large Metro	0.07	<i>0.11</i>	0.23	<i>0.39</i>	0.07	<i>0.11</i>	0.24	<i>0.29</i>	0.07	<i>0.13</i>
Small Metro	0.06	<i>0.11</i>	0.29	<i>0.61</i>	0.06	<i>0.10</i>	0.18	<i>0.22</i>	0.06	<i>0.11</i>
Nonmetro	0.07	<i>0.12</i>	0.38	<i>1.13</i>	0.07	<i>0.08</i>	0.29	<i>0.44</i>	0.06	<i>0.07</i>
Urbanized	0.13	<i>0.20</i>	0.55	<i>1.98</i>	0.13	<i>0.13</i>	0.46	<i>0.80</i>	0.13	<i>0.07</i>
Less Urbanized	0.08	<i>0.17</i>	0.60	<i>1.44</i>	0.07	<i>0.13</i>	0.34	<i>0.50</i>	0.06	<i>0.14</i>
Completely Rural	0.18	<i>0.14</i>	0.99	*	0.18	<i>0.04</i>	1.22	<i>0.50</i>	0.15	*
POVERTY LEVEL¹										
Less Than 100%	0.13	<i>0.19</i>	0.37	<i>0.90</i>	0.13	<i>0.18</i>	0.34	<i>0.50</i>	0.13	<i>0.18</i>
100-199%	0.08	<i>0.11</i>	0.35	<i>0.59</i>	0.08	<i>0.11</i>	0.27	<i>0.37</i>	0.08	<i>0.11</i>
200% or More	0.06	<i>0.09</i>	0.23	<i>0.40</i>	0.06	<i>0.09</i>	0.19	<i>0.21</i>	0.06	<i>0.10</i>
HEALTH INSURANCE²										
Private	0.05	<i>0.08</i>	0.23	<i>0.40</i>	0.05	<i>0.08</i>	0.19	<i>0.23</i>	0.05	<i>0.08</i>
Medicaid/CHIP	0.10	<i>0.23</i>	0.25	<i>0.65</i>	0.10	<i>0.25</i>	0.27	<i>0.29</i>	0.11	<i>0.30</i>
Other ³	0.09	<i>0.05</i>	0.73	<i>1.76</i>	0.09	<i>0.04</i>	0.52	<i>0.28</i>	0.09	<i>0.04</i>
No Coverage	0.10	<i>0.26</i>	0.62	<i>0.45</i>	0.10	<i>0.27</i>	0.29	<i>0.66</i>	0.10	<i>0.29</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.76C Inhalant Use in Past Month: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Standard Errors of Numbers in Thousands, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	84	<i>129</i>	24	<i>22</i>	80	<i>127</i>	26	<i>41</i>	75	<i>125</i>
GEOGRAPHIC REGION										
Northeast	31	<i>68</i>	9	<i>12</i>	29	<i>66</i>	11	<i>19</i>	27	<i>69</i>
Midwest	30	<i>53</i>	10	<i>9</i>	28	<i>52</i>	9	<i>8</i>	27	<i>52</i>
South	47	<i>73</i>	14	<i>16</i>	45	<i>71</i>	13	<i>29</i>	43	<i>66</i>
West	54	<i>66</i>	13	<i>6</i>	52	<i>66</i>	17	<i>19</i>	47	<i>63</i>
COUNTY TYPE										
Large Metro	79	<i>116</i>	20	<i>19</i>	77	<i>114</i>	23	<i>30</i>	71	<i>115</i>
Small Metro	28	<i>49</i>	13	<i>10</i>	24	<i>48</i>	11	<i>22</i>	22	<i>43</i>
Nonmetro	13	<i>29</i>	6	<i>6</i>	12	<i>27</i>	5	<i>17</i>	11	<i>21</i>
Urbanized	10	<i>18</i>	4	<i>4</i>	9	<i>17</i>	5	<i>*</i>	8	<i>6</i>
Less Urbanized	8	<i>22</i>	4	<i>1</i>	8	<i>22</i>	1	<i>6</i>	8	<i>21</i>
Completely Rural	2	<i>4</i>	2	<i>*</i>	<i>*</i>	<i>1</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>
POVERTY LEVEL¹										
Less Than 100%	22	<i>35</i>	10	<i>10</i>	20	<i>34</i>	14	<i>26</i>	15	<i>22</i>
100-199%	30	<i>44</i>	13	<i>4</i>	27	<i>44</i>	14	<i>19</i>	23	<i>39</i>
200% or More	76	<i>121</i>	18	<i>20</i>	74	<i>118</i>	17	<i>24</i>	70	<i>117</i>
HEALTH INSURANCE²										
Private	61	<i>101</i>	18	<i>20</i>	58	<i>97</i>	22	<i>30</i>	52	<i>94</i>
Medicaid/CHIP	29	<i>71</i>	15	<i>13</i>	25	<i>70</i>	10	<i>9</i>	21	<i>69</i>
Other ³	48	<i>24</i>	4	<i>*</i>	48	<i>24</i>	6	<i>1</i>	48	<i>24</i>
No Coverage	17	<i>49</i>	3	<i>*</i>	17	<i>49</i>	8	<i>26</i>	15	<i>41</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.76D Inhalant Use in Past Month: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Standard Errors of Percentages, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	0.03	<i>0.05</i>	0.10	<i>0.09</i>	0.03	<i>0.05</i>	0.08	<i>0.12</i>	0.03	<i>0.06</i>
GEOGRAPHIC REGION										
Northeast	0.06	<i>0.14</i>	0.23	<i>0.31</i>	0.07	<i>0.15</i>	0.20	<i>0.33</i>	0.07	<i>0.18</i>
Midwest	0.05	<i>0.09</i>	0.20	<i>0.17</i>	0.05	<i>0.10</i>	0.12	<i>0.11</i>	0.06	<i>0.12</i>
South	0.04	<i>0.07</i>	0.15	<i>0.16</i>	0.05	<i>0.07</i>	0.10	<i>0.23</i>	0.05	<i>0.08</i>
West	0.08	<i>0.10</i>	0.22	<i>0.09</i>	0.09	<i>0.11</i>	0.21	<i>0.24</i>	0.09	<i>0.12</i>
COUNTY TYPE										
Large Metro	0.05	<i>0.08</i>	0.14	<i>0.14</i>	0.05	<i>0.08</i>	0.12	<i>0.17</i>	0.06	<i>0.10</i>
Small Metro	0.03	<i>0.06</i>	0.16	<i>0.13</i>	0.03	<i>0.06</i>	0.11	<i>0.20</i>	0.03	<i>0.07</i>
Nonmetro	0.03	<i>0.07</i>	0.16	<i>0.19</i>	0.03	<i>0.07</i>	0.11	<i>0.36</i>	0.04	<i>0.06</i>
Urbanized	0.06	<i>0.11</i>	0.28	<i>0.33</i>	0.06	<i>0.11</i>	0.24	*	0.06	<i>0.04</i>
Less Urbanized	0.05	<i>0.11</i>	0.21	<i>0.06</i>	0.05	<i>0.12</i>	0.05	<i>0.27</i>	0.05	<i>0.13</i>
Completely Rural	0.04	<i>0.08</i>	0.39	*	*	<i>0.03</i>	*	*	*	*
POVERTY LEVEL¹										
Less Than 100%	0.06	<i>0.09</i>	0.19	<i>0.21</i>	0.06	<i>0.09</i>	0.17	<i>0.32</i>	0.06	<i>0.07</i>
100-199%	0.06	<i>0.09</i>	0.25	<i>0.10</i>	0.06	<i>0.09</i>	0.18	<i>0.25</i>	0.06	<i>0.10</i>
200% or More	0.04	<i>0.07</i>	0.12	<i>0.12</i>	0.04	<i>0.07</i>	0.10	<i>0.14</i>	0.05	<i>0.08</i>
HEALTH INSURANCE²										
Private	0.03	<i>0.06</i>	0.12	<i>0.13</i>	0.04	<i>0.06</i>	0.11	<i>0.15</i>	0.04	<i>0.07</i>
Medicaid/CHIP	0.06	<i>0.15</i>	0.16	<i>0.15</i>	0.07	<i>0.17</i>	0.14	<i>0.12</i>	0.08	<i>0.21</i>
Other ³	0.07	<i>0.03</i>	0.34	*	0.07	<i>0.03</i>	0.22	<i>0.03</i>	0.07	<i>0.03</i>
No Coverage	0.06	<i>0.17</i>	0.23	*	0.07	<i>0.18</i>	0.16	<i>0.49</i>	0.07	<i>0.19</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.