

Table 1.79A Any Use of Prescription Pain Relievers in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Numbers in Thousands, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	82,559	75,198	3,980	3,079	78,579	72,118	8,357	6,858	70,222	65,261
GEOGRAPHIC REGION										
Northeast	12,396	11,685	552	357	11,844	11,328	1,238	1,094	10,606	10,235
Midwest	17,550	15,012	795	666	16,755	14,347	1,822	1,316	14,933	13,030
South	32,479	30,506	1,612	1,403	30,867	29,103	3,209	2,801	27,658	26,302
West	20,134	17,994	1,021	653	19,112	17,341	2,087	1,647	17,025	15,693
COUNTY TYPE										
Large Metro	43,623	38,079	2,052	1,521	41,571	36,558	4,543	3,409	37,028	33,149
Small Metro	26,925	25,231	1,324	1,127	25,600	24,104	2,691	2,324	22,909	21,780
Nonmetro	12,011	11,888	604	431	11,408	11,456	1,123	1,124	10,285	10,332
Urbanized	5,037	4,966	255	178	4,783	4,789	558	550	4,225	4,239
Less Urbanized	5,560	5,433	275	193	5,285	5,240	467	538	4,817	4,702
Completely Rural	1,414	1,488	74	*	1,340	1,428	98	37	1,242	1,391
POVERTY LEVEL¹										
Less Than 100%	11,567	13,045	907	636	10,660	12,408	1,883	1,586	8,777	10,822
100-199%	16,296	14,097	882	545	15,414	13,552	1,944	1,701	13,470	11,851
200% or More	54,407	47,937	2,191	1,898	52,216	46,039	4,241	3,451	47,975	42,588
HEALTH INSURANCE²										
Private	52,152	44,978	2,161	1,729	49,991	43,249	4,988	3,956	45,003	39,294
Medicaid/CHIP	15,284	15,295	1,629	1,061	13,655	14,233	1,909	1,854	11,746	12,380
Other ³	26,242	26,624	245	186	25,998	26,438	625	554	25,373	25,884
No Coverage	6,636	6,925	143	285	6,494	6,640	1,189	929	5,305	5,711

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.79B Any Use of Prescription Pain Relievers in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Percentages, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	30.0	27.2	16.0	12.3	31.4	28.6	24.8	20.5	32.4	29.9
GEOGRAPHIC REGION										
Northeast	26.0	24.5	13.9	9.1	27.0	25.9	21.5	19.3	27.9	26.9
Midwest	30.6	26.2	15.1	12.6	32.2	27.5	25.4	18.6	33.3	28.9
South	31.1	28.8	16.7	14.4	32.5	30.3	25.2	22.0	33.6	31.6
West	30.7	27.2	17.0	10.8	32.1	28.9	25.8	20.5	33.1	30.2
COUNTY TYPE										
Large Metro	28.4	25.2	15.1	10.8	29.7	26.7	24.2	19.2	30.5	27.8
Small Metro	32.3	29.8	17.1	14.7	33.8	31.3	25.4	21.1	35.2	33.0
Nonmetro	31.6	28.8	16.8	13.4	33.1	30.1	25.9	23.6	34.1	31.0
Urbanized	31.7	30.1	16.6	13.4	33.3	31.5	27.5	25.0	34.3	32.7
Less Urbanized	31.2	27.6	16.6	12.0	32.7	29.0	24.2	24.3	33.8	29.6
Completely Rural	32.5	29.3	18.0	*	34.1	29.7	26.1	10.4	34.9	31.2
POVERTY LEVEL¹										
Less Than 100%	30.9	31.1	17.8	13.8	33.0	33.3	23.9	19.5	35.9	37.1
100-199%	30.8	27.9	16.5	12.1	32.4	29.5	25.1	22.1	33.8	31.0
200% or More	29.6	26.0	15.1	12.0	30.9	27.4	25.0	20.0	31.5	28.2
HEALTH INSURANCE²										
Private	29.0	25.9	15.2	11.5	30.2	27.2	24.4	20.2	31.0	28.2
Medicaid/CHIP	33.8	31.4	17.3	12.7	38.2	35.3	26.3	23.8	41.2	38.1
Other ³	37.4	34.9	20.0	11.2	37.7	35.4	24.6	22.8	38.2	35.8
No Coverage	24.5	24.3	12.3	19.5	25.1	24.6	24.7	17.6	25.2	26.3

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.80A Misuse of Prescription Pain Relievers in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Numbers in Thousands, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	9,724	<i>9,254</i>	567	<i>396</i>	9,158	<i>8,858</i>	1,754	<i>1,361</i>	7,404	<i>7,498</i>
GEOGRAPHIC REGION										
Northeast	1,352	<i>1,631</i>	57	<i>65</i>	1,295	<i>1,566</i>	234	<i>206</i>	1,062	<i>1,360</i>
Midwest	2,072	<i>1,686</i>	126	<i>84</i>	1,945	<i>1,602</i>	408	<i>313</i>	1,537	<i>1,290</i>
South	3,703	<i>3,307</i>	232	<i>126</i>	3,472	<i>3,181</i>	655	<i>501</i>	2,817	<i>2,680</i>
West	2,597	<i>2,630</i>	152	<i>121</i>	2,445	<i>2,509</i>	457	<i>341</i>	1,988	<i>2,167</i>
COUNTY TYPE										
Large Metro	5,240	<i>4,684</i>	286	<i>190</i>	4,954	<i>4,494</i>	969	<i>679</i>	3,985	<i>3,815</i>
Small Metro	3,326	<i>3,198</i>	215	<i>148</i>	3,110	<i>3,050</i>	579	<i>419</i>	2,532	<i>2,631</i>
Nonmetro	1,159	<i>1,372</i>	66	<i>58</i>	1,093	<i>1,314</i>	206	<i>262</i>	887	<i>1,051</i>
Urbanized	537	<i>584</i>	28	<i>24</i>	509	<i>560</i>	121	<i>120</i>	388	<i>440</i>
Less Urbanized	495	<i>754</i>	24	<i>33</i>	471	<i>721</i>	64	<i>132</i>	407	<i>589</i>
Completely Rural	127	<i>34</i>	14	<i>1</i>	113	<i>33</i>	22	<i>11</i>	92	<i>22</i>
POVERTY LEVEL¹										
Less Than 100%	1,926	<i>2,436</i>	129	<i>141</i>	1,797	<i>2,296</i>	434	<i>344</i>	1,363	<i>1,952</i>
100-199%	2,171	<i>1,775</i>	144	<i>55</i>	2,028	<i>1,721</i>	419	<i>336</i>	1,609	<i>1,385</i>
200% or More	5,597	<i>5,003</i>	294	<i>201</i>	5,303	<i>4,803</i>	871	<i>642</i>	4,432	<i>4,161</i>
HEALTH INSURANCE²										
Private	5,055	<i>4,578</i>	288	<i>170</i>	4,767	<i>4,408</i>	973	<i>748</i>	3,794	<i>3,660</i>
Medicaid/CHIP	2,429	<i>2,724</i>	240	<i>173</i>	2,189	<i>2,551</i>	401	<i>352</i>	1,788	<i>2,199</i>
Other ³	1,741	<i>1,927</i>	43	<i>20</i>	1,698	<i>1,906</i>	114	<i>106</i>	1,584	<i>1,800</i>
No Coverage	1,605	<i>1,494</i>	10	<i>*</i>	1,595	<i>1,437</i>	333	<i>233</i>	1,262	<i>1,203</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.80B Misuse of Prescription Pain Relievers in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Percentages, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	3.5	3.3	2.3	1.6	3.7	3.5	5.2	4.1	3.4	3.4
GEOGRAPHIC REGION										
Northeast	2.8	3.4	1.4	1.7	3.0	3.6	4.1	3.6	2.8	3.6
Midwest	3.6	2.9	2.4	1.6	3.7	3.1	5.7	4.4	3.4	2.9
South	3.5	3.1	2.4	1.3	3.7	3.3	5.2	3.9	3.4	3.2
West	4.0	4.0	2.5	2.0	4.1	4.2	5.7	4.3	3.9	4.2
COUNTY TYPE										
Large Metro	3.4	3.1	2.1	1.3	3.5	3.3	5.2	3.8	3.3	3.2
Small Metro	4.0	3.8	2.8	1.9	4.1	4.0	5.5	3.8	3.9	4.0
Nonmetro	3.0	3.3	1.8	1.8	3.2	3.4	4.8	5.5	2.9	3.2
Urbanized	3.4	3.5	1.8	1.8	3.5	3.7	6.0	5.4	3.1	3.4
Less Urbanized	2.8	3.8	1.5	2.0	2.9	4.0	3.3	6.0	2.9	3.7
Completely Rural	2.9	0.7	3.4	0.6	2.9	0.7	5.8	3.0	2.6	0.5
POVERTY LEVEL¹										
Less Than 100%	5.1	5.8	2.5	3.1	5.6	6.2	5.5	4.2	5.6	6.7
100-199%	4.1	3.5	2.7	1.2	4.3	3.7	5.4	4.4	4.0	3.6
200% or More	3.0	2.7	2.0	1.3	3.1	2.9	5.1	3.7	2.9	2.8
HEALTH INSURANCE²										
Private	2.8	2.6	2.0	1.1	2.9	2.8	4.8	3.8	2.6	2.6
Medicaid/CHIP	5.4	5.6	2.6	2.1	6.1	6.3	5.5	4.5	6.3	6.8
Other ³	2.5	2.5	3.5	1.2	2.5	2.6	4.5	4.4	2.4	2.5
No Coverage	5.9	5.3	0.8	*	6.2	5.3	6.9	4.4	6.0	5.5

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.81A Misuse of Prescription Pain Relievers in Past Month: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Numbers in Thousands, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	2,819	2,536	165	96	2,654	2,440	420	288	2,234	2,152
GEOGRAPHIC REGION										
Northeast	409	386	13	19	395	368	58	54	337	313
Midwest	626	459	44	27	581	432	95	69	486	362
South	1,045	877	63	36	982	841	154	111	828	730
West	739	814	44	14	695	800	112	54	583	746
COUNTY TYPE										
Large Metro	1,541	1,253	79	51	1,462	1,203	225	143	1,237	1,060
Small Metro	922	969	65	34	857	935	127	79	730	855
Nonmetro	355	313	21	11	334	303	67	66	267	237
Urbanized	133	118	7	*	125	113	38	*	87	100
Less Urbanized	181	195	8	6	172	190	24	53	148	137
Completely Rural	42	*	5	*	37	*	5	*	31	*
POVERTY LEVEL¹										
Less Than 100%	538	794	32	16	505	778	98	105	407	673
100-199%	584	479	41	12	543	467	101	58	442	409
200% or More	1,685	1,244	91	68	1,594	1,176	208	106	1,385	1,070
HEALTH INSURANCE²										
Private	1,354	988	95	50	1,259	938	221	135	1,038	803
Medicaid/CHIP	766	911	62	38	705	872	120	117	584	756
Other ³	474	562	7	7	467	555	12	16	455	539
No Coverage	565	459	2	2	563	457	80	41	484	416

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.81B Misuse of Prescription Pain Relievers in Past Month: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Percentages, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	1.0	<i>0.9</i>	0.7	<i>0.4</i>	1.1	<i>1.0</i>	1.2	<i>0.9</i>	1.0	<i>1.0</i>
GEOGRAPHIC REGION										
Northeast	0.9	<i>0.8</i>	0.3	<i>0.5</i>	0.9	<i>0.8</i>	1.0	<i>1.0</i>	0.9	<i>0.8</i>
Midwest	1.1	<i>0.8</i>	0.8	<i>0.5</i>	1.1	<i>0.8</i>	1.3	<i>1.0</i>	1.1	<i>0.8</i>
South	1.0	<i>0.8</i>	0.7	<i>0.4</i>	1.0	<i>0.9</i>	1.2	<i>0.9</i>	1.0	<i>0.9</i>
West	1.1	<i>1.2</i>	0.7	<i>0.2</i>	1.2	<i>1.3</i>	1.4	<i>0.7</i>	1.1	<i>1.4</i>
COUNTY TYPE										
Large Metro	1.0	<i>0.8</i>	0.6	<i>0.4</i>	1.0	<i>0.9</i>	1.2	<i>0.8</i>	1.0	<i>0.9</i>
Small Metro	1.1	<i>1.1</i>	0.8	<i>0.4</i>	1.1	<i>1.2</i>	1.2	<i>0.7</i>	1.1	<i>1.3</i>
Nonmetro	0.9	<i>0.8</i>	0.6	<i>0.3</i>	1.0	<i>0.8</i>	1.6	<i>1.4</i>	0.9	<i>0.7</i>
Urbanized	0.8	<i>0.7</i>	0.5	*	0.9	<i>0.7</i>	1.9	*	0.7	<i>0.8</i>
Less Urbanized	1.0	<i>1.0</i>	0.5	<i>0.3</i>	1.1	<i>1.0</i>	1.2	<i>2.4</i>	1.0	<i>0.9</i>
Completely Rural	1.0	*	1.3	*	0.9	*	1.4	*	0.9	*
POVERTY LEVEL¹										
Less Than 100%	1.4	<i>1.9</i>	0.6	<i>0.4</i>	1.6	<i>2.1</i>	1.2	<i>1.3</i>	1.7	<i>2.3</i>
100-199%	1.1	<i>0.9</i>	0.8	<i>0.3</i>	1.1	<i>1.0</i>	1.3	<i>0.8</i>	1.1	<i>1.1</i>
200% or More	0.9	<i>0.7</i>	0.6	<i>0.4</i>	0.9	<i>0.7</i>	1.2	<i>0.6</i>	0.9	<i>0.7</i>
HEALTH INSURANCE²										
Private	0.8	<i>0.6</i>	0.7	<i>0.3</i>	0.8	<i>0.6</i>	1.1	<i>0.7</i>	0.7	<i>0.6</i>
Medicaid/CHIP	1.7	<i>1.9</i>	0.7	<i>0.5</i>	2.0	<i>2.2</i>	1.7	<i>1.5</i>	2.0	<i>2.3</i>
Other ³	0.7	<i>0.7</i>	0.5	<i>0.4</i>	0.7	<i>0.7</i>	0.5	<i>0.7</i>	0.7	<i>0.7</i>
No Coverage	2.1	<i>1.6</i>	0.2	<i>0.1</i>	2.2	<i>1.7</i>	1.6	<i>0.8</i>	2.3	<i>1.9</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.