

**Table 1.98C Misuse of Central Nervous System Stimulants in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Standard Errors of Numbers in Thousands, 2019 and 2020**

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>TOTAL</b>	276	459	39	55	270	454	116	159	240	417
<b>GEOGRAPHIC REGION</b>										
Northeast	104	189	10	17	104	188	40	56	88	179
Midwest	116	165	17	17	113	164	43	67	101	143
South	183	287	27	34	178	283	82	96	146	263
West	158	264	19	37	156	258	66	89	143	239
<b>COUNTY TYPE</b>										
Large Metro	241	354	32	42	236	351	119	129	200	322
Small Metro	155	280	21	35	151	274	66	104	134	250
Nonmetro	83	122	9	12	82	120	32	41	75	111
Urbanized	64	81	7	7	63	81	26	23	59	77
Less Urbanized	52	91	4	10	52	89	18	34	46	80
Completely Rural	21	12	2	0	21	12	7	*	17	9
<b>POVERTY LEVEL<sup>1</sup></b>										
Less Than 100%	137	221	14	18	135	219	78	85	106	199
100-199%	128	192	15	11	128	193	47	75	121	177
200% or More	204	347	34	48	200	341	80	115	178	323
<b>HEALTH INSURANCE<sup>2</sup></b>										
Private	211	309	34	49	207	304	116	140	169	273
Medicaid/CHIP	150	233	21	30	148	231	45	51	141	226
Other <sup>3</sup>	112	183	8	5	112	183	26	31	109	180
No Coverage	108	184	6	6	107	184	34	63	103	170

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

<sup>1</sup> Respondents aged 18 to 22 who were living in a college dormitory were excluded.

<sup>2</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>3</sup> Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 1.98D Misuse of Central Nervous System Stimulants in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Standard Errors of Percentages, 2019 and 2020**

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>TOTAL</b>	0.10	<i>0.17</i>	0.16	<i>0.22</i>	0.11	<i>0.18</i>	0.34	<i>0.48</i>	0.11	<i>0.19</i>
<b>GEOGRAPHIC REGION</b>										
Northeast	0.22	<i>0.40</i>	0.26	<i>0.43</i>	0.24	<i>0.43</i>	0.69	<i>0.99</i>	0.23	<i>0.47</i>
Midwest	0.20	<i>0.29</i>	0.33	<i>0.33</i>	0.22	<i>0.31</i>	0.60	<i>0.94</i>	0.23	<i>0.32</i>
South	0.18	<i>0.27</i>	0.28	<i>0.34</i>	0.19	<i>0.29</i>	0.65	<i>0.75</i>	0.18	<i>0.32</i>
West	0.24	<i>0.40</i>	0.32	<i>0.61</i>	0.26	<i>0.43</i>	0.81	<i>1.11</i>	0.28	<i>0.46</i>
<b>COUNTY TYPE</b>										
Large Metro	0.15	<i>0.23</i>	0.23	<i>0.29</i>	0.16	<i>0.25</i>	0.53	<i>0.68</i>	0.16	<i>0.27</i>
Small Metro	0.17	<i>0.33</i>	0.27	<i>0.45</i>	0.19	<i>0.35</i>	0.53	<i>0.83</i>	0.20	<i>0.37</i>
Nonmetro	0.21	<i>0.29</i>	0.24	<i>0.38</i>	0.23	<i>0.31</i>	0.67	<i>0.84</i>	0.25	<i>0.33</i>
Urbanized	0.36	<i>0.49</i>	0.46	<i>0.54</i>	0.40	<i>0.53</i>	1.15	<i>0.96</i>	0.45	<i>0.60</i>
Less Urbanized	0.29	<i>0.44</i>	0.25	<i>0.62</i>	0.31	<i>0.47</i>	0.84	<i>1.53</i>	0.32	<i>0.49</i>
Completely Rural	0.43	<i>0.25</i>	0.54	<i>0.17</i>	0.47	<i>0.26</i>	1.67	*	0.44	<i>0.21</i>
<b>POVERTY LEVEL<sup>1</sup></b>										
Less Than 100%	0.35	<i>0.51</i>	0.28	<i>0.40</i>	0.40	<i>0.57</i>	0.80	<i>0.97</i>	0.43	<i>0.67</i>
100-199%	0.24	<i>0.37</i>	0.29	<i>0.24</i>	0.26	<i>0.41</i>	0.57	<i>0.93</i>	0.30	<i>0.46</i>
200% or More	0.11	<i>0.18</i>	0.22	<i>0.30</i>	0.11	<i>0.20</i>	0.44	<i>0.64</i>	0.11	<i>0.21</i>
<b>HEALTH INSURANCE<sup>2</sup></b>										
Private	0.11	<i>0.17</i>	0.23	<i>0.32</i>	0.12	<i>0.19</i>	0.47	<i>0.65</i>	0.11	<i>0.19</i>
Medicaid/CHIP	0.32	<i>0.47</i>	0.22	<i>0.35</i>	0.40	<i>0.56</i>	0.58	<i>0.64</i>	0.48	<i>0.69</i>
Other <sup>3</sup>	0.16	<i>0.24</i>	0.61	<i>0.29</i>	0.16	<i>0.24</i>	1.01	<i>1.23</i>	0.16	<i>0.25</i>
No Coverage	0.39	<i>0.64</i>	0.52	<i>0.43</i>	0.40	<i>0.67</i>	0.68	<i>1.17</i>	0.47	<i>0.77</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

<sup>1</sup> Respondents aged 18 to 22 who were living in a college dormitory were excluded.

<sup>2</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>3</sup> Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 1.99C Misuse of Central Nervous System Stimulants in Past Month: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Standard Errors of Numbers in Thousands, 2019 and 2020**

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>TOTAL</b>	184	<i>316</i>	26	<i>22</i>	182	<i>314</i>	66	<i>81</i>	169	<i>296</i>
<b>GEOGRAPHIC REGION</b>										
Northeast	67	<i>133</i>	6	<i>8</i>	67	<i>134</i>	26	<i>30</i>	62	<i>131</i>
Midwest	66	<i>113</i>	12	<i>10</i>	65	<i>113</i>	28	<i>28</i>	58	<i>108</i>
South	114	<i>184</i>	19	<i>14</i>	112	<i>182</i>	41	<i>54</i>	98	<i>169</i>
West	111	<i>192</i>	10	<i>10</i>	110	<i>191</i>	37	<i>46</i>	108	<i>179</i>
<b>COUNTY TYPE</b>										
Large Metro	166	<i>239</i>	19	<i>17</i>	165	<i>238</i>	60	<i>64</i>	149	<i>224</i>
Small Metro	85	<i>191</i>	16	<i>14</i>	84	<i>190</i>	33	<i>51</i>	78	<i>179</i>
Nonmetro	49	<i>88</i>	6	<i>5</i>	49	<i>87</i>	15	<i>17</i>	48	<i>85</i>
Urbanized	42	<i>65</i>	6	<i>4</i>	41	<i>64</i>	12	<i>5</i>	41	<i>64</i>
Less Urbanized	25	<i>59</i>	1	<i>2</i>	25	<i>58</i>	9	<i>16</i>	23	<i>56</i>
Completely Rural	12	<i>8</i>	1	<i>*</i>	12	<i>8</i>	3	<i>*</i>	11	<i>8</i>
<b>POVERTY LEVEL<sup>1</sup></b>										
Less Than 100%	97	<i>175</i>	10	<i>2</i>	96	<i>175</i>	40	<i>39</i>	87	<i>168</i>
100-199%	79	<i>117</i>	7	<i>1</i>	79	<i>117</i>	27	<i>35</i>	73	<i>112</i>
200% or More	138	<i>234</i>	23	<i>22</i>	137	<i>233</i>	45	<i>56</i>	127	<i>221</i>
<b>HEALTH INSURANCE<sup>2</sup></b>										
Private	128	<i>210</i>	22	<i>21</i>	127	<i>209</i>	60	<i>63</i>	114	<i>197</i>
Medicaid/CHIP	97	<i>177</i>	12	<i>7</i>	97	<i>177</i>	22	<i>19</i>	94	<i>174</i>
Other <sup>3</sup>	68	<i>109</i>	4	<i>2</i>	68	<i>109</i>	17	<i>18</i>	66	<i>107</i>
No Coverage	82	<i>131</i>	0	<i>2</i>	82	<i>131</i>	22	<i>37</i>	80	<i>124</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

<sup>1</sup> Respondents aged 18 to 22 who were living in a college dormitory were excluded.

<sup>2</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>3</sup> Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 1.99D Misuse of Central Nervous System Stimulants in Past Month: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Standard Errors of Percentages, 2019 and 2020**

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>TOTAL</b>	0.07	<i>0.11</i>	0.10	<i>0.09</i>	0.07	<i>0.12</i>	0.20	<i>0.24</i>	0.08	<i>0.14</i>
<b>GEOGRAPHIC REGION</b>										
Northeast	0.14	<i>0.28</i>	0.15	<i>0.20</i>	0.15	<i>0.31</i>	0.45	<i>0.53</i>	0.16	<i>0.34</i>
Midwest	0.12	<i>0.20</i>	0.23	<i>0.20</i>	0.12	<i>0.22</i>	0.39	<i>0.40</i>	0.13	<i>0.24</i>
South	0.11	<i>0.17</i>	0.19	<i>0.15</i>	0.12	<i>0.19</i>	0.32	<i>0.43</i>	0.12	<i>0.20</i>
West	0.17	<i>0.29</i>	0.17	<i>0.16</i>	0.19	<i>0.32</i>	0.46	<i>0.57</i>	0.21	<i>0.34</i>
<b>COUNTY TYPE</b>										
Large Metro	0.11	<i>0.16</i>	0.14	<i>0.12</i>	0.11	<i>0.17</i>	0.30	<i>0.35</i>	0.12	<i>0.19</i>
Small Metro	0.10	<i>0.22</i>	0.20	<i>0.19</i>	0.11	<i>0.24</i>	0.30	<i>0.44</i>	0.12	<i>0.27</i>
Nonmetro	0.13	<i>0.21</i>	0.16	<i>0.15</i>	0.14	<i>0.23</i>	0.33	<i>0.36</i>	0.16	<i>0.25</i>
Urbanized	0.25	<i>0.39</i>	0.35	<i>0.33</i>	0.28	<i>0.42</i>	0.54	<i>0.21</i>	0.32	<i>0.49</i>
Less Urbanized	0.14	<i>0.29</i>	0.07	<i>0.12</i>	0.15	<i>0.32</i>	0.45	<i>0.73</i>	0.16	<i>0.35</i>
Completely Rural	0.27	<i>0.16</i>	0.20	*	0.30	<i>0.17</i>	0.75	*	0.30	<i>0.18</i>
<b>POVERTY LEVEL<sup>1</sup></b>										
Less Than 100%	0.25	<i>0.41</i>	0.19	<i>0.05</i>	0.29	<i>0.45</i>	0.47	<i>0.47</i>	0.35	<i>0.56</i>
100-199%	0.15	<i>0.23</i>	0.13	<i>0.03</i>	0.17	<i>0.25</i>	0.34	<i>0.44</i>	0.18	<i>0.29</i>
200% or More	0.07	<i>0.13</i>	0.15	<i>0.14</i>	0.08	<i>0.14</i>	0.27	<i>0.33</i>	0.08	<i>0.14</i>
<b>HEALTH INSURANCE<sup>2</sup></b>										
Private	0.07	<i>0.12</i>	0.16	<i>0.14</i>	0.08	<i>0.13</i>	0.28	<i>0.32</i>	0.08	<i>0.14</i>
Medicaid/CHIP	0.21	<i>0.36</i>	0.13	<i>0.09</i>	0.27	<i>0.43</i>	0.30	<i>0.24</i>	0.32	<i>0.53</i>
Other <sup>3</sup>	0.10	<i>0.14</i>	0.36	<i>0.13</i>	0.10	<i>0.14</i>	0.66	<i>0.73</i>	0.10	<i>0.15</i>
No Coverage	0.30	<i>0.45</i>	0.03	<i>0.13</i>	0.31	<i>0.48</i>	0.45	<i>0.70</i>	0.37	<i>0.56</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

<sup>1</sup> Respondents aged 18 to 22 who were living in a college dormitory were excluded.

<sup>2</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>3</sup> Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.