

**Table 10.30A Had Serious Thoughts of Suicide in Past Year: Among People Aged 18 or Older; by Detailed Age Category, Numbers in Thousands, 2008-2020**

Age Category	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>TOTAL</b>	8,335	8,467	8,724	8,540	9,031	9,292	9,436	9,767	9,829	10,642	10,744	11,951	<i>12,197</i>
<b>18-25</b>	2,227	2,027	2,257	2,329	2,489	2,568	2,586	2,878	3,022	3,555	3,709	3,926	<i>3,771</i>
18-20	1,022	950	1,067	1,104	1,161	1,260	1,167	1,294	1,408	1,596	1,571	1,747	<i>1,552</i>
21-25	1,205	1,077	1,191	1,225	1,328	1,308	1,420	1,584	1,614	1,958	2,138	2,179	<i>2,219</i>
<b>26 or Older</b>	6,107	6,440	6,467	6,211	6,542	6,723	6,850	6,890	6,807	7,087	7,035	8,025	<i>8,427</i>
26-49	3,951	4,290	4,006	3,587	4,064	3,875	3,953	4,072	4,173	4,292	4,657	5,331	<i>5,317</i>
26-29	776	754	742	765	759	808	796	990	944	1,175	1,284	1,522	<i>1,279</i>
30-34	698	812	755	536	777	888	894	936	1,068	922	1,165	1,276	<i>1,422</i>
35-39	767	954	593	822	726	693	727	720	855	753	883	990	<i>932</i>
40-44	811	832	790	680	838	680	850	709	585	800	697	755	<i>832</i>
45-49	899	938	1,125	784	965	805	685	716	721	642	627	787	<i>852</i>
50 or Older	2,157	2,150	2,461	2,624	2,478	2,849	2,897	2,817	2,634	2,795	2,379	2,693	<i>3,110</i>
50-54	695	721	668	931	602	1,119	873	806	796	523	603	583	<i>905</i>
55-59	599	693	698	656	528	721	832	647	641	749	584	704	<i>712</i>
60-64	296	280	435	401	548	376	461	519	413	484	427	463	<i>431</i>
65 or Older	567	456	660	636	800	632	731	846	783	1,039	766	943	<i>1,061</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019, and Quarters 1 and 4, 2020.

**Table 10.30B Had Serious Thoughts of Suicide in Past Year: Among People Aged 18 or Older; by Detailed Age Category, Percentages, 2008-2020**

Age Category	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>TOTAL</b>	3.7	3.7	3.8	3.7	3.9	3.9	3.9	4.0	4.0	4.3	4.3	4.8	<i>4.9</i>
<b>18-25</b>	6.8	6.1	6.7	6.8	7.2	7.4	7.5	8.3	8.8	10.5	11.0	11.8	<i>11.3</i>
18-20	7.8	7.0	7.9	8.2	8.8	9.6	9.0	10.0	11.2	12.5	12.2	13.4	<i>12.5</i>
21-25	6.1	5.4	5.8	5.9	6.3	6.1	6.5	7.3	7.5	9.2	10.3	10.7	<i>10.6</i>
<b>26 or Older</b>	3.2	3.3	3.3	3.1	3.3	3.3	3.4	3.3	3.3	3.3	3.3	3.7	<i>3.9</i>
26-49	4.0	4.3	4.1	3.7	4.2	4.0	4.0	4.1	4.2	4.3	4.7	5.3	<i>5.3</i>
26-29	4.6	4.5	4.5	4.5	4.4	4.7	4.7	5.8	5.5	6.6	7.3	8.5	<i>7.2</i>
30-34	3.8	4.2	3.8	2.8	3.9	4.4	4.3	4.4	5.0	4.3	5.3	5.8	<i>6.3</i>
35-39	3.8	4.8	3.1	4.2	3.7	3.6	3.7	3.7	4.2	3.6	4.1	4.6	<i>4.2</i>
40-44	3.8	4.0	3.9	3.4	4.2	3.2	4.2	3.5	3.0	4.1	3.6	3.9	<i>4.3</i>
45-49	4.0	4.2	5.1	3.6	4.5	4.0	3.3	3.5	3.6	3.2	3.2	3.9	<i>4.4</i>
50 or Older	2.3	2.3	2.6	2.6	2.4	2.7	2.7	2.6	2.4	2.5	2.1	2.4	<i>2.7</i>
50-54	3.2	3.2	3.0	4.0	2.6	4.9	3.7	3.5	3.7	2.5	2.9	2.9	<i>4.6</i>
55-59	3.4	3.7	3.6	3.5	2.6	3.5	4.0	3.1	2.9	3.4	2.7	3.3	<i>3.3</i>
60-64	2.0	1.8	2.6	2.2	3.3	2.1	2.6	2.8	2.2	2.4	2.1	2.3	<i>2.1</i>
65 or Older	1.5	1.2	1.7	1.6	1.9	1.5	1.6	1.8	1.6	2.1	1.5	1.8	<i>2.0</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019, and Quarters 1 and 4, 2020.

**Table 10.33A Made Any Suicide Plans in Past Year: Among People Aged 18 or Older; by Detailed Age Category, Numbers in Thousands, 2008-2020**

Age Category	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>TOTAL</b>	2,299	2,253	2,542	2,406	2,681	2,714	2,667	2,702	2,756	3,195	3,312	3,455	3,222
<b>18-25</b>	643	654	661	663	813	852	806	934	981	1,266	1,160	1,313	<i>1,343</i>
18-20	347	335	324	320	433	434	388	468	469	647	523	608	<i>659</i>
21-25	296	320	338	343	380	418	419	466	512	618	637	705	<i>684</i>
<b>26 or Older</b>	1,656	1,598	1,880	1,743	1,868	1,862	1,860	1,768	1,775	1,930	2,153	2,142	<i>1,879</i>
26-49	1,056	1,016	1,013	1,076	1,272	1,249	1,107	1,033	1,271	1,230	1,450	1,466	<i>1,360</i>
26-29	200	158	147	277	232	255	211	272	253	291	442	412	<i>359</i>
30-34	209	152	225	124	157	274	212	242	344	247	316	361	<i>243</i>
35-39	221	213	142	264	267	250	252	171	282	192	314	229	<i>266</i>
40-44	202	265	266	202	288	245	221	169	186	244	200	234	<i>186</i>
45-49	224	227	233	208	329	226	211	179	206	256	180	231	<i>306</i>
50 or Older	600	583	867	667	596	614	754	734	504	700	702	676	<i>519</i>
50-54	259	212	298	295	157	163	159	295	185	168	210	123	<i>304</i>
55-59	142	169	173	195	113	204	192	151	104	166	222	204	<i>65</i>
60-64	87	100	108	70	143	90	161	94	116	133	116	133	<i>78</i>
65 or Older	111	101	288	107	184	157	242	194	99	233	153	216	<i>72</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents who had no suicide thoughts were categorized as not making any suicide plans in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019, and Quarters 1 and 4, 2020.

**Table 10.33B Made Any Suicide Plans in Past Year: Among People Aged 18 or Older; by Detailed Age Category, Percentages, 2008-2020**

Age Category	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>TOTAL</b>	1.0	1.0	1.1	1.0	1.1	1.1	1.1	1.1	1.1	1.3	1.3	1.4	<i>1.3</i>
<b>18-25</b>	2.0	2.0	1.9	1.9	2.4	2.5	2.3	2.7	2.9	3.7	3.4	3.9	<i>4.0</i>
18-20	2.7	2.5	2.4	2.4	3.3	3.3	3.0	3.6	3.7	5.1	4.1	4.7	<i>5.3</i>
21-25	1.5	1.6	1.7	1.7	1.8	1.9	1.9	2.1	2.4	2.9	3.1	3.5	<i>3.3</i>
<b>26 or Older</b>	0.9	0.8	1.0	0.9	0.9	0.9	0.9	0.9	0.9	0.9	1.0	1.0	<i>0.9</i>
26-49	1.1	1.0	1.0	1.1	1.3	1.3	1.1	1.1	1.3	1.2	1.4	1.5	<i>1.3</i>
26-29	1.2	0.9	0.9	1.6	1.4	1.5	1.2	1.6	1.5	1.6	2.5	2.3	<i>2.0</i>
30-34	1.1	0.8	1.1	0.6	0.8	1.4	1.0	1.1	1.6	1.1	1.4	1.6	<i>1.1</i>
35-39	1.1	1.1	0.7	1.3	1.4	1.3	1.3	0.9	1.4	0.9	1.4	1.1	<i>1.2</i>
40-44	0.9	1.3	1.3	1.0	1.4	1.1	1.1	0.8	1.0	1.2	1.0	1.2	<i>1.0</i>
45-49	1.0	1.0	1.1	1.0	1.5	1.1	1.0	0.9	1.0	1.3	0.9	1.1	<i>1.6</i>
50 or Older	0.7	0.6	0.9	0.7	0.6	0.6	0.7	0.7	0.5	0.6	0.6	0.6	<i>0.4</i>
50-54	1.2	1.0	1.4	1.3	0.7	0.7	0.7	1.3	0.8	0.8	1.0	0.6	<i>1.5</i>
55-59	0.8	0.9	0.9	1.0	0.5	1.0	0.9	0.7	0.5	0.8	1.0	1.0	<i>0.3</i>
60-64	0.6	0.6	0.7	0.4	0.9	0.5	0.9	0.5	0.6	0.7	0.6	0.7	<i>0.4</i>
65 or Older	0.3	0.3	0.7	0.3	0.4	0.4	0.5	0.4	0.2	0.5	0.3	0.4	<i>0.1</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents who had no suicide thoughts were categorized as not making any suicide plans in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019, and Quarters 1 and 4, 2020.

**Table 10.35A Attempted Suicide in Past Year: Among People Aged 18 or Older; by Detailed Age Category, Numbers in Thousands, 2008-2020**

Age Category	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>TOTAL</b>	1,088	1,040	1,122	1,135	1,290	1,312	1,120	1,399	1,319	1,388	1,442	1,379	<i>1,203</i>
<b>18-25</b>	395	372	393	415	508	434	433	571	616	648	647	604	<i>627</i>
18-20	242	190	205	221	282	255	190	274	334	364	316	300	<i>336</i>
21-25	153	182	189	194	226	180	244	298	281	284	332	305	<i>292</i>
<b>26 or Older</b>	693	668	729	721	782	878	687	827	704	739	795	775	<i>576</i>
26-49	446	465	399	443	502	596	491	506	496	435	521	579	<i>452</i>
26-29	106	83	44	119	100	98	99	115	131	114	105	156	<i>184</i>
30-34	117	81	105	73	85	158	91	131	114	83	121	154	<i>98</i>
35-39	58	121	48	102	87	126	79	77	114	88	157	91	<i>57</i>
40-44	61	62	86	93	126	135	100	107	60	52	79	105	<i>61</i>
45-49	103	118	115	55	105	80	123	76	77	97	59	74	<i>51</i>
50 or Older	247	203	329	278	279	283	196	321	208	304	274	195	<i>124</i>
50-54	151	51	62	136	56	43	62	136	47	105	83	31	<i>84</i>
55-59	42	50	60	112	53	120	62	67	65	58	118	58	<i>9</i>
60-64	29	35	47	2	57	59	17	46	25	29	14	13	<i>26</i>
65 or Older	25	66	160	29	114	60	54	72	71	113	58	94	<i>6</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents who had no suicide thoughts were categorized as not attempting suicide in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019, and Quarters 1 and 4, 2020.

**Table 10.35B Attempted Suicide in Past Year: Among People Aged 18 or Older; by Detailed Age Category, Percentages, 2008-2020**

Age Category	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>TOTAL</b>	0.5	0.5	0.5	0.5	0.6	0.6	0.5	0.6	0.5	0.6	0.6	0.6	<i>0.5</i>
<b>18-25</b>	1.2	1.1	1.2	1.2	1.5	1.3	1.2	1.6	1.8	1.9	1.9	1.8	<i>1.9</i>
18-20	1.8	1.4	1.5	1.6	2.1	1.9	1.5	2.1	2.7	2.9	2.4	2.3	<i>2.7</i>
21-25	0.8	0.9	0.9	0.9	1.1	0.8	1.1	1.4	1.3	1.3	1.6	1.5	<i>1.4</i>
<b>26 or Older</b>	0.4	0.3	0.4	0.4	0.4	0.4	0.3	0.4	0.3	0.3	0.4	0.4	<i>0.3</i>
26-49	0.4	0.5	0.4	0.5	0.5	0.6	0.5	0.5	0.5	0.4	0.5	0.6	<i>0.4</i>
26-29	0.6	0.5	0.3	0.7	0.6	0.6	0.6	0.7	0.8	0.6	0.6	0.9	<i>1.0</i>
30-34	0.6	0.4	0.5	0.4	0.4	0.8	0.4	0.6	0.5	0.4	0.6	0.7	<i>0.4</i>
35-39	0.3	0.6	0.2	0.5	0.4	0.7	0.4	0.4	0.6	0.4	0.7	0.4	<i>0.3</i>
40-44	0.3	0.3	0.4	0.5	0.6	0.6	0.5	0.5	0.3	0.3	0.4	0.5	<i>0.3</i>
45-49	0.5	0.5	0.5	0.3	0.5	0.4	0.6	0.4	0.4	0.5	0.3	0.4	<i>0.3</i>
50 or Older	0.3	0.2	0.3	0.3	0.3	0.3	0.2	0.3	0.2	0.3	0.2	0.2	<i>0.1</i>
50-54	0.7	0.2	0.3	0.6	0.2	0.2	0.3	0.6	0.2	0.5	0.4	0.2	<i>0.4</i>
55-59	0.2	0.3	0.3	0.6	0.3	0.6	0.3	0.3	0.3	0.3	0.5	0.3	<i>0.0</i>
60-64	0.2	0.2	0.3	0.0	0.3	0.3	0.1	0.3	0.1	0.1	0.1	0.1	<i>0.1</i>
65 or Older	0.1	0.2	0.4	0.1	0.3	0.1	0.1	0.2	0.1	0.2	0.1	0.2	<i>0.0</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents who had no suicide thoughts were categorized as not attempting suicide in the past 12 months. Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019, and Quarters 1 and 4, 2020.