

Table 11.3A Major Depressive Episode with Severe Impairment in Past Year: Among People Aged 12 to 17; by Demographic Characteristics, Numbers in Thousands, 2006-2020

Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
TOTAL	1,358	1,371	1,460	1,404	1,350	1,388	1,544	1,868	1,990	2,129	2,168	2,265	2,423	2,671	<i>2,919</i>
AGE GROUP															
12-13	211	200	239	235	232	218	285	314	375	388	354	332	382	530	<i>641</i>
14-15	518	500	505	521	479	487	590	752	707	826	789	861	891	970	<i>1,094</i>
16-17	629	671	716	648	639	683	669	801	909	915	1,025	1,072	1,150	1,171	<i>1,184</i>
GENDER															
Male	335	386	359	391	395	397	373	435	461	477	539	581	628	736	<i>732</i>
Female	1,023	986	1,101	1,013	954	991	1,172	1,432	1,529	1,652	1,629	1,684	1,795	1,935	<i>2,187</i>
HISPANIC ORIGIN AND RACE															
Not Hispanic or Latino	1,118	1,141	1,226	1,150	1,093	1,113	1,152	1,425	1,540	1,651	1,703	1,694	1,841	1,981	<i>2,230</i>
White	871	873	944	858	853	799	883	1,046	1,167	1,258	1,290	1,256	1,354	1,415	<i>1,616</i>
Black or African American	150	193	171	204	157	183	164	207	214	198	196	233	225	256	<i>294</i>
AIAN	9	4	8	5	7	13	4	6	8	*	8	6	*	19	*
NHOPI	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Asian	54	39	44	48	44	60	30	98	80	67	118	101	127	142	<i>186</i>
Two or More Races	32	32	50	31	30	57	64	61	66	98	86	89	109	134	<i>118</i>
Hispanic or Latino	240	230	233	254	257	275	392	442	450	477	465	571	582	690	<i>689</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2019, and Quarters 1 and 4, 2020.

Table 11.3B Major Depressive Episode with Severe Impairment in Past Year: Among People Aged 12 to 17; by Demographic Characteristics, Percentages, 2006-2020

Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
TOTAL	5.5	5.5	6.0	5.8	5.7	5.7	6.3	7.7	8.2	8.8	9.0	9.4	10.0	11.1	<i>12.0</i>
AGE GROUP															
12-13	2.7	2.5	3.2	3.2	3.0	2.8	3.7	4.1	4.9	5.1	4.7	4.4	5.1	6.8	<i>7.9</i>
14-15	6.0	6.0	6.1	6.2	6.1	5.9	7.1	9.1	8.5	9.8	9.4	10.4	10.9	11.9	<i>13.0</i>
16-17	7.5	7.9	8.4	7.7	7.7	8.1	8.0	9.7	10.9	11.1	12.4	12.7	13.7	14.5	<i>15.3</i>
GENDER															
Male	2.6	3.0	2.9	3.2	3.2	3.2	3.0	3.5	3.7	3.8	4.4	4.7	5.1	6.0	<i>5.9</i>
Female	8.4	8.2	9.3	8.6	8.2	8.3	9.8	12.0	13.0	14.0	13.7	14.2	15.2	16.5	<i>18.4</i>
HISPANIC ORIGIN AND RACE															
Not Hispanic or Latino	5.5	5.7	6.2	5.9	5.7	5.8	6.1	7.6	8.2	8.9	9.2	9.2	10.1	11.0	<i>12.3</i>
White	5.8	5.9	6.5	6.1	6.2	5.9	6.5	7.8	8.9	9.7	10.0	9.8	10.7	11.4	<i>13.0</i>
Black or African American	3.9	5.1	4.6	5.7	4.5	5.4	4.8	6.2	6.4	5.9	6.0	7.1	6.9	8.0	<i>9.7</i>
AIAN	6.6	2.6	6.5	4.3	5.4	9.8	2.6	3.8	4.9	*	5.7	3.9	*	11.5	*
NHOPI	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Asian	5.3	3.9	4.7	5.0	4.3	5.0	2.6	8.1	6.6	5.5	9.3	7.9	9.2	11.3	<i>10.3</i>
Two or More Races	8.0	7.8	10.2	6.0	5.9	8.1	9.0	8.4	8.9	12.6	10.9	10.9	14.2	14.9	<i>19.7</i>
Hispanic or Latino	5.4	5.1	5.1	5.4	5.4	5.2	7.3	8.2	8.2	8.5	8.2	9.9	10.0	11.7	<i>11.3</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2019, and Quarters 1 and 4, 2020.

Table 11.5A Receipt of Treatment for Depression in Past Year: Among People Aged 12 to 17 with Major Depressive Episode with Severe Impairment in Past Year; by Demographic Characteristics, Numbers in Thousands, 2006-2020

Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
TOTAL	627	601	622	542	554	602	633	832	882	945	1,001	1,074	1,133	1,321	<i>1,361</i>
AGE GROUP															
12-13	92	85	91	78	66	91	91	150	143	142	131	153	177	218	*
14-15	232	216	194	184	215	200	239	320	307	374	369	364	406	492	<i>521</i>
16-17	304	300	336	280	273	311	302	363	433	429	501	558	550	612	<i>527</i>
GENDER															
Male	134	154	132	132	132	147	127	157	189	206	208	208	274	300	<i>341</i>
Female	493	447	489	410	422	455	506	675	693	739	793	866	859	1,021	<i>1,020</i>
HISPANIC ORIGIN AND RACE															
Not Hispanic or Latino	535	534	550	443	448	502	506	645	725	755	827	868	882	1,018	<i>1,088</i>
White	432	439	461	352	380	379	403	504	572	563	668	679	695	800	<i>917</i>
Black or African American	53	72	65	58	42	82	56	72	99	97	71	97	90	108	*
AIAN	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
NHOPI	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Asian	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Two or More Races	*	*	*	*	*	*	*	*	*	*	*	*	*	63	*
Hispanic or Latino	92	67	72	99	106	100	127	187	158	190	174	206	251	304	<i>273</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown past year treatment for depression data were excluded.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2019, and Quarters 1 and 4, 2020.

Table 11.5B Receipt of Treatment for Depression in Past Year: Among People Aged 12 to 17 with Major Depressive Episode with Severe Impairment in Past Year; by Demographic Characteristics, Percentages, 2006-2020

Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
TOTAL	46.5	43.9	42.6	38.8	41.1	43.5	41.0	45.0	44.7	44.6	46.7	47.5	46.9	49.7	<i>46.9</i>
AGE GROUP															
12-13	44.1	42.3	38.3	33.8	28.7	42.3	32.0	48.5	38.7	37.0	37.6	46.2	47.2	41.9	*
14-15	44.8	43.2	38.6	35.6	45.0	41.0	40.5	43.2	43.4	45.4	47.5	42.3	45.6	50.8	<i>48.1</i>
16-17	48.8	44.9	46.9	43.2	42.7	45.5	45.4	45.3	48.1	47.1	49.2	52.0	47.8	52.3	<i>44.8</i>
GENDER															
Male	40.1	40.2	36.9	34.0	33.4	37.3	34.4	36.4	41.2	43.1	39.0	35.8	43.7	41.2	<i>46.6</i>
Female	48.6	45.4	44.5	40.7	44.3	45.9	43.2	47.6	45.7	45.1	49.3	51.5	48.1	52.9	<i>47.0</i>
HISPANIC ORIGIN AND RACE															
Not Hispanic or Latino	48.0	46.8	44.9	38.7	41.1	45.2	43.9	45.6	47.3	46.0	48.9	51.3	48.1	51.7	<i>49.2</i>
White	49.8	50.4	48.9	41.1	44.5	47.4	45.7	48.6	49.2	45.0	52.1	54.1	51.6	57.0	<i>57.3</i>
Black or African American	35.5	37.4	38.0	28.9	26.9	44.6	34.4	34.9	46.9	48.8	37.1	41.6	40.1	42.3	*
AIAN	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
NHOPI	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Asian	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Two or More Races	*	*	*	*	*	*	*	*	*	*	*	*	*	47.3	*
Hispanic or Latino	39.4	29.4	30.8	39.3	41.3	36.4	32.6	43.1	35.6	40.0	38.4	36.1	43.2	44.0	<i>39.6</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown past year treatment for depression data were excluded.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2019, and Quarters 1 and 4, 2020.