

**Table 13.11C Perceived COVID-19 Pandemic Effect on Access to Medical Care: Among People Aged 12 or Older; by Age Group, Standard Errors of Numbers in Thousands, Quarter 4, 2020**

<b>Perceived Effect on Access to Medical Care</b>	<b>Aged 12+</b>	<b>Aged 12-17</b>	<b>Aged 18+</b>	<b>Aged 18-25</b>	<b>Aged 26+</b>	<b>Aged 26-49</b>	<b>Aged 50+</b>
<b>PERCEIVED EFFECT AMONG TOTAL POPULATION</b>							
Appointments moved from in-person to telehealth	1,794	311	1,738	360	1,694	838	1,491
Delays or cancellations in appointments or preventative services	1,673	316	1,651	321	1,590	755	1,443
Delays in getting prescriptions	1,145	204	1,133	252	1,084	545	922
Unable to access needed care resulting in moderate to severe impact on health	877	83	872	194	838	509	670

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents could report that the questions about access to medical care did not apply to them. Respondents who reported that the respective question did not apply to them were classified as not having experienced that effect.

NOTE: Respondents with unknown information on their perception of the COVID-19 pandemic's effect on their medical care were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, Quarter 4, 2020.

**Table 13.11D Perceived COVID-19 Pandemic Effect on Access to Medical Care: Among People Aged 12 or Older; by Age Group, Standard Errors of Percentages, Quarter 4, 2020**

<b>Perceived Effect on Access to Medical Care</b>	<b>Aged 12+</b>	<b>Aged 12-17</b>	<b>Aged 18+</b>	<b>Aged 18-25</b>	<b>Aged 26+</b>	<b>Aged 26-49</b>	<b>Aged 50+</b>
<b>PERCEIVED EFFECT AMONG TOTAL POPULATION</b>							
Appointments moved from in-person to telehealth	0.66	1.28	0.71	1.10	0.80	0.85	1.30
Delays or cancellations in appointments or preventative services	0.62	1.31	0.67	0.98	0.75	0.77	1.26
Delays in getting prescriptions	0.42	0.84	0.46	0.77	0.51	0.55	0.80
Unable to access needed care resulting in moderate to severe impact on health	0.33	0.34	0.36	0.59	0.39	0.52	0.58

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents could report that the questions about access to medical care did not apply to them. Respondents who reported that the respective question did not apply to them were classified as not having experienced that effect.

NOTE: Respondents with unknown information on their perception of the COVID-19 pandemic's effect on their medical care were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, Quarter 4, 2020.