

**Table 2.3C Tobacco Product Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020**

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
<b>TOTAL</b>	894	<i>1,431</i>	723	<i>1,188</i>	686	<i>1,117</i>
12-20	201	<i>296</i>	163	<i>233</i>	117	<i>182</i>
21 or Older	1,640	<i>2,327</i>	873	<i>1,350</i>	787	<i>1,244</i>
<b>12-17</b>	83	<i>144</i>	67	<i>107</i>	48	<i>72</i>
12-13	26	<i>30</i>	22	<i>30</i>	14	<i>22</i>
14-15	56	<i>86</i>	42	<i>58</i>	26	<i>39</i>
16-17	69	<i>109</i>	57	<i>80</i>	39	<i>52</i>
<b>18 or Older</b>	853	<i>1,375</i>	713	<i>1,171</i>	678	<i>1,109</i>
<b>18-25</b>	180	<i>333</i>	171	<i>278</i>	153	<i>249</i>
18-20	176	<i>239</i>	144	<i>196</i>	106	<i>161</i>
21-25	254	<i>386</i>	195	<i>286</i>	157	<i>236</i>
<b>26 or Older</b>	818	<i>1,287</i>	692	<i>1,144</i>	658	<i>1,093</i>
26-49	416	<i>645</i>	410	<i>687</i>	391	<i>624</i>
26-29	307	<i>446</i>	229	<i>332</i>	198	<i>280</i>
30-34	340	<i>544</i>	231	<i>372</i>	204	<i>334</i>
35-39	346	<i>536</i>	228	<i>380</i>	206	<i>324</i>
40-44	318	<i>480</i>	203	<i>321</i>	187	<i>304</i>
45-49	310	<i>472</i>	202	<i>324</i>	181	<i>299</i>
50 or Older	677	<i>1,098</i>	533	<i>875</i>	512	<i>870</i>
50-54	487	<i>753</i>	315	<i>515</i>	287	<i>497</i>
55-59	509	<i>722</i>	323	<i>434</i>	301	<i>402</i>
60-64	492	<i>793</i>	270	<i>472</i>	250	<i>457</i>
65 or Older	500	<i>729</i>	314	<i>434</i>	290	<i>436</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Information about past year use of pipe tobacco was not collected. The estimate for tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 2.3D Tobacco Product Use in Lifetime, Past Year, and Past Month: Among People Aged 12 or Older; by Detailed Age Category, Standard Errors of Percentages, 2019 and 2020**

Age Category	Lifetime (2019)	Lifetime (2020)	Past Year (2019)	Past Year (2020)	Past Month (2019)	Past Month (2020)
<b>TOTAL</b>	0.32	<i>0.52</i>	0.26	<i>0.43</i>	0.25	<i>0.40</i>
12-20	0.40	<i>0.72</i>	0.35	<i>0.58</i>	0.27	<i>0.47</i>
21 or Older	0.35	<i>0.55</i>	0.30	<i>0.48</i>	0.28	<i>0.46</i>
<b>12-17</b>	0.33	<i>0.58</i>	0.27	<i>0.43</i>	0.19	<i>0.29</i>
12-13	0.31	<i>0.35</i>	0.26	<i>0.36</i>	0.16	<i>0.26</i>
14-15	0.59	<i>0.93</i>	0.47	<i>0.65</i>	0.30	<i>0.44</i>
16-17	0.71	<i>1.27</i>	0.62	<i>0.96</i>	0.45	<i>0.64</i>
<b>18 or Older</b>	0.34	<i>0.55</i>	0.28	<i>0.46</i>	0.27	<i>0.44</i>
<b>18-25</b>	0.53	<i>0.99</i>	0.51	<i>0.83</i>	0.45	<i>0.74</i>
18-20	0.83	<i>1.51</i>	0.75	<i>1.31</i>	0.63	<i>1.14</i>
21-25	0.67	<i>1.20</i>	0.65	<i>1.07</i>	0.59	<i>0.97</i>
<b>26 or Older</b>	0.38	<i>0.59</i>	0.32	<i>0.52</i>	0.30	<i>0.50</i>
26-49	0.41	<i>0.64</i>	0.40	<i>0.68</i>	0.39	<i>0.61</i>
26-29	0.95	<i>1.48</i>	0.96	<i>1.50</i>	0.91	<i>1.32</i>
30-34	0.83	<i>1.33</i>	0.83	<i>1.30</i>	0.77	<i>1.23</i>
35-39	0.86	<i>1.29</i>	0.87	<i>1.38</i>	0.83	<i>1.26</i>
40-44	0.88	<i>1.49</i>	0.89	<i>1.41</i>	0.86	<i>1.36</i>
45-49	0.91	<i>1.48</i>	0.87	<i>1.46</i>	0.79	<i>1.39</i>
50 or Older	0.59	<i>0.94</i>	0.46	<i>0.75</i>	0.44	<i>0.74</i>
50-54	1.38	<i>2.15</i>	1.34	<i>2.17</i>	1.25	<i>2.15</i>
55-59	1.26	<i>2.27</i>	1.28	<i>1.79</i>	1.22	<i>1.70</i>
60-64	1.22	<i>1.88</i>	1.12	<i>1.95</i>	1.08	<i>1.92</i>
65 or Older	0.95	<i>1.34</i>	0.59	<i>0.79</i>	0.55	<i>0.80</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Information about past year use of pipe tobacco was not collected. The estimate for tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.