

**Table 5.1C Substance Use Disorder for Specific Substances in Past Year: Among People Aged 12 or Older; by Age Group, Standard Errors of Numbers in Thousands, 2019 and 2020**

Substance Use Disorder	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>ILLICIT DRUGS</b>	nc	586	nc	104	nc	580	nc	210	nc	523
Marijuana	nc	493	nc	95	nc	487	nc	206	nc	426
Cocaine	nc	166	nc	14	nc	165	nc	44	nc	159
Heroin	nc	151	nc	*	nc	151	nc	16	nc	142
Hallucinogens	nc	59	nc	20	nc	54	nc	31	nc	45
Inhalants	nc	66	nc	35	nc	56	nc	19	nc	53
Methamphetamine	nc	192	nc	12	nc	192	nc	23	nc	191
Misuse of Prescription Psychotherapeutics	nc	309	nc	30	nc	309	nc	69	nc	296
Pain Relievers	nc	256	nc	24	nc	255	nc	48	nc	247
Stimulants	nc	112	nc	14	nc	111	nc	33	nc	106
Tranquilizers or Sedatives	nc	167	nc	23	nc	166	nc	44	nc	162
Tranquilizers	nc	135	nc	22	nc	133	nc	40	nc	128
Sedatives	nc	116	nc	5	nc	116	nc	20	nc	114
Benzodiazepines	--	--	--	--	--	--	--	--	--	--
Opioids	nc	268	nc	24	nc	267	nc	48	nc	259
Central Nervous System Stimulants	nc	265	nc	21	nc	264	nc	60	nc	258
Illicit Drugs Other Than Marijuana <sup>1</sup>	nc	390	nc	48	nc	389	nc	83	nc	376
<b>ALCOHOL</b>	nc	797	nc	86	nc	798	nc	241	nc	750
<b>BOTH ILLICIT DRUGS AND ALCOHOL</b>	nc	342	nc	56	nc	340	nc	140	nc	306
<b>ILLICIT DRUGS OR ALCOHOL</b>	nc	915	nc	123	nc	913	nc	279	nc	846
<b>ILLICIT DRUGS ONLY</b>	nc	495	nc	89	nc	487	nc	173	nc	446
<b>ALCOHOL ONLY</b>	nc	710	nc	66	nc	709	nc	180	nc	684

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

<sup>1</sup> Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 5.1D Substance Use Disorder for Specific Substances in Past Year: Among People Aged 12 or Older; by Age Group, Standard Errors of Percentages, 2019 and 2020**

Substance Use Disorder	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>ILLCIT DRUGS</b>	nc	0.21	nc	0.42	nc	0.23	nc	0.63	nc	0.24
Marijuana	nc	0.18	nc	0.38	nc	0.19	nc	0.61	nc	0.19
Cocaine	nc	0.06	nc	0.06	nc	0.07	nc	0.13	nc	0.07
Heroin	nc	0.05	nc	*	nc	0.06	nc	0.05	nc	0.06
Hallucinogens	nc	0.02	nc	0.08	nc	0.02	nc	0.09	nc	0.02
Inhalants	nc	0.02	nc	0.14	nc	0.02	nc	0.06	nc	0.02
Methamphetamine	nc	0.07	nc	0.05	nc	0.08	nc	0.07	nc	0.09
Misuse of Prescription Psychotherapeutics	nc	0.11	nc	0.12	nc	0.12	nc	0.21	nc	0.14
Pain Relievers	nc	0.09	nc	0.10	nc	0.10	nc	0.14	nc	0.11
Stimulants	nc	0.04	nc	0.05	nc	0.04	nc	0.10	nc	0.05
Tranquilizers or Sedatives	nc	0.06	nc	0.09	nc	0.07	nc	0.13	nc	0.07
Tranquilizers	nc	0.05	nc	0.09	nc	0.05	nc	0.12	nc	0.06
Sedatives	nc	0.04	nc	0.02	nc	0.05	nc	0.06	nc	0.05
Benzodiazepines	--	--	--	--	--	--	--	--	--	--
Opioids	nc	0.10	nc	0.10	nc	0.11	nc	0.14	nc	0.12
Central Nervous System Stimulants	nc	0.10	nc	0.09	nc	0.10	nc	0.18	nc	0.12
Illicit Drugs Other Than Marijuana <sup>1</sup>	nc	0.14	nc	0.19	nc	0.15	nc	0.25	nc	0.17
<b>ALCOHOL</b>	nc	0.29	nc	0.34	nc	0.32	nc	0.72	nc	0.34
<b>BOTH ILLICIT DRUGS AND ALCOHOL</b>	nc	0.12	nc	0.22	nc	0.13	nc	0.42	nc	0.14
<b>ILLCIT DRUGS OR ALCOHOL</b>	nc	0.33	nc	0.49	nc	0.36	nc	0.83	nc	0.39
<b>ILLCIT DRUGS ONLY</b>	nc	0.18	nc	0.36	nc	0.19	nc	0.52	nc	0.20
<b>ALCOHOL ONLY</b>	nc	0.26	nc	0.26	nc	0.28	nc	0.54	nc	0.31

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

<sup>1</sup> Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.