

Table 5.6C Illicit Drug Use Disorder in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Standard Errors of Numbers in Thousands, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	nc	586	nc	104	nc	580	nc	210	nc	523
GEOGRAPHIC REGION										
Northeast	nc	266	nc	36	nc	263	nc	82	nc	245
Midwest	nc	244	nc	39	nc	240	nc	87	nc	207
South	nc	368	nc	67	nc	362	nc	123	nc	331
West	nc	288	nc	61	nc	287	nc	118	nc	267
COUNTY TYPE										
Large Metro	nc	463	nc	81	nc	453	nc	186	nc	411
Small Metro	nc	360	nc	60	nc	355	nc	137	nc	314
Nonmetro	nc	240	nc	46	nc	234	nc	79	nc	217
Urbanized	nc	149	nc	22	nc	145	nc	58	nc	129
Less Urbanized	nc	188	nc	39	nc	183	nc	53	nc	170
Completely Rural	nc	23	nc	*	nc	23	nc	10	nc	19
POVERTY LEVEL¹										
Less Than 100%	nc	311	nc	48	nc	305	nc	124	nc	275
100-199%	nc	279	nc	56	nc	274	nc	130	nc	239
200% or More	nc	471	nc	83	nc	461	nc	151	nc	422
HEALTH INSURANCE²										
Private	nc	428	nc	80	nc	422	nc	184	nc	372
Medicaid/CHIP	nc	356	nc	71	nc	349	nc	104	nc	322
Other ³	nc	293	nc	19	nc	292	nc	79	nc	284
No Coverage	nc	225	nc	20	nc	223	nc	86	nc	198

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Illicit drug use disorder (IDUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). IDUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 5.6D Illicit Drug Use Disorder in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Standard Errors of Percentages, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	nc	0.21	nc	0.42	nc	0.23	nc	0.63	nc	0.24
GEOGRAPHIC REGION										
Northeast	nc	0.56	nc	0.91	nc	0.60	nc	1.44	nc	0.64
Midwest	nc	0.42	nc	0.74	nc	0.46	nc	1.23	nc	0.46
South	nc	0.35	nc	0.68	nc	0.38	nc	0.97	nc	0.40
West	nc	0.44	nc	1.02	nc	0.48	nc	1.48	nc	0.51
COUNTY TYPE										
Large Metro	nc	0.29	nc	0.54	nc	0.31	nc	0.90	nc	0.33
Small Metro	nc	0.41	nc	0.77	nc	0.44	nc	1.05	nc	0.46
Nonmetro	nc	0.54	nc	1.28	nc	0.58	nc	1.57	nc	0.62
Urbanized	nc	0.86	nc	1.54	nc	0.93	nc	2.56	nc	0.97
Less Urbanized	nc	0.85	nc	2.12	nc	0.92	nc	2.30	nc	0.99
Completely Rural	nc	0.46	nc	*	nc	0.48	nc	2.30	nc	0.44
POVERTY LEVEL¹										
Less Than 100%	nc	0.70	nc	1.01	nc	0.77	nc	1.36	nc	0.90
100-199%	nc	0.53	nc	1.20	nc	0.57	nc	1.49	nc	0.60
200% or More	nc	0.24	nc	0.50	nc	0.26	nc	0.81	nc	0.27
HEALTH INSURANCE²										
Private	nc	0.24	nc	0.50	nc	0.26	nc	0.81	nc	0.26
Medicaid/CHIP	nc	0.67	nc	0.83	nc	0.79	nc	1.22	nc	0.91
Other ³	nc	0.38	nc	1.17	nc	0.39	nc	2.91	nc	0.39
No Coverage	nc	0.75	nc	1.43	nc	0.78	nc	1.50	nc	0.87

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Illicit drug use disorder (IDUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). IDUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 5.7C Alcohol Use Disorder in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Standard Errors of Numbers in Thousands, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)	Aged 12-20 (2019)	Aged 12-20 (2020)	Aged 21+ (2019)	Aged 21+ (2020)
TOTAL	nc	797	nc	86	nc	798	nc	241	nc	750	nc	153	nc	821
GEOGRAPHIC REGION														
Northeast	nc	298	nc	36	nc	303	nc	96	nc	276	nc	56	nc	309
Midwest	nc	390	nc	32	nc	390	nc	109	nc	368	nc	67	nc	406
South	nc	407	nc	45	nc	409	nc	145	nc	383	nc	76	nc	430
West	nc	490	nc	53	nc	489	nc	115	nc	464	nc	87	nc	528
COUNTY TYPE														
Large Metro	nc	666	nc	75	nc	660	nc	194	nc	621	nc	113	nc	649
Small Metro	nc	457	nc	40	nc	459	nc	183	nc	411	nc	85	nc	453
Nonmetro	nc	286	nc	23	nc	286	nc	64	nc	274	nc	37	nc	283
Urbanized	nc	192	nc	15	nc	191	nc	39	nc	185	nc	22	nc	188
Less Urbanized	nc	211	nc	17	nc	214	nc	50	nc	198	nc	30	nc	212
Completely Rural	nc	79	nc	1	nc	79	nc	8	nc	78	nc	3	nc	79
POVERTY LEVEL¹														
Less Than 100%	nc	331	nc	49	nc	330	nc	152	nc	299	nc	97	nc	313
100-199%	nc	350	nc	24	nc	348	nc	121	nc	320	nc	40	nc	344
200% or More	nc	704	nc	69	nc	701	nc	165	nc	670	nc	94	nc	701
HEALTH INSURANCE²														
Private	nc	644	nc	67	nc	640	nc	235	nc	599	nc	114	nc	630
Medicaid/CHIP	nc	348	nc	37	nc	351	nc	86	nc	339	nc	57	nc	352
Other ³	nc	415	nc	20	nc	415	nc	65	nc	407	nc	44	nc	414
No Coverage	nc	329	nc	*	nc	326	nc	83	nc	303	nc	58	nc	323

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Alcohol use disorder (AUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). AUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 5.7D Alcohol Use Disorder in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Standard Errors of Percentages, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)	Aged 12-20 (2019)	Aged 12-20 (2020)	Aged 21+ (2019)	Aged 21+ (2020)
TOTAL	nc	0.29	nc	0.34	nc	0.32	nc	0.72	nc	0.34	nc	0.39	nc	0.33
GEOGRAPHIC REGION														
Northeast	nc	0.63	nc	0.92	nc	0.69	nc	1.69	nc	0.72	nc	0.92	nc	0.71
Midwest	nc	0.68	nc	0.60	nc	0.75	nc	1.54	nc	0.82	nc	0.79	nc	0.78
South	nc	0.38	nc	0.46	nc	0.43	nc	1.14	nc	0.46	nc	0.51	nc	0.45
West	nc	0.74	nc	0.87	nc	0.82	nc	1.43	nc	0.89	nc	0.92	nc	0.84
COUNTY TYPE														
Large Metro	nc	0.41	nc	0.51	nc	0.45	nc	0.95	nc	0.49	nc	0.52	nc	0.47
Small Metro	nc	0.52	nc	0.52	nc	0.57	nc	1.38	nc	0.60	nc	0.70	nc	0.59
Nonmetro	nc	0.63	nc	0.68	nc	0.69	nc	1.25	nc	0.76	nc	0.67	nc	0.72
Urbanized	nc	1.05	nc	1.07	nc	1.15	nc	1.71	nc	1.32	nc	1.00	nc	1.19
Less Urbanized	nc	0.96	nc	1.02	nc	1.04	nc	2.10	nc	1.12	nc	1.01	nc	1.10
Completely Rural	nc	1.43	nc	0.56	nc	1.52	nc	2.11	nc	1.63	nc	0.75	nc	1.56
POVERTY LEVEL¹														
Less Than 100%	nc	0.75	nc	1.04	nc	0.84	nc	1.57	nc	0.99	nc	1.18	nc	0.88
100-199%	nc	0.64	nc	0.53	nc	0.70	nc	1.38	nc	0.78	nc	0.55	nc	0.74
200% or More	nc	0.36	nc	0.42	nc	0.40	nc	0.87	nc	0.42	nc	0.41	nc	0.41
HEALTH INSURANCE²														
Private	nc	0.34	nc	0.43	nc	0.38	nc	1.01	nc	0.40	nc	0.50	nc	0.39
Medicaid/CHIP	nc	0.68	nc	0.43	nc	0.82	nc	1.08	nc	0.98	nc	0.46	nc	0.90
Other ³	nc	0.53	nc	1.19	nc	0.54	nc	2.47	nc	0.55	nc	1.64	nc	0.55
No Coverage	nc	1.04	nc	*	nc	1.09	nc	1.48	nc	1.27	nc	1.81	nc	1.14

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Alcohol use disorder (AUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). AUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 5.8C Substance Use Disorder in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Standard Errors of Numbers in Thousands, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	nc	915	nc	123	nc	913	nc	279	nc	846
GEOGRAPHIC REGION										
Northeast	nc	386	nc	42	nc	388	nc	111	nc	355
Midwest	nc	433	nc	45	nc	434	nc	124	nc	403
South	nc	502	nc	76	nc	499	nc	161	nc	471
West	nc	531	nc	73	nc	524	nc	146	nc	496
COUNTY TYPE										
Large Metro	nc	806	nc	100	nc	795	nc	255	nc	739
Small Metro	nc	555	nc	69	nc	549	nc	220	nc	494
Nonmetro	nc	359	nc	46	nc	353	nc	95	nc	330
Urbanized	nc	238	nc	23	nc	234	nc	66	nc	218
Less Urbanized	nc	270	nc	40	nc	267	nc	70	nc	243
Completely Rural	nc	85	nc	6	nc	82	nc	14	nc	80
POVERTY LEVEL¹										
Less Than 100%	nc	431	nc	64	nc	424	nc	180	nc	384
100-199%	nc	422	nc	58	nc	417	nc	158	nc	385
200% or More	nc	817	nc	98	nc	807	nc	208	nc	762
HEALTH INSURANCE²										
Private	nc	739	nc	92	nc	734	nc	283	nc	681
Medicaid/CHIP	nc	470	nc	73	nc	463	nc	124	nc	434
Other ³	nc	498	nc	27	nc	497	nc	89	nc	487
No Coverage	nc	386	nc	46	nc	379	nc	107	nc	348

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 5.8D Substance Use Disorder in Past Year: Among People Aged 12 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Standard Errors of Percentages, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
TOTAL	nc	0.33	nc	0.49	nc	0.36	nc	0.83	nc	0.39
GEOGRAPHIC REGION										
Northeast	nc	0.81	nc	1.08	nc	0.89	nc	1.97	nc	0.93
Midwest	nc	0.75	nc	0.86	nc	0.83	nc	1.75	nc	0.89
South	nc	0.47	nc	0.78	nc	0.52	nc	1.27	nc	0.57
West	nc	0.80	nc	1.21	nc	0.87	nc	1.82	nc	0.95
COUNTY TYPE										
Large Metro	nc	0.48	nc	0.67	nc	0.52	nc	1.15	nc	0.56
Small Metro	nc	0.60	nc	0.88	nc	0.66	nc	1.52	nc	0.69
Nonmetro	nc	0.76	nc	1.29	nc	0.82	nc	1.78	nc	0.89
Urbanized	nc	1.24	nc	1.57	nc	1.35	nc	2.81	nc	1.49
Less Urbanized	nc	1.16	nc	2.13	nc	1.26	nc	2.77	nc	1.34
Completely Rural	nc	1.54	nc	2.05	nc	1.57	nc	3.29	nc	1.67
POVERTY LEVEL¹										
Less Than 100%	nc	0.95	nc	1.34	nc	1.04	nc	1.75	nc	1.23
100-199%	nc	0.76	nc	1.24	nc	0.82	nc	1.65	nc	0.91
200% or More	nc	0.40	nc	0.58	nc	0.44	nc	1.05	nc	0.47
HEALTH INSURANCE²										
Private	nc	0.38	nc	0.57	nc	0.42	nc	1.12	nc	0.44
Medicaid/CHIP	nc	0.87	nc	0.86	nc	1.03	nc	1.44	nc	1.20
Other ³	nc	0.63	nc	1.63	nc	0.64	nc	3.19	nc	0.65
No Coverage	nc	1.19	nc	3.01	nc	1.23	nc	1.78	nc	1.44

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

¹ Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.