

Table 6.1A Drugs Used with Alcohol or within 2 Hours of Alcohol Use on Most Recent Use of Alcohol in Past Month: Among Past Month Alcohol Users Aged 12 or Older; by Age Group and Underage and Legal Drinking Age Groups, Numbers in Thousands, 2019 and 2020

Drug	AGE GROUP										UNDERAGE AND LEGAL DRINKING AGE GROUP			
	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)	Aged 12-20 (2019)	Aged 12-20 (2020)	Aged 21+ (2019)	Aged 21+ (2020)
Marijuana	11,886	<i>12,402</i>	432	<i>369</i>	11,454	<i>12,033</i>	3,178	<i>3,323</i>	8,276	<i>8,710</i>	1,441	<i>1,192</i>	10,445	<i>11,210</i>
Cocaine (Including Crack)	583	<i>740</i>	7	*	575	<i>732</i>	141	<i>142</i>	435	<i>590</i>	36	<i>21</i>	547	<i>719</i>
Heroin	98	<i>156</i>	*	*	98	<i>156</i>	3	<i>1</i>	94	<i>154</i>	*	*	98	<i>156</i>
Hallucinogens	146	<i>102</i>	3	*	143	<i>102</i>	34	<i>59</i>	109	<i>43</i>	12	<i>1</i>	134	<i>101</i>
Inhalants	45	<i>61</i>	7	*	38	<i>61</i>	13	<i>4</i>	25	<i>57</i>	7	<i>3</i>	38	<i>58</i>
Methamphetamine	372	<i>417</i>	1	*	372	<i>417</i>	10	<i>13</i>	362	<i>404</i>	7	*	366	<i>417</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates are based on nonimputed responses to multiple questions; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 6.1B Drugs Used with Alcohol or within 2 Hours of Alcohol Use on Most Recent Use of Alcohol in Past Month: Among Past Month Alcohol Users Aged 12 or Older; by Age Group and Underage and Legal Drinking Age Groups, Percentages, 2019 and 2020

Drug	AGE GROUP										UNDERAGE AND LEGAL DRINKING AGE GROUP			
	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)	Aged 12-20 (2019)	Aged 12-20 (2020)	Aged 21+ (2019)	Aged 21+ (2020)
Marijuana	8.6	<i>9.0</i>	19.3	<i>18.5</i>	8.4	<i>8.9</i>	17.6	<i>19.5</i>	7.0	<i>7.3</i>	21.1	<i>20.2</i>	7.9	<i>8.5</i>
Cocaine (Including Crack)	0.4	<i>0.5</i>	0.3	*	0.4	<i>0.5</i>	0.8	<i>0.8</i>	0.4	<i>0.5</i>	0.5	<i>0.3</i>	0.4	<i>0.5</i>
Heroin	0.1	<i>0.1</i>	*	*	0.1	<i>0.1</i>	0.0	<i>0.0</i>	0.1	<i>0.1</i>	*	*	0.1	<i>0.1</i>
Hallucinogens	0.1	<i>0.1</i>	0.1	*	0.1	<i>0.1</i>	0.2	<i>0.3</i>	0.1	<i>0.0</i>	0.2	<i>0.0</i>	0.1	<i>0.1</i>
Inhalants	0.0	<i>0.0</i>	0.3	*	0.0	<i>0.0</i>	0.1	<i>0.0</i>	0.0	<i>0.0</i>	0.1	<i>0.0</i>	0.0	<i>0.0</i>
Methamphetamine	0.3	<i>0.3</i>	0.0	*	0.3	<i>0.3</i>	0.1	<i>0.1</i>	0.3	<i>0.3</i>	0.1	*	0.3	<i>0.3</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates are based on nonimputed responses to multiple questions; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.