

Table 6.20C Types of Illicit Drug, Tobacco Product, and Alcohol Use in Past Month: Among Females Aged 15 to 44; by Pregnancy Status, Standard Errors of Numbers in Thousands, 2019 and 2020

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2019	2020	2019	2020	2019	2020
ILLCIT DRUGS	256	<i>412</i>	22	<i>45</i>	250	<i>399</i>
Marijuana	239	<i>394</i>	21	<i>45</i>	233	<i>381</i>
Cocaine	54	<i>87</i>	3	<i>6</i>	51	<i>86</i>
Crack	15	<i>24</i>	0	<i>*</i>	15	<i>24</i>
Heroin	28	<i>61</i>	*	<i>*</i>	28	<i>34</i>
Hallucinogens	56	<i>71</i>	0	<i>6</i>	54	<i>70</i>
LSD	27	<i>33</i>	0	<i>6</i>	27	<i>32</i>
PCP	*	<i>*</i>	0	<i>*</i>	*	<i>*</i>
Ecstasy	36	<i>39</i>	*	<i>*</i>	36	<i>39</i>
Inhalants	22	<i>48</i>	0	<i>*</i>	22	<i>48</i>
Methamphetamine	41	<i>105</i>	1	<i>*</i>	41	<i>92</i>
Misuse of Prescription Psychotherapeutics	108	<i>144</i>	6	<i>8</i>	108	<i>134</i>
Pain Relievers	75	<i>102</i>	5	<i>6</i>	75	<i>88</i>
Stimulants	62	<i>81</i>	5	<i>5</i>	62	<i>81</i>
Tranquilizers or Sedatives	66	<i>112</i>	5	<i>5</i>	66	<i>98</i>
Tranquilizers	65	<i>106</i>	5	<i>5</i>	65	<i>92</i>
Sedatives	18	<i>43</i>	*	<i>5</i>	18	<i>43</i>
Benzodiazepines	--	<i>--</i>	--	<i>--</i>	--	<i>--</i>
Opioids	77	<i>107</i>	5	<i>6</i>	77	<i>93</i>
Illicit Drugs Other Than Marijuana ²	131	<i>193</i>	6	<i>10</i>	130	<i>185</i>
TOBACCO PRODUCTS	278	<i>437</i>	27	<i>37</i>	271	<i>424</i>
Cigarettes	263	<i>410</i>	25	<i>37</i>	255	<i>401</i>
Daily Cigarettes	189	<i>288</i>	19	<i>27</i>	185	<i>278</i>
Smokeless Tobacco	47	<i>78</i>	8	<i>1</i>	47	<i>69</i>
Cigars	102	<i>182</i>	13	<i>3</i>	101	<i>180</i>
Pipe Tobacco	38	<i>71</i>	1	<i>2</i>	38	<i>70</i>
ALCOHOL	518	<i>802</i>	30	<i>46</i>	515	<i>785</i>
Binge Alcohol Use ³	362	<i>532</i>	21	<i>33</i>	358	<i>519</i>
Heavy Alcohol Use	149	<i>250</i>	4	<i>3</i>	147	<i>244</i>
GENERAL SUBSTANCES						
Illicit Drugs, Tobacco Products, or Alcohol	563	<i>864</i>	38	<i>64</i>	557	<i>844</i>
Illicit Drugs or Alcohol	539	<i>826</i>	34	<i>59</i>	535	<i>807</i>
Tobacco Products or Alcohol	547	<i>855</i>	37	<i>54</i>	541	<i>839</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ The Total column includes respondents with unknown pregnancy status.

² Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

³ Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 6.20D Types of Illicit Drug, Tobacco Product, and Alcohol Use in Past Month: Among Females Aged 15 to 44; by Pregnancy Status, Standard Errors of Percentages, 2019 and 2020

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2019	2020	2019	2020	2019	2020
ILLICIT DRUGS	0.34	<i>0.58</i>	1.04	<i>2.05</i>	0.35	<i>0.60</i>
Marijuana	0.32	<i>0.56</i>	1.01	<i>2.04</i>	0.33	<i>0.58</i>
Cocaine	0.08	<i>0.13</i>	0.13	<i>0.27</i>	0.08	<i>0.14</i>
Crack	0.02	<i>0.04</i>	0.01	*	0.02	<i>0.04</i>
Heroin	0.04	<i>0.09</i>	*	*	0.05	<i>0.06</i>
Hallucinogens	0.09	<i>0.11</i>	0.01	<i>0.27</i>	0.09	<i>0.11</i>
LSD	0.04	<i>0.05</i>	0.01	<i>0.27</i>	0.04	<i>0.05</i>
PCP	*	*	0.01	*	*	*
Ecstasy	0.06	<i>0.06</i>	*	*	0.06	<i>0.06</i>
Inhalants	0.03	<i>0.07</i>	0.01	*	0.04	<i>0.08</i>
Methamphetamine	0.06	<i>0.16</i>	0.03	*	0.07	<i>0.15</i>
Misuse of Prescription Psychotherapeutics	0.16	<i>0.22</i>	0.29	<i>0.39</i>	0.17	<i>0.21</i>
Pain Relievers	0.12	<i>0.16</i>	0.22	<i>0.30</i>	0.12	<i>0.14</i>
Stimulants	0.10	<i>0.12</i>	0.25	<i>0.26</i>	0.10	<i>0.13</i>
Tranquilizers or Sedatives	0.10	<i>0.17</i>	0.22	<i>0.24</i>	0.11	<i>0.16</i>
Tranquilizers	0.10	<i>0.16</i>	0.22	<i>0.24</i>	0.10	<i>0.15</i>
Sedatives	0.03	<i>0.07</i>	*	<i>0.24</i>	0.03	<i>0.07</i>
Benzodiazepines	--	--	--	--	--	--
Opioids	0.12	<i>0.16</i>	0.22	<i>0.30</i>	0.12	<i>0.15</i>
Illicit Drugs Other Than Marijuana ²	0.19	<i>0.29</i>	0.29	<i>0.47</i>	0.20	<i>0.29</i>
TOBACCO PRODUCTS	0.37	<i>0.59</i>	1.25	<i>1.82</i>	0.37	<i>0.61</i>
Cigarettes	0.36	<i>0.56</i>	1.16	<i>1.82</i>	0.36	<i>0.58</i>
Daily Cigarettes	0.28	<i>0.42</i>	0.91	<i>1.34</i>	0.28	<i>0.43</i>
Smokeless Tobacco	0.07	<i>0.12</i>	0.37	<i>0.03</i>	0.08	<i>0.11</i>
Cigars	0.16	<i>0.27</i>	0.63	<i>0.14</i>	0.16	<i>0.29</i>
Pipe Tobacco	0.06	<i>0.11</i>	0.03	<i>0.10</i>	0.06	<i>0.11</i>
ALCOHOL	0.48	<i>0.81</i>	1.40	<i>2.16</i>	0.49	<i>0.82</i>
Binge Alcohol Use ³	0.43	<i>0.66</i>	0.97	<i>1.55</i>	0.44	<i>0.68</i>
Heavy Alcohol Use	0.22	<i>0.36</i>	0.19	<i>0.14</i>	0.23	<i>0.38</i>
GENERAL SUBSTANCES						
Illicit Drugs, Tobacco Products, or Alcohol	0.46	<i>0.81</i>	1.70	<i>2.77</i>	0.47	<i>0.82</i>
Illicit Drugs or Alcohol	0.47	<i>0.80</i>	1.55	<i>2.58</i>	0.49	<i>0.81</i>
Tobacco Products or Alcohol	0.46	<i>0.81</i>	1.65	<i>2.52</i>	0.47	<i>0.82</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ The Total column includes respondents with unknown pregnancy status.

² Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

³ Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.