

**Table 6.6C Source Where Friend or Relative Obtained Prescription Pain Relievers: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Pain Relievers from a Friend or Relative for Free in Past Year; by Age Group, Standard Errors of Numbers in Thousands, 2019 and 2020**

Source Where Friend or Relative Obtained Pain Relievers	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	153	<i>244</i>	*	*	152	<i>232</i>	39	<i>62</i>	148	<i>225</i>
Prescription from One Doctor	151	<i>236</i>	*	*	150	<i>224</i>	39	<i>62</i>	147	<i>217</i>
Prescriptions from More Than One Doctor	20	<i>15</i>	*	*	20	<i>15</i>	7	*	19	<i>14</i>
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	8	*	*	*	8	*	3	*	7	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	48	<i>86</i>	*	*	48	<i>86</i>	13	<i>21</i>	47	*
From Friend or Relative for Free	38	<i>80</i>	*	*	38	<i>80</i>	12	*	37	*
Bought from Friend or Relative	24	<i>35</i>	*	*	24	<i>34</i>	5	*	23	*
Took from Friend or Relative without Asking	19	<i>12</i>	*	*	18	<i>11</i>	4	*	17	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	24	<i>31</i>	*	*	23	<i>30</i>	6	<i>6</i>	22	<i>29</i>
<b>SOME OTHER WAY<sup>1</sup></b>	16	<i>29</i>	*	*	15	<i>29</i>	5	<i>5</i>	14	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 6.6D Source Where Friend or Relative Obtained Prescription Pain Relievers: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Pain Relievers from a Friend or Relative for Free in Past Year; by Age Group, Standard Errors of Percentages, 2019 and 2020**

Source Where Friend or Relative Obtained Pain Relievers	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	1.66	<i>3.08</i>	*	*	1.68	<i>3.20</i>	2.56	<i>4.85</i>	2.02	<i>3.75</i>
Prescription from One Doctor	1.76	<i>3.51</i>	*	*	1.78	<i>3.63</i>	2.79	<i>4.90</i>	2.14	<i>4.28</i>
Prescriptions from More Than One Doctor	0.63	<i>0.51</i>	*	*	0.65	<i>0.53</i>	1.15	*	0.75	<i>0.62</i>
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.25	*	*	*	0.26	*	0.51	*	0.29	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	1.44	<i>2.82</i>	*	*	1.47	<i>2.93</i>	2.16	<i>4.51</i>	1.78	*
From Friend or Relative for Free	1.18	<i>2.67</i>	*	*	1.22	<i>2.78</i>	2.00	*	1.44	*
Bought from Friend or Relative	0.74	<i>1.19</i>	*	*	0.76	<i>1.24</i>	0.90	*	0.91	*
Took from Friend or Relative without Asking	0.59	<i>0.41</i>	*	*	0.57	<i>0.42</i>	0.79	*	0.68	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.75	<i>1.06</i>	*	*	0.73	<i>1.09</i>	1.03	<i>1.40</i>	0.86	<i>1.28</i>
<b>SOME OTHER WAY<sup>1</sup></b>	0.51	<i>1.02</i>	*	*	0.49	<i>1.07</i>	0.96	<i>1.14</i>	0.56	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 6.8C Source Where Friend or Relative Obtained Prescription Stimulants: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Stimulants from a Friend or Relative for Free in Past Year; by Age Group, Standard Errors of Numbers in Thousands, 2019 and 2020**

Source Where Friend or Relative Obtained Stimulants	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	111	<i>182</i>	*	*	108	<i>178</i>	53	<i>73</i>	89	<i>162</i>
Prescription from One Doctor	111	<i>178</i>	*	*	108	<i>177</i>	53	<i>72</i>	89	<i>160</i>
Prescriptions from More Than One Doctor	12	<i>22</i>	*	*	9	<i>19</i>	8	<i>4</i>	4	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	<i>3</i>	*	*	*	<i>3</i>	*	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	47	<i>57</i>	*	*	46	<i>56</i>	20	<i>27</i>	42	<i>50</i>
From Friend or Relative for Free	38	<i>45</i>	*	*	37	<i>45</i>	16	<i>11</i>	34	*
Bought from Friend or Relative	27	<i>35</i>	*	*	27	<i>34</i>	12	<i>24</i>	24	<i>24</i>
Took from Friend or Relative without Asking	0	<i>3</i>	*	*	*	<i>3</i>	*	<i>0</i>	*	<i>3</i>
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	15	<i>30</i>	*	*	13	<i>21</i>	2	*	13	<i>19</i>
<b>SOME OTHER WAY<sup>1</sup></b>	14	<i>7</i>	*	*	14	<i>1</i>	4	<i>1</i>	13	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 6.8D Source Where Friend or Relative Obtained Prescription Stimulants: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Stimulants from a Friend or Relative for Free in Past Year; by Age Group, Standard Errors of Percentages, 2019 and 2020**

Source Where Friend or Relative Obtained Stimulants	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	1.94	<i>2.49</i>	*	*	2.04	<i>2.45</i>	2.14	<i>3.58</i>	3.04	<i>3.22</i>
Prescription from One Doctor	1.98	<i>2.57</i>	*	*	2.07	<i>2.56</i>	2.31	<i>3.65</i>	3.04	<i>3.38</i>
Prescriptions from More Than One Doctor	0.53	<i>0.90</i>	*	*	0.41	<i>0.83</i>	0.91	<i>0.59</i>	0.32	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	<i>0.14</i>	*	*	*	<i>0.15</i>	*	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	1.82	<i>2.23</i>	*	*	1.92	<i>2.31</i>	2.10	<i>3.45</i>	2.83	<i>3.02</i>
From Friend or Relative for Free	1.55	<i>1.80</i>	*	*	1.63	<i>1.87</i>	1.78	<i>1.57</i>	2.43	*
Bought from Friend or Relative	1.10	<i>1.41</i>	*	*	1.18	<i>1.47</i>	1.28	<i>3.18</i>	1.75	<i>1.57</i>
Took from Friend or Relative without Asking	0.01	<i>0.11</i>	*	*	*	<i>0.11</i>	*	<i>0.06</i>	*	<i>0.16</i>
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.62	<i>1.20</i>	*	*	0.62	<i>0.90</i>	0.27	*	0.99	<i>1.21</i>
<b>SOME OTHER WAY<sup>1</sup></b>	0.57	<i>0.31</i>	*	*	0.62	<i>0.05</i>	0.50	<i>0.17</i>	0.96	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 6.10C Source Where Friend or Relative Obtained Prescription Tranquilizers: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Tranquilizers from a Friend or Relative for Free in Past Year; by Age Group, Standard Errors of Numbers in Thousands, 2019 and 2020**

Source Where Friend or Relative Obtained Tranquilizers	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	112	<i>213</i>	*	*	111	<i>213</i>	37	*	107	<i>199</i>
Prescription from One Doctor	110	<i>212</i>	*	*	109	<i>213</i>	37	*	105	<i>199</i>
Prescriptions from More Than One Doctor	14	<i>15</i>	*	*	14	<i>15</i>	2	*	14	<i>15</i>
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	10	*	*	*	10	*	3	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	40	<i>59</i>	*	*	37	<i>58</i>	20	*	32	<i>52</i>
From Friend or Relative for Free	28	<i>55</i>	*	*	27	<i>55</i>	16	*	22	<i>52</i>
Bought from Friend or Relative	28	<i>11</i>	*	*	26	<i>10</i>	13	7	23	7
Took from Friend or Relative without Asking	6	<i>15</i>	*	*	5	<i>14</i>	*	*	*	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	22	<i>33</i>	*	*	20	<i>26</i>	14	*	15	<i>14</i>
<b>SOME OTHER WAY<sup>1</sup></b>	15	<i>29</i>	*	*	14	<i>29</i>	1	0	14	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 6.10D Source Where Friend or Relative Obtained Prescription Tranquilizers: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Tranquilizers from a Friend or Relative for Free in Past Year; by Age Group, Standard Errors of Percentages, 2019 and 2020**

Source Where Friend or Relative Obtained Tranquilizers	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	2.02	2.62	*	*	2.06	2.55	3.68	*	2.42	2.96
Prescription from One Doctor	2.19	2.68	*	*	2.25	2.62	3.73	*	2.72	3.05
Prescriptions from More Than One Doctor	0.67	0.58	*	*	0.70	0.59	0.45	*	0.95	0.74
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.46	*	*	*	0.48	*	0.59	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	1.76	2.18	*	*	1.77	2.22	3.26	*	2.07	2.50
From Friend or Relative for Free	1.26	2.07	*	*	1.31	2.11	2.72	*	1.47	2.48
Bought from Friend or Relative	1.26	0.45	*	*	1.28	0.41	2.35	1.44	1.52	0.37
Took from Friend or Relative without Asking	0.27	0.56	*	*	0.25	0.57	*	*	*	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	1.01	1.30	*	*	1.00	1.04	2.48	*	1.02	0.68
<b>SOME OTHER WAY<sup>1</sup></b>	0.67	1.12	*	*	0.70	1.15	0.18	0.02	0.96	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 6.12C Source Where Friend or Relative Obtained Prescription Sedatives: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Sedatives from a Friend or Relative for Free in Past Year; by Age Group, Standard Errors of Numbers in Thousands, 2019 and 2020**

Source Where Friend or Relative Obtained Sedatives	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	64	*	*	*	*	*	*	*	*	*
Prescription from One Doctor	64	*	*	*	64	*	*	*	*	*
Prescriptions from More Than One Doctor	5	*	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	17	*	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*	*	*
Bought from Friend or Relative	7	*	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	3	*	*	*	*	*	*	*	*	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0	*	*	*	0	*	*	*	*	*
<b>SOME OTHER WAY<sup>1</sup></b>	*	*	*	*	*	*	*	*	*	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 6.12D Source Where Friend or Relative Obtained Prescription Sedatives: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Sedatives from a Friend or Relative for Free in Past Year; by Age Group, Standard Errors of Percentages, 2019 and 2020**

Source Where Friend or Relative Obtained Sedatives	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	3.46	*	*	*	*	*	*	*	*	*
Prescription from One Doctor	3.62	*	*	*	3.69	*	*	*	*	*
Prescriptions from More Than One Doctor	1.05	*	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	3.46	*	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*	*	*
Bought from Friend or Relative	1.49	*	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	0.70	*	*	*	*	*	*	*	*	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.03	*	*	*	0.04	*	*	*	*	*
<b>SOME OTHER WAY<sup>1</sup></b>	*	*	*	*	*	*	*	*	*	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.