

**Table 7.19C Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 12 or Older and among Past Year Users of Specific Substances Aged 12 or Older; Standard Errors of Numbers in Thousands, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																			
Marijuana	146	142	163	165	143	170	182	176	218	219	225	259	225	218	232	252	256	286	<i>498</i>
Cocaine	35	22	36	47	56	39	26	16	29	18	33	50	22	32	21	37	42	34	<i>30</i>
Crack	16	10	32	35	14	29	10	14	19	14	13	28	20	25	11	33	20	12	<i>22</i>
Heroin	28	13	22	7	58	14	23	19	18	29	44	27	29	27	38	33	36	35	<i>112</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	8	15	13	31	<i>19</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	18	5	12	15	13	<i>54</i>
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	30	33	39	40	54	<i>116</i>
Alcohol	426	362	419	409	435	409	440	450	413	445	415	401	358	369	381	375	368	366	<i>645</i>
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																			
Marijuana	145	144	163	165	145	171	187	175	221	219	227	261	231	218	236	258	260	296	<i>513</i>
Cocaine	35	21	36	47	56	39	26	16	29	18	33	50	22	32	21	37	42	34	<i>30</i>
Crack	16	10	32	35	14	29	10	14	*	*	13	*	20	25	11	*	20	12	<i>*</i>
Heroin	*	*	*	7	*	*	*	19	18	29	*	27	29	27	38	33	36	35	<i>*</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	8	15	13	31	<i>19</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	18	5	12	15	13	<i>*</i>
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	30	33	39	40	55	<i>116</i>
Alcohol	442	377	430	425	452	424	458	475	419	457	436	417	377	387	391	388	380	386	<i>664</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.19D Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 12 or Older and among Past Year Users of Specific Substances Aged 12 or Older; Standard Errors of Percentages, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																			
Marijuana	0.06	0.06	0.07	0.07	0.06	0.07	0.07	0.07	0.09	0.08	0.09	0.10	0.08	0.08	0.09	0.09	0.09	0.10	<i>0.18</i>
Cocaine	0.01	0.01	0.01	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.02	0.01	<i>0.01</i>
Crack	0.01	0.00	0.01	0.01	0.01	0.01	0.00	0.01	0.01	0.01	0.00	0.01	0.01	0.01	0.00	0.01	0.01	0.00	<i>0.01</i>
Heroin	0.01	0.01	0.01	0.00	0.02	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	<i>0.04</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.00	0.00	0.01	0.00	0.01	<i>0.01</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.00	0.00	0.01	0.00	<i>0.02</i>
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.01	0.01	0.01	0.02	<i>0.04</i>
Alcohol	0.18	0.15	0.17	0.17	0.18	0.16	0.18	0.18	0.16	0.17	0.16	0.15	0.13	0.14	0.14	0.14	0.13	0.13	<i>0.23</i>
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																			
Marijuana	0.51	0.53	0.58	0.60	0.52	0.62	0.64	0.55	0.65	0.65	0.62	0.71	0.56	0.54	0.56	0.54	0.53	0.52	<i>0.89</i>
Cocaine	0.59	0.36	0.62	0.83	0.91	0.66	0.48	0.34	0.63	0.47	0.71	1.16	0.47	0.65	0.41	0.60	0.74	0.60	<i>0.58</i>
Crack	1.02	0.71	2.30	2.41	0.97	1.94	0.91	1.40	*	*	1.39	*	2.48	2.83	1.30	*	2.47	1.58	*
Heroin	*	*	*	1.81	*	*	*	3.12	2.92	4.11	*	3.62	2.89	3.03	3.55	3.35	3.48	4.42	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.28	0.17	0.29	0.24	0.51	<i>0.27</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.02	0.33	0.68	0.75	0.61	*
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.68	2.25	2.23	2.03	2.46	<i>4.08</i>
Alcohol	0.27	0.23	0.27	0.25	0.26	0.25	0.26	0.26	0.24	0.25	0.24	0.23	0.20	0.21	0.21	0.21	0.20	0.20	<i>0.35</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.20C Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 12 to 17 and among Past Year Users of Specific Substances Aged 12 to 17; Standard Errors of Numbers in Thousands, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																			
Marijuana	29	21	25	23	23	22	22	22	25	22	22	24	24	25	23	23	23	25	56
Cocaine	4	5	2	4	3	4	2	1	1	2	*	*	*	*	2	*	2	*	*
Crack	2	*	*	3	*	3	*	*	*	2	*	*	*	*	*	*	*	*	*
Heroin	2	1	2	*	*	*	1	2	*	*	*	*	2	*	2	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	2	*	2	*	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	3	5	8	3	4	2
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	2	2	2	*	*
Alcohol	16	13	13	12	13	11	10	9	10	10	8	10	14	13	9	10	6	7	5
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																			
Marijuana	29	21	25	23	23	22	22	23	26	22	22	24	24	25	23	23	24	25	57
Cocaine	4	5	2	4	3	4	2	1	1	*	0	*	*	*	*	*	*	*	*
Crack	2	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1	2	0	2	1	0
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	3	5	8	3	4	2
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*	*
Alcohol	16	13	14	12	13	11	10	8	10	10	8	10	14	13	9	10	6	7	5

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.20D Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 12 to 17 and among Past Year Users of Specific Substances Aged 12 to 17; Standard Errors of Percentages, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																			
Marijuana	0.12	0.09	0.10	0.09	0.09	0.09	0.09	0.09	0.10	0.09	0.09	0.10	0.10	0.10	0.09	0.09	0.09	0.10	<i>0.23</i>
Cocaine	0.02	0.02	0.01	0.02	0.01	0.02	0.01	0.01	0.01	0.01	*	*	*	*	0.01	*	0.01	*	*
Crack	0.01	*	*	0.01	*	0.01	*	*	*	0.01	*	*	*	*	*	*	*	*	*
Heroin	0.01	0.01	0.01	*	*	*	0.01	0.01	*	*	*	*	0.01	*	0.01	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.01	*	0.01	*	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.02	0.03	0.01	0.02	<i>0.01</i>
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.01	0.01	0.01	*	*
Alcohol	0.06	0.05	0.05	0.05	0.05	0.05	0.04	0.03	0.04	0.04	0.03	0.04	0.05	0.05	0.03	0.04	0.03	0.03	<i>0.02</i>
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																			
Marijuana	0.68	0.55	0.65	0.67	0.67	0.68	0.64	0.64	0.69	0.59	0.63	0.67	0.71	0.74	0.75	0.72	0.71	0.72	<i>2.09</i>
Cocaine	0.74	1.06	0.51	0.94	0.65	0.97	0.80	0.60	0.55	*	0.10	*	*	*	*	*	*	*	*
Crack	2.40	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.13	0.50	0.07	0.64	0.14	<i>0.09</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.38	0.89	1.41	0.42	0.52	<i>0.37</i>
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*	*
Alcohol	0.18	0.15	0.16	0.14	0.15	0.14	0.13	0.11	0.14	0.15	0.13	0.16	0.23	0.22	0.16	0.18	0.12	0.14	<i>0.11</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.21C Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 18 or Older and among Past Year Users of Specific Substances Aged 18 or Older; Standard Errors of Numbers in Thousands, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																			
Marijuana	142	140	162	165	140	169	181	175	217	217	223	258	222	214	230	251	254	282	<i>499</i>
Cocaine	35	21	35	46	56	39	26	16	29	18	33	50	22	32	21	37	42	34	<i>30</i>
Crack	16	10	32	35	14	29	10	14	19	14	13	28	20	25	11	33	20	12	<i>22</i>
Heroin	28	13	22	7	58	14	23	18	18	29	44	27	29	27	38	33	36	35	<i>112</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	8	15	13	31	<i>19</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	18	*	9	15	13	<i>54</i>
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	30	33	39	40	55	<i>116</i>
Alcohol	424	362	417	408	434	408	439	448	412	444	413	400	356	368	380	374	367	366	<i>643</i>
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																			
Marijuana	142	141	162	164	142	170	185	174	220	217	225	260	228	215	234	257	258	293	<i>513</i>
Cocaine	35	21	36	46	56	39	26	16	29	18	33	50	22	32	21	37	42	34	<i>30</i>
Crack	16	10	32	35	14	29	10	14	*	*	13	*	20	25	11	*	20	12	<i>*</i>
Heroin	*	*	*	7	*	*	*	18	18	29	*	27	29	27	38	33	36	35	<i>*</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	8	15	13	31	<i>19</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	2	9	*	13	<i>*</i>
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	30	33	39	40	55	<i>116</i>
Alcohol	441	377	429	424	452	423	458	474	419	457	436	417	375	386	391	388	380	386	<i>664</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.21D Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 18 or Older and among Past Year Users of Specific Substances Aged 18 or Older; Standard Errors of Percentages, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																			
Marijuana	0.07	0.07	0.08	0.08	0.06	0.08	0.08	0.08	0.09	0.09	0.09	0.11	0.09	0.09	0.09	0.10	0.10	0.11	<i>0.20</i>
Cocaine	0.02	0.01	0.02	0.02	0.03	0.02	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.02	0.01	<i>0.01</i>
Crack	0.01	0.00	0.01	0.02	0.01	0.01	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.00	0.01	0.01	0.00	<i>0.01</i>
Heroin	0.01	0.01	0.01	0.00	0.03	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.02	0.01	0.01	0.01	<i>0.04</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.00	0.01	0.01	0.01	<i>0.01</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	*	0.00	0.01	0.01	<i>0.02</i>
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.01	0.02	0.02	0.02	<i>0.05</i>
Alcohol	0.20	0.17	0.19	0.19	0.20	0.18	0.20	0.20	0.18	0.19	0.18	0.17	0.15	0.15	0.16	0.15	0.15	0.15	<i>0.26</i>
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																			
Marijuana	0.59	0.62	0.67	0.69	0.59	0.70	0.73	0.63	0.73	0.73	0.69	0.78	0.61	0.58	0.61	0.59	0.57	0.55	<i>0.94</i>
Cocaine	0.64	0.38	0.66	0.88	0.97	0.71	0.51	0.36	0.66	0.50	0.74	1.20	0.49	0.67	0.42	0.62	0.76	0.61	<i>0.59</i>
Crack	1.08	0.76	2.41	2.50	1.02	2.03	0.94	1.45	*	*	1.42	*	2.54	2.84	1.32	*	2.49	1.61	*
Heroin	*	*	*	2.00	*	*	*	3.29	3.06	4.47	*	3.76	2.97	3.11	3.60	3.40	3.52	4.42	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.31	0.18	0.32	0.25	0.55	<i>0.28</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.20	0.73	*	0.90	*
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.72	2.30	2.29	2.08	2.51	<i>4.12</i>
Alcohol	0.28	0.24	0.28	0.26	0.27	0.26	0.27	0.28	0.25	0.27	0.25	0.24	0.21	0.21	0.22	0.21	0.21	0.21	<i>0.36</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.22C Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 18 to 25 and among Past Year Users of Specific Substances Aged 18 to 25; Standard Errors of Numbers in Thousands, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																			
Marijuana	57	58	59	66	64	64	65	67	73	74	87	84	91	90	88	97	92	86	<i>173</i>
Cocaine	5	8	6	8	8	7	5	7	3	5	8	3	4	9	5	10	4	9	<i>20</i>
Crack	2	3	4	2	3	5	3	2	2	*	3	*	2	5	*	*	*	5	*
Heroin	3	4	4	4	5	6	8	7	9	9	15	6	8	11	10	12	8	8	<i>14</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	7	3	6	3	4	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	2	5	2	*	*
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	7	9	9	8	11	<i>4</i>
Alcohol	41	41	40	38	41	44	43	44	43	40	43	39	47	47	50	52	44	46	<i>57</i>
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																			
Marijuana	58	60	61	67	65	67	67	71	76	77	96	88	94	96	93	102	103	93	<i>183</i>
Cocaine	5	8	6	8	8	7	5	7	3	5	8	3	4	9	5	10	4	9	*
Crack	2	3	4	2	3	5	3	2	2	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	4	*	*	8	*	9	9	*	6	8	*	10	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	7	3	6	3	4	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	0	*
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	7	9	9	8	*	*
Alcohol	42	42	41	39	40	45	44	44	43	42	43	40	48	47	51	53	45	45	<i>58</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.22D Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 18 to 25 and among Past Year Users of Specific Substances Aged 18 to 25; Standard Errors of Percentages, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																			
Marijuana	0.18	0.18	0.18	0.20	0.20	0.20	0.20	0.20	0.21	0.21	0.25	0.24	0.26	0.26	0.25	0.28	0.27	0.26	<i>0.52</i>
Cocaine	0.02	0.02	0.02	0.03	0.03	0.02	0.02	0.02	0.01	0.02	0.02	0.01	0.01	0.03	0.01	0.03	0.01	0.03	<i>0.06</i>
Crack	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01	*	0.01	*	0.01	0.01	*	*	*	0.02	*
Heroin	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.03	0.03	0.04	0.02	0.02	0.03	0.03	0.03	0.02	0.02	<i>0.04</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.02	0.01	0.02	0.01	0.01	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.01	0.02	0.01	*	*
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.02	0.03	0.03	0.02	0.03	<i>0.01</i>
Alcohol	0.13	0.13	0.12	0.12	0.12	0.13	0.13	0.13	0.13	0.12	0.12	0.11	0.14	0.13	0.15	0.15	0.13	0.14	<i>0.17</i>
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																			
Marijuana	0.57	0.62	0.61	0.66	0.64	0.66	0.68	0.61	0.66	0.66	0.72	0.72	0.74	0.73	0.71	0.73	0.69	0.69	<i>1.34</i>
Cocaine	0.25	0.36	0.27	0.37	0.37	0.34	0.28	0.41	0.20	0.35	0.49	0.17	0.24	0.47	0.26	0.47	0.19	0.52	*
Crack	0.78	1.00	1.53	0.75	1.20	1.95	1.29	1.18	1.04	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	2.36	*	*	4.46	*	4.17	3.48	*	2.44	2.64	*	4.05	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.28	0.14	0.27	0.12	0.17	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	0.05	*
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2.25	3.31	2.31	2.71	*	*
Alcohol	0.17	0.16	0.16	0.15	0.16	0.17	0.17	0.17	0.16	0.15	0.16	0.15	0.18	0.18	0.19	0.21	0.18	0.19	<i>0.24</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.23C Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 26 or Older and among Past Year Users of Specific Substances Aged 26 or Older; Standard Errors of Numbers in Thousands, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																			
Marijuana	127	127	149	149	125	149	167	156	199	202	198	239	194	184	202	224	229	262	<i>454</i>
Cocaine	35	20	35	45	55	38	25	15	28	17	33	50	21	31	20	35	42	32	<i>22</i>
Crack	16	10	31	35	14	29	10	14	19	14	12	28	20	24	11	33	20	11	<i>22</i>
Heroin	28	12	21	*	58	13	22	17	16	28	41	26	27	24	36	30	35	34	<i>99</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	11	7	13	13	31	<i>19</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	18	*	*	15	13	<i>54</i>
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	29	32	38	40	53	<i>116</i>
Alcohol	421	360	414	404	433	404	438	441	408	442	409	396	352	363	375	369	360	361	<i>643</i>
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																			
Marijuana	127	128	149	149	126	150	170	156	201	202	199	241	198	185	204	229	230	267	<i>468</i>
Cocaine	35	20	35	45	55	38	25	15	28	17	32	50	21	31	20	35	42	32	<i>22</i>
Crack	16	10	32	35	14	29	10	*	*	*	*	*	20	*	11	*	20	11	<i>*</i>
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	27	24	37	30	35	*	<i>*</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	7	13	13	31	<i>19</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	1	*	*	*	<i>*</i>
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	29	32	38	40	53	<i>116</i>
Alcohol	438	377	428	423	452	421	458	470	416	456	433	413	374	381	388	385	374	383	<i>663</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.23D Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 26 or Older and among Past Year Users of Specific Substances Aged 26 or Older; Standard Errors of Percentages, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																			
Marijuana	0.07	0.07	0.08	0.08	0.07	0.08	0.09	0.08	0.10	0.10	0.10	0.12	0.09	0.09	0.10	0.11	0.11	0.12	<i>0.21</i>
Cocaine	0.02	0.01	0.02	0.02	0.03	0.02	0.01	0.01	0.01	0.01	0.02	0.02	0.01	0.01	0.01	0.02	0.02	0.01	<i>0.01</i>
Crack	0.01	0.01	0.02	0.02	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.00	<i>0.01</i>
Heroin	0.02	0.01	0.01	*	0.03	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.02	0.01	0.02	0.02	<i>0.05</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.00	0.01	0.01	0.01	<i>0.01</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	*	*	0.01	0.01	<i>0.02</i>
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.02	0.02	0.02	0.02	<i>0.05</i>
Alcohol	0.23	0.20	0.23	0.22	0.23	0.21	0.23	0.23	0.21	0.22	0.20	0.20	0.17	0.17	0.18	0.17	0.17	0.17	<i>0.29</i>
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																			
Marijuana	0.93	0.98	1.08	1.08	0.92	1.07	1.13	0.96	1.12	1.13	1.02	1.17	0.82	0.78	0.80	0.78	0.74	0.70	<i>1.14</i>
Cocaine	1.02	0.58	1.09	1.54	1.59	1.13	0.79	0.53	1.04	0.82	1.12	1.93	0.76	1.07	0.67	0.93	1.16	0.88	<i>0.61</i>
Crack	1.31	0.92	2.99	3.27	1.25	2.44	1.12	*	*	*	*	*	3.03	*	1.47	*	2.80	1.52	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	4.07	3.87	4.54	4.17	4.22	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.36	0.60	0.44	0.96	<i>0.44</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.17	*	*	*	*
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2.07	2.74	2.90	2.40	2.79	<i>4.43</i>
Alcohol	0.34	0.29	0.33	0.31	0.33	0.31	0.33	0.32	0.30	0.31	0.29	0.28	0.24	0.25	0.25	0.25	0.24	0.24	<i>0.42</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.24C Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 12 to 20 and among Past Year Users of Specific Substances Aged 12 to 20; Standard Errors of Numbers in Thousands, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																				
Marijuana	50	49	43	53	50	47	51	50	52	52	66	58	62	61	56	62	60	60	<i>107</i>	
Cocaine	5	6	3	5	6	6	4	6	2	6	7	2	2	6	4	5	2	4	<i>20</i>	
Crack	3	3	2	3	1	3	2	*	2	2	3	*	*	4	*	*	*	4	*	
Heroin	2	4	2	2	*	3	3	3	8	6	5	2	7	4	3	3	3	4	*	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	3	2	2	3	*	*	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	3	5	10	3	4	2	
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	6	7	7	5	6	*	
Alcohol	27	28	24	22	22	25	25	23	24	23	18	23	25	27	18	26	21	20	22	
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																				
Marijuana	50	49	43	53	50	47	51	50	52	52	66	58	62	61	56	62	60	60	<i>107</i>	
Cocaine	5	6	3	5	6	6	4	6	2	6	7	2	2	6	4	5	2	4	*	
Crack	3	*	*	*	1	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	2	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	3	2	2	3	1	<i>0</i>	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	3	5	10	3	4	2	
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*	*	
Alcohol	27	28	24	22	22	25	25	23	24	23	18	23	25	27	18	26	21	20	22	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2002 to 2019 standard errors of numbers in thousands may differ from previously published estimates because of a change to the standard error estimation method. See the Introduction to these tables for more details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.24D Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 12 to 20 and among Past Year Users of Specific Substances Aged 12 to 20; Standard Errors of Percentages, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																				
Marijuana	0.13	0.13	0.11	0.13	0.13	0.12	0.13	0.13	0.13	0.13	0.16	0.15	0.16	0.16	0.15	0.16	0.15	0.15	<i>0.28</i>	
Cocaine	0.01	0.02	0.01	0.01	0.02	0.01	0.01	0.02	0.01	0.01	0.02	0.00	0.01	0.02	0.01	0.01	0.01	0.01	<i>0.05</i>	
Crack	0.01	0.01	0.01	0.01	0.00	0.01	0.01	*	0.00	0.01	0.01	*	*	0.01	*	*	*	0.01	*	
Heroin	0.00	0.01	0.01	0.01	*	0.01	0.01	0.01	0.02	0.01	0.01	0.00	0.02	0.01	0.01	0.01	0.01	0.01	*	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.01	0.01	0.01	*	*	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.01	0.03	0.01	0.01	<i>0.01</i>	
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.02	0.02	0.02	0.01	0.02	*	
Alcohol	0.07	0.07	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.05	0.06	0.06	0.07	0.05	0.07	0.05	0.05	<i>0.06</i>	
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																				
Marijuana	0.57	0.57	0.52	0.64	0.61	0.59	0.63	0.57	0.60	0.59	0.70	0.66	0.74	0.73	0.71	0.76	0.69	0.67	<i>1.53</i>	
Cocaine	0.39	0.50	0.28	0.39	0.45	0.45	0.39	0.65	0.29	0.68	0.89	0.31	0.35	0.79	0.60	0.73	0.31	0.79	*	
Crack	1.30	*	*	*	0.64	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	1.49	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.17	0.16	0.18	0.24	0.05	<i>0.03</i>	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.29	0.68	1.25	0.32	0.40	<i>0.29</i>	
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*	*	*
Alcohol	0.16	0.16	0.13	0.12	0.13	0.14	0.14	0.14	0.15	0.14	0.12	0.15	0.17	0.19	0.13	0.20	0.16	0.15	<i>0.20</i>	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.25C Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 21 or Older and among Past Year Users of Specific Substances Aged 21 or Older; Standard Errors of Numbers in Thousands, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																			
Marijuana	137	133	156	156	135	161	178	165	214	212	215	251	215	204	224	250	250	284	<i>505</i>
Cocaine	35	20	35	46	56	39	25	15	29	17	33	50	21	31	21	36	42	33	<i>22</i>
Crack	16	10	32	35	14	29	10	14	19	14	13	28	20	24	11	33	20	11	<i>22</i>
Heroin	28	12	22	6	58	14	23	18	17	28	44	27	28	26	38	33	36	35	<i>112</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	8	14	13	31	<i>19</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	18	*	*	15	13	<i>54</i>
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	29	33	38	40	54	<i>116</i>
Alcohol	441	378	429	424	452	422	457	473	418	456	437	415	376	385	390	386	379	386	<i>663</i>
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																			
Marijuana	137	133	156	156	135	161	178	165	214	212	215	251	215	204	224	250	250	284	<i>505</i>
Cocaine	35	20	35	46	56	39	25	15	29	17	33	50	21	31	21	36	42	33	<i>22</i>
Crack	16	10	32	35	14	29	10	14	*	*	*	*	20	24	11	*	20	11	<i>*</i>
Heroin	*	*	*	6	*	*	*	*	17	*	*	27	28	26	38	33	36	35	<i>*</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	8	14	13	31	<i>19</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	1	7	*	13	<i>*</i>
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	29	33	38	40	54	<i>116</i>
Alcohol	441	378	429	424	452	422	457	473	418	456	437	415	376	385	390	386	379	386	<i>663</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2002 to 2019 standard errors of numbers in thousands may differ from previously published estimates because of a change to the standard error estimation method. See the Introduction to these tables for more details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.25D Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 21 or Older and among Past Year Users of Specific Substances Aged 21 or Older; Standard Errors of Percentages, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION</b>																			
Marijuana	0.07	0.07	0.08	0.08	0.06	0.08	0.08	0.08	0.10	0.10	0.10	0.11	0.09	0.09	0.10	0.10	0.10	0.12	<i>0.20</i>
Cocaine	0.02	0.01	0.02	0.02	0.03	0.02	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.02	0.02	0.01	<i>0.01</i>
Crack	0.01	0.00	0.02	0.02	0.01	0.01	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.00	0.01	0.01	0.00	<i>0.01</i>
Heroin	0.01	0.01	0.01	0.00	0.03	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.02	0.01	0.02	0.01	<i>0.05</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.00	0.01	0.01	0.01	<i>0.01</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	*	*	0.01	0.01	<i>0.02</i>
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.01	0.02	0.02	0.02	<i>0.05</i>
Alcohol	0.21	0.18	0.21	0.20	0.21	0.19	0.21	0.21	0.19	0.20	0.19	0.18	0.16	0.16	0.16	0.16	0.16	0.15	<i>0.27</i>
<b>DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS</b>																			
Marijuana	0.70	0.73	0.81	0.81	0.70	0.82	0.85	0.73	0.86	0.86	0.80	0.90	0.66	0.65	0.67	0.65	0.63	0.60	<i>1.01</i>
Cocaine	0.75	0.44	0.79	1.04	1.17	0.83	0.59	0.39	0.76	0.57	0.83	1.35	0.54	0.75	0.47	0.68	0.84	0.66	<i>0.46</i>
Crack	1.16	0.81	2.61	2.79	1.13	2.17	1.01	1.53	*	*	*	*	2.67	2.90	1.36	*	2.59	1.55	*
Heroin	*	*	*	2.38	*	*	*	*	3.34	*	*	4.07	3.14	3.29	3.79	3.52	3.69	4.52	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.41	0.24	0.39	0.30	0.65	<i>0.32</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.12	0.70	*	1.07	*
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.79	2.41	2.42	2.19	2.58	<i>4.19</i>
Alcohol	0.30	0.26	0.30	0.28	0.29	0.27	0.29	0.29	0.26	0.28	0.26	0.25	0.22	0.22	0.23	0.22	0.22	0.22	<i>0.38</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.26C Daily or Almost Daily Use of Specific Substances in Past Month: Among All People Aged 12 or Older and among Past Month Users of Specific Substances Aged 12 or Older; Standard Errors of Numbers in Thousands, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																				
Marijuana	171	175	200	194	196	204	218	237	275	265	269	311	262	277	269	300	313	340	<i>566</i>	
Cocaine	27	30	49	45	75	41	37	39	47	25	36	55	26	32	43	38	28	38	<i>34</i>	
Crack	14	26	33	28	49	30	27	31	18	14	29	35	20	22	35	20	21	29	<i>25</i>	
Heroin	37	22	31	11	70	29	28	18	22	43	42	33	33	30	41	38	28	38	<i>114</i>	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	14	7	11	15	6	<i>31</i>	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	11	11	19	18	<i>78</i>	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	43	49	50	57	57	<i>157</i>	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	54	57	48	61	62	<i>72</i>	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	26	28	28	21	26	<i>41</i>	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	25	31	20	29	30	<i>44</i>	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	15	24	17	*	13	<i>29</i>	
Cigarettes (Daily Only) <sup>2</sup>	702	655	646	626	637	655	688	659	665	675	691	656	528	546	542	523	537	517	<i>853</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	215	210	198	202	205	<i>267</i>	
Cigars	121	143	136	143	142	149	152	147	164	155	177	155	113	157	150	140	154	138	<i>321</i>	
Alcohol	577	563	525	555	578	551	582	638	561	585	605	590	497	487	508	491	534	514	<i>774</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	173	168	163	179	163	<i>378</i>	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																				
Marijuana	173	174	199	192	198	205	221	236	281	267	278	318	273	281	273	308	319	356	<i>583</i>	
Cocaine	27	30	49	45	75	41	37	39	47	25	36	55	26	32	43	38	28	38	<i>34</i>	
Crack	14	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	<i>*</i>
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	33	*	*	39	*	*	*	<i>*</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	14	7	11	15	6	<i>31</i>	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	11	11	19	18	<i>*</i>	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	43	49	50	57	57	<i>158</i>	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	54	57	48	61	62	<i>72</i>	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	27	28	28	21	26	<i>41</i>	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	25	31	20	29	30	<i>44</i>	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	<i>*</i>	
Cigarettes (Daily Only) <sup>2</sup>	774	714	693	664	690	727	730	726	737	709	740	707	578	580	586	561	559	553	<i>898</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	219	208	201	203	205	<i>267</i>	
Cigars	121	145	137	142	143	149	152	146	165	154	177	156	113	158	150	141	154	139	<i>323</i>	
Alcohol	619	617	561	604	617	590	629	694	607	623	654	643	547	529	532	526	581	559	<i>812</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	175	169	164	180	164	<i>381</i>	
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>	538	525	505	489	482	509	512	503	486	441	455	466	365	367	379	369	342	348	<i>568</i>	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.26D Daily or Almost Daily Use of Specific Substances in Past Month: Among All People Aged 12 or Older and among Past Month Users of Specific Substances Aged 12 or Older; Standard Errors of Percentages, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																					
Marijuana	0.07	0.07	0.08	0.08	0.08	0.08	0.09	0.09	0.11	0.10	0.10	0.12	0.10	0.10	0.10	0.11	0.11	0.12	<i>0.20</i>		
Cocaine	0.01	0.01	0.02	0.02	0.03	0.02	0.01	0.02	0.02	0.01	0.01	0.02	0.01	0.01	0.02	0.01	0.01	0.01	0.01	<i>0.01</i>	
Crack	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	<i>0.01</i>	
Heroin	0.02	0.01	0.01	0.00	0.03	0.01	0.01	0.01	0.01	0.02	0.02	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01	<i>0.04</i>	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.00	0.00	0.01	0.00	0.00	<i>0.01</i>	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.00	0.00	0.00	0.01	0.01	0.01	<i>0.03</i>	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.02	0.02	0.02	0.02	0.02	0.02	<i>0.06</i>	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.02	0.02	0.02	0.02	0.02	0.02	<i>0.03</i>	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.01	0.01	0.01	0.01	0.01	0.01	<i>0.01</i>	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.01	0.01	0.01	0.01	0.01	0.01	<i>0.02</i>	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	0.01	0.01	0.01	*	0.00	0.00	<i>0.01</i>	
Cigarettes (Daily Only) <sup>2</sup>	0.30	0.28	0.27	0.26	0.26	0.26	0.28	0.26	0.26	0.26	0.27	0.25	0.20	0.20	0.20	0.19	0.20	0.19	0.19	<i>0.31</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	0.08	0.08	0.07	0.07	0.07	0.07	<i>0.10</i>	
Cigars	0.05	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.07	0.06	0.04	0.06	0.06	0.05	0.06	0.05	0.05	<i>0.12</i>	
Alcohol	0.25	0.24	0.22	0.23	0.23	0.22	0.23	0.25	0.22	0.23	0.23	0.22	0.19	0.18	0.19	0.18	0.20	0.19	0.19	<i>0.28</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.06	0.06	0.06	0.07	0.06	0.06	<i>0.14</i>	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																					
Marijuana	0.95	0.98	1.01	1.05	1.06	1.09	1.16	1.06	1.17	1.08	1.04	1.13	0.86	0.91	0.81	0.89	0.87	0.80	0.80	<i>1.23</i>	
Cocaine	1.31	1.29	2.17	1.78	2.78	1.85	1.91	2.31	2.96	1.80	2.07	3.28	1.70	1.63	2.14	1.68	1.42	1.79	1.79	<i>1.80</i>	
Crack	2.39	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	<i>*</i>
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	5.89	*	*	5.88	*	*	*	*	<i>*</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.12	0.48	0.79	0.91	0.30	0.30	<i>1.75</i>	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	2.45	1.92	1.96	3.00	2.16	2.16	<i>*</i>	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	4.06	5.83	4.90	4.64	4.24	4.24	<i>5.97</i>	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	1.34	1.59	1.42	1.95	2.05	2.05	<i>2.73</i>	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	1.54	1.55	1.49	1.22	1.58	1.58	<i>2.64</i>	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	1.31	1.50	1.15	1.67	1.57	1.57	<i>2.27</i>	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*	<i>*</i>	
Cigarettes (Daily Only) <sup>2</sup>	0.66	0.67	0.63	0.62	0.59	0.65	0.70	0.68	0.71	0.71	0.71	0.73	0.59	0.64	0.66	0.69	0.71	0.71	0.71	<i>1.25</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	1.39	1.38	1.41	1.44	1.48	1.48	<i>2.79</i>	
Cigars	0.89	1.03	0.93	0.94	0.94	1.04	1.06	1.01	1.09	1.09	1.19	1.13	0.88	1.10	1.04	1.02	1.13	1.08	1.08	<i>2.53</i>	
Alcohol	0.44	0.44	0.41	0.41	0.43	0.41	0.42	0.46	0.40	0.41	0.42	0.41	0.33	0.33	0.34	0.32	0.35	0.35	0.35	<i>0.52</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.26	0.25	0.24	0.26	0.24	0.24	<i>0.59</i>	
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																					
	0.91	0.82	0.87	0.86	0.85	0.88	0.94	0.98	0.94	0.90	0.94	1.00	0.83	0.87	0.89	0.92	0.94	0.95	0.95	<i>1.78</i>	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.27C Daily or Almost Daily Use of Specific Substances in Past Month: Among All People Aged 12 to 17 and among Past Month Users of Specific Substances Aged 12 to 17; Standard Errors of Numbers in Thousands, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																			
Marijuana	31	28	31	26	29	27	27	29	28	28	28	28	30	32	29	30	29	33	67
Cocaine	4	4	5	4	*	4	4	3	1	*	2	*	5	2	2	*	2	*	*
Crack	3	4	3	*	*	3	*	2	*	*	*	*	*	*	*	*	*	*	*
Heroin	2	2	4	*	*	*	2	1	*	7	*	*	3	*	*	*	3	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1	3	2	2	*	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	4	6	5	3	8	4
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	2	3	2	2	5
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	3	3	4	*	2	*
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	5	5	3	*	5	3
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	7	3	4	5	1	*
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	42	38	42	33	35	34	28	29	30	28	25	21	27	22	15	14	14	12	12
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	16	15	12	13	12	12
Cigars	14	11	12	11	11	9	9	10	9	8	7	6	9	9	10	10	8	12	15
Alcohol	15	15	17	19	13	14	12	15	13	11	12	9	11	12	8	9	7	11	7
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	5	2	6	5	7	*
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																			
Marijuana	32	29	32	26	29	27	28	29	29	29	29	28	31	32	29	30	29	33	67
Cocaine	4	4	*	4	1	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	3	8	*
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	3	3	4	1	*	*
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	41	39	43	34	35	35	29	30	30	28	25	21	27	22	15	13	14	12	*
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	16	15	12	13	12	*
Cigars	14	11	12	11	11	9	9	10	9	8	7	6	9	9	10	10	8	12	*
Alcohol	15	15	17	20	13	14	12	15	13	11	12	9	11	12	8	9	7	11	7
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	5	2	6	5	7	*
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>	19	17	17	15	15	16	12	12	11	10	7	7	8	5	*	*	*	*	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.27D Daily or Almost Daily Use of Specific Substances in Past Month: Among All People Aged 12 to 17 and among Past Month Users of Specific Substances Aged 12 to 17; Standard Errors of Percentages, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																				
Marijuana	0.13	0.11	0.12	0.10	0.11	0.11	0.11	0.12	0.12	0.11	0.11	0.11	0.12	0.13	0.11	0.12	0.12	0.13	<i>0.27</i>	
Cocaine	0.01	0.02	0.02	0.02	*	0.02	0.02	0.01	0.01	*	0.01	*	0.02	0.01	0.01	*	0.01	*	*	
Crack	0.01	0.01	0.01	*	*	0.01	*	0.01	*	*	*	*	*	*	*	*	*	*	*	
Heroin	0.01	0.01	0.01	*	*	*	0.01	0.01	*	0.03	*	*	0.01	*	*	*	0.01	*	*	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.00	0.01	0.01	0.01	*	*	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.02	0.02	0.02	0.01	0.03	<i>0.01</i>	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	0.01	0.01	0.01	0.01	<i>0.02</i>	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.01	0.01	0.02	*	0.01	*	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.02	0.02	0.01	*	0.02	<i>0.01</i>	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.03	0.01	0.01	0.02	0.01	*	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*	
Cigarettes (Daily Only) <sup>2</sup>	0.17	0.15	0.17	0.13	0.14	0.14	0.11	0.12	0.12	0.11	0.10	0.08	0.11	0.09	0.06	0.05	0.06	0.05	<i>0.05</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	0.06	0.06	0.05	0.05	0.05	<i>0.05</i>	
Cigars	0.06	0.04	0.05	0.04	0.04	0.04	0.03	0.04	0.04	0.03	0.03	0.02	0.04	0.03	0.04	0.04	0.03	0.05	<i>0.06</i>	
Alcohol	0.06	0.06	0.07	0.08	0.05	0.06	0.05	0.06	0.05	0.04	0.05	0.04	0.04	0.05	0.03	0.03	0.03	0.04	<i>0.03</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.02	0.01	0.02	0.02	0.03	*	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																				
Marijuana	1.32	1.26	1.38	1.38	1.44	1.41	1.39	1.38	1.41	1.27	1.38	1.35	1.41	1.60	1.54	1.70	1.54	1.58	<i>3.80</i>	
Cocaine	2.37	2.54	*	2.81	0.53	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*	*	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	1.42	3.43	*	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	1.16	1.35	1.85	0.49	*	*	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*	
Cigarettes (Daily Only) <sup>2</sup>	1.03	1.06	1.13	1.12	1.19	1.16	1.11	1.17	1.29	1.28	1.33	1.35	1.89	1.84	1.64	1.64	2.04	2.04	*	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	3.58	3.47	3.05	4.09	4.22	*	
Cigars	1.16	0.94	0.97	0.96	1.03	0.83	0.87	0.98	1.10	0.93	1.07	1.05	1.71	1.58	2.11	1.90	1.95	3.09	*	
Alcohol	0.34	0.33	0.38	0.46	0.31	0.35	0.33	0.40	0.39	0.32	0.38	0.31	0.37	0.47	0.36	0.35	0.31	0.47	<i>0.32</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.32	0.20	0.45	0.39	0.59	*	
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																				
	1.61	1.68	1.80	1.87	1.94	2.14	2.08	2.12	2.24	1.97	1.88	2.47	2.52	2.51	*	*	*	*	*	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.28C Daily or Almost Daily Use of Specific Substances in Past Month: Among All People Aged 18 or Older and among Past Month Users of Specific Substances Aged 18 or Older; Standard Errors of Numbers in Thousands, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																				
Marijuana	169	172	198	193	192	204	215	233	273	264	268	310	259	273	267	300	309	336	<i>564</i>	
Cocaine	27	30	48	45	75	41	37	39	47	25	36	55	26	32	43	38	28	38	<i>34</i>	
Crack	13	26	33	28	48	30	27	31	18	14	29	35	20	22	35	20	21	29	<i>25</i>	
Heroin	37	22	31	11	70	29	28	18	22	42	42	33	33	30	41	38	28	38	<i>114</i>	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	14	*	11	15	6	<i>31</i>	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	10	10	19	16	<i>78</i>	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	43	49	49	57	57	<i>157</i>	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	54	57	48	61	62	<i>72</i>	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	26	28	28	21	25	<i>41</i>	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	24	30	20	28	30	<i>44</i>	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	15	24	17	*	13	<i>29</i>	
Cigarettes (Daily Only) <sup>2</sup>	703	644	645	628	634	650	685	653	659	674	688	657	526	545	540	521	533	516	<i>844</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	215	210	198	201	203	<i>267</i>	
Cigars	120	143	136	143	142	148	152	147	164	154	177	154	113	157	150	139	154	139	<i>321</i>	
Alcohol	573	559	524	552	576	549	581	633	560	584	604	588	492	487	507	490	531	512	<i>771</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	173	168	162	178	163	<i>378</i>	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																				
Marijuana	170	171	198	191	194	204	218	233	279	266	276	316	269	277	272	306	316	353	<i>582</i>	
Cocaine	27	30	48	45	75	41	37	39	47	25	36	55	26	32	43	38	28	38	<i>34</i>	
Crack	13	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	<i>*</i>
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	39	*	*	*	<i>*</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	14	6	11	15	6	<i>31</i>	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	10	10	*	16	<i>*</i>	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	43	*	49	57	57	<i>158</i>	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	54	57	48	61	62	<i>72</i>	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	26	28	28	21	25	<i>41</i>	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	24	30	20	29	30	<i>44</i>	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	<i>*</i>	
Cigarettes (Daily Only) <sup>2</sup>	773	710	689	660	688	725	728	723	734	707	739	705	577	577	586	560	558	553	<i>898</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	219	207	200	202	203	<i>267</i>	
Cigars	121	145	136	142	143	148	151	146	165	154	177	156	113	158	150	140	155	139	<i>323</i>	
Alcohol	618	616	561	604	617	589	628	693	607	623	654	643	547	528	532	525	581	558	<i>812</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	175	169	163	180	164	<i>381</i>	
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																				
	538	524	506	489	481	509	511	502	485	441	455	465	366	366	379	369	342	348	<i>568</i>	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.28D Daily or Almost Daily Use of Specific Substances in Past Month: Among All People Aged 18 or Older and among Past Month Users of Specific Substances Aged 18 or Older; Standard Errors of Percentages, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																			
Marijuana	0.08	0.08	0.09	0.09	0.09	0.09	0.10	0.10	0.12	0.11	0.11	0.13	0.11	0.11	0.11	0.12	0.12	0.13	<i>0.22</i>
Cocaine	0.01	0.01	0.02	0.02	0.03	0.02	0.02	0.02	0.02	0.01	0.02	0.02	0.01	0.01	0.02	0.02	0.01	0.02	<i>0.01</i>
Crack	0.01	0.01	0.02	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	<i>0.01</i>
Heroin	0.02	0.01	0.01	0.01	0.03	0.01	0.01	0.01	0.01	0.02	0.02	0.01	0.01	0.01	0.02	0.02	0.01	0.02	<i>0.05</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	*	0.00	0.01	0.00	<i>0.01</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.00	0.00	0.01	0.01	<i>0.03</i>
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.02	0.02	0.02	0.02	0.02	<i>0.06</i>
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.02	0.02	0.02	0.02	0.02	<i>0.03</i>
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.01	0.01	0.01	0.01	0.01	<i>0.02</i>
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.01	0.01	0.01	0.01	0.01	<i>0.02</i>
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	0.01	0.01	0.01	*	0.01	<i>0.01</i>
Cigarettes (Daily Only) <sup>2</sup>	0.33	0.30	0.30	0.29	0.29	0.29	0.30	0.29	0.29	0.29	0.29	0.28	0.22	0.22	0.22	0.21	0.21	0.21	<i>0.34</i>
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	0.09	0.09	0.08	0.08	0.08	<i>0.11</i>
Cigars	0.06	0.07	0.06	0.07	0.06	0.07	0.07	0.06	0.07	0.07	0.08	0.07	0.05	0.06	0.06	0.06	0.06	0.06	<i>0.13</i>
Alcohol	0.27	0.26	0.24	0.25	0.26	0.25	0.26	0.28	0.24	0.25	0.26	0.25	0.20	0.20	0.21	0.20	0.21	0.20	<i>0.31</i>
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.07	0.07	0.07	0.07	0.07	<i>0.15</i>
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																			
Marijuana	1.09	1.11	1.17	1.18	1.17	1.22	1.29	1.18	1.28	1.20	1.14	1.23	0.92	0.96	0.87	0.94	0.91	0.84	<i>1.28</i>
Cocaine	1.40	1.36	2.30	1.89	2.91	1.94	2.00	2.40	3.08	1.89	2.11	3.37	1.72	1.68	2.17	1.70	1.43	1.82	<i>1.81</i>
Crack	2.42	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	5.92	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.24	0.48	0.87	0.97	0.32	<i>1.84</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	2.17	2.43	*	2.64	*
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	4.11	*	5.00	4.71	4.29	<i>5.99</i>
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	1.44	1.70	1.52	2.06	2.17	<i>2.85</i>
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	1.63	1.62	1.59	1.31	1.68	<i>2.75</i>
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	1.38	1.57	1.22	1.74	1.67	<i>2.30</i>
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	0.70	0.70	0.66	0.65	0.61	0.68	0.72	0.70	0.72	0.72	0.73	0.74	0.61	0.65	0.66	0.70	0.72	0.72	<i>1.26</i>
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	1.42	1.43	1.46	1.48	1.51	<i>2.84</i>
Cigars	0.97	1.12	1.01	1.02	1.02	1.12	1.14	1.09	1.16	1.15	1.25	1.18	0.91	1.14	1.08	1.06	1.17	1.12	<i>2.57</i>
Alcohol	0.45	0.46	0.42	0.42	0.44	0.42	0.43	0.47	0.41	0.42	0.43	0.41	0.34	0.33	0.35	0.33	0.36	0.35	<i>0.53</i>
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.26	0.25	0.25	0.26	0.25	<i>0.60</i>
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																			
	0.93	0.83	0.88	0.88	0.86	0.89	0.95	1.00	0.94	0.91	0.95	1.01	0.84	0.88	0.90	0.92	0.94	0.95	<i>1.78</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.29C Daily or Almost Daily Use of Specific Substances in Past Month: Among All People Aged 18 to 25 and among Past Month Users of Specific Substances Aged 18 to 25; Standard Errors of Numbers in Thousands, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																				
Marijuana	63	71	74	75	77	74	78	81	86	88	93	93	107	105	102	107	105	100	<i>194</i>	
Cocaine	6	7	11	7	12	7	7	10	7	5	9	4	4	7	6	7	5	9	*	
Crack	3	3	5	*	6	7	3	9	6	*	3	*	2	4	*	3	2	4	*	
Heroin	4	4	4	7	5	7	7	4	10	14	16	12	11	13	10	17	8	9	<i>14</i>	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	6	2	5	4	4	6	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	4	7	4	*	*	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	12	7	11	9	12	<i>12</i>	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	14	9	14	10	8	6	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	8	4	8	5	7	*	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	10	7	8	7	8	<i>20</i>	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	5	*	2	*	*	
Cigarettes (Daily Only) <sup>2</sup>	122	125	126	119	124	123	121	118	116	125	124	109	119	119	106	101	87	85	<i>128</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	53	53	47	42	48	<i>58</i>	
Cigars	24	28	28	30	29	32	29	29	36	33	36	31	34	36	36	39	38	36	<i>68</i>	
Alcohol	62	67	71	68	77	70	70	76	74	71	69	70	75	65	68	67	71	68	<i>126</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	28	26	25	26	26	<i>52</i>	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																				
Marijuana	65	74	79	80	82	79	81	87	93	95	104	99	114	115	111	115	122	112	<i>205</i>	
Cocaine	6	7	11	7	12	7	7	10	7	5	9	4	4	7	6	7	5	9	*	
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	6	2	5	4	4	*	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*	*	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	14	9	14	10	8	*	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	8	4	8	5	7	*	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	10	7	8	7	8	*	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*	
Cigarettes (Daily Only) <sup>2</sup>	135	144	146	135	134	135	129	130	120	134	129	115	127	130	109	105	92	85	<i>129</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	54	54	47	42	48	<i>58</i>	
Cigars	24	28	28	30	29	33	29	30	37	33	36	31	34	37	37	39	38	37	<i>69</i>	
Alcohol	67	69	77	75	82	76	77	82	79	74	73	74	79	68	70	69	75	69	<i>131</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	29	26	25	27	26	<i>52</i>	
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																				
	79	81	76	77	74	71	67	64	60	62	54	49	57	53	57	51	42	41	<i>51</i>	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.29D Daily or Almost Daily Use of Specific Substances in Past Month: Among All People Aged 18 to 25 and among Past Month Users of Specific Substances Aged 18 to 25; Standard Errors of Percentages, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																				
Marijuana	0.20	0.22	0.23	0.23	0.24	0.23	0.24	0.24	0.25	0.26	0.27	0.27	0.31	0.30	0.29	0.31	0.31	0.30	<i>0.58</i>	
Cocaine	0.02	0.02	0.03	0.02	0.04	0.02	0.02	0.03	0.02	0.01	0.03	0.01	0.01	0.02	0.02	0.02	0.02	0.03	*	
Crack	0.01	0.01	0.02	*	0.02	0.02	0.01	0.03	0.02	*	0.01	*	0.01	0.01	*	0.01	0.01	0.01	*	
Heroin	0.01	0.01	0.01	0.02	0.01	0.02	0.02	0.01	0.03	0.04	0.04	0.04	0.03	0.04	0.03	0.05	0.02	0.03	<i>0.04</i>	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.02	0.01	0.01	0.01	0.01	<i>0.02</i>	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.01	0.02	0.01	*	*	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.03	0.02	0.03	0.03	0.04	<i>0.04</i>	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.04	0.03	0.04	0.03	0.02	<i>0.02</i>	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.02	0.01	0.02	0.01	0.02	*	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.03	0.02	0.02	0.02	0.02	<i>0.06</i>	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	0.01	*	0.01	*	*	
Cigarettes (Daily Only) <sup>2</sup>	0.39	0.40	0.39	0.37	0.38	0.38	0.37	0.35	0.34	0.36	0.36	0.31	0.34	0.34	0.31	0.29	0.25	0.25	<i>0.38</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	0.15	0.15	0.14	0.12	0.14	<i>0.17</i>	
Cigars	0.08	0.09	0.09	0.09	0.09	0.10	0.09	0.09	0.11	0.10	0.10	0.09	0.10	0.10	0.11	0.11	0.11	0.11	<i>0.20</i>	
Alcohol	0.20	0.21	0.22	0.21	0.23	0.21	0.21	0.23	0.22	0.21	0.20	0.20	0.22	0.19	0.20	0.20	0.21	0.20	<i>0.38</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.08	0.07	0.07	0.08	0.08	<i>0.15</i>	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																				
Marijuana	0.94	1.04	1.06	1.07	1.11	1.13	1.11	1.05	1.09	1.05	1.06	1.09	1.19	1.19	1.10	1.14	1.07	1.06	<i>1.85</i>	
Cocaine	0.99	0.99	1.55	0.78	1.64	1.22	1.40	2.03	1.32	1.00	2.18	1.06	0.84	1.12	1.10	1.08	1.01	1.62	*	
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.00	0.33	0.80	0.71	0.69	*	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*	*	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	1.66	1.35	2.18	2.02	1.83	*	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	1.01	0.58	1.10	0.85	1.15	*	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	1.63	1.29	1.45	1.84	2.45	*	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*	
Cigarettes (Daily Only) <sup>2</sup>	0.72	0.69	0.72	0.73	0.77	0.76	0.81	0.80	0.80	0.86	0.88	0.83	0.91	1.02	1.03	1.02	1.11	1.13	<i>2.24</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	2.00	2.05	2.25	2.21	2.13	<i>4.18</i>	
Cigars	0.66	0.74	0.66	0.74	0.71	0.81	0.73	0.73	0.89	0.84	0.92	0.86	0.95	1.09	1.09	1.11	1.21	1.27	<i>3.13</i>	
Alcohol	0.32	0.34	0.34	0.33	0.36	0.33	0.33	0.35	0.34	0.33	0.32	0.32	0.35	0.31	0.34	0.34	0.36	0.37	<i>0.70</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.21	0.19	0.20	0.22	0.23	<i>0.48</i>	
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																				
	0.93	0.88	0.86	0.93	0.93	0.92	0.91	0.92	0.94	0.97	0.90	0.90	1.16	1.11	1.42	1.45	1.49	1.60	<i>3.03</i>	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.30C Daily or Almost Daily Use of Specific Substances in Past Month: Among All People Aged 26 or Older and among Past Month Users of Specific Substances Aged 26 or Older; Standard Errors of Numbers in Thousands, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																				
Marijuana	153	157	178	176	172	185	199	210	249	246	241	286	228	238	240	273	283	310	<i>513</i>	
Cocaine	26	29	47	44	74	40	36	38	47	25	35	55	25	31	42	37	27	37	<i>34</i>	
Crack	13	26	33	28	48	29	27	29	17	14	29	35	20	22	35	19	21	29	<i>25</i>	
Heroin	37	21	30	9	70	29	26	18	20	40	40	30	31	27	40	34	26	37	<i>102</i>	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	*	10	14	*	<i>31</i>	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	9	7	18	16	<i>78</i>	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	42	48	48	56	56	<i>156</i>	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	52	56	45	60	62	<i>71</i>	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	25	27	27	20	24	<i>41</i>	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	21	30	18	28	29	<i>39</i>	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	14	23	17	*	13	<i>29</i>	
Cigarettes (Daily Only) <sup>2</sup>	681	618	611	603	618	630	662	632	635	650	662	648	502	524	520	501	520	501	<i>836</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	207	203	196	194	193	<i>259</i>	
Cigars	117	140	131	140	139	144	151	144	158	150	172	151	107	150	147	132	147	136	<i>310</i>	
Alcohol	568	549	521	545	564	546	579	622	555	577	596	583	483	473	494	485	522	505	<i>762</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	169	165	162	176	160	<i>374</i>	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																				
Marijuana	154	157	178	174	174	185	202	210	253	247	246	292	235	243	243	279	284	322	<i>532</i>	
Cocaine	26	29	47	45	75	40	36	38	47	25	35	*	25	31	42	37	27	37	<i>34</i>	
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	<i>*</i>
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	<i>*</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	6	10	14	4	<i>31</i>	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*	<i>*</i>	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	41	*	48	56	56	<i>*</i>	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	52	56	45	60	62	<i>71</i>	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	25	27	27	20	24	<i>*</i>	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	22	30	18	28	29	<i>39</i>	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	<i>*</i>	
Cigarettes (Daily Only) <sup>2</sup>	738	678	649	632	672	696	705	696	709	681	709	691	545	546	564	537	544	538	<i>887</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	212	201	197	195	193	<i>259</i>	
Cigars	118	141	132	140	139	144	151	143	159	149	172	152	108	150	147	133	148	137	<i>312</i>	
Alcohol	616	608	557	598	605	584	628	687	600	615	648	638	541	517	524	524	574	553	<i>804</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	171	166	163	178	161	<i>377</i>	
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																				
	528	508	491	476	473	500	501	495	479	435	447	458	355	356	370	362	338	344	<i>563</i>	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.30D Daily or Almost Daily Use of Specific Substances in Past Month: Among All People Aged 26 or Older and among Past Month Users of Specific Substances Aged 26 or Older; Standard Errors of Percentages, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																				
Marijuana	0.09	0.09	0.10	0.09	0.09	0.10	0.10	0.11	0.13	0.12	0.12	0.14	0.11	0.11	0.11	0.13	0.13	0.14	<i>0.24</i>	
Cocaine	0.01	0.02	0.03	0.02	0.04	0.02	0.02	0.02	0.02	0.01	0.02	0.03	0.01	0.01	0.02	0.02	0.01	0.02	<i>0.02</i>	
Crack	0.01	0.01	0.02	0.01	0.03	0.02	0.01	0.02	0.01	0.01	0.01	0.02	0.01	0.01	0.02	0.01	0.01	0.01	<i>0.01</i>	
Heroin	0.02	0.01	0.02	0.00	0.04	0.02	0.01	0.01	0.01	0.02	0.02	0.01	0.02	0.01	0.02	0.02	0.01	0.02	<i>0.05</i>	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	*	0.00	0.01	*	<i>0.01</i>	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.00	0.00	0.01	0.01	<i>0.04</i>	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.02	0.02	0.02	0.03	0.03	<i>0.07</i>	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.02	0.03	0.02	0.03	0.03	<i>0.03</i>	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.01	0.01	0.01	0.01	0.01	<i>0.02</i>	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.01	0.01	0.01	0.01	0.01	<i>0.02</i>	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	0.01	0.01	0.01	*	0.01	<i>0.01</i>	
Cigarettes (Daily Only) <sup>2</sup>	0.38	0.34	0.33	0.33	0.33	0.33	0.35	0.33	0.33	0.33	0.33	0.32	0.24	0.25	0.25	0.24	0.24	0.23	<i>0.38</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	0.10	0.10	0.09	0.09	0.09	<i>0.12</i>	
Cigars	0.07	0.08	0.07	0.08	0.07	0.08	0.08	0.07	0.08	0.08	0.09	0.07	0.05	0.07	0.07	0.06	0.07	0.06	<i>0.14</i>	
Alcohol	0.32	0.30	0.28	0.29	0.30	0.29	0.30	0.32	0.28	0.29	0.30	0.29	0.24	0.23	0.24	0.23	0.24	0.23	<i>0.35</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.08	0.08	0.08	0.08	0.07	<i>0.17</i>	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																				
Marijuana	1.77	1.79	1.86	1.89	1.80	1.98	1.99	1.86	2.00	1.86	1.72	1.80	1.25	1.29	1.21	1.23	1.24	1.04	<i>1.58</i>	
Cocaine	2.05	1.98	3.48	2.94	4.09	2.68	2.74	3.30	4.64	2.88	2.71	*	2.50	2.40	3.04	2.40	1.90	2.46	<i>2.37</i>	
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	0.94	1.46	1.53	0.33	<i>2.89</i>	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*	*	*
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	4.63	*	5.86	5.14	4.57	*	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	1.84	2.11	1.81	2.44	2.52	<i>3.21</i>	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	3.02	2.92	2.58	2.03	2.72	*	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	1.86	2.17	1.71	2.24	1.99	<i>2.53</i>	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*	
Cigarettes (Daily Only) <sup>2</sup>	0.87	0.86	0.80	0.79	0.74	0.83	0.86	0.84	0.86	0.88	0.85	0.88	0.72	0.76	0.75	0.79	0.79	0.79	<i>1.36</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	1.69	1.72	1.77	1.73	1.76	<i>3.34</i>	
Cigars	1.35	1.58	1.44	1.43	1.43	1.60	1.62	1.54	1.62	1.62	1.70	1.64	1.23	1.46	1.42	1.36	1.48	1.44	<i>3.05</i>	
Alcohol	0.54	0.54	0.50	0.49	0.52	0.50	0.51	0.55	0.49	0.49	0.51	0.49	0.39	0.38	0.40	0.37	0.40	0.40	<i>0.60</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.32	0.31	0.30	0.32	0.30	<i>0.72</i>	
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																				
	1.12	0.99	1.05	1.02	1.00	1.06	1.10	1.16	1.09	1.05	1.09	1.15	0.93	0.98	1.00	1.02	1.03	1.01	<i>1.86</i>	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.31C Daily or Almost Daily Use of Specific Substances in Past Month: Among All People Aged 12 to 20 and among Past Month Users of Specific Substances Aged 12 to 20; Standard Errors of Numbers in Thousands, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																				
Marijuana	59	59	56	62	65	58	61	63	66	66	77	66	74	71	69	71	73	74	<i>124</i>	
Cocaine	5	6	9	6	5	6	6	6	6	4	4	3	6	6	5	2	3	4	*	
Crack	4	4	4	*	*	4	*	4	6	*	3	*	*	4	*	2	*	4	*	
Heroin	4	4	4	3	2	4	4	3	5	10	6	9	8	3	2	*	3	4	*	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	6	3	2	4	2	*	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	4	6	7	3	8	4	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	8	*	8	6	6	9	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	5	6	8	6	3	*	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	7	5	7	2	6	3	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	8	7	4	8	3	20	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*	
Cigarettes (Daily Only) <sup>2</sup>	85	87	87	85	86	83	80	75	75	75	66	63	68	66	55	46	44	40	65	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	36	34	27	25	27	37	
Cigars	23	22	22	25	19	21	21	25	25	24	19	22	22	23	24	27	24	26	51	
Alcohol	35	38	43	41	39	38	36	40	40	32	31	32	36	30	26	29	26	26	33	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	15	10	13	11	9	13	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																				
Marijuana	59	59	56	62	65	58	61	63	66	66	77	66	74	71	69	71	73	74	<i>124</i>	
Cocaine	5	6	9	6	5	6	6	6	6	4	4	*	*	6	*	2	*	*	*	
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	6	*	2	4	2	*	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	4	6	7	3	8	*	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	5	6	8	6	3	*	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	7	5	7	*	6	*	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	8	7	4	8	3	*	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*	
Cigarettes (Daily Only) <sup>2</sup>	85	87	87	85	86	83	80	75	75	75	66	63	68	66	55	46	44	40	65	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	36	34	27	25	27	*	
Cigars	23	22	22	25	19	21	21	25	25	24	19	22	22	23	24	27	24	26	*	
Alcohol	35	38	43	41	39	38	36	40	40	32	31	32	36	30	26	29	26	26	33	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	15	10	13	11	9	13	
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																				
	44	42	43	45	43	38	39	33	35	36	25	24	29	21	28	25	19	14	*	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2002 to 2019 standard errors of numbers in thousands may differ from previously published estimates because of a change to the standard error estimation method. See the Introduction to these tables for more details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.31D Daily or Almost Daily Use of Specific Substances in Past Month: Among All People Aged 12 to 20 and among Past Month Users of Specific Substances Aged 12 to 20; Standard Errors of Percentages, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																			
Marijuana	0.15	0.16	0.14	0.16	0.16	0.15	0.15	0.16	0.17	0.16	0.19	0.17	0.19	0.18	0.18	0.18	0.19	0.18	<i>0.33</i>
Cocaine	0.01	0.02	0.02	0.01	0.01	0.02	0.02	0.01	0.02	0.01	0.01	0.01	0.02	0.02	0.01	0.01	0.01	0.01	*
Crack	0.01	0.01	0.01	*	*	0.01	*	0.01	0.01	*	0.01	*	*	0.01	*	0.01	*	0.01	*
Heroin	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.03	0.02	0.02	0.02	0.01	0.01	*	0.01	0.01	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.01	0.01	0.01	0.00	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.02	0.02	0.01	0.02	<i>0.01</i>
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.02	*	0.02	0.02	0.02	<i>0.03</i>
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.01	0.02	0.02	0.02	0.01	*
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.02	0.01	0.02	0.01	0.01	<i>0.01</i>
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.02	0.02	0.01	0.02	0.01	<i>0.05</i>
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	0.22	0.22	0.23	0.21	0.22	0.20	0.20	0.19	0.19	0.19	0.17	0.16	0.18	0.17	0.15	0.12	0.11	0.11	<i>0.17</i>
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	0.09	0.09	0.07	0.07	0.07	<i>0.10</i>
Cigars	0.06	0.06	0.06	0.06	0.05	0.06	0.05	0.06	0.07	0.06	0.05	0.06	0.06	0.06	0.06	0.07	0.06	0.07	<i>0.14</i>
Alcohol	0.09	0.10	0.11	0.10	0.10	0.10	0.09	0.10	0.10	0.08	0.08	0.08	0.09	0.08	0.07	0.08	0.07	0.07	<i>0.09</i>
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.04	0.03	0.03	0.03	0.02	<i>0.03</i>
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																			
Marijuana	1.02	1.01	0.99	1.14	1.16	1.13	1.10	1.07	1.16	1.04	1.15	1.15	1.27	1.24	1.30	1.32	1.24	1.17	<i>2.41</i>
Cocaine	1.36	1.48	2.27	1.26	1.21	1.71	1.95	2.52	2.58	1.49	2.32	*	*	2.56	*	1.15	*	*	*
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.47	*	0.62	1.17	0.42	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.61	2.95	3.19	1.13	2.89	*
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.93	1.32	2.05	1.67	0.92	*
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	1.66	1.30	1.83	*	1.84	*
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	1.95	1.96	1.32	3.27	1.46	*
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	0.82	0.83	0.85	0.86	0.90	0.88	0.96	0.84	0.97	1.00	1.04	1.07	1.24	1.41	1.49	1.31	1.46	1.63	<i>3.66</i>
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	2.71	2.72	2.47	2.54	2.83	*
Cigars	0.78	0.76	0.69	0.81	0.68	0.70	0.71	0.84	0.95	0.88	0.83	1.01	1.12	1.24	1.38	1.49	1.46	1.85	*
Alcohol	0.31	0.34	0.36	0.36	0.34	0.33	0.34	0.36	0.39	0.32	0.32	0.36	0.40	0.37	0.35	0.38	0.35	0.36	<i>0.52</i>
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.29	0.23	0.27	0.26	0.22	<i>0.37</i>
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																			
	1.11	1.11	1.17	1.29	1.31	1.25	1.34	1.25	1.36	1.43	1.29	1.40	1.75	1.70	2.68	2.67	2.59	2.43	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.32C Daily or Almost Daily Use of Specific Substances in Past Month: Among All People Aged 21 or Older and among Past Month Users of Specific Substances Aged 21 or Older; Standard Errors of Numbers in Thousands, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																				
Marijuana	166	165	189	185	185	197	212	221	267	259	265	307	255	263	258	298	303	343	<i>569</i>	
Cocaine	26	29	48	45	75	40	36	39	47	25	36	55	25	31	43	38	28	38	<i>34</i>	
Crack	13	26	33	28	48	30	27	31	17	14	29	35	20	22	35	19	21	29	<i>25</i>	
Heroin	37	21	31	11	70	29	27	18	22	41	42	32	32	30	41	39	28	38	<i>114</i>	
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	6	11	14	5	<i>31</i>	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	10	8	19	16	<i>78</i>	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	42	49	48	56	57	<i>157</i>	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	54	56	47	61	62	<i>72</i>	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	26	27	27	21	25	<i>41</i>	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	24	30	20	28	30	<i>39</i>	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	14	24	17	*	13	<i>29</i>	
Cigarettes (Daily Only) <sup>2</sup>	758	702	680	654	682	712	716	714	726	700	727	703	565	566	579	554	556	549	<i>897</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	218	205	198	201	201	<i>263</i>	
Cigars	119	143	134	142	141	147	151	144	162	152	175	154	111	155	149	138	151	138	<i>318</i>	
Alcohol	619	614	562	603	612	588	626	693	604	620	653	640	547	527	531	525	580	557	<i>813</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	175	169	163	180	164	<i>380</i>	
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																				
Marijuana	166	165	189	185	185	197	212	221	267	259	265	307	255	263	258	298	303	343	<i>569</i>	
Cocaine	26	29	48	45	75	40	36	39	47	25	36	55	25	31	43	38	28	38	<i>34</i>	
Crack	13	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	<i>*</i>
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	39	*	*	*	<i>*</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	13	6	11	14	5	<i>31</i>	
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*	<i>*</i>	
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	42	*	48	56	57	<i>*</i>	
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	54	56	47	61	62	<i>72</i>	
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	26	27	27	21	25	<i>41</i>	
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	24	30	20	28	30	<i>39</i>	
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	<i>*</i>	
Cigarettes (Daily Only) <sup>2</sup>	758	702	680	654	682	712	716	714	726	700	727	703	565	566	579	554	556	549	<i>897</i>	
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	218	205	198	201	201	<i>263</i>	
Cigars	119	143	134	142	141	147	151	144	162	152	175	154	111	155	149	138	151	138	<i>318</i>	
Alcohol	619	614	562	603	612	588	626	693	604	620	653	640	547	527	531	525	580	557	<i>813</i>	
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	175	169	163	180	164	<i>380</i>	
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>	535	522	504	485	480	505	505	501	483	439	452	464	363	364	378	367	341	347	<i>568</i>	

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2002 to 2019 standard errors of numbers in thousands may differ from previously published estimates because of a change to the standard error estimation method. See the Introduction to these tables for more details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

**Table 7.32D Daily or Almost Daily Use of Specific Substances in Past Month: Among All People Aged 21 or Older and among Past Month Users of Specific Substances Aged 21 or Older; Standard Errors of Percentages, 2002-2020**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG TOTAL POPULATION</b>																			
Marijuana	0.08	0.08	0.09	0.09	0.09	0.09	0.10	0.10	0.12	0.12	0.12	0.13	0.11	0.11	0.11	0.12	0.13	0.14	<i>0.23</i>
Cocaine	0.01	0.01	0.02	0.02	0.04	0.02	0.02	0.02	0.02	0.01	0.02	0.02	0.01	0.01	0.02	0.02	0.01	0.02	<i>0.01</i>
Crack	0.01	0.01	0.02	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.02	0.01	0.01	0.01	<i>0.01</i>
Heroin	0.02	0.01	0.02	0.01	0.03	0.01	0.01	0.01	0.01	0.02	0.02	0.01	0.01	0.01	0.02	0.02	0.01	0.02	<i>0.05</i>
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.00	0.00	0.01	0.00	<i>0.01</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.00	0.00	0.01	0.01	<i>0.03</i>
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	0.02	0.02	0.02	0.02	0.02	<i>0.07</i>
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	0.02	0.02	0.02	0.03	0.03	<i>0.03</i>
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	0.01	0.01	0.01	0.01	0.01	<i>0.02</i>
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	0.01	0.01	0.01	0.01	0.01	<i>0.02</i>
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	0.01	0.01	0.01	*	0.01	<i>0.01</i>
Cigarettes (Daily Only) <sup>2</sup>	0.35	0.32	0.31	0.30	0.30	0.31	0.32	0.30	0.30	0.30	0.31	0.29	0.23	0.23	0.23	0.22	0.22	0.22	<i>0.35</i>
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	0.09	0.09	0.08	0.08	0.08	<i>0.11</i>
Cigars	0.06	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.08	0.07	0.05	0.07	0.06	0.06	0.06	0.06	<i>0.13</i>
Alcohol	0.29	0.28	0.26	0.27	0.28	0.26	0.27	0.30	0.26	0.26	0.27	0.26	0.22	0.21	0.22	0.21	0.22	0.21	<i>0.32</i>
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.08	0.07	0.07	0.08	0.07	<i>0.16</i>
<b>DAILY/ALMOST DAILY USE IN PAST MONTH AMONG PAST MONTH USERS</b>																			
Marijuana	1.31	1.35	1.38	1.43	1.38	1.47	1.53	1.39	1.52	1.43	1.34	1.41	1.02	1.05	0.96	1.03	1.02	0.90	<i>1.37</i>
Cocaine	1.59	1.55	2.62	2.16	3.36	2.17	2.25	2.64	3.44	2.22	2.29	3.66	1.90	1.83	2.35	1.85	1.56	1.94	<i>1.91</i>
Crack	2.50	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	5.96	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	1.48	0.61	1.05	1.10	0.35	<i>2.19</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*	*
Methamphetamine	--	--	--	--	--	--	--	--	--	--	--	--	--	4.24	*	5.23	4.83	4.35	*
Misuse of Prescription Pain Relievers <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	--	1.54	1.81	1.59	2.19	2.28	<i>2.94</i>
Misuse of Prescription Stimulants	--	--	--	--	--	--	--	--	--	--	--	--	--	1.97	1.97	1.80	1.42	1.91	<i>2.89</i>
Misuse of Prescription Tranquilizers	--	--	--	--	--	--	--	--	--	--	--	--	--	1.56	1.74	1.37	1.89	1.77	<i>2.14</i>
Misuse of Prescription Sedatives	--	--	--	--	--	--	--	--	--	--	--	--	--	*	*	*	*	*	*
Cigarettes (Daily Only) <sup>2</sup>	0.75	0.76	0.71	0.69	0.65	0.72	0.77	0.74	0.77	0.76	0.76	0.78	0.64	0.68	0.69	0.73	0.74	0.73	<i>1.29</i>
Smokeless Tobacco	--	--	--	--	--	--	--	--	--	--	--	--	--	1.51	1.51	1.55	1.56	1.58	<i>2.93</i>
Cigars	1.12	1.29	1.17	1.18	1.16	1.31	1.32	1.25	1.33	1.32	1.41	1.33	1.01	1.24	1.18	1.16	1.26	1.21	<i>2.72</i>
Alcohol	0.48	0.48	0.44	0.44	0.46	0.44	0.45	0.49	0.43	0.44	0.45	0.43	0.35	0.35	0.36	0.34	0.37	0.36	<i>0.54</i>
Binge Alcohol <sup>3</sup>	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.28	0.27	0.26	0.28	0.26	<i>0.63</i>
<b>SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS<sup>4</sup></b>																			
	0.99	0.88	0.94	0.92	0.90	0.94	0.99	1.05	0.99	0.95	0.98	1.05	0.86	0.90	0.92	0.95	0.96	0.96	<i>1.80</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

<sup>1</sup> Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

<sup>2</sup> Includes only respondents who smoked cigarettes on each of the past 30 days.

<sup>3</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

<sup>4</sup> Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.