

Table 8.1C Any Mental Illness in Past Year: Among People Aged 18 or Older; by Gender and Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
TOTAL	631	981	404	583	467	754
18-25	163	282	102	177	117	202
18-20	144	219	76	124	105	171
21-25	165	287	97	163	122	215
26 or Older	599	937	385	558	441	729
26-49	366	552	232	371	266	400
26-29	194	288	120	179	145	220
30-34	213	345	135	224	155	238
35-39	190	298	114	188	148	233
40-44	175	268	102	172	135	208
45-49	160	266	100	181	125	199
50 or Older	460	737	297	431	339	595
50-54	230	370	158	170	171	326
55-59	254	407	164	206	183	347
60-64	207	346	119	242	172	254
65 or Older	295	458	172	276	232	365

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Any Mental Illness (AMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 8.1D Any Mental Illness in Past Year: Among People Aged 18 or Older; by Gender and Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
TOTAL	0.25	<i>0.39</i>	0.33	<i>0.48</i>	0.36	<i>0.58</i>
18-25	0.48	<i>0.84</i>	0.60	<i>1.06</i>	0.70	<i>1.21</i>
18-20	0.78	<i>1.40</i>	0.91	<i>1.67</i>	1.15	<i>2.11</i>
21-25	0.60	<i>1.07</i>	0.79	<i>1.36</i>	0.90	<i>1.54</i>
26 or Older	0.28	<i>0.43</i>	0.37	<i>0.53</i>	0.39	<i>0.64</i>
26-49	0.36	<i>0.54</i>	0.46	<i>0.74</i>	0.52	<i>0.78</i>
26-29	0.87	<i>1.35</i>	1.16	<i>1.80</i>	1.27	<i>1.98</i>
30-34	0.78	<i>1.24</i>	1.04	<i>1.71</i>	1.13	<i>1.73</i>
35-39	0.78	<i>1.16</i>	1.02	<i>1.64</i>	1.12	<i>1.66</i>
40-44	0.80	<i>1.24</i>	0.99	<i>1.63</i>	1.17	<i>1.82</i>
45-49	0.74	<i>1.24</i>	0.92	<i>1.73</i>	1.11	<i>1.80</i>
50 or Older	0.40	<i>0.63</i>	0.55	<i>0.79</i>	0.55	<i>0.96</i>
50-54	1.04	<i>1.64</i>	1.45	<i>1.85</i>	1.47	<i>2.58</i>
55-59	1.08	<i>1.68</i>	1.49	<i>1.82</i>	1.47	<i>2.65</i>
60-64	0.94	<i>1.50</i>	1.18	<i>2.10</i>	1.40	<i>2.21</i>
65 or Older	0.56	<i>0.84</i>	0.73	<i>1.13</i>	0.80	<i>1.22</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Any Mental Illness (AMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 8.4C Serious Mental Illness in Past Year: Among People Aged 18 or Older; by Gender and Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020

Age Category	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
TOTAL	328	<i>571</i>	203	<i>332</i>	257	<i>437</i>
18-25	90	<i>176</i>	57	<i>107</i>	70	<i>130</i>
18-20	63	<i>101</i>	35	<i>68</i>	48	<i>83</i>
21-25	83	<i>159</i>	48	<i>87</i>	64	<i>125</i>
26 or Older	315	<i>525</i>	193	<i>312</i>	250	<i>413</i>
26-49	195	<i>316</i>	117	<i>210</i>	163	<i>231</i>
26-29	98	<i>147</i>	63	<i>91</i>	74	<i>114</i>
30-34	101	<i>158</i>	65	<i>98</i>	76	<i>119</i>
35-39	96	<i>153</i>	57	<i>97</i>	79	<i>111</i>
40-44	88	<i>129</i>	51	<i>79</i>	71	<i>103</i>
45-49	83	<i>154</i>	48	<i>113</i>	66	<i>105</i>
50 or Older	226	<i>405</i>	148	<i>224</i>	177	<i>346</i>
50-54	125	<i>171</i>	90	<i>95</i>	84	<i>142</i>
55-59	127	<i>247</i>	89	<i>92</i>	93	<i>229</i>
60-64	97	<i>208</i>	41	<i>151</i>	85	<i>143</i>
65 or Older	112	<i>200</i>	73	<i>112</i>	86	<i>165</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Serious Mental Illness (SMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of SMI are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 8.4D Serious Mental Illness in Past Year: Among People Aged 18 or Older; by Gender and Detailed Age Category, Standard Errors of Percentages, 2019 and 2020

Age Category	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
TOTAL	0.13	<i>0.23</i>	0.17	<i>0.27</i>	0.20	<i>0.34</i>
18-25	0.27	<i>0.53</i>	0.34	<i>0.64</i>	0.42	<i>0.78</i>
18-20	0.44	<i>0.77</i>	0.50	<i>1.03</i>	0.69	<i>1.26</i>
21-25	0.37	<i>0.68</i>	0.45	<i>0.80</i>	0.56	<i>1.05</i>
26 or Older	0.15	<i>0.24</i>	0.19	<i>0.30</i>	0.22	<i>0.36</i>
26-49	0.19	<i>0.31</i>	0.23	<i>0.42</i>	0.32	<i>0.45</i>
26-29	0.52	<i>0.79</i>	0.67	<i>1.01</i>	0.80	<i>1.21</i>
30-34	0.43	<i>0.66</i>	0.58	<i>0.82</i>	0.65	<i>1.01</i>
35-39	0.44	<i>0.66</i>	0.55	<i>0.89</i>	0.68	<i>0.92</i>
40-44	0.44	<i>0.66</i>	0.53	<i>0.81</i>	0.69	<i>1.03</i>
45-49	0.40	<i>0.77</i>	0.48	<i>1.12</i>	0.62	<i>1.04</i>
50 or Older	0.20	<i>0.35</i>	0.27	<i>0.41</i>	0.29	<i>0.56</i>
50-54	0.59	<i>0.84</i>	0.87	<i>1.06</i>	0.79	<i>1.28</i>
55-59	0.59	<i>1.08</i>	0.85	<i>0.85</i>	0.84	<i>1.93</i>
60-64	0.46	<i>0.97</i>	0.42	<i>1.39</i>	0.77	<i>1.36</i>
65 or Older	0.21	<i>0.37</i>	0.31	<i>0.46</i>	0.30	<i>0.55</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Serious Mental Illness (SMI) aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of SMI are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.