

**Table 8.36C Major Depressive Episode in Past Year: Among People Aged 18 or Older; by Gender and Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020**

Age Category	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
<b>TOTAL</b>	385	674	240	358	310	528
<b>18-25</b>	124	228	73	125	96	171
18-20	102	153	53	87	76	122
21-25	108	203	62	104	85	166
<b>26 or Older</b>	360	616	230	340	289	492
26-49	221	366	138	250	176	263
26-29	115	161	77	106	84	116
30-34	123	194	73	135	94	132
35-39	110	174	58	114	94	119
40-44	100	146	61	105	77	102
45-49	96	181	60	127	73	136
50 or Older	275	479	183	238	214	415
50-54	146	206	100	116	105	174
55-59	137	272	85	115	108	241
60-64	128	213	69	106	104	185
65 or Older	159	286	111	156	120	236

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 8.36D Major Depressive Episode in Past Year: Among People Aged 18 or Older; by Gender and Detailed Age Category, Standard Errors of Percentages, 2019 and 2020**

Age Category	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
<b>TOTAL</b>	0.16	<i>0.27</i>	0.20	<i>0.30</i>	0.24	<i>0.41</i>
<b>18-25</b>	0.37	<i>0.69</i>	0.44	<i>0.76</i>	0.58	<i>1.04</i>
18-20	0.64	<i>1.11</i>	0.69	<i>1.28</i>	1.01	<i>1.75</i>
21-25	0.47	<i>0.84</i>	0.58	<i>0.94</i>	0.73	<i>1.31</i>
<b>26 or Older</b>	0.17	<i>0.28</i>	0.22	<i>0.33</i>	0.26	<i>0.44</i>
26-49	0.22	<i>0.36</i>	0.28	<i>0.50</i>	0.35	<i>0.52</i>
26-29	0.60	<i>0.87</i>	0.81	<i>1.17</i>	0.89	<i>1.27</i>
30-34	0.53	<i>0.82</i>	0.64	<i>1.14</i>	0.80	<i>1.14</i>
35-39	0.49	<i>0.75</i>	0.56	<i>1.04</i>	0.79	<i>1.00</i>
40-44	0.50	<i>0.74</i>	0.63	<i>1.06</i>	0.76	<i>1.02</i>
45-49	0.47	<i>0.88</i>	0.59	<i>1.25</i>	0.70	<i>1.30</i>
50 or Older	0.24	<i>0.41</i>	0.34	<i>0.44</i>	0.35	<i>0.68</i>
50-54	0.70	<i>1.02</i>	0.97	<i>1.29</i>	0.98	<i>1.57</i>
55-59	0.64	<i>1.20</i>	0.84	<i>1.07</i>	0.97	<i>2.06</i>
60-64	0.60	<i>1.00</i>	0.70	<i>0.99</i>	0.92	<i>1.74</i>
65 or Older	0.30	<i>0.53</i>	0.47	<i>0.64</i>	0.42	<i>0.80</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 8.37C Major Depressive Episode with Severe Impairment in Past Year: Among People Aged 18 or Older; by Gender and Detailed Age Category, Standard Errors of Numbers in Thousands, 2019 and 2020**

Age Category	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
<b>TOTAL</b>	331	578	200	317	259	448
<b>18-25</b>	101	190	60	107	79	151
18-20	73	119	39	73	57	98
21-25	86	170	50	88	69	141
<b>26 or Older</b>	316	518	189	297	246	407
26-49	184	320	116	210	148	235
26-29	95	137	61	85	70	104
30-34	91	154	54	98	75	116
35-39	91	152	50	108	76	99
40-44	85	113	53	73	64	85
45-49	81	165	49	113	63	119
50 or Older	236	400	146	219	184	339
50-54	114	183	79	112	83	146
55-59	119	215	74	111	90	178
60-64	108	188	52	100	92	159
65 or Older	134	242	87	136	103	199

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 8.37D Major Depressive Episode with Severe Impairment in Past Year: Among People Aged 18 or Older; by Gender and Detailed Age Category, Standard Errors of Percentages, 2019 and 2020**

Age Category	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
<b>TOTAL</b>	0.13	<i>0.23</i>	0.17	<i>0.26</i>	0.20	<i>0.35</i>
<b>18-25</b>	0.30	<i>0.57</i>	0.36	<i>0.65</i>	0.48	<i>0.92</i>
18-20	0.50	<i>0.90</i>	0.56	<i>1.11</i>	0.80	<i>1.46</i>
21-25	0.39	<i>0.74</i>	0.48	<i>0.81</i>	0.62	<i>1.17</i>
<b>26 or Older</b>	0.15	<i>0.24</i>	0.18	<i>0.29</i>	0.22	<i>0.36</i>
26-49	0.18	<i>0.32</i>	0.23	<i>0.42</i>	0.29	<i>0.46</i>
26-29	0.51	<i>0.76</i>	0.65	<i>0.96</i>	0.77	<i>1.16</i>
30-34	0.40	<i>0.66</i>	0.49	<i>0.84</i>	0.65	<i>1.00</i>
35-39	0.42	<i>0.67</i>	0.48	<i>1.00</i>	0.66	<i>0.84</i>
40-44	0.43	<i>0.58</i>	0.56	<i>0.74</i>	0.64	<i>0.86</i>
45-49	0.40	<i>0.82</i>	0.49	<i>1.13</i>	0.61	<i>1.15</i>
50 or Older	0.21	<i>0.35</i>	0.27	<i>0.40</i>	0.30	<i>0.55</i>
50-54	0.56	<i>0.91</i>	0.78	<i>1.24</i>	0.79	<i>1.35</i>
55-59	0.56	<i>0.98</i>	0.73	<i>1.03</i>	0.82	<i>1.58</i>
60-64	0.51	<i>0.89</i>	0.54	<i>0.94</i>	0.82	<i>1.52</i>
65 or Older	0.26	<i>0.45</i>	0.37	<i>0.56</i>	0.36	<i>0.67</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.