

Table 8.42A Substance Use in Past Year and Past Month: Among People Aged 18 or Older; by Past Year Level of Mental Illness and Past Year Major Depressive Episode (MDE) and MDE with Severe Impairment Status, Numbers in Thousands, 2019 and 2020

Substance	Total		Any Mental Illness ¹		Serious Mental Illness ¹		Any Mental Illness Excluding Serious Mental Illness ¹		No Mental Illness ¹		MDE ²		MDE with Severe Impairment ^{2,3}		No MDE ²	
	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)
PAST YEAR USE																
Illicit Drugs	52,924	54,982	19,998	21,062	6,475	6,786	13,524	14,276	32,926	33,920	8,555	9,220	6,112	6,671	43,776	45,178
Marijuana	44,958	46,338	16,741	17,358	5,225	5,572	11,516	11,786	28,217	28,980	7,200	7,787	5,080	5,613	37,318	38,046
Cocaine	5,371	5,007	2,523	2,330	974	1,027	1,549	1,303	2,848	2,677	1,148	1,266	809	866	4,179	3,724
Crack	766	621	477	477	287	187	190	291	289	143	256	180	225	156	510	440
Heroin	745	719	423	602	260	251	163	350	322	117	223	175	182	149	511	542
Hallucinogens	5,571	6,614	2,616	3,289	863	1,375	1,753	1,915	2,954	3,325	1,223	1,709	854	1,175	4,252	4,848
LSD	2,191	2,364	1,069	1,303	326	627	743	675	1,122	1,061	529	816	341	572	1,613	1,510
PCP	52	94	17	84	10	60	7	24	35	10	8	64	7	53	31	30
Ecstasy	2,395	2,464	1,114	1,134	344	577	770	556	1,281	1,331	496	718	341	480	1,866	1,744
Inhalants	1,399	1,712	696	865	234	452	462	413	703	847	357	529	259	368	1,003	1,162
Methamphetamine	1,958	2,365	1,150	1,568	534	745	616	823	808	796	471	792	418	648	1,462	1,505
Misuse of Prescription Psychotherapeutics	15,242	15,218	7,347	7,364	2,799	2,646	4,548	4,719	7,895	7,854	3,329	3,227	2,465	2,483	11,706	11,884
Pain Relievers	9,158	8,567	4,396	4,127	1,746	1,577	2,651	2,550	4,761	4,440	2,017	1,761	1,514	1,390	7,031	6,747
Stimulants	4,499	4,830	2,259	2,203	885	870	1,374	1,333	2,240	2,626	1,101	1,128	785	801	3,358	3,673
Tranquilizers or Sedatives	5,459	5,971	3,175	3,751	1,366	1,509	1,809	2,242	2,285	2,220	1,530	1,810	1,204	1,444	3,834	4,140
Tranquilizers	4,829	5,238	2,864	3,304	1,272	1,363	1,592	1,941	1,965	1,934	1,417	1,629	1,108	1,323	3,318	3,588
Sedatives	1,016	1,253	574	862	250	417	324	445	442	391	282	461	223	391	728	779
Benzodiazepines	4,460	4,646	2,639	2,999	1,167	1,193	1,472	1,806	1,821	1,647	1,338	1,517	1,054	1,225	3,037	3,108
Opioids	9,499	8,789	4,534	4,292	1,813	1,645	2,720	2,647	4,965	4,497	2,066	1,787	1,550	1,416	7,318	6,943
Central Nervous System Stimulants	9,638	9,801	4,704	4,658	1,844	1,971	2,860	2,687	4,933	5,143	2,129	2,342	1,542	1,708	7,407	7,357
Illicit Drugs Other Than Marijuana ⁴	21,727	22,174	9,969	10,604	3,696	3,960	6,273	6,644	11,758	11,570	4,456	4,804	3,269	3,559	16,960	17,159
PAST MONTH USE																
Alcohol	137,381	136,476	29,216	29,505	7,399	7,665	21,817	21,841	108,165	106,971	11,058	11,566	7,170	8,035	124,902	124,020
Binge Alcohol Use ⁵	64,633	60,376	15,895	15,084	4,283	4,388	11,612	10,697	48,738	45,291	6,205	6,165	3,970	4,340	57,705	53,823
Heavy Alcohol Use	15,835	17,364	4,436	4,629	1,297	1,445	3,138	3,184	11,399	12,735	1,808	1,922	1,107	1,359	13,853	15,287
Cigarette Use	45,313	41,051	13,991	12,225	4,322	3,900	9,669	8,326	31,322	28,825	5,328	4,961	3,866	3,566	39,260	35,500
Daily Cigarette Use	26,721	25,032	8,283	7,304	2,560	2,306	5,724	4,998	18,437	17,728	3,164	2,856	2,334	2,040	23,173	21,871

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. Therefore, some estimates in this table could differ from corresponding estimates in other tables that did not undergo sample size adjustments. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

² Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

³ Respondents with unknown impairment data were excluded.

⁴ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

⁵ Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 8.42B Substance Use in Past Year and Past Month: Among People Aged 18 or Older; by Past Year Level of Mental Illness and Past Year Major Depressive Episode (MDE) and MDE with Severe Impairment Status, Percentages, 2019 and 2020

Substance	Total (2019)	Total (2020)	Any Mental Illness ¹ (2019)	Any Mental Illness ¹ (2020)	Serious Mental Illness ¹ (2019)	Serious Mental Illness ¹ (2020)	Any Mental Illness Excluding Serious Mental Illness ¹	Any Mental Illness Excluding Serious Mental Illness ¹	No Mental Illness ¹	No Mental Illness ¹	MDE ² (2019)	MDE ² (2020)	MDE with Severe Impairment ^{2,3}	MDE with Severe Impairment ^{2,3}	No MDE ²	No MDE ²
							(2019)	(2020)	(2019)	(2020)			(2019)	(2020)	(2019)	(2020)
PAST YEAR USE																
Illicit Drugs	21.1	<i>21.8</i>	38.8	<i>39.8</i>	49.4	<i>47.8</i>	35.2	<i>36.9</i>	16.6	<i>17.0</i>	44.2	<i>44.0</i>	46.6	<i>44.9</i>	19.2	<i>19.8</i>
Marijuana	18.0	<i>18.4</i>	32.5	<i>32.8</i>	39.8	<i>39.2</i>	30.0	<i>30.5</i>	14.2	<i>14.6</i>	37.2	<i>37.1</i>	38.7	<i>37.8</i>	16.4	<i>16.7</i>
Cocaine	2.1	<i>2.0</i>	4.9	<i>4.4</i>	7.4	<i>7.2</i>	4.0	<i>3.4</i>	1.4	<i>1.3</i>	5.9	<i>6.0</i>	6.2	<i>5.8</i>	1.8	<i>1.6</i>
Crack	0.3	<i>0.2</i>	0.9	<i>0.9</i>	2.2	<i>1.3</i>	0.5	<i>0.8</i>	0.1	<i>0.1</i>	1.3	<i>0.9</i>	1.7	<i>1.0</i>	0.2	<i>0.2</i>
Heroin	0.3	<i>0.3</i>	0.8	<i>1.1</i>	2.0	<i>1.8</i>	0.4	<i>0.9</i>	0.2	<i>0.1</i>	1.2	<i>0.8</i>	1.4	<i>1.0</i>	0.2	<i>0.2</i>
Hallucinogens	2.2	<i>2.6</i>	5.1	<i>6.2</i>	6.6	<i>9.7</i>	4.6	<i>5.0</i>	1.5	<i>1.7</i>	6.3	<i>8.1</i>	6.5	<i>7.9</i>	1.9	<i>2.1</i>
LSD	0.9	<i>0.9</i>	2.1	<i>2.5</i>	2.5	<i>4.4</i>	1.9	<i>1.7</i>	0.6	<i>0.5</i>	2.7	<i>3.9</i>	2.6	<i>3.9</i>	0.7	<i>0.7</i>
PCP	0.0	<i>0.0</i>	0.0	<i>0.2</i>	0.1	<i>0.4</i>	0.0	<i>0.1</i>	0.0	<i>0.0</i>	0.0	<i>0.3</i>	0.1	<i>0.4</i>	0.0	<i>0.0</i>
Ecstasy	1.0	<i>1.0</i>	2.2	<i>2.1</i>	2.6	<i>4.1</i>	2.0	<i>1.4</i>	0.6	<i>0.7</i>	2.6	<i>3.4</i>	2.6	<i>3.2</i>	0.8	<i>0.8</i>
Inhalants	0.6	<i>0.7</i>	1.4	<i>1.6</i>	1.8	<i>3.2</i>	1.2	<i>1.1</i>	0.4	<i>0.4</i>	1.8	<i>2.5</i>	2.0	<i>2.5</i>	0.4	<i>0.5</i>
Methamphetamine	0.8	<i>0.9</i>	2.2	<i>3.0</i>	4.1	<i>5.2</i>	1.6	<i>2.1</i>	0.4	<i>0.4</i>	2.4	<i>3.8</i>	3.2	<i>4.4</i>	0.6	<i>0.7</i>
Misuse of Prescription Psychotherapeutics	6.1	<i>6.0</i>	14.3	<i>13.9</i>	21.3	<i>18.6</i>	11.8	<i>12.2</i>	4.0	<i>3.9</i>	17.2	<i>15.4</i>	18.8	<i>16.7</i>	5.1	<i>5.2</i>
Pain Relievers	3.7	<i>3.4</i>	8.5	<i>7.8</i>	13.3	<i>11.1</i>	6.9	<i>6.6</i>	2.4	<i>2.2</i>	10.4	<i>8.4</i>	11.6	<i>9.4</i>	3.1	<i>3.0</i>
Stimulants	1.8	<i>1.9</i>	4.4	<i>4.2</i>	6.7	<i>6.1</i>	3.6	<i>3.4</i>	1.1	<i>1.3</i>	5.7	<i>5.4</i>	6.0	<i>5.4</i>	1.5	<i>1.6</i>
Tranquilizers or Sedatives	2.2	<i>2.4</i>	6.2	<i>7.1</i>	10.4	<i>10.6</i>	4.7	<i>5.8</i>	1.1	<i>1.1</i>	7.9	<i>8.6</i>	9.2	<i>9.7</i>	1.7	<i>1.8</i>
Tranquilizers	1.9	<i>2.1</i>	5.6	<i>6.3</i>	9.7	<i>9.6</i>	4.1	<i>5.0</i>	1.0	<i>1.0</i>	7.3	<i>7.8</i>	8.5	<i>8.9</i>	1.5	<i>1.6</i>
Sedatives	0.4	<i>0.5</i>	1.1	<i>1.6</i>	1.9	<i>2.9</i>	0.8	<i>1.2</i>	0.2	<i>0.2</i>	1.5	<i>2.2</i>	1.7	<i>2.6</i>	0.3	<i>0.3</i>
Benzodiazepines	1.8	<i>1.8</i>	5.1	<i>5.7</i>	8.9	<i>8.4</i>	3.8	<i>4.7</i>	0.9	<i>0.8</i>	6.9	<i>7.2</i>	8.0	<i>8.3</i>	1.3	<i>1.4</i>
Opioids	3.8	<i>3.5</i>	8.8	<i>8.1</i>	13.8	<i>11.6</i>	7.1	<i>6.8</i>	2.5	<i>2.3</i>	10.7	<i>8.5</i>	11.8	<i>9.5</i>	3.2	<i>3.0</i>
Central Nervous System Stimulants	3.9	<i>3.9</i>	9.1	<i>8.8</i>	14.1	<i>13.9</i>	7.5	<i>7.0</i>	2.5	<i>2.6</i>	11.0	<i>11.2</i>	11.8	<i>11.5</i>	3.2	<i>3.2</i>
Illicit Drugs Other Than Marijuana ⁴	8.7	<i>8.8</i>	19.4	<i>20.1</i>	28.2	<i>27.9</i>	16.3	<i>17.2</i>	5.9	<i>5.8</i>	23.0	<i>22.9</i>	24.9	<i>24.0</i>	7.4	<i>7.5</i>
PAST MONTH USE																
Alcohol	54.9	<i>54.2</i>	56.7	<i>55.8</i>	56.4	<i>54.0</i>	56.8	<i>56.5</i>	54.4	<i>53.7</i>	57.1	<i>55.1</i>	54.7	<i>54.1</i>	54.8	<i>54.3</i>
Binge Alcohol Use ⁵	25.8	<i>24.0</i>	30.9	<i>28.5</i>	32.7	<i>30.9</i>	30.3	<i>27.7</i>	24.5	<i>22.8</i>	32.1	<i>29.4</i>	30.3	<i>29.2</i>	25.3	<i>23.6</i>
Heavy Alcohol Use	6.3	<i>6.9</i>	8.6	<i>8.8</i>	9.9	<i>10.2</i>	8.2	<i>8.2</i>	5.7	<i>6.4</i>	9.3	<i>9.2</i>	8.4	<i>9.2</i>	6.1	<i>6.7</i>
Cigarette Use	18.1	<i>16.3</i>	27.2	<i>23.1</i>	33.0	<i>27.5</i>	25.2	<i>21.5</i>	15.8	<i>14.5</i>	27.5	<i>23.6</i>	29.5	<i>24.0</i>	17.2	<i>15.5</i>
Daily Cigarette Use	10.7	<i>9.9</i>	16.1	<i>13.8</i>	19.5	<i>16.2</i>	14.9	<i>12.9</i>	9.3	<i>8.9</i>	16.3	<i>13.6</i>	17.8	<i>13.7</i>	10.2	<i>9.6</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. Therefore, some estimates in this table could differ from corresponding estimates in other tables that did not undergo sample size adjustments. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

² Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

³ Respondents with unknown impairment data were excluded.

⁴ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

⁵ Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 8.43A Substance Use in Past Year and Past Month: Among People Aged 18 to 25; by Past Year Level of Mental Illness and Past Year Major Depressive Episode (MDE) and MDE with Severe Impairment Status, Numbers in Thousands, 2019 and 2020

Substance	Total		Any Mental Illness ¹		Serious Mental Illness ¹		Any Mental Illness Excluding Serious Mental Illness ¹		Any Mental Illness Excluding Serious Mental Illness ¹		No Mental Illness ¹		No Mental Illness ¹		MDE ²		MDE ²		MDE with Severe Impairment ^{2,3}		MDE with Severe Impairment ^{2,3}		No MDE ²		No MDE ²	
	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)
PAST YEAR USE																										
Illicit Drugs	13,193	12,279	5,357	5,219	1,761	1,848	3,597	3,372	7,836	7,060	2,750	2,931	1,953	2,076	10,214	9,208										
Marijuana	11,952	11,443	4,866	4,923	1,614	1,767	3,252	3,156	7,085	6,520	2,504	2,750	1,755	1,956	9,258	8,556										
Cocaine	1,789	1,469	873	699	305	330	568	369	915	770	451	506	308	340	1,306	962										
Crack	61	52	47	38	19	12	28	26	13	14	23	20	23	17	37	32										
Heroin	87	66	65	34	30	17	34	17	22	32	39	20	35	16	46	45										
Hallucinogens	2,425	2,444	1,167	1,211	398	605	769	607	1,258	1,233	604	810	426	552	1,761	1,604										
LSD	1,202	1,241	595	686	188	340	407	346	607	555	293	433	213	310	878	785										
PCP	30	63	11	58	10	53	2	5	19	5	8	57	7	53	10	6										
Ecstasy	1,089	861	474	427	141	207	333	220	615	435	235	283	170	171	835	578										
Inhalants	577	513	333	328	125	208	208	120	244	185	195	236	160	162	350	272										
Methamphetamine	275	180	146	114	50	52	96	62	130	66	59	85	49	74	214	94										
Misuse of Prescription Psychotherapeutics	3,868	3,189	1,878	1,552	731	696	1,147	856	1,990	1,637	1,042	1,034	791	742	2,750	2,117										
Pain Relievers	1,754	1,350	893	731	389	305	503	426	861	619	514	461	409	326	1,180	878										
Stimulants	1,951	1,627	957	799	371	301	587	498	994	828	510	512	373	343	1,419	1,094										
Tranquilizers or Sedatives	1,409	1,250	775	798	345	434	430	364	634	452	485	590	396	416	903	654										
Tranquilizers	1,313	1,122	712	737	305	408	407	329	601	385	437	531	355	386	856	586										
Sedatives	175	256	120	185	78	98	42	87	55	72	91	161	82	123	82	95										
Benzodiazepines	1,266	1,111	694	736	292	414	402	321	572	375	428	524	347	384	827	581										
Opioids	1,776	1,367	904	739	389	307	515	432	871	627	517	463	409	328	1,199	893										
Central Nervous System Stimulants	3,205	2,604	1,554	1,234	556	501	997	733	1,651	1,369	797	816	557	568	2,355	1,767										
Illicit Drugs Other Than Marijuana ⁴	5,791	5,161	2,700	2,434	988	1,060	1,712	1,374	3,091	2,726	1,438	1,557	1,062	1,098	4,217	3,544										
PAST MONTH USE																										
Alcohol	18,323	17,212	6,031	6,099	1,804	1,976	4,227	4,123	12,292	11,113	3,051	3,374	2,022	2,373	15,012	13,662										
Binge Alcohol Use ⁵	11,559	10,421	3,792	3,738	1,169	1,329	2,624	2,408	7,767	6,683	1,956	2,092	1,313	1,442	9,454	8,206										
Heavy Alcohol Use	2,844	2,842	944	1,098	350	423	594	675	1,900	1,745	522	584	358	395	2,269	2,200										
Cigarette Use	5,917	4,745	2,263	1,824	781	750	1,482	1,075	3,654	2,921	1,112	1,131	831	829	4,665	3,529										
Daily Cigarette Use	2,097	1,578	794	576	286	242	508	334	1,303	1,003	345	367	286	270	1,715	1,177										

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. Therefore, some estimates in this table could differ from corresponding estimates in other tables that did not undergo sample size adjustments. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

² Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

³ Respondents with unknown impairment data were excluded.

⁴ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

⁵ Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 8.43B Substance Use in Past Year and Past Month: Among People Aged 18 to 25; by Past Year Level of Mental Illness and Past Year Major Depressive Episode (MDE) and MDE with Severe Impairment Status, Percentages, 2019 and 2020

Substance	Total (2019)	Total (2020)	Any Mental Illness ¹ (2019)	Any Mental Illness ¹ (2020)	Serious Mental Illness ¹ (2019)	Serious Mental Illness ¹ (2020)	Any Mental Illness Excluding Serious Mental Illness ¹ (2019)	Any Mental Illness Excluding Serious Mental Illness ¹ (2020)	No Mental Illness ¹ (2019)	No Mental Illness ¹ (2020)	MDE ² (2019)	MDE ² (2020)	MDE with Severe Impairment ^{2,3} (2019)	MDE with Severe Impairment ^{2,3} (2020)	No MDE ² (2019)	No MDE ² (2020)
PAST YEAR USE																
Illicit Drugs	39.1	36.7	54.0	51.0	60.7	56.7	51.2	48.3	32.9	30.4	54.5	52.2	57.3	52.2	36.3	33.6
Marijuana	35.4	34.2	49.0	48.1	55.7	54.2	46.3	45.2	29.8	28.0	49.6	49.0	51.5	49.2	32.9	31.2
Cocaine	5.3	4.4	8.8	6.8	10.5	10.1	8.1	5.3	3.8	3.3	8.9	9.0	9.0	8.5	4.6	3.5
Crack	0.2	0.2	0.5	0.4	0.7	0.4	0.4	0.4	0.1	0.1	0.5	0.3	0.7	0.4	0.1	0.1
Heroin	0.3	0.2	0.7	0.3	1.0	0.5	0.5	0.2	0.1	0.1	0.8	0.3	1.0	0.4	0.2	0.2
Hallucinogens	7.2	7.3	11.8	11.8	13.7	18.6	10.9	8.7	5.3	5.3	12.0	14.4	12.5	13.9	6.3	5.9
LSD	3.6	3.7	6.0	6.7	6.5	10.4	5.8	5.0	2.6	2.4	5.8	7.7	6.3	7.8	3.1	2.9
PCP	0.1	0.2	0.1	0.6	0.3	1.6	0.0	0.1	0.1	0.0	0.2	1.0	0.2	1.3	0.0	0.0
Ecstasy	3.2	2.6	4.8	4.2	4.9	6.3	4.7	3.2	2.6	1.9	4.7	5.0	5.0	4.3	3.0	2.1
Inhalants	1.7	1.5	3.4	3.2	4.3	6.4	3.0	1.7	1.0	0.8	3.9	4.2	4.7	4.1	1.2	1.0
Methamphetamine	0.8	0.5	1.5	1.1	1.7	1.6	1.4	0.9	0.5	0.3	1.2	1.5	1.4	1.9	0.8	0.3
Misuse of Prescription Psychotherapeutics	11.5	9.5	18.9	15.2	25.2	21.4	16.3	12.3	8.4	7.0	20.7	18.4	23.2	18.6	9.8	7.7
Pain Relievers	5.2	4.0	9.0	7.1	13.4	9.4	7.2	6.1	3.6	2.7	10.2	8.2	12.0	8.2	4.2	3.2
Stimulants	5.8	4.9	9.6	7.8	12.8	9.2	8.3	7.1	4.2	3.6	10.1	9.1	11.0	8.6	5.0	4.0
Tranquilizers or Sedatives	4.2	3.7	7.8	7.8	11.9	13.3	6.1	5.2	2.7	1.9	9.6	10.5	11.6	10.5	3.2	2.4
Tranquilizers	3.9	3.4	7.2	7.2	10.5	12.5	5.8	4.7	2.5	1.7	8.7	9.5	10.4	9.7	3.0	2.1
Sedatives	0.5	0.8	1.2	1.8	2.7	3.0	0.6	1.2	0.2	0.3	1.8	2.9	2.4	3.1	0.3	0.3
Benzodiazepines	3.8	3.3	7.0	7.2	10.1	12.7	5.7	4.6	2.4	1.6	8.5	9.3	10.2	9.6	2.9	2.1
Opioids	5.3	4.1	9.1	7.2	13.4	9.4	7.3	6.2	3.7	2.7	10.2	8.3	12.0	8.2	4.3	3.3
Central Nervous System Stimulants	9.5	7.8	15.6	12.1	19.2	15.4	14.2	10.5	6.9	5.9	15.8	14.5	16.3	14.3	8.4	6.4
Illicit Drugs Other Than Marijuana ⁴	17.2	15.4	27.2	23.8	34.1	32.5	24.4	19.7	13.0	11.7	28.5	27.8	31.2	27.6	15.0	12.9
PAST MONTH USE																
Alcohol	54.3	51.4	60.7	59.6	62.2	60.6	60.1	59.1	51.6	47.8	60.5	60.1	59.3	59.6	53.3	49.9
Binge Alcohol Use ⁵	34.3	31.1	38.2	36.5	40.3	40.8	37.3	34.5	32.6	28.7	38.8	37.3	38.5	36.2	33.6	29.9
Heavy Alcohol Use	8.4	8.5	9.5	10.7	12.1	13.0	8.5	9.7	8.0	7.5	10.3	10.4	10.5	9.9	8.1	8.0
Cigarette Use	17.5	14.2	22.8	17.8	26.9	23.0	21.1	15.4	15.4	12.6	22.0	20.2	24.4	20.8	16.6	12.9
Daily Cigarette Use	6.2	4.7	8.0	5.6	9.9	7.4	7.2	4.8	5.5	4.3	6.8	6.5	8.4	6.8	6.1	4.3

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. Therefore, some estimates in this table could differ from corresponding estimates in other tables that did not undergo sample size adjustments. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

² Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

³ Respondents with unknown impairment data were excluded.

⁴ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

⁵ Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 8.44A Substance Use in Past Year and Past Month: Among People Aged 26 or Older; by Past Year Level of Mental Illness and Past Year Major Depressive Episode (MDE) and MDE with Severe Impairment Status, Numbers in Thousands, 2019 and 2020

Substance	Total		Any Mental Illness ¹		Serious Mental Illness ¹		Any Mental Illness Excluding Serious Mental Illness ¹		No Mental Illness ¹		MDE ²		MDE with Severe Impairment ^{2,3}		No MDE ²	
	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)
PAST YEAR USE																
Illicit Drugs	39,731	42,703	14,641	15,842	4,714	4,938	9,927	10,904	25,090	26,860	5,805	6,290	4,159	4,595	33,562	35,970
Marijuana	33,007	34,895	11,875	12,435	3,611	3,805	8,263	8,630	21,132	22,460	4,696	5,037	3,325	3,658	28,059	29,490
Cocaine	3,582	3,539	1,649	1,632	669	697	981	934	1,933	1,907	697	760	501	526	2,873	2,762
Crack	706	569	430	440	268	175	162	265	276	129	233	160	203	139	473	408
Heroin	658	653	358	568	229	235	129	333	300	85	185	156	147	133	465	497
Hallucinogens	3,146	4,170	1,449	2,078	465	770	985	1,308	1,697	2,092	618	899	428	623	2,492	3,245
LSD	989	1,123	474	617	138	288	336	329	515	506	236	383	127	262	735	725
PCP	22	31	5	26	1	7	5	19	16	*	*	7	*	*	21	24
Ecstasy	1,306	1,603	639	707	203	371	436	336	667	896	262	435	171	309	1,031	1,166
Inhalants	822	1,199	363	538	109	244	254	293	459	662	162	292	99	206	653	890
Methamphetamine	1,682	2,185	1,004	1,454	484	693	520	761	678	731	412	707	369	574	1,248	1,410
Misuse of Prescription Psychotherapeutics	11,374	12,029	5,468	5,812	2,068	1,949	3,401	3,863	5,906	6,217	2,287	2,193	1,674	1,741	8,957	9,767
Pain Relievers	7,404	7,217	3,503	3,396	1,356	1,272	2,147	2,124	3,900	3,820	1,503	1,300	1,106	1,064	5,851	5,869
Stimulants	2,548	3,202	1,301	1,405	514	569	787	836	1,247	1,798	591	616	412	459	1,939	2,579
Tranquilizers or Sedatives	4,050	4,721	2,400	2,953	1,021	1,075	1,379	1,878	1,650	1,769	1,045	1,220	808	1,028	2,931	3,486
Tranquilizers	3,516	4,116	2,152	2,567	967	955	1,186	1,612	1,364	1,549	980	1,099	753	937	2,461	3,002
Sedatives	841	996	454	677	173	319	281	358	387	319	191	300	141	267	646	684
Benzodiazepines	3,194	3,535	1,946	2,263	876	779	1,070	1,485	1,249	1,272	910	993	707	842	2,210	2,527
Opioids	7,723	7,422	3,629	3,553	1,424	1,338	2,205	2,215	4,094	3,870	1,549	1,324	1,141	1,088	6,119	6,050
Central Nervous System Stimulants	6,433	7,197	3,151	3,423	1,288	1,469	1,863	1,954	3,282	3,773	1,331	1,526	985	1,139	5,053	5,590
Illicit Drugs Other Than Marijuana ⁴	15,936	17,014	7,269	8,170	2,709	2,900	4,560	5,269	8,666	8,844	3,017	3,247	2,207	2,461	12,743	13,614
PAST MONTH USE																
Alcohol	119,058	119,265	23,185	23,407	5,595	5,689	17,590	17,718	95,873	95,858	8,007	8,193	5,149	5,662	109,890	110,359
Binge Alcohol Use ⁵	53,073	49,955	12,103	11,347	3,114	3,058	8,989	8,288	40,971	38,608	4,249	4,074	2,657	2,899	48,251	45,617
Heavy Alcohol Use	12,991	14,521	3,492	3,531	947	1,022	2,544	2,509	9,499	10,990	1,286	1,339	749	964	11,584	13,087
Cigarette Use	39,396	36,306	11,728	10,401	3,542	3,150	8,187	7,251	27,668	25,905	4,216	3,829	3,036	2,737	34,595	31,971
Daily Cigarette Use	24,624	23,454	7,490	6,729	2,273	2,064	5,216	4,665	17,134	16,725	2,819	2,489	2,048	1,771	21,458	20,694

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. Therefore, some estimates in this table could differ from corresponding estimates in other tables that did not undergo sample size adjustments. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

² Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

³ Respondents with unknown impairment data were excluded.

⁴ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

⁵ Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 8.44B Substance Use in Past Year and Past Month: Among People Aged 26 or Older; by Past Year Level of Mental Illness and Past Year Major Depressive Episode (MDE) and MDE with Severe Impairment Status, Percentages, 2019 and 2020

Substance	Total (2019)	Total (2020)	Any Mental Illness ¹ (2019)	Any Mental Illness ¹ (2020)	Serious Mental Illness ¹ (2019)	Serious Mental Illness ¹ (2020)	Any Mental Illness Excluding Serious Mental Illness ¹	Any Mental Illness Excluding Serious Mental Illness ¹	No Mental Illness ¹	No Mental Illness ¹	MDE ² (2019)	MDE ² (2020)	MDE with Severe Impairment ^{2,3}	MDE with Severe Impairment ^{2,3}	No MDE ² (2019)	No MDE ² (2020)
							(2019)	(2020)	(2019)	(2020)			(2019)	(2020)	(2019)	(2020)
PAST YEAR USE																
Illicit Drugs	18.3	<i>19.5</i>	35.2	<i>37.2</i>	46.1	<i>45.1</i>	31.7	<i>34.4</i>	14.3	<i>15.3</i>	40.6	<i>40.9</i>	42.9	<i>42.3</i>	16.8	<i>17.9</i>
Marijuana	15.2	<i>16.0</i>	28.6	<i>29.2</i>	35.3	<i>34.8</i>	26.4	<i>27.2</i>	12.1	<i>12.8</i>	32.8	<i>32.8</i>	34.3	<i>33.7</i>	14.0	<i>14.7</i>
Cocaine	1.7	<i>1.6</i>	4.0	<i>3.8</i>	6.5	<i>6.4</i>	3.1	<i>2.9</i>	1.1	<i>1.1</i>	4.9	<i>4.9</i>	5.2	<i>4.8</i>	1.4	<i>1.4</i>
Crack	0.3	<i>0.3</i>	1.0	<i>1.0</i>	2.6	<i>1.6</i>	0.5	<i>0.8</i>	0.2	<i>0.1</i>	1.6	<i>1.0</i>	2.1	<i>1.3</i>	0.2	<i>0.2</i>
Heroin	0.3	<i>0.3</i>	0.9	<i>1.3</i>	2.2	<i>2.1</i>	0.4	<i>1.1</i>	0.2	<i>0.0</i>	1.3	<i>1.0</i>	1.5	<i>1.2</i>	0.2	<i>0.2</i>
Hallucinogens	1.5	<i>1.9</i>	3.5	<i>4.9</i>	4.5	<i>7.0</i>	3.1	<i>4.1</i>	1.0	<i>1.2</i>	4.3	<i>5.8</i>	4.4	<i>5.7</i>	1.2	<i>1.6</i>
LSD	0.5	<i>0.5</i>	1.1	<i>1.4</i>	1.4	<i>2.6</i>	1.1	<i>1.0</i>	0.3	<i>0.3</i>	1.7	<i>2.5</i>	1.3	<i>2.4</i>	0.4	<i>0.4</i>
PCP	0.0	<i>0.0</i>	0.0	<i>0.1</i>	0.0	<i>0.1</i>	0.0	<i>0.1</i>	0.0	<i>*</i>	<i>*</i>	<i>0.0</i>	<i>*</i>	<i>*</i>	0.0	<i>0.0</i>
Ecstasy	0.6	<i>0.7</i>	1.5	<i>1.7</i>	2.0	<i>3.4</i>	1.4	<i>1.1</i>	0.4	<i>0.5</i>	1.8	<i>2.8</i>	1.8	<i>2.8</i>	0.5	<i>0.6</i>
Inhalants	0.4	<i>0.5</i>	0.9	<i>1.3</i>	1.1	<i>2.2</i>	0.8	<i>0.9</i>	0.3	<i>0.4</i>	1.1	<i>1.9</i>	1.0	<i>1.9</i>	0.3	<i>0.4</i>
Methamphetamine	0.8	<i>1.0</i>	2.4	<i>3.4</i>	4.7	<i>6.3</i>	1.7	<i>2.4</i>	0.4	<i>0.4</i>	2.9	<i>4.6</i>	3.8	<i>5.3</i>	0.6	<i>0.7</i>
Misuse of Prescription Psychotherapeutics	5.3	<i>5.5</i>	13.2	<i>13.6</i>	20.2	<i>17.8</i>	10.8	<i>12.2</i>	3.4	<i>3.5</i>	16.0	<i>14.3</i>	17.2	<i>16.0</i>	4.5	<i>4.9</i>
Pain Relievers	3.4	<i>3.3</i>	8.4	<i>8.0</i>	13.3	<i>11.6</i>	6.8	<i>6.7</i>	2.2	<i>2.2</i>	10.5	<i>8.5</i>	11.4	<i>9.8</i>	2.9	<i>2.9</i>
Stimulants	1.2	<i>1.5</i>	3.1	<i>3.3</i>	5.0	<i>5.2</i>	2.5	<i>2.6</i>	0.7	<i>1.0</i>	4.1	<i>4.0</i>	4.2	<i>4.2</i>	1.0	<i>1.3</i>
Tranquilizers or Sedatives	1.9	<i>2.2</i>	5.8	<i>6.9</i>	10.0	<i>9.8</i>	4.4	<i>5.9</i>	0.9	<i>1.0</i>	7.3	<i>7.9</i>	8.3	<i>9.5</i>	1.5	<i>1.7</i>
Tranquilizers	1.6	<i>1.9</i>	5.2	<i>6.0</i>	9.5	<i>8.7</i>	3.8	<i>5.1</i>	0.8	<i>0.9</i>	6.8	<i>7.1</i>	7.8	<i>8.6</i>	1.2	<i>1.5</i>
Sedatives	0.4	<i>0.5</i>	1.1	<i>1.6</i>	1.7	<i>2.9</i>	0.9	<i>1.1</i>	0.2	<i>0.2</i>	1.3	<i>2.0</i>	1.5	<i>2.5</i>	0.3	<i>0.3</i>
Benzodiazepines	1.5	<i>1.6</i>	4.7	<i>5.3</i>	8.6	<i>7.1</i>	3.4	<i>4.7</i>	0.7	<i>0.7</i>	6.4	<i>6.5</i>	7.3	<i>7.7</i>	1.1	<i>1.3</i>
Opioids	3.6	<i>3.4</i>	8.7	<i>8.3</i>	13.9	<i>12.2</i>	7.0	<i>7.0</i>	2.3	<i>2.2</i>	10.8	<i>8.6</i>	11.8	<i>10.0</i>	3.1	<i>3.0</i>
Central Nervous System Stimulants	3.0	<i>3.3</i>	7.6	<i>8.0</i>	12.6	<i>13.4</i>	5.9	<i>6.2</i>	1.9	<i>2.1</i>	9.3	<i>9.9</i>	10.1	<i>10.5</i>	2.5	<i>2.8</i>
Illicit Drugs Other Than Marijuana ⁴	7.4	<i>7.8</i>	17.5	<i>19.2</i>	26.5	<i>26.5</i>	14.5	<i>16.6</i>	5.0	<i>5.0</i>	21.1	<i>21.1</i>	22.7	<i>22.7</i>	6.4	<i>6.8</i>
PAST MONTH USE																
Alcohol	55.0	<i>54.6</i>	55.8	<i>54.9</i>	54.8	<i>52.0</i>	56.1	<i>55.9</i>	54.8	<i>54.5</i>	55.9	<i>53.3</i>	53.0	<i>52.1</i>	55.0	<i>54.9</i>
Binge Alcohol Use ⁵	24.5	<i>22.9</i>	29.1	<i>26.6</i>	30.5	<i>28.0</i>	28.7	<i>26.2</i>	23.4	<i>22.0</i>	29.7	<i>26.5</i>	27.4	<i>26.7</i>	24.1	<i>22.7</i>
Heavy Alcohol Use	6.0	<i>6.6</i>	8.4	<i>8.3</i>	9.3	<i>9.3</i>	8.1	<i>7.9</i>	5.4	<i>6.3</i>	9.0	<i>8.7</i>	7.7	<i>8.9</i>	5.8	<i>6.5</i>
Cigarette Use	18.2	<i>16.6</i>	28.2	<i>24.4</i>	34.7	<i>28.8</i>	26.1	<i>22.9</i>	15.8	<i>14.7</i>	29.5	<i>24.9</i>	31.3	<i>25.2</i>	17.3	<i>15.9</i>
Daily Cigarette Use	11.4	<i>10.7</i>	18.0	<i>15.8</i>	22.3	<i>18.9</i>	16.6	<i>14.7</i>	9.8	<i>9.5</i>	19.7	<i>16.2</i>	21.1	<i>16.3</i>	10.7	<i>10.3</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. Therefore, some estimates in this table could differ from corresponding estimates in other tables that did not undergo sample size adjustments. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

² Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

³ Respondents with unknown impairment data were excluded.

⁴ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

⁵ Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 8.45A Substance Use in Past Year and Past Month: Among People Aged 26 to 49; by Past Year Level of Mental Illness and Past Year Major Depressive Episode (MDE) and MDE with Severe Impairment Status, Numbers in Thousands, 2019 and 2020

Substance	Total		Any Mental Illness ¹		Serious Mental Illness ¹		Any Mental Illness Excluding Serious Mental Illness ¹		No Mental Illness ¹		MDE ²		MDE with Severe Impairment ^{2,3}		No MDE ²	
	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)
PAST YEAR USE																
Illicit Drugs	26,092	27,233	10,401	10,943	3,400	3,547	7,001	7,395	15,692	16,290	4,101	4,259	2,871	3,121	21,773	22,811
Marijuana	22,016	23,038	8,513	9,176	2,724	2,927	5,789	6,249	13,504	13,862	3,348	3,559	2,319	2,592	18,488	19,316
Cocaine	2,863	2,921	1,444	1,204	559	493	886	711	1,418	1,717	594	565	423	361	2,256	2,350
Crack	343	356	269	255	171	118	98	136	74	101	138	104	131	97	205	251
Heroin	474	440	318	377	214	195	103	182	156	63	161	116	141	93	304	323
Hallucinogens	2,742	3,611	1,339	1,781	438	709	901	1,072	1,403	1,830	560	806	382	583	2,156	2,779
LSD	890	1,049	471	567	135	281	336	286	419	482	234	354	124	262	639	680
PCP	19	10	5	8	1	7	5	1	13	*	1	7	*	*	18	*
Ecstasy	1,261	1,565	632	686	195	371	436	315	630	879	254	435	163	309	994	1,128
Inhalants	623	1,056	292	451	109	203	183	248	331	606	135	251	99	206	481	790
Methamphetamine	1,262	1,355	809	893	437	385	373	509	453	461	354	408	321	303	895	916
Misuse of Prescription Psychotherapeutics	7,601	7,630	3,897	3,902	1,458	1,427	2,439	2,475	3,704	3,728	1,626	1,612	1,154	1,253	5,920	6,008
Pain Relievers	4,797	4,323	2,470	2,288	953	955	1,517	1,333	2,327	2,034	1,030	930	732	754	3,735	3,392
Stimulants	2,247	2,675	1,138	1,280	465	463	673	816	1,109	1,395	517	562	366	404	1,712	2,105
Tranquilizers or Sedatives	2,582	3,182	1,661	1,940	688	751	973	1,190	921	1,242	733	893	522	734	1,831	2,288
Tranquilizers	2,342	2,862	1,521	1,738	635	660	885	1,078	821	1,124	671	810	467	658	1,653	2,051
Sedatives	423	530	280	340	129	188	151	152	144	191	147	161	99	152	273	369
Benzodiazepines	2,169	2,555	1,401	1,608	579	588	823	1,021	768	946	635	757	455	615	1,516	1,795
Opioids	4,971	4,477	2,573	2,416	1,019	1,021	1,553	1,394	2,398	2,061	1,066	954	768	778	3,869	3,522
Central Nervous System Stimulants	5,196	5,547	2,699	2,584	1,106	1,012	1,593	1,572	2,497	2,963	1,132	1,129	838	785	4,026	4,375
Illicit Drugs Other Than Marijuana ⁴	11,025	11,387	5,416	5,475	1,983	2,010	3,433	3,464	5,609	5,912	2,197	2,285	1,584	1,694	8,747	9,049
PAST MONTH USE																
Alcohol	62,340	60,652	16,174	15,839	4,278	4,081	11,896	11,758	46,166	44,813	5,611	5,377	3,723	3,750	56,256	54,946
Binge Alcohol Use ⁵	33,413	30,700	9,051	8,172	2,393	2,147	6,658	6,025	24,362	22,528	3,053	2,834	1,985	1,953	30,131	27,714
Heavy Alcohol Use	7,816	8,251	2,530	2,594	745	820	1,785	1,774	5,286	5,656	902	1,084	595	788	6,873	7,102
Cigarette Use	22,628	20,772	7,686	6,854	2,547	2,376	5,139	4,478	14,942	13,918	2,864	2,693	2,095	1,820	19,430	17,919
Daily Cigarette Use	13,106	12,536	4,641	4,414	1,537	1,525	3,103	2,888	8,465	8,122	1,848	1,704	1,367	1,102	11,078	10,753

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. Therefore, some estimates in this table could differ from corresponding estimates in other tables that did not undergo sample size adjustments. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

² Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

³ Respondents with unknown impairment data were excluded.

⁴ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

⁵ Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 8.45B Substance Use in Past Year and Past Month: Among People Aged 26 to 49; by Past Year Level of Mental Illness and Past Year Major Depressive Episode (MDE) and MDE with Severe Impairment Status, Percentages, 2019 and 2020

Substance	Total		Any Mental Illness ¹		Serious Mental Illness ¹		Any Mental Illness Excluding Serious Mental Illness ¹		No Mental Illness ¹		MDE ²		MDE with Severe Impairment ^{2,3}		No MDE ²	
	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)
PAST YEAR USE																
Illicit Drugs	25.8	26.8	41.1	42.6	49.7	50.8	38.0	39.6	20.6	21.5	46.0	46.5	47.1	47.9	23.8	24.9
Marijuana	21.7	22.7	33.7	35.7	39.8	41.9	31.4	33.4	17.8	18.3	37.6	38.8	38.0	39.8	20.2	21.1
Cocaine	2.8	2.9	5.7	4.7	8.2	7.1	4.8	3.8	1.9	2.3	6.7	6.2	6.9	5.5	2.5	2.6
Crack	0.3	0.4	1.1	1.0	2.5	1.7	0.5	0.7	0.1	0.1	1.5	1.1	2.2	1.5	0.2	0.3
Heroin	0.5	0.4	1.3	1.5	3.1	2.8	0.6	1.0	0.2	0.1	1.8	1.3	2.3	1.4	0.3	0.4
Hallucinogens	2.7	3.6	5.3	6.9	6.4	10.2	4.9	5.7	1.8	2.4	6.3	8.8	6.3	9.0	2.4	3.0
LSD	0.9	1.0	1.9	2.2	2.0	4.0	1.8	1.5	0.6	0.6	2.6	3.9	2.0	4.0	0.7	0.7
PCP	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	*	0.0	0.1	*	*	0.0	*
Ecstasy	1.2	1.5	2.5	2.7	2.8	5.3	2.4	1.7	0.8	1.2	2.8	4.7	2.7	4.8	1.1	1.2
Inhalants	0.6	1.0	1.2	1.8	1.6	2.9	1.0	1.3	0.4	0.8	1.5	2.7	1.6	3.2	0.5	0.9
Methamphetamine	1.2	1.3	3.2	3.5	6.4	5.5	2.0	2.7	0.6	0.6	4.0	4.5	5.3	4.7	1.0	1.0
Misuse of Prescription Psychotherapeutics	7.5	7.5	15.4	15.2	21.3	20.4	13.2	13.2	4.9	4.9	18.2	17.6	18.9	19.3	6.5	6.6
Pain Relievers	4.7	4.3	9.8	8.9	13.9	13.7	8.2	7.1	3.1	2.7	11.6	10.1	12.0	11.6	4.1	3.7
Stimulants	2.2	2.6	4.5	5.0	6.8	6.6	3.7	4.4	1.5	1.8	5.8	6.1	6.0	6.2	1.9	2.3
Tranquilizers or Sedatives	2.5	3.1	6.6	7.6	10.0	10.8	5.3	6.4	1.2	1.6	8.2	9.7	8.6	11.3	2.0	2.5
Tranquilizers	2.3	2.8	6.0	6.8	9.3	9.5	4.8	5.8	1.1	1.5	7.5	8.8	7.7	10.1	1.8	2.2
Sedatives	0.4	0.5	1.1	1.3	1.9	2.7	0.8	0.8	0.2	0.3	1.6	1.8	1.6	2.3	0.3	0.4
Benzodiazepines	2.1	2.5	5.5	6.3	8.4	8.4	4.5	5.5	1.0	1.2	7.1	8.3	7.5	9.4	1.7	2.0
Opioids	4.9	4.4	10.2	9.4	14.9	14.6	8.4	7.5	3.2	2.7	12.0	10.4	12.6	12.0	4.2	3.8
Central Nervous System Stimulants	5.1	5.5	10.7	10.1	16.1	14.5	8.6	8.4	3.3	3.9	12.7	12.3	13.7	12.1	4.4	4.8
Illicit Drugs Other Than Marijuana ⁴	10.9	11.2	21.4	21.3	29.0	28.8	18.6	18.5	7.4	7.8	24.6	24.9	26.0	26.0	9.6	9.9
PAST MONTH USE																
Alcohol	61.6	59.7	64.0	61.7	62.5	58.5	64.5	62.9	60.7	59.1	62.9	58.7	61.0	57.6	61.6	59.9
Binge Alcohol Use ⁵	33.0	30.2	35.8	31.8	34.9	30.8	36.1	32.2	32.1	29.7	34.2	30.9	32.5	30.0	33.0	30.2
Heavy Alcohol Use	7.7	8.1	10.0	10.1	10.9	11.8	9.7	9.5	7.0	7.5	10.1	11.8	9.8	12.1	7.5	7.7
Cigarette Use	22.3	20.5	30.4	26.7	37.2	34.0	27.9	24.0	19.7	18.3	32.1	29.4	34.3	28.0	21.3	19.5
Daily Cigarette Use	12.9	12.3	18.4	17.2	22.5	21.9	16.8	15.5	11.1	10.7	20.7	18.6	22.4	16.9	12.1	11.7

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. Therefore, some estimates in this table could differ from corresponding estimates in other tables that did not undergo sample size adjustments. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

² Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

³ Respondents with unknown impairment data were excluded.

⁴ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

⁵ Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 8.46A Substance Use in Past Year and Past Month: Among People Aged 50 or Older; by Past Year Level of Mental Illness and Past Year Major Depressive Episode (MDE) and MDE with Severe Impairment Status, Numbers in Thousands, 2019 and 2020

Substance	Total		Any Mental Illness ¹		Serious Mental Illness ¹		Any Mental Illness Excluding Serious Mental Illness ¹		No Mental Illness ¹		MDE ²		MDE with Severe Impairment ^{2,3}		No MDE ²	
	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)
PAST YEAR USE																
Illicit Drugs	13,639	15,470	4,240	4,900	1,314	1,391	2,926	3,509	9,399	10,570	1,704	2,031	1,288	1,474	11,789	13,159
Marijuana	10,990	11,857	3,362	3,259	887	879	2,475	2,381	7,628	8,598	1,348	1,478	1,006	1,065	9,572	10,175
Cocaine	720	618	205	428	110	205	95	223	515	190	103	195	79	165	617	412
Crack	363	213	161	185	96	*	65	129	202	28	95	56	71	*	268	157
Heroin	184	213	41	191	15	*	26	151	144	22	23	*	6	*	161	174
Hallucinogens	404	559	110	297	26	61	83	235	294	262	59	92	46	*	336	466
LSD	99	74	3	50	3	6	*	43	96	24	3	29	3	*	96	45
PCP	*	22	*	18	*	*	*	18	*	*	*	*	*	*	*	22
Ecstasy	45	38	8	21	8	*	*	21	37	16	8	*	8	*	37	38
Inhalants	199	143	71	87	*	42	71	45	128	56	*	42	*	*	172	100
Methamphetamine	420	830	195	561	48	*	147	252	225	269	58	298	48	*	353	494
Misuse of Prescription Psychotherapeutics	3,773	4,400	1,572	1,910	610	522	962	1,388	2,201	2,490	662	581	520	488	3,036	3,759
Pain Relievers	2,607	2,894	1,034	1,108	403	317	630	792	1,573	1,786	473	370	374	310	2,115	2,477
Stimulants	301	528	163	125	50	105	113	19	138	403	74	54	46	54	227	473
Tranquilizers or Sedatives	1,468	1,539	739	1,013	333	325	406	688	729	526	311	328	286	294	1,101	1,199
Tranquilizers	1,174	1,254	632	829	331	295	300	534	543	425	309	289	286	279	808	952
Sedatives	417	466	174	338	44	131	130	207	243	128	44	139	42	115	373	315
Benzodiazepines	1,026	981	544	655	297	191	248	464	481	326	275	237	251	227	694	732
Opioids	2,752	2,945	1,057	1,137	405	317	652	820	1,695	1,808	483	370	374	310	2,250	2,528
Central Nervous System Stimulants	1,236	1,650	452	840	182	457	269	382	785	810	199	397	147	354	1,027	1,215
Illicit Drugs Other Than Marijuana ⁴	4,911	5,626	1,853	2,695	726	890	1,127	1,805	3,057	2,931	821	962	623	768	3,996	4,565
PAST MONTH USE																
Alcohol	56,719	58,612	7,012	7,568	1,317	1,608	5,694	5,960	49,707	51,044	2,396	2,815	1,426	1,912	53,634	55,412
Binge Alcohol Use ⁵	19,660	19,255	3,052	3,175	721	911	2,330	2,263	16,609	16,081	1,196	1,240	672	945	18,120	17,903
Heavy Alcohol Use	5,174	6,271	962	937	203	202	759	735	4,213	5,333	385	255	154	177	4,711	5,985
Cigarette Use	16,768	15,534	4,042	3,547	995	774	3,047	2,773	12,726	11,986	1,352	1,137	941	916	15,165	14,051
Daily Cigarette Use	11,518	10,918	2,849	2,315	736	539	2,113	1,776	8,669	8,603	971	784	680	669	10,379	9,941

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. Therefore, some estimates in this table could differ from corresponding estimates in other tables that did not undergo sample size adjustments. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

² Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

³ Respondents with unknown impairment data were excluded.

⁴ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

⁵ Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 8.46B Substance Use in Past Year and Past Month: Among People Aged 50 or Older; by Past Year Level of Mental Illness and Past Year Major Depressive Episode (MDE) and MDE with Severe Impairment Status, Percentages, 2019 and 2020

Substance	Total		Any Mental Illness ¹		Serious Mental Illness ¹		Any Mental Illness Excluding Serious Mental Illness ¹		Any Mental Illness Excluding Serious Mental Illness ¹		MDE ²		MDE with Severe Impairment ^{2,3}		No MDE ²	
	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)
PAST YEAR USE																
Illicit Drugs	11.8	<i>13.2</i>	26.0	<i>28.9</i>	39.0	<i>35.1</i>	22.7	<i>27.0</i>	9.5	<i>10.6</i>	31.6	<i>32.8</i>	35.7	<i>33.8</i>	10.9	<i>12.0</i>
Marijuana	9.5	<i>10.1</i>	20.6	<i>19.2</i>	26.3	<i>22.2</i>	19.2	<i>18.3</i>	7.7	<i>8.6</i>	25.0	<i>23.8</i>	27.9	<i>24.5</i>	8.8	<i>9.3</i>
Cocaine	0.6	<i>0.5</i>	1.3	<i>2.5</i>	3.3	<i>5.2</i>	0.7	<i>1.7</i>	0.5	<i>0.2</i>	1.9	<i>3.1</i>	2.2	<i>3.8</i>	0.6	<i>0.4</i>
Crack	0.3	<i>0.2</i>	1.0	<i>1.1</i>	2.9	<i>*</i>	0.5	<i>1.0</i>	0.2	<i>0.0</i>	1.8	<i>0.9</i>	2.0	<i>*</i>	0.2	<i>0.1</i>
Heroin	0.2	<i>0.2</i>	0.2	<i>1.1</i>	0.4	<i>*</i>	0.2	<i>1.2</i>	0.1	<i>0.0</i>	0.4	<i>*</i>	0.2	<i>*</i>	0.1	<i>0.2</i>
Hallucinogens	0.3	<i>0.5</i>	0.7	<i>1.8</i>	0.8	<i>1.5</i>	0.6	<i>1.8</i>	0.3	<i>0.3</i>	1.1	<i>1.5</i>	1.3	<i>*</i>	0.3	<i>0.4</i>
LSD	0.1	<i>0.1</i>	0.0	<i>0.3</i>	0.1	<i>0.2</i>	<i>*</i>	<i>0.3</i>	0.1	<i>0.0</i>	0.0	<i>0.5</i>	0.1	<i>*</i>	0.1	<i>0.0</i>
PCP	<i>*</i>	<i>0.0</i>	<i>*</i>	<i>0.1</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>0.1</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>0.0</i>
Ecstasy	0.0	<i>0.0</i>	0.0	<i>0.1</i>	0.2	<i>*</i>	<i>*</i>	<i>0.2</i>	0.0	<i>0.0</i>	0.1	<i>*</i>	0.2	<i>*</i>	0.0	<i>0.0</i>
Inhalants	0.2	<i>0.1</i>	0.4	<i>0.5</i>	<i>*</i>	<i>1.0</i>	0.6	<i>0.3</i>	0.1	<i>0.1</i>	<i>*</i>	<i>0.7</i>	<i>*</i>	<i>*</i>	0.2	<i>0.1</i>
Methamphetamine	0.4	<i>0.7</i>	1.2	<i>3.3</i>	1.4	<i>*</i>	1.1	<i>1.9</i>	0.2	<i>0.3</i>	1.1	<i>4.8</i>	1.3	<i>*</i>	0.3	<i>0.5</i>
Misuse of Prescription Psychotherapeutics	3.3	<i>3.8</i>	9.7	<i>11.3</i>	18.1	<i>13.2</i>	7.4	<i>10.7</i>	2.2	<i>2.5</i>	12.3	<i>9.4</i>	14.4	<i>11.2</i>	2.8	<i>3.4</i>
Pain Relievers	2.3	<i>2.5</i>	6.3	<i>6.5</i>	12.0	<i>8.0</i>	4.9	<i>6.1</i>	1.6	<i>1.8</i>	8.8	<i>6.0</i>	10.4	<i>7.1</i>	1.9	<i>2.3</i>
Stimulants	0.3	<i>0.5</i>	1.0	<i>0.7</i>	1.5	<i>2.7</i>	0.9	<i>0.2</i>	0.1	<i>0.4</i>	1.4	<i>0.9</i>	1.3	<i>1.2</i>	0.2	<i>0.4</i>
Tranquilizers or Sedatives	1.3	<i>1.3</i>	4.5	<i>6.0</i>	9.9	<i>8.2</i>	3.1	<i>5.3</i>	0.7	<i>0.5</i>	5.8	<i>5.3</i>	7.9	<i>6.7</i>	1.0	<i>1.1</i>
Tranquilizers	1.0	<i>1.1</i>	3.9	<i>4.9</i>	9.8	<i>7.4</i>	2.3	<i>4.1</i>	0.5	<i>0.4</i>	5.7	<i>4.7</i>	7.9	<i>6.4</i>	0.7	<i>0.9</i>
Sedatives	0.4	<i>0.4</i>	1.1	<i>2.0</i>	1.3	<i>3.3</i>	1.0	<i>1.6</i>	0.2	<i>0.1</i>	0.8	<i>2.2</i>	1.2	<i>2.7</i>	0.3	<i>0.3</i>
Benzodiazepines	0.9	<i>0.8</i>	3.3	<i>3.9</i>	8.8	<i>4.8</i>	1.9	<i>3.6</i>	0.5	<i>0.3</i>	5.1	<i>3.8</i>	7.0	<i>5.2</i>	0.6	<i>0.7</i>
Opioids	2.4	<i>2.5</i>	6.5	<i>6.7</i>	12.0	<i>8.0</i>	5.0	<i>6.3</i>	1.7	<i>1.8</i>	8.9	<i>6.0</i>	10.4	<i>7.1</i>	2.1	<i>2.3</i>
Central Nervous System Stimulants	1.1	<i>1.4</i>	2.8	<i>5.0</i>	5.4	<i>11.5</i>	2.1	<i>2.9</i>	0.8	<i>0.8</i>	3.7	<i>6.4</i>	4.1	<i>8.1</i>	0.9	<i>1.1</i>
Illicit Drugs Other Than Marijuana ⁴	4.3	<i>4.8</i>	11.4	<i>15.9</i>	21.5	<i>22.5</i>	8.7	<i>13.9</i>	3.1	<i>2.9</i>	15.2	<i>15.5</i>	17.3	<i>17.6</i>	3.7	<i>4.2</i>
PAST MONTH USE																
Alcohol	49.2	<i>50.1</i>	43.1	<i>44.7</i>	39.1	<i>40.6</i>	44.1	<i>45.9</i>	50.2	<i>51.1</i>	44.4	<i>45.4</i>	39.6	<i>43.9</i>	49.4	<i>50.7</i>
Binge Alcohol Use ⁵	17.1	<i>16.5</i>	18.7	<i>18.7</i>	21.4	<i>23.0</i>	18.0	<i>17.4</i>	16.8	<i>16.1</i>	22.2	<i>20.0</i>	18.6	<i>21.7</i>	16.7	<i>16.4</i>
Heavy Alcohol Use	4.5	<i>5.4</i>	5.9	<i>5.5</i>	6.0	<i>5.1</i>	5.9	<i>5.7</i>	4.3	<i>5.3</i>	7.1	<i>4.1</i>	4.3	<i>4.1</i>	4.3	<i>5.5</i>
Cigarette Use	14.5	<i>13.3</i>	24.8	<i>20.9</i>	29.5	<i>19.5</i>	23.6	<i>21.4</i>	12.9	<i>12.0</i>	25.1	<i>18.3</i>	26.1	<i>21.0</i>	14.0	<i>12.8</i>
Daily Cigarette Use	10.0	<i>9.3</i>	17.5	<i>13.7</i>	21.8	<i>13.6</i>	16.4	<i>13.7</i>	8.8	<i>8.6</i>	18.0	<i>12.7</i>	18.9	<i>15.4</i>	9.6	<i>9.1</i>

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. Therefore, some estimates in this table could differ from corresponding estimates in other tables that did not undergo sample size adjustments. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

¹ Mental Illness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

² Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

³ Respondents with unknown impairment data were excluded.

⁴ Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

⁵ Estimates of binge alcohol use include use by those who were heavy alcohol users.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.