

**Table 8.62C Substance Use Disorder Status and Substance Use Disorder for Specific Substances in Past Year: Among People Aged 18 or Older; by Past Year Suicide Behavior, Standard Errors of Numbers in Thousands, 2019 and 2020**

Substance Use Disorder Status	Had Serious Thoughts of Suicide (2019)	Had Serious Thoughts of Suicide (2020)	No Serious Thoughts of Suicide (2019)	No Serious Thoughts of Suicide (2020)	Made Any Suicide Plans <sup>1</sup> (2019)	Made Any Suicide Plans <sup>1</sup> (2020)	Did Not Make Any Suicide Plans <sup>1</sup> (2019)	Did Not Make Any Suicide Plans <sup>1</sup> (2020)	Attempted Suicide <sup>1</sup> (2019)	Attempted Suicide <sup>1</sup> (2020)	Did Not Attempt Suicide <sup>1</sup> (2019)	Did Not Attempt Suicide <sup>1</sup> (2020)
<b>TOTAL POPULATION</b>	308	510	2,077	2,884	153	239	2,118	2,937	93	131	2,127	2,952
<b>ILLICIT DRUGS</b>	nc	257	nc	517	nc	129	nc	570	nc	77	nc	587
Marijuana	nc	187	nc	448	nc	102	nc	480	nc	57	nc	491
Cocaine	nc	101	nc	140	nc	84	nc	153	nc	*	nc	173
Heroin	nc	40	nc	130	nc	13	nc	136	nc	26	nc	134
Hallucinogens	nc	24	nc	50	nc	10	nc	54	nc	8	nc	55
Inhalants	nc	20	nc	49	nc	6	nc	52	nc	5	nc	53
Methamphetamine	nc	81	nc	155	nc	58	nc	168	nc	*	nc	173
Misuse of Prescription Psychotherapeutics	nc	154	nc	247	nc	47	nc	288	nc	31	nc	291
Pain Relievers	nc	124	nc	206	nc	42	nc	236	nc	27	nc	239
Stimulants	nc	58	nc	95	nc	15	nc	113	nc	4	nc	114
Tranquilizers or Sedatives	nc	99	nc	124	nc	30	nc	156	nc	22	nc	158
Tranquilizers	nc	97	nc	99	nc	30	nc	136	nc	22	nc	138
Sedatives	nc	69	nc	79	nc	6	nc	104	nc	4	nc	104
Benzodiazepines	--	--	--	--	--	--	--	--	--	--	--	--
Opioids	nc	128	nc	215	nc	43	nc	248	nc	37	nc	248
Central Nervous System Stimulants	nc	138	nc	213	nc	97	nc	240	nc	49	nc	257
Illicit Drugs Other Than Marijuana <sup>2</sup>	nc	200	nc	312	nc	104	nc	360	nc	58	nc	372
<b>ALCOHOL</b>	nc	251	nc	750	nc	107	nc	804	nc	80	nc	812
<b>BOTH ILLICIT DRUGS AND ALCOHOL</b>	nc	167	nc	287	nc	81	nc	329	nc	52	nc	341
<b>ILLICIT DRUGS OR ALCOHOL</b>	nc	314	nc	887	nc	147	nc	948	nc	96	nc	954
<b>ILLICIT DRUGS ONLY</b>	nc	192	nc	434	nc	97	nc	468	nc	57	nc	480
<b>ALCOHOL ONLY</b>	nc	192	nc	679	nc	69	nc	710	nc	61	nc	714
<b>NO SUBSTANCE USE DISORDER</b>	nc	386	nc	2,718	nc	182	nc	2,749	nc	84	nc	2,759

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

NOTE: Respondents with unknown suicide information were excluded.

<sup>1</sup> Respondents who had no suicide thoughts were categorized as not making any suicide plans and not attempting suicide in the past 12 months.

<sup>2</sup> Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 8.62D Substance Use Disorder Status and Substance Use Disorder for Specific Substances in Past Year: Among People Aged 18 or Older; by Past Year Suicide Behavior, Standard Errors of Percentages, 2019 and 2020**

Substance Use Disorder Status	Had Serious Thoughts of Suicide (2019)	Had Serious Thoughts of Suicide (2020)	No Serious Thoughts of Suicide (2019)	No Serious Thoughts of Suicide (2020)	Made Any Suicide Plans <sup>1</sup> (2019)	Made Any Suicide Plans <sup>1</sup> (2020)	Did Not Make Any Suicide Plans <sup>1</sup> (2019)	Did Not Make Any Suicide Plans <sup>1</sup> (2020)	Attempted Suicide <sup>1</sup> (2019)	Attempted Suicide <sup>1</sup> (2020)	Did Not Attempt Suicide <sup>1</sup> (2019)	Did Not Attempt Suicide <sup>1</sup> (2020)
<b>TOTAL POPULATION</b>	0.00	<i>0.00</i>	0.00	<i>0.00</i>	0.00	<i>0.00</i>	0.00	<i>0.00</i>	0.00	<i>0.00</i>	0.00	<i>0.00</i>
<b>ILLICIT DRUGS</b>	nc	<i>1.81</i>	nc	<i>0.21</i>	nc	<i>3.23</i>	nc	<i>0.22</i>	nc	<i>4.99</i>	nc	<i>0.23</i>
Marijuana	nc	<i>1.42</i>	nc	<i>0.18</i>	nc	<i>2.76</i>	nc	<i>0.19</i>	nc	<i>4.12</i>	nc	<i>0.19</i>
Cocaine	nc	<i>0.81</i>	nc	<i>0.06</i>	nc	<i>2.46</i>	nc	<i>0.06</i>	nc	*	nc	<i>0.07</i>
Heroin	nc	<i>0.33</i>	nc	<i>0.05</i>	nc	<i>0.40</i>	nc	<i>0.06</i>	nc	<i>2.14</i>	nc	<i>0.05</i>
Hallucinogens	nc	<i>0.20</i>	nc	<i>0.02</i>	nc	<i>0.32</i>	nc	<i>0.02</i>	nc	<i>0.66</i>	nc	<i>0.02</i>
Inhalants	nc	<i>0.16</i>	nc	<i>0.02</i>	nc	<i>0.19</i>	nc	<i>0.02</i>	nc	<i>0.39</i>	nc	<i>0.02</i>
Methamphetamine	nc	<i>0.66</i>	nc	<i>0.06</i>	nc	<i>1.76</i>	nc	<i>0.07</i>	nc	*	nc	<i>0.07</i>
Misuse of Prescription Psychotherapeutics	nc	<i>1.20</i>	nc	<i>0.10</i>	nc	<i>1.47</i>	nc	<i>0.12</i>	nc	<i>2.55</i>	nc	<i>0.12</i>
Pain Relievers	nc	<i>0.98</i>	nc	<i>0.09</i>	nc	<i>1.30</i>	nc	<i>0.10</i>	nc	<i>2.21</i>	nc	<i>0.10</i>
Stimulants	nc	<i>0.47</i>	nc	<i>0.04</i>	nc	<i>0.47</i>	nc	<i>0.05</i>	nc	<i>0.30</i>	nc	<i>0.05</i>
Tranquilizers or Sedatives	nc	<i>0.80</i>	nc	<i>0.05</i>	nc	<i>0.93</i>	nc	<i>0.06</i>	nc	<i>1.82</i>	nc	<i>0.06</i>
Tranquilizers	nc	<i>0.78</i>	nc	<i>0.04</i>	nc	<i>0.94</i>	nc	<i>0.05</i>	nc	<i>1.84</i>	nc	<i>0.06</i>
Sedatives	nc	<i>0.56</i>	nc	<i>0.03</i>	nc	<i>0.18</i>	nc	<i>0.04</i>	nc	<i>0.33</i>	nc	<i>0.04</i>
Benzodiazepines	--	--	--	--	--	--	--	--	--	--	--	--
Opioids	nc	<i>1.01</i>	nc	<i>0.09</i>	nc	<i>1.33</i>	nc	<i>0.10</i>	nc	<i>2.94</i>	nc	<i>0.10</i>
Central Nervous System Stimulants	nc	<i>1.10</i>	nc	<i>0.09</i>	nc	<i>2.78</i>	nc	<i>0.10</i>	nc	<i>3.79</i>	nc	<i>0.10</i>
Illicit Drugs Other Than Marijuana <sup>2</sup>	nc	<i>1.52</i>	nc	<i>0.13</i>	nc	<i>2.93</i>	nc	<i>0.15</i>	nc	<i>4.33</i>	nc	<i>0.15</i>
<b>ALCOHOL</b>	nc	<i>1.76</i>	nc	<i>0.30</i>	nc	<i>2.99</i>	nc	<i>0.31</i>	nc	<i>5.38</i>	nc	<i>0.31</i>
<b>BOTH ILLICIT DRUGS AND ALCOHOL</b>	nc	<i>1.28</i>	nc	<i>0.12</i>	nc	<i>2.34</i>	nc	<i>0.13</i>	nc	<i>3.94</i>	nc	<i>0.14</i>
<b>ILLICIT DRUGS OR ALCOHOL</b>	nc	<i>1.95</i>	nc	<i>0.35</i>	nc	<i>3.47</i>	nc	<i>0.36</i>	nc	<i>5.09</i>	nc	<i>0.36</i>
<b>ILLICIT DRUGS ONLY</b>	nc	<i>1.45</i>	nc	<i>0.18</i>	nc	<i>2.64</i>	nc	<i>0.18</i>	nc	<i>4.14</i>	nc	<i>0.19</i>
<b>ALCOHOL ONLY</b>	nc	<i>1.46</i>	nc	<i>0.27</i>	nc	<i>2.07</i>	nc	<i>0.28</i>	nc	<i>4.56</i>	nc	<i>0.27</i>
<b>NO SUBSTANCE USE DISORDER</b>	nc	<i>1.95</i>	nc	<i>0.35</i>	nc	<i>3.47</i>	nc	<i>0.36</i>	nc	<i>5.09</i>	nc	<i>0.36</i>

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

NOTE: Respondents with unknown suicide information were excluded.

<sup>1</sup> Respondents who had no suicide thoughts were categorized as not making any suicide plans and not attempting suicide in the past 12 months.

<sup>2</sup> Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.