

**Table 9.10A Co-Occurring Substance Use Disorder, Illicit Drug Use Disorder, or Alcohol Use Disorder and Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year: Among People Aged 12 to 17; by Gender, Numbers in Thousands, 2019 and 2020**

<b>Substance Use Disorder (SUD), Illicit Drug Use Disorder (IDUD), or Alcohol Use Disorder (AUD) and MDE</b>	<b>Total (2019)</b>	<b>Total (2020)</b>	<b>Male (2019)</b>	<b>Male (2020)</b>	<b>Female (2019)</b>	<b>Female (2020)</b>
<b>SUD and MDE</b>	nc	644	nc	110	nc	533
SUD and MDE with Severe Impairment	nc	447	nc	70	nc	377
<b>IDUD and MDE</b>	nc	487	nc	84	nc	403
IDUD and MDE with Severe Impairment	nc	315	nc	54	nc	261
<b>AUD and MDE</b>	nc	328	nc	51	nc	276
AUD and MDE with Severe Impairment	nc	260	nc	38	nc	222

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 9.10B Co-Occurring Substance Use Disorder, Illicit Drug Use Disorder, or Alcohol Use Disorder and Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year: Among People Aged 12 to 17; by Gender, Percentages, 2019 and 2020**

<b>Substance Use Disorder (SUD), Illicit Drug Use Disorder (IDUD), or Alcohol Use Disorder (AUD) and MDE</b>	<b>Total (2019)</b>	<b>Total (2020)</b>	<b>Male (2019)</b>	<b>Male (2020)</b>	<b>Female (2019)</b>	<b>Female (2020)</b>
<b>SUD and MDE</b>	nc	2.7	nc	0.9	nc	4.5
SUD and MDE with Severe Impairment	nc	1.8	nc	0.6	nc	3.2
<b>IDUD and MDE</b>	nc	2.0	nc	0.7	nc	3.4
IDUD and MDE with Severe Impairment	nc	1.3	nc	0.4	nc	2.2
<b>AUD and MDE</b>	nc	1.3	nc	0.4	nc	2.3
AUD and MDE with Severe Impairment	nc	1.1	nc	0.3	nc	1.9

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.