

**Table 9.10C Co-Occurring Substance Use Disorder, Illicit Drug Use Disorder, or Alcohol Use Disorder and Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year: Among People Aged 12 to 17; by Gender, Standard Errors of Numbers in Thousands, 2019 and 2020**

| <b>Substance Use Disorder (SUD), Illicit Drug Use Disorder (IDUD), or Alcohol Use Disorder (AUD) and MDE</b> | <b>Total (2019)</b> | <b>Total (2020)</b> | <b>Male (2019)</b> | <b>Male (2020)</b> | <b>Female (2019)</b> | <b>Female (2020)</b> |
|--|---------------------|---------------------|--------------------|--------------------|----------------------|----------------------|
| <b>SUD and MDE</b>   | nc                  | 79                  | nc                 | 25                 | nc                   | 74                   |
| SUD and MDE with Severe Impairment   | nc                  | 60                  | nc                 | 19                 | nc                   | 55                   |
| <b>IDUD and MDE</b>  | nc                  | 69                  | nc                 | 21                 | nc                   | 65                   |
| IDUD and MDE with Severe Impairment  | nc                  | 49                  | nc                 | 17                 | nc                   | 45                   |
| <b>AUD and MDE</b>   | nc                  | 56                  | nc                 | 18                 | nc                   | 53                   |
| AUD and MDE with Severe Impairment   | nc                  | 49                  | nc                 | 15                 | nc                   | 47                   |

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 9.10D Co-Occurring Substance Use Disorder, Illicit Drug Use Disorder, or Alcohol Use Disorder and Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year: Among People Aged 12 to 17; by Gender, Standard Errors of Percentages, 2019 and 2020**

| <b>Substance Use Disorder (SUD), Illicit Drug Use Disorder (IDUD), or Alcohol Use Disorder (AUD) and MDE</b> | <b>Total (2019)</b> | <b>Total (2020)</b> | <b>Male (2019)</b> | <b>Male (2020)</b> | <b>Female (2019)</b> | <b>Female (2020)</b> |
|--|---------------------|---------------------|--------------------|--------------------|----------------------|----------------------|
| <b>SUD and MDE</b>   | nc                  | 0.33                | nc                 | 0.20               | nc                   | 0.62                 |
| SUD and MDE with Severe Impairment   | nc                  | 0.25                | nc                 | 0.15               | nc                   | 0.47                 |
| <b>IDUD and MDE</b>  | nc                  | 0.28                | nc                 | 0.17               | nc                   | 0.54                 |
| IDUD and MDE with Severe Impairment  | nc                  | 0.20                | nc                 | 0.13               | nc                   | 0.38                 |
| <b>AUD and MDE</b>   | nc                  | 0.23                | nc                 | 0.15               | nc                   | 0.45                 |
| AUD and MDE with Severe Impairment   | nc                  | 0.20                | nc                 | 0.12               | nc                   | 0.39                 |

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.