

Table 9.5A Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year: Among People Aged 12 to 17; Receipt of Treatment for Depression in Past Year: Among People Aged 12 to 17 with MDE and among People Aged 12 to 17 with MDE with Severe Impairment in Past Year; by Demographic Characteristics, Numbers in Thousands, 2019 and 2020

Demographic Characteristic	Had MDE (2019)	Had MDE (2020)	Had MDE with Severe Impairment ¹ (2019)	Had MDE with Severe Impairment ¹ (2020)	Received	Received	Received	Received
					Treatment for Depression in Past Year among People with MDE ² (2019)	Treatment for Depression in Past Year among People with MDE ² (2020)	Treatment for Depression in Past Year among People with Severe Impairment ^{1,2} (2019)	Treatment for Depression in Past Year among People with Severe Impairment ^{1,2} (2020)
TOTAL	3,783	4,131	2,671	2,919	1,627	1,710	1,321	1,361
AGE GROUP								
12-13	816	902	530	641	302	*	218	*
14-15	1,341	1,533	970	1,094	585	629	492	521
16-17	1,626	1,696	1,171	1,184	740	675	612	527
GENDER								
Male	1,076	1,135	736	732	391	448	300	341
Female	2,708	2,996	1,935	2,187	1,236	1,262	1,021	1,020
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	2,755	3,176	1,981	2,230	1,251	1,357	1,018	1,088
White	1,975	2,330	1,415	1,616	985	1,133	800	917
Black or African American	367	392	256	294	130	*	108	*
AIAN	20	*	19	*	*	*	*	*
NHOPI	*	*	*	*	*	*	*	*
Asian	190	252	142	186	*	*	*	*
Two or More Races	189	179	134	118	83	*	63	*
Hispanic or Latino	1,028	956	690	689	376	354	304	273

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

¹ Respondents with unknown impairment data were excluded.

² Respondents with unknown past year treatment for depression data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 9.5B Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year: Among People Aged 12 to 17; Receipt of Treatment for Depression in Past Year: Among People Aged 12 to 17 with MDE and among People Aged 12 to 17 with MDE with Severe Impairment in Past Year; by Demographic Characteristics, Percentages, 2019 and 2020

Demographic Characteristic	Had MDE (2019)	Had MDE (2020)	Had MDE with Severe Impairment ¹ (2019)	Had MDE with Severe Impairment ¹ (2020)	Received	Received	Received	Received
					Treatment for Depression in Past Year among People with MDE ² (2019)	Treatment for Depression in Past Year among People with MDE ² (2020)	Treatment for Depression in Past Year among People with MDE with Severe Impairment ^{1,2} (2019)	Treatment for Depression in Past Year among People with MDE with Severe Impairment ^{1,2} (2020)
TOTAL	15.7	17.0	11.1	12.0	43.3	41.6	49.7	46.9
AGE GROUP								
12-13	10.5	11.2	6.8	7.9	37.5	*	41.9	*
14-15	16.4	18.2	11.9	13.0	43.9	41.5	50.8	48.1
16-17	20.1	21.9	14.5	15.3	45.6	40.0	52.3	44.8
GENDER								
Male	8.8	9.2	6.0	5.9	36.8	39.7	41.2	46.6
Female	23.0	25.2	16.5	18.4	45.8	42.4	52.9	47.0
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	15.2	17.5	11.0	12.3	45.7	43.1	51.7	49.2
White	15.9	18.7	11.4	13.0	50.3	49.1	57.0	57.3
Black or African American	11.4	12.9	8.0	9.7	35.6	*	42.3	*
AIAN	12.2	*	11.5	*	*	*	*	*
NHOPI	*	*	*	*	*	*	*	*
Asian	15.1	13.9	11.3	10.3	*	*	*	*
Two or More Races	20.9	29.9	14.9	19.7	44.4	*	47.3	*
Hispanic or Latino	17.3	15.7	11.7	11.3	36.8	37.0	44.0	39.6

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

¹ Respondents with unknown impairment data were excluded.

² Respondents with unknown past year treatment for depression data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.