

Table 9.5C Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year: Among People Aged 12 to 17; Receipt of Treatment for Depression in Past Year: Among People Aged 12 to 17 with MDE and among People Aged 12 to 17 with MDE with Severe Impairment in Past Year; by Demographic Characteristics, Standard Errors of Numbers in Thousands, 2019 and 2020

Demographic Characteristic	Had MDE (2019)	Had MDE (2020)	Had MDE with Severe Impairment ¹ (2019)	Had MDE with Severe Impairment ¹ (2020)	Received	Received	Received	Received
					Treatment for Depression in Past Year among People with MDE ² (2019)	Treatment for Depression in Past Year among People with MDE ² (2020)	Treatment for Depression in Past Year among People with Severe Impairment ^{1,2} (2019)	Treatment for Depression in Past Year among People with Severe Impairment ^{1,2} (2020)
TOTAL	90	199	77	160	64	138	58	112
AGE GROUP								
12-13	40	127	34	102	24	*	21	*
14-15	61	124	53	99	40	73	37	65
16-17	63	130	54	102	42	69	38	59
GENDER								
Male	47	105	42	89	29	73	26	64
Female	75	154	65	133	57	105	52	93
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	73	172	65	130	55	124	50	97
White	72	166	60	120	50	118	45	93
Black or African American	29	60	25	48	18	*	17	*
AIAN	5	*	5	*	*	*	*	*
NHOPI	*	*	*	*	*	*	*	*
Asian	30	58	26	47	*	*	*	*
Two or More Races	19	41	16	26	13	*	11	*
Hispanic or Latino	52	102	44	92	33	61	29	56

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

¹ Respondents with unknown impairment data were excluded.

² Respondents with unknown past year treatment for depression data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 9.5D Major Depressive Episode (MDE) or MDE with Severe Impairment in Past Year: Among People Aged 12 to 17; Receipt of Treatment for Depression in Past Year: Among People Aged 12 to 17 with MDE and among People Aged 12 to 17 with MDE with Severe Impairment in Past Year; by Demographic Characteristics, Standard Errors of Percentages, 2019 and 2020

Demographic Characteristic	Had MDE (2019)	Had MDE (2020)	Had MDE with Severe Impairment ¹ (2019)	Had MDE with Severe Impairment ¹ (2020)	Received	Received	Received	Received
					Treatment for Depression in Past Year among People with MDE ² (2019)	Treatment for Depression in Past Year among People with MDE ² (2020)	Treatment for Depression in Past Year among People with Severe Impairment ^{1,2} (2019)	Treatment for Depression in Past Year among People with Severe Impairment ^{1,2} (2020)
TOTAL	0.37	<i>0.82</i>	0.32	<i>0.66</i>	1.34	2.45	1.63	2.69
AGE GROUP								
12-13	0.49	<i>1.42</i>	0.43	<i>1.16</i>	2.50	*	3.08	*
14-15	0.68	<i>1.35</i>	0.60	<i>1.09</i>	2.27	3.92	2.73	4.46
16-17	0.69	<i>1.55</i>	0.62	<i>1.26</i>	1.99	3.32	2.29	4.02
GENDER								
Male	0.38	<i>0.85</i>	0.34	<i>0.72</i>	2.21	4.49	2.82	6.03
Female	0.63	<i>1.30</i>	0.55	<i>1.13</i>	1.62	2.73	1.89	3.00
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.41	<i>0.95</i>	0.36	<i>0.71</i>	1.45	2.78	1.76	2.95
White	0.50	<i>1.18</i>	0.44	<i>0.88</i>	1.64	3.36	1.88	3.60
Black or African American	0.85	<i>1.81</i>	0.73	<i>1.44</i>	3.89	*	4.93	*
AIAN	3.19	*	3.12	*	*	*	*	*
NHOPI	*	*	*	*	*	*	*	*
Asian	2.02	<i>2.80</i>	1.85	<i>2.26</i>	*	*	*	*
Two or More Races	1.82	<i>5.11</i>	1.67	<i>3.49</i>	4.80	*	5.79	*
Hispanic or Latino	0.87	<i>1.67</i>	0.74	<i>1.51</i>	2.67	4.83	3.51	5.82

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

¹ Respondents with unknown impairment data were excluded.

² Respondents with unknown past year treatment for depression data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.