

Table 9.7A Major Depressive Episode in Past Year: Among People Aged 12 to 17; by Past Year Substance Use Disorder Status and Demographic Characteristics, Numbers in Thousands, 2019 and 2020

Substance Use Disorder Status/Age Group	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
SUBSTANCE USE DISORDER	nc	644	nc	110	nc	533
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	314	nc	*	nc	*
Illicit Drug Use Disorder	nc	487	nc	84	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	212	nc	*	nc	*
Alcohol Use Disorder	nc	*	nc	*	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	*	nc	*	nc	*
NO SUBSTANCE USE DISORDER	nc	3,488	nc	1,025	nc	2,463
12-13	nc	852	nc	190	nc	661
14-15	nc	1,254	nc	392	nc	862
16-17	nc	1,382	nc	443	nc	939

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 9.7B Major Depressive Episode in Past Year: Among People Aged 12 to 17; by Past Year Substance Use Disorder Status and Demographic Characteristics, Percentages, 2019 and 2020

Substance Use Disorder Status/Age Group	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
SUBSTANCE USE DISORDER	nc	41.7	nc	16.1	nc	62.1
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	36.9	nc	*	nc	*
Illicit Drug Use Disorder	nc	41.5	nc	15.1	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	31.8	nc	*	nc	*
Alcohol Use Disorder	nc	*	nc	*	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	*	nc	*	nc	*
NO SUBSTANCE USE DISORDER	nc	15.3	nc	8.7	nc	22.4
12-13	nc	10.7	nc	4.5	nc	17.7
14-15	nc	16.0	nc	9.5	nc	23.1
16-17	nc	20.0	nc	13.2	nc	26.6

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 9.8A Major Depressive Episode with Severe Impairment in Past Year: Among People Aged 12 to 17; by Past Year Substance Use Disorder Status and Demographic Characteristics, Numbers in Thousands, 2019 and 2020

Substance Use Disorder Status/Age Group	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
SUBSTANCE USE DISORDER	nc	447	nc	70	nc	377
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	191	nc	41	nc	*
Illicit Drug Use Disorder	nc	315	nc	54	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	114	nc	*	nc	*
Alcohol Use Disorder	nc	*	nc	*	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	*	nc	*	nc	*
NO SUBSTANCE USE DISORDER	nc	2,473	nc	662	nc	1,811
12-13	nc	603	nc	118	nc	486
14-15	nc	876	nc	237	nc	640
16-17	nc	993	nc	307	nc	686

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 9.8B Major Depressive Episode with Severe Impairment in Past Year: Among People Aged 12 to 17; by Past Year Substance Use Disorder Status and Demographic Characteristics, Percentages, 2019 and 2020

Substance Use Disorder Status/Age Group	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
SUBSTANCE USE DISORDER	nc	29.1	nc	10.2	nc	44.3
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	22.7	nc	10.4	nc	*
Illicit Drug Use Disorder	nc	27.1	nc	9.7	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	17.3	nc	*	nc	*
Alcohol Use Disorder	nc	*	nc	*	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	*	nc	*	nc	*
NO SUBSTANCE USE DISORDER	nc	10.9	nc	5.7	nc	16.4
12-13	nc	7.6	nc	2.8	nc	13.0
14-15	nc	11.2	nc	5.8	nc	17.1
16-17	nc	14.4	nc	9.1	nc	19.4

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.