

**Table 9.7C Major Depressive Episode in Past Year: Among People Aged 12 to 17; by Past Year Substance Use Disorder Status and Demographic Characteristics, Standard Errors of Numbers in Thousands, 2019 and 2020**

<b>Substance Use Disorder Status/Age Group</b>	<b>Total (2019)</b>	<b>Total (2020)</b>	<b>Male (2019)</b>	<b>Male (2020)</b>	<b>Female (2019)</b>	<b>Female (2020)</b>
<b>SUBSTANCE USE DISORDER</b>	nc	81	nc	25	nc	76
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	55	nc	*	nc	*
<b>Illicit Drug Use Disorder</b>	nc	70	nc	21	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	42	nc	*	nc	*
<b>Alcohol Use Disorder</b>	nc	*	nc	*	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	*	nc	*	nc	*
<b>NO SUBSTANCE USE DISORDER</b>	nc	212	nc	110	nc	166
12-13	nc	126	nc	58	nc	99
14-15	nc	111	nc	54	nc	94
16-17	nc	117	nc	77	nc	86

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 9.7D Major Depressive Episode in Past Year: Among People Aged 12 to 17; by Past Year Substance Use Disorder Status and Demographic Characteristics, Standard Errors of Percentages, 2019 and 2020**

<b>Substance Use Disorder Status/Age Group</b>	<b>Total (2019)</b>	<b>Total (2020)</b>	<b>Male (2019)</b>	<b>Male (2020)</b>	<b>Female (2019)</b>	<b>Female (2020)</b>
<b>SUBSTANCE USE DISORDER</b>	nc	4.05	nc	3.43	nc	5.39
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	5.18	nc	*	nc	*
<b>Illicit Drug Use Disorder</b>	nc	4.56	nc	3.46	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	5.49	nc	*	nc	*
<b>Alcohol Use Disorder</b>	nc	*	nc	*	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	*	nc	*	nc	*
<b>NO SUBSTANCE USE DISORDER</b>	nc	0.82	nc	0.88	nc	1.31
12-13	nc	1.43	nc	1.33	nc	2.29
14-15	nc	1.33	nc	1.28	nc	2.27
16-17	nc	1.59	nc	2.05	nc	2.28

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 9.8C Major Depressive Episode with Severe Impairment in Past Year: Among People Aged 12 to 17; by Past Year Substance Use Disorder Status and Demographic Characteristics, Standard Errors of Numbers in Thousands, 2019 and 2020**

Substance Use Disorder Status/Age Group	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
<b>SUBSTANCE USE DISORDER</b>	nc	61	nc	19	nc	57
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	39	nc	14	nc	*
<b>Illicit Drug Use Disorder</b>	nc	50	nc	17	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	24	nc	*	nc	*
<b>Alcohol Use Disorder</b>	nc	*	nc	*	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	*	nc	*	nc	*
<b>NO SUBSTANCE USE DISORDER</b>	nc	168	nc	92	nc	138
12-13	nc	101	nc	51	nc	87
14-15	nc	87	nc	47	nc	72
16-17	nc	95	nc	60	nc	72

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 9.8D Major Depressive Episode with Severe Impairment in Past Year: Among People Aged 12 to 17; by Past Year Substance Use Disorder Status and Demographic Characteristics, Standard Errors of Percentages, 2019 and 2020**

<b>Substance Use Disorder Status/Age Group</b>	<b>Total (2019)</b>	<b>Total (2020)</b>	<b>Male (2019)</b>	<b>Male (2020)</b>	<b>Female (2019)</b>	<b>Female (2020)</b>
<b>SUBSTANCE USE DISORDER</b>	nc	3.52	nc	2.74	nc	5.40
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	4.23	nc	3.51	nc	*
<b>Illicit Drug Use Disorder</b>	nc	3.87	nc	2.94	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	3.70	nc	*	nc	*
<b>Alcohol Use Disorder</b>	nc	*	nc	*	nc	*
12-13	nc	*	nc	*	nc	*
14-15	nc	*	nc	*	nc	*
16-17	nc	*	nc	*	nc	*
<b>NO SUBSTANCE USE DISORDER</b>	nc	0.67	nc	0.75	nc	1.12
12-13	nc	1.18	nc	1.18	nc	2.09
14-15	nc	1.05	nc	1.13	nc	1.75
16-17	nc	1.32	nc	1.67	nc	1.96

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details on these changes.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.