


Results from the 2020 National Survey on Drug Use and Health: Graphics from the Key Findings Report

Center for Behavioral Health Statistics and Quality
Substance Abuse and Mental Health Services Administration
U.S. Department of Health and Human Services



Past Month General Substance Use and Nicotine Vaping: Among People Aged 12 or Older; 2020




Rx = prescription.

Note: General Substance Use includes any illicit drug, alcohol, and tobacco product use. Tobacco products are defined as cigarettes, smokeless tobacco, cigars, and pipe tobacco.


Note: The estimated numbers of current users of different substances are not mutually exclusive because people could have used more than one type of substance in the past month.

Past Month Tobacco Use and Nicotine Vaping: Among People Aged 12 or Older; 2020



Note: The estimated numbers of current users of different tobacco products or nicotine vaping are not mutually exclusive because people could have used more than one type of tobacco product or used tobacco products and vaped nicotine in the past month.

Past Month Cigarette Use: Among People Aged 12 or Older; 2002-2020




Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.


Age	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
12 or Older	26.0	25.4	24.9	24.9	25.0	24.3	24.0	23.3	23.0	22.1	22.1	21.3	20.8	19.4	19.1	17.9	17.2	16.7	15.0
12 to 17	13.0	12.2	11.9	10.8	10.4	9.9	9.2	9.0	8.4	7.8	6.6	5.6	4.9	4.2	3.4	3.2	2.7	2.3	1.4
18 to 25	40.8	40.2	39.5	39.0	38.5	36.2	35.7	35.8	34.4	33.5	31.8	30.6	28.4	26.7	23.5	22.3	19.1	17.5	13.9
26 or Older	25.2	24.7	24.1	24.3	24.7	24.1	23.8	23.0	22.8	21.9	22.4	21.6	21.5	20.0	20.2	18.9	18.5	18.2	16.7

Daily Cigarette Use: Among Past Month Cigarette Smokers Aged 12 or Older; Smoking of One or More Packs of Cigarettes per Day: Among Current Daily Smokers; 2020




Note: Current daily smokers with unknown data about the number of cigarettes smoked per day were excluded from the pie chart on the right.

Past Month Nicotine Vaping: Among People Aged 12 or Older; 2020




Current, Binge, and Heavy Alcohol Use: Among People Aged 12 or Older; 2020



Note: Binge Alcohol Use is defined as drinking five or more drinks (for males) or four or more drinks (for females) on the same occasion on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as binge drinking on the same occasion on 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.


Past Month Binge Alcohol Use: Among People Aged 12 or Older; 2015-2020



Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.


Past Month Heavy Alcohol Use: Among People Aged 12 or Older; 2015-2020



Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.


Past Year Illicit Drug Use: Among People Aged 12 or Older; 2020



Rx = prescription.

Note: The estimated numbers of past year users of different illicit drugs are not mutually exclusive because people could have used more than one type of illicit drug in the past year.


Past Year Illicit Drug Use: Among People Aged 12 or Older; 2015-2020



Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.


Past Year Marijuana Use: Among People Aged 12 or Older; 2002-2020




Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.


Past Year Methamphetamine Use: Among People Aged 12 or Older; 2015-2020




Past Year Hallucinogen Use: Among People Aged 12 or Older; 2015-2020



Past Year Inhalant Use: Among People Aged 12 or Older; 2015-2020




Past Year Prescription Stimulant Misuse: Among People Aged 12 or Older; 2015-2020



Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.


Past Year Prescription Pain Reliever Misuse: Among People Aged 12 or Older; 2015-2020



Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.


Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Past Year Prescription Pain Reliever Misuse: Among People Aged 12 or Older; by Selected Pain Reliever Subtype, 2020



* Low precision; no estimate reported.

Source Where Pain Relievers Were Obtained for Most Recent Misuse: Among People Aged 12 or Older Who Misused Pain Relievers in the Past Year; 2020




9.3 Million People Aged 12 or Older Who Misused Pain Relievers in the Past Year


Note: Respondents with unknown data for the Source for Most Recent Misuse or who reported Some Other Way but did not specify a valid way were excluded.

Note: The percentages do not add to 100 percent due to rounding.


Past Year Opioid Misuse: Among People Aged 12 or Older; 2020



Past Year Central Nervous System (CNS) Stimulant Misuse: Among People Aged 12 or Older; 2020




Past Year Initiates of Substances: Among People Aged 12 or Older; 2020



Rx = prescription.

Note: Estimates for prescription pain relievers, prescription tranquilizers, prescription stimulants, and prescription sedatives are for the initiation of misuse.


Past Year Alcohol Initiates: Among People Aged 12 or Older; 2002-2020



Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Past Year Marijuana Initiates: Among People Aged 12 or Older; 2002-2020




Note: Estimates of less than 0.05 million round to 0.0 million when shown to the nearest tenth of a million.

Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Past Year Prescription Pain Reliever Misuse Initiates: Among People Aged 12 or Older; 2015-2020




Age Category: ▲ 12 or Older ○ 12 to 17 □ 18 to 25 ◇ 26 or Older

Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.


Perceived Great Risk from Substance Use: Among People Aged 12 or Older; 2015-2020




Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.


Substance Use Disorder, Alcohol Use Disorder, and Illicit Drug Use Disorder in the Past Year: Among People Aged 12 or Older; 2020




People Aged 12 or Older with a Past Year Substance Use Disorder (SUD); 2020




Alcohol Use Disorder (AUD) and Illicit Drug Use Disorder (IDUD) in the Past Year: Among People Aged 12 or Older with a Past Year Substance Use Disorder (SUD); 2020



Marijuana Use Disorder, Pain Reliever Use Disorder, and Methamphetamine Use Disorder in the Past Year: Among People Aged 12 or Older; 2020



Major Depressive Episode (MDE) and MDE with Severe Impairment in the Past Year: Among Youths Aged 12 to 17; 2004-2020




N/A = not available.

Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

MDE Status	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
MDE	9.0	8.8	7.9	8.2	8.3	8.1	8.0	8.2	9.1	10.7	11.4	12.5	12.8	13.3	14.4	15.7	17.0
MDE with Severe Impairment	N/A	N/A	5.5	5.5	6.0	5.8	5.7	5.7	6.3	7.7	8.2	8.8	9.0	9.4	10.0	11.1	12.0


Major Depressive Episode with Severe Impairment in the Past Year: Among Adults Aged 18 or Older; 2009-2020



Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.


Serious Mental Illness in the Past Year: Among Adults Aged 18 or Older; 2008-2020



Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.


Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Past Year Substance Use Disorder (SUD) and Major Depressive Episode (MDE): Among Youths Aged 12 to 17; 2020



Note: Youth respondents with unknown MDE data were excluded.


Substance Use: Among Youths Aged 12 to 17; by Past Year Major Depressive Episode (MDE) Status, 2020




⁺ Difference between this estimate and the estimate for youths without MDE is statistically significant at the .05 level.

Note: Youth respondents with unknown MDE data were excluded.


Past Year Substance Use Disorder (SUD) and Any Mental Illness (AMI): Among Adults Aged 18 or Older; 2020



Past Year Substance Use Disorder (SUD) and Serious Mental Illness (SMI): Among Adults Aged 18 or Older; 2020




Substance Use: Among Adults Aged 18 or Older; by Mental Illness Status, 2020




* Difference between this estimate and the estimate for adults without mental illness is statistically significant at the .05 level.


Adults Aged 18 or Older with Serious Thoughts of Suicide, Suicide Plans, or Suicide Attempts in the Past Year; 2020



Had Serious Thoughts of Suicide, Made a Suicide Plan, or Attempted Suicide in the Past Year: Among Adults Aged 18 or Older; 2020




Youths Aged 12 to 17 with Serious Thoughts of Suicide, Suicide Plans, or Suicide Attempts in the Past Year; 2020




3.0 Million Youths Aged 12 to 17 Had Serious Thoughts of Suicide, Made Suicide Plans, or Attempted Suicide in the Past Year

Need for Substance Use Treatment in the Past Year: Among People Aged 12 or Older; 2020




Locations Where Substance Use Treatment in the Past Year Was Received: Among People Aged 12 or Older; 2020




Note: Locations where people received substance use treatment are not mutually exclusive because respondents could report that they received treatment in more than one location in the past year.


Received Any Substance Use Treatment in the Past Year: Among People Aged 12 or Older Who Had a Substance Use Disorder in the Past Year; 2020



Received Substance Use Treatment at a Specialty Facility in the Past Year: Among People Aged 12 or Older Who Needed Substance Use Treatment in the Past Year; 2020



Perceived Need for Substance Use Treatment: Among People Aged 12 or Older with a Past Year Substance Use Disorder (SUD) Who Did Not Receive Substance Use Treatment at a Specialty Facility in the Past Year; 2020




38.4 Million People with an SUD Who Did Not Receive Substance Use Treatment at a Specialty Facility

Note: People who had an SUD were classified as needing substance use treatment.

Note: The percentages do not add to 100 percent due to rounding.

Received Treatment in the Past Year for Depression: Among Youths Aged 12 to 17 with a Past Year Major Depressive Episode (MDE) or MDE with Severe Impairment; 2004-2020




N/A = not available.

Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

MDE Status	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
MDE	40.3	37.8	38.8	39.0	37.7	34.6	37.8	38.4	37.0	38.1	41.2	39.3	40.9	41.5	41.4	43.3	41.6
MDE with Severe Impairment	N/A	N/A	46.5	43.9	42.6	38.8	41.1	43.5	41.0	45.0	44.7	44.6	46.7	47.5	46.9	49.7	46.9

Received Treatment in the Past Year for Depression: Among Adults Aged 18 or Older with a Past Year Major Depressive Episode; 2009-2020




Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Age	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
18 or Older	64.3	68.2	68.1	68.0	68.6	68.6	67.2	65.3	66.8	64.8	66.3	66.0
18 to 25	47.0	48.7	47.8	49.8	50.8	49.5	46.8	44.1	50.7	49.6	50.9	57.6
26 to 49	64.8	68.1	68.1	68.8	66.7	67.9	67.4	67.4	67.3	64.4	68.9	64.8
50 or Older	73.8	78.4	80.0	76.8	81.3	80.8	80.9	77.3	79.7	78.9	76.5	75.3

Sources of Mental Health Services in the Past Year: Among Youths Aged 12 to 17; 2002-2020




N/A = not available.

Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Source	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Specialty Mental Health Setting	11.8	12.4	13.4	13.4	13.0	12.4	12.7	12.0	12.1	12.6	12.7	13.6	13.7	13.3	14.7	14.8	16.0	16.7	17.3
Education Setting	N/A	12.1	12.4	11.9	12.9	13.0	13.2	13.2	13.1	13.3	14.2	15.4	12.8						
General Medical Setting	2.7	2.9	3.4	3.2	2.8	2.8	2.9	2.5	2.5	2.5	2.5	2.8	2.9	2.7	2.9	3.3	3.1	3.7	3.1
Child Welfare Setting	N/A	0.4	0.4	0.6	0.4	0.4	0.4	0.3	0.4	0.4	0.4	0.4	0.3						
Juvenile Justice Setting	N/A	0.4	0.3	0.4	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.1						

Type of Mental Health Services Received in the Past Year: Among Adults Aged 18 or Older; 2002-2020




Note: Mental Health Services include any combination of inpatient or outpatient services or receipt of prescription medication.

Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Service Type	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Mental Health Services	13.0	13.2	12.8	13.0	12.9	13.3	13.5	13.4	13.8	13.6	14.5	14.6	14.8	14.2	14.4	14.8	15.0	16.1	16.9
Inpatient	0.7	0.8	0.9	1.0	0.7	1.0	0.9	0.8	0.8	0.8	0.8	0.9	1.0	0.9	0.9	1.0	1.0	1.0	0.9
Outpatient	7.4	7.1	7.1	6.8	6.7	7.0	6.8	6.4	6.6	6.7	6.6	6.6	6.7	7.1	6.9	7.5	7.9	8.3	8.8
Prescription Medication	10.5	10.9	10.5	10.7	10.9	11.2	11.4	11.3	11.7	11.5	12.4	12.5	12.6	11.8	12.0	12.1	12.2	13.1	13.8

Mental Health Services Received in the Past Year: Among Adults Aged 18 or Older with Any Mental Illness in the Past Year; 2008-2020




Note: Mental Health Services include any combination of inpatient or outpatient services or receipt of prescription medication.

Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Mental Health Services Received in the Past Year: Among Adults Aged 18 or Older with Serious Mental Illness in the Past Year; 2008-2020




Note: Mental Health Services include any combination of inpatient or outpatient services or receipt of prescription medication.

Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.


Perceived Unmet Need for Mental Health Services in the Past Year: Among Adults Aged 18 or Older with Any Mental Illness in the Past Year; 2008-2020



Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Perceived Unmet Need for Mental Health Services in the Past Year: Among Adults Aged 18 or Older with Serious Mental Illness in the Past Year; 2008-2020




Note: There is no connecting line between 2019 and 2020 to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Note: The estimate in 2020 is italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed.

Age	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
18 or Older	43.7	46.3	42.0	43.1	41.6	38.6	42.9	38.2	39.7	44.2	45.1	47.7	49.7
18 to 25	50.0	52.2	53.1	55.0	49.8	51.5	53.6	50.3	53.7	55.9	59.5	62.5	63.4
26 to 49	44.8	49.2	44.3	45.2	46.2	42.4	45.4	43.3	39.7	45.2	45.2	47.6	48.0
50 or Older	38.2	37.5	32.7	33.9	30.1	27.1	33.9	23.2	30.4	32.5	31.9	35.1	41.4


Receipt of Substance Use Treatment at a Specialty Facility and Mental Health Services in the Past Year: Among Adults Aged 18 or Older with Past Year Substance Use Disorder and Any Mental Illness; 2020



Note: Mental Health Services include any combination of inpatient or outpatient services or receipt of prescription medication.

MH = mental health; SU Tx = substance use treatment.

Receipt of Substance Use Treatment at a Specialty Facility and Mental Health Services in the Past Year: Among Adults Aged 18 or Older with Past Year Substance Use Disorder and Serious Mental Illness; 2020




Note: Mental Health Services include any combination of inpatient or outpatient services or receipt of prescription medication.

MH = mental health; SU Tx = substance use treatment.


Note: The percentages do not add to 100 percent due to rounding.

Perceived COVID-19 Pandemic Negative Effect on Emotional or Mental Health: Among Youths Aged 12 to 17; by Past Year Major Depressive Episode (MDE) Status, Quarter 4, 2020




Note: The percentages do not add to 100 percent due to rounding.

Perceived COVID-19 Pandemic Negative Effect on Emotional or Mental Health: Among Adults Aged 18 or Older; by Past Year Mental Illness Status, Quarter 4, 2020




Note: The percentages do not add to 100 percent due to rounding.

Perceived COVID-19 Pandemic Effect on Alcohol Use: Among Past Year Alcohol Users Aged 12 or Older; Quarter 4, 2020



Note: The percentages do not add to 100 percent due to rounding.


Perceived COVID-19 Pandemic Effect on Drug Use: Among Past Year Users of Drugs Other than Alcohol Aged 12 or Older; Quarter 4, 2020



Note: Use of drugs other than alcohol included the use of marijuana, cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine in the past year or any use (i.e., not necessarily misuse) of prescription pain relievers, tranquilizers, stimulants, or sedatives in the past year. Drugs other than alcohol did not include tobacco products or nicotine vaping.

Note: The percentages do not add to 100 percent due to rounding.

Perceived COVID-19 Pandemic Effect on Mental Health Services: Among Adults Aged 18 or Older Who Received Services; Quarter 4, 2020



Perceived COVID-19 Pandemic Effect on Access to Medical Care: Among People Aged 12 or Older; Quarter 4, 2020

