

## CALIFORNIA

**Table 19A Substance Use, Perceptions of Great Risk, and Mental Health Measures: Among People Aged 12 or Older in California; by Age Group, Estimated Numbers (in Thousands), 2021**

Measure	12+	12-17	18-25	26+	18+
<b>ILLICIT DRUGS</b>					
Illicit Drug Use in the Past Month <sup>1,2</sup>	5,207	215	1,123	3,869	4,993
Marijuana Use in the Past Year	6,886	316	1,541	5,028	6,569
Marijuana Use in the Past Month	4,704	161	1,030	3,513	4,543
Perceptions of Great Risk from Smoking Marijuana Once a Month	7,974	699	588	6,687	7,275
First Use of Marijuana in the Past Year among Those at Risk for Initiation of Marijuana Use <sup>3,4</sup>	401	102	167	132	299
Illicit Drug Use Other Than Marijuana in the Past Month <sup>1,2</sup>	1,363	49	169	1,145	1,314
Cocaine Use in the Past Year	662	3	180	480	659
Perceptions of Great Risk from Using Cocaine Once a Month	20,806	1,635	2,266	16,905	19,170
Heroin Use in the Past Year <sup>5</sup>	--	--	5	99	103
Perceptions of Great Risk from Trying Heroin Once or Twice	26,701	1,843	3,128	21,729	24,858
Methamphetamine Use in the Past Year	421	3	23	394	418
Prescription Pain Reliever Misuse in the Past Year <sup>2</sup>	1,054	54	135	865	1,000
Opioid Misuse in the Past Year <sup>2,6</sup>	1,116	54	129	933	1,062
<b>ALCOHOL</b>					
Alcohol Use in the Past Month	15,040	196	1,769	13,075	14,844
Binge Alcohol Use in the Past Month <sup>7</sup>	6,804	126	1,107	5,570	6,677
Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week	15,851	1,471	1,768	12,611	14,380
Alcohol Use in the Past Month <sup>8</sup> (People Aged 12 to 20)	652	--	--	--	--
Binge Alcohol Use in the Past Month <sup>7,8</sup> (People Aged 12 to 20)	386	--	--	--	--
<b>TOBACCO PRODUCTS</b>					
Tobacco Product Use in the Past Month <sup>9</sup>	4,306	58	460	3,788	4,247
Cigarette Use in the Past Month	3,671	46	364	3,261	3,625
Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day	24,007	2,102	2,716	19,188	21,905
<b>MENTAL HEALTH MEASURES IN THE PAST YEAR</b>					
Any Mental Illness <sup>4,10</sup>	--	--	1,330	5,516	6,846
Serious Mental Illness <sup>4,10</sup>	--	--	479	1,351	1,830
Received Mental Health Services <sup>11</sup>	--	--	570	3,776	4,346
Major Depressive Episode <sup>4,12</sup>	--	613	816	1,685	2,501
Had Serious Thoughts of Suicide	--	--	558	977	1,535
Made Any Suicide Plans	--	--	206	233	439
Attempted Suicide	--	--	120	91	212

-- = not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes small area estimation approach. For confidence intervals, see Tables 1 to 35 in *2021 National Survey on Drug Use and Health: Model-Based Estimated Totals (in Thousands) (50 States and the District of Columbia)* at <https://www.samhsa.gov/data/report/2021-nsduh-estimated-totals-state>.

NOTE: Estimated numbers appearing as 0 in this table mean that the estimate is greater than 0 but less than 500 because estimated numbers are shown in thousands.

<sup>1</sup> Illicit Drug Use includes the misuse of prescription psychotherapeutics or the use of marijuana (including vaping), cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine. Illicit Drug Use Other Than Marijuana includes the misuse of prescription psychotherapeutics or the use of cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine. Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.<sup>2</sup> Prescription pain relievers are a type of prescription psychotherapeutic. Misuse of prescription psychotherapeutics is defined as use in any way not directed by a doctor, including use without a prescription of one's own; use in greater amounts, more often, or longer than told; or use in any other way not directed by a doctor. Prescription psychotherapeutics do not include over-the-counter drugs.<sup>3</sup> *First use of marijuana in the past year among those at risk for initiation* =  $X_1 \div 2$ , where  $X_1$  is the number of marijuana initiates in the past 24 months.<sup>4</sup> For details, see Section B of *2021 National Survey on Drug Use and Health: Guide to State Tables and Summary of Small Area Estimation Methodology* at <https://www.samhsa.gov/data/report/2021-nsduh-guide-state-tables-and-summary-sae-methodology>.<sup>5</sup> Estimates for youths aged 12 to 17 are not available for past year heroin use because past year heroin use was extremely rare among youths aged 12 to 17 in the 2021 NSDUH. As a result, estimates for people aged 12 or older are also not produced.<sup>6</sup> Respondents were classified as misusing opioids in the past year if they reported using heroin or misusing prescription pain relievers in the past year.<sup>7</sup> Binge Alcohol Use is defined as drinking five or more drinks (for males) or four or more drinks (for females) on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.<sup>8</sup> Underage drinking is defined for individuals aged 12 to 20; therefore, the "12+" estimate reflects that age group and not individuals aged 12 or older.<sup>9</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., snuff, dip, chewing tobacco, or snus), cigars, or pipe tobacco.<sup>10</sup> Mental Illness aligns with *Diagnostic and Statistical Manual of Mental Disorders*, 4th Edition (DSM-IV) criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These estimates are based on indicators of AMI and SMI rather than direct measures of diagnostic status.<sup>11</sup> Mental Health Services for adults includes inpatient treatment/counseling; outpatient treatment/counseling; or use of prescription medication for problems with emotions, nerves, or mental health.<sup>12</sup> Major depressive episode (MDE) is based on the *Diagnostic and Statistical Manual of Mental Disorders*, 5th edition (DSM-5) definition, which specifies a period of at least 2 weeks when an individual experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from adults aged 18 or older to produce an estimate for those aged 12 or older.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2021.

## CALIFORNIA

**Table 19B Substance Use Disorder and Treatment in the Past Year: Among People Aged 12 or Older in California; by Age Group, Estimated Numbers (in Thousands), 2021**

Measure	12+	12-17	18-25	26+	18+
<b>SUBSTANCE USE DISORDER AND TREATMENT</b>					
Drug Use Disorder <sup>1,2</sup>	2,729	194	666	1,868	2,535
Pain Reliever Use Disorder <sup>1,3</sup>	504	36	59	410	469
Opioid Use Disorder <sup>1,3,4</sup>	521	36	60	426	485
Alcohol Use Disorder <sup>1</sup>	3,650	103	608	2,940	3,547
Substance Use Disorder <sup>1,2</sup>	5,362	235	954	4,172	5,127
Needing But Not Receiving Treatment at a Specialty Facility for Illicit Drug Use <sup>5,6,7</sup>	2,327	144	616	1,567	2,183
Needing But Not Receiving Treatment at a Specialty Facility for Alcohol Use <sup>5</sup>	3,371	94	609	2,669	3,277
Needing But Not Receiving Treatment at a Specialty Facility for Substance Use <sup>5,6,7</sup>	4,817	200	942	3,675	4,617

NOTE: Estimates are based on a survey-weighted hierarchical Bayes small area estimation approach. For confidence intervals, see Tables 1 to 35 in *2021 National Survey on Drug Use and Health: Model-Based Estimated Totals (in Thousands) (50 States and the District of Columbia)* at <https://www.samhsa.gov/data/report/2021-nsduh-estimated-totals-state>.

NOTE: Estimated numbers appearing as 0 in this table mean that the estimate is greater than 0 but less than 500 because estimated numbers are shown in thousands.

<sup>1</sup> Substance Use Disorder (SUD) estimates are based on *Diagnostic and Statistical Manual of Mental Disorders*, 5th edition (DSM-5) criteria. SUD is defined as meeting the criteria for drug or alcohol use disorder. Beginning with the 2021 NSDUH, questions on prescription drug use disorder were asked of all past year users of prescription drugs, regardless of whether they misused prescription drugs. The estimates in these rows include prescription drug use disorder data from all past year users of prescription drugs.

<sup>2</sup> Drug use includes the use of marijuana (including vaping), cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine in the past year or any use (i.e., not necessarily misuse) of prescription pain relievers, tranquilizers, stimulants, or sedatives in the past year.

<sup>3</sup> Pain relievers are a type of prescription drug.

<sup>4</sup> Opioid Use Disorder is defined as meeting the criteria for heroin or pain reliver use disorder.

<sup>5</sup> Respondents were classified as needing substance use treatment if they met the DSM-5 criteria for an illicit drug or alcohol use disorder or received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Substance use treatment questions are asked of respondents who used alcohol or illicit drugs in their lifetime. Respondents who used prescription drugs but who did not misuse prescription drugs in their lifetime may not receive these questions. Needing But Not Receiving Substance Use Treatment refers to respondents who are classified as needing illicit drug or alcohol use treatment but who did not receive illicit drug or alcohol use treatment at a specialty facility.

<sup>6</sup> Illicit drug or alcohol use disorder estimates are based on DSM-5 criteria. Beginning with the 2021 NSDUH, questions on prescription drug use disorder were asked of all past year users of prescription drugs, regardless of whether they misused prescription drugs. The estimates in these rows do not include prescription drug use disorder data from the past year users of prescription drugs who were not also misusers of prescription drugs.

<sup>7</sup> Illicit Drug Use includes the misuse of prescription psychotherapeutics or the use of marijuana (including vaping), cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2021.

## CALIFORNIA

**Table 20A Substance Use, Perceptions of Great Risk, and Mental Health Measures: Among People Aged 12 or Older in California; by Age Group, Percentages, 2021**

Measure	12+	12-17	18-25	26+	18+
<b>ILLICIT DRUGS</b>					
Illicit Drug Use in the Past Month <sup>1,2</sup>	15.73	6.90	28.24	14.87	16.65
Marijuana Use in the Past Year	20.80	10.16	38.76	19.33	21.90
Marijuana Use in the Past Month	14.21	5.18	25.91	13.50	15.15
Perceptions of Great Risk from Smoking Marijuana Once a Month	24.08	22.47	14.77	25.71	24.25
First Use of Marijuana in the Past Year among Those at Risk for Initiation of Marijuana Use <sup>3,4</sup>	2.27	3.65	7.99	1.02	2.02
Illicit Drug Use Other Than Marijuana in the Past Month <sup>1,2</sup>	4.12	1.58	4.25	4.40	4.38
Cocaine Use in the Past Year	2.00	0.09	4.52	1.84	2.20
Perceptions of Great Risk from Using Cocaine Once a Month	62.86	52.58	56.98	64.98	63.91
Heroin Use in the Past Year <sup>5</sup>	--	--	0.12	0.38	0.34
Perceptions of Great Risk from Trying Heroin Once or Twice	80.69	59.25	78.66	83.53	82.88
Methamphetamine Use in the Past Year	1.27	0.11	0.58	1.52	1.39
Prescription Pain Reliever Misuse in the Past Year <sup>2</sup>	3.18	1.73	3.40	3.32	3.33
Opioid Misuse in the Past Year <sup>2,6</sup>	3.37	1.72	3.25	3.59	3.54
<b>ALCOHOL</b>					
Alcohol Use in the Past Month	45.44	6.32	44.48	50.26	49.50
Binge Alcohol Use in the Past Month <sup>7</sup>	20.55	4.06	27.84	21.41	22.26
Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week	47.88	47.30	44.46	48.48	47.94
Alcohol Use in the Past Month <sup>8</sup> (People Aged 12 to 20)	13.82	--	--	--	--
Binge Alcohol Use in the Past Month <sup>7,8</sup> (People Aged 12 to 20)	8.18	--	--	--	--
<b>TOBACCO PRODUCTS</b>					
Tobacco Product Use in the Past Month <sup>9</sup>	13.01	1.87	11.56	14.56	14.16
Cigarette Use in the Past Month	11.09	1.47	9.15	12.54	12.09
Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day	72.53	67.59	68.31	73.76	73.04
<b>MENTAL HEALTH MEASURES IN THE PAST YEAR</b>					
Any Mental Illness <sup>4,10</sup>	--	--	33.44	21.21	22.83
Serious Mental Illness <sup>4,10</sup>	--	--	12.05	5.19	6.10
Received Mental Health Services <sup>11</sup>	--	--	14.33	14.52	14.49
Major Depressive Episode <sup>4,12</sup>	--	19.72	20.52	6.48	8.34
Had Serious Thoughts of Suicide	--	--	14.04	3.76	5.12
Made Any Suicide Plans	--	--	5.18	0.89	1.46
Attempted Suicide	--	--	3.03	0.35	0.71

-- = not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes small area estimation approach. For confidence intervals, see Tables 1 to 35 in *2021 National Survey on Drug Use and Health: Model-Based Prevalence Estimates (50 States and the District of Columbia)* at <https://www.samhsa.gov/data/report/2021-nsduh-state-prevalence-estimates>.<sup>1</sup> Illicit Drug Use includes the misuse of prescription psychotherapeutics or the use of marijuana (including vaping), cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine. Illicit Drug Use Other Than Marijuana includes the misuse of prescription psychotherapeutics or the use of cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine. Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.<sup>2</sup> Prescription pain relievers are a type of prescription psychotherapeutic. Misuse of prescription psychotherapeutics is defined as use in any way not directed by a doctor, including use without a prescription of one's own; use in greater amounts, more often, or longer than told; or use in any other way not directed by a doctor. Prescription psychotherapeutics do not include over-the-counter drugs.<sup>3</sup> *First use of marijuana in the past year among those at risk for initiation (%) = 100 \* [(X<sub>1</sub> + (0.5 \* X<sub>1</sub> + X<sub>2</sub>)) ÷ 2]*, where X<sub>1</sub> is the number of marijuana initiates in the past 24 months and X<sub>2</sub> is the number of individuals who never used marijuana (with the at-risk population defined as 0.5 \* X<sub>1</sub> + X<sub>2</sub>). Both of the computation components, X<sub>1</sub> and X<sub>2</sub>, are based on a survey-weighted hierarchical Bayes small area estimation approach. The age group shown is based on a respondent's age at the time of the interview, not his or her age at first use.<sup>4</sup> For details, see Section B of *2021 National Survey on Drug Use and Health: Guide to State Tables and Summary of Small Area Estimation Methodology* at <https://www.samhsa.gov/data/report/2021-nsduh-guide-state-tables-and-summary-sae-methodology>.<sup>5</sup> Estimates for youths aged 12 to 17 are not available for past year heroin use because past year heroin use was extremely rare among youths aged 12 to 17 in the 2021 NSDUH. As a result, estimates for people aged 12 or older are also not produced.<sup>6</sup> Respondents were classified as misusing opioids in the past year if they reported using heroin or misusing prescription pain relievers in the past year.<sup>7</sup> Binge Alcohol Use is defined as drinking five or more drinks (for males) or four or more drinks (for females) on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.<sup>8</sup> Underage drinking is defined for individuals aged 12 to 20; therefore, the "12+" estimate reflects that age group and not individuals aged 12 or older.<sup>9</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., snuff, dip, chewing tobacco, or snus), cigars, or pipe tobacco.<sup>10</sup> Mental Illness aligns with *Diagnostic and Statistical Manual of Mental Disorders*, 4th Edition (DSM-IV) criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These estimates are based on indicators of AMI and SMI rather than direct measures of diagnostic status.<sup>11</sup> Mental Health Services for adults includes inpatient treatment/counseling; outpatient treatment/counseling; or use of prescription medication for problems with emotions, nerves, or mental health.<sup>12</sup> Major depressive episode (MDE) is based on the *Diagnostic and Statistical Manual of Mental Disorders*, 5th edition (DSM-5) definition, which specifies a period of at least 2 weeks when an individual experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from adults aged 18 or older to produce an estimate for those aged 12 or older.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2021.

## CALIFORNIA

**Table 20B Substance Use Disorder and Treatment in the Past Year: Among People Aged 12 or Older in California; by Age Group, Percentages, 2021**

Measure	12+	12-17	18-25	26+	18+
<b>SUBSTANCE USE DISORDER AND TREATMENT</b>					
Drug Use Disorder <sup>1,2</sup>	8.24	6.25	16.75	7.18	8.45
Pain Reliever Use Disorder <sup>1,3</sup>	1.52	1.15	1.48	1.58	1.56
Opioid Use Disorder <sup>1,3,4</sup>	1.58	1.17	1.50	1.64	1.62
Alcohol Use Disorder <sup>1</sup>	11.03	3.31	15.28	11.30	11.83
Substance Use Disorder <sup>1,2</sup>	16.20	7.56	24.00	16.04	17.09
Needing But Not Receiving Treatment at a Specialty Facility for Illicit Drug Use <sup>5,6,7</sup>	7.03	4.61	15.49	6.03	7.28
Needing But Not Receiving Treatment at a Specialty Facility for Alcohol Use <sup>5</sup>	10.18	3.01	15.31	10.26	10.93
Needing But Not Receiving Treatment at a Specialty Facility for Substance Use <sup>5,6,7</sup>	14.55	6.45	23.69	14.13	15.39

NOTE: Estimates are based on a survey-weighted hierarchical Bayes small area estimation approach. For confidence intervals, see Tables 1 to 35 in *2021 National Survey on Drug Use and Health: Model-Based Prevalence Estimates (50 States and the District of Columbia)* at <https://www.samhsa.gov/data/report/2021-nsduh-state-prevalence-estimates>.

<sup>1</sup> Substance Use Disorder (SUD) estimates are based on *Diagnostic and Statistical Manual of Mental Disorders*, 5th edition (DSM-5) criteria. SUD is defined as meeting the criteria for drug or alcohol use disorder. Beginning with the 2021 NSDUH, questions on prescription drug use disorder were asked of all past year users of prescription drugs, regardless of whether they misused prescription drugs. The estimates in these rows include prescription drug use disorder data from all past year users of prescription drugs.

<sup>2</sup> Drug use includes the use of marijuana (including vaping), cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine in the past year or any use (i.e., not necessarily misuse) of prescription pain relievers, tranquilizers, stimulants, or sedatives in the past year.

<sup>3</sup> Pain relievers are a type of prescription drug.

<sup>4</sup> Opioid Use Disorder is defined as meeting the criteria for heroin or pain reliver use disorder.

<sup>5</sup> Respondents were classified as needing substance use treatment if they met the DSM-5 criteria for an illicit drug or alcohol use disorder or received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Substance use treatment questions are asked of respondents who used alcohol or illicit drugs in their lifetime. Respondents who used prescription drugs but who did not misuse prescription drugs in their lifetime may not receive these questions. Needing But Not Receiving Substance Use Treatment refers to respondents who are classified as needing illicit drug or alcohol use treatment but who did not receive illicit drug or alcohol use treatment at a specialty facility.

<sup>6</sup> Illicit drug or alcohol use disorder estimates are based on DSM-5 criteria. Beginning with the 2021 NSDUH, questions on prescription drug use disorder were asked of all past year users of prescription drugs, regardless of whether they misused prescription drugs. The estimates in these rows do not include prescription drug use disorder data from the past year users of prescription drugs who were not also misusers of prescription drugs.

<sup>7</sup> Illicit Drug Use includes the misuse of prescription psychotherapeutics or the use of marijuana (including vaping), cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2021.